

Changes in macrobenthos communities during the invasive *Spartina alterniflora* removal and mangrove restoration

Lin Lin ^a, Yuqing Guo ^a, Guanzhao Wang ^a, Shilong Feng ^a, Kai Liu ^a, Mingcheng Hu ^a, Mouxin Ye ^b, Chunxi Cao ^b, Ruohai Chen ^c, Silong Ding ^c, Zenghui Peng ^d, Fenfen Ji ^{a*}, Yi-Jia Shih ^{e*}

^a *Fisheries College, Jimei University, Xiamen 361021, China*

^b *Mangrove Conservation Foundation, Shenzhen, Guangdong, China*

^c *Quanzhou Bay Estuary Wetland Nature Reserve Development Center, Quanzhou, Fujian, China*

^d *Hunan Shuitiandi Environmental Technology Co., Ltd., Changsha 410205, China*

^e *Sustainable Ocean Governance Centre, National Sun Yat-sen University, Kaohsiung 80424, Taiwan*

* Corresponding author.

E-mail address: ffji1213266226@163.com (F. Ji); cja0313@mail.nsysu.edu.tw (YJ SHIH).

Table S1 the number of OTU sequences obtained through eDNA metabarcoding

Month	The number of sequences
2022.9	57886
2022.10	37299
2022.12	56975




Figure S1 the species-filtering steps to identify target species




Figure S2 the changes in sediment environmental factors (TOC, TN and TP)