



Hepatic Stellate Cells and Hepatocarcinogenesis

Anna E. Barry^{1,2†}, Rajkumar Baldeosingh^{1,2†}, Ryan Lamm¹, Keyur Patel¹, Kai Zhang^{1,2}, Dana A. Dominguez³, Kayla J. Kirton¹, Ashesh P. Shah¹ and Hien Dang^{1,2*}

¹ Department of Surgery, Thomas Jefferson University, Philadelphia, PA, United States, ² Sidney Kimmel Cancer Center, Philadelphia, PA, United States, ³ Department of General Surgery, UCSF East Bay, Oakland, CA, United States

OPEN ACCESS

Edited by:

Tomer Cooks,
Ben-Gurion University of the Negev,
Israel

Reviewed by:

Ana I. Robles,
National Cancer Institute (NCI),
United States
Oleg N. Demidov,
INSERM U1231 Lipides, Nutrition,
Cancer (LNC), France

*Correspondence:

Hien Dang
Hien.Dang@jefferson.edu

† These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Molecular Medicine,
a section of the journal
Frontiers in Cell and Developmental
Biology

Received: 28 May 2020

Accepted: 13 July 2020

Published: 05 August 2020

Citation:

Barry AE, Baldeosingh R,
Lamm R, Patel K, Zhang K,
Dominguez DA, Kirton KJ, Shah AP
and Dang H (2020) Hepatic Stellate
Cells and Hepatocarcinogenesis.
Front. Cell Dev. Biol. 8:709.
doi: 10.3389/fcell.2020.00709

Hepatic stellate cells (HSCs) are a significant component of the hepatocellular carcinoma (HCC) tumor microenvironment (TME). Activated HSCs transform into myofibroblast-like cells to promote fibrosis in response to liver injury or chronic inflammation, leading to cirrhosis and HCC. The hepatic TME is comprised of cellular components, including activated HSCs, tumor-associated macrophages, endothelial cells, immune cells, and non-cellular components, such as growth factors, proteolytic enzymes and their inhibitors, and other extracellular matrix (ECM) proteins. Interactions between HCC cells and their microenvironment have become topics under active investigation. These interactions within the hepatic TME have the potential to drive carcinogenesis and create challenges in generating effective therapies. Current studies reveal potential mechanisms through which activated HSCs drive hepatocarcinogenesis utilizing matricellular proteins and paracrine crosstalk within the TME. Since activated HSCs are primary secretors of ECM proteins during liver injury and inflammation, they help promote fibrogenesis, infiltrate the HCC stroma, and contribute to HCC development. In this review, we examine several recent studies revealing the roles of HSCs and their clinical implications in the development of fibrosis and cirrhosis within the hepatic TME.

Keywords: hepatic stellate cells, hepatocytes, hepatocellular carcinoma, tumor microenvironment, inflammation, fibrosis

INTRODUCTION

Hepatocellular carcinoma (HCC) is the sixth leading cause of cancer related deaths in western countries (Choo et al., 2016; Bray et al., 2018), accounting for up to 90% of all primary liver cancers (Lozano et al., 2012). Although the percentage of HCC cases is dramatically higher in eastern Asia and most African countries, HCC is on the rise in the United States (U.S.) (Rawla et al., 2018). This trend is primarily due to increases in the incidences of chronic hepatic inflammation including fatty liver disease (FLD) (Nordenstedt et al., 2010) and chronic hepatitis C (HCV) infection (Kanwal et al., 2011). Although the distribution of HCC etiology varies between geographic regions, the most common etiology worldwide is viral hepatitis. In developing areas, such as Sub-Saharan Africa and Eastern Asia, hepatitis B virus (HBV) passed through vertical transmission at birth is the most common etiology (El-Serag, 2012). In developed worlds such as North America, HCV infection acquired later in life is endemic (Armstrong et al., 2006; Barazani et al., 2007; Wasley et al., 2008). In addition to viral hepatitis and FLD, alcoholic cirrhosis is a major contributing etiology of HCC; all of which can induce fibrosis, cirrhosis, and ultimately lead to development of HCC. This pathway

to malignancy, driven largely through fibrosis, is supported by the fact that 90% of HCC patients have cirrhosis (Parkin et al., 2005; Ananthakrishnan et al., 2006).

One of the main components in the development of fibrosis, cirrhosis and HCC (Wynn, 2008; Sokolovic et al., 2010) are the liver-specific pericytes, known as hepatic stellate cells (HSCs), which are located in the perisinusoidal space of the liver (Tsuchida and Friedman, 2017). Under normal conditions, HSCs exist in a quiescent state containing abundant lipid droplets of vitamin A (Blaner et al., 2009), and are highly sensitive to extracellular signals from fibrotic stimuli (Yin et al., 2013) including hepatitis, inflammation, or tissue injury (Wynn, 2008). In the presence of liver injury, HSCs are activated, transitioning from a quiescent to a myofibroblast phenotype with proliferative, migratory and invasive capabilities (Coulouarn et al., 2012; Carloni et al., 2014; Novikova et al., 2017). The well-known sequence of HSC activation can be split into two pathways. The first includes 'initiation,' which describes changes in HSC gene expression and phenotype rendering them more sensitive to paracrine stimuli. The second pathway includes 'perpetuation,' which amplifies the HSC-initiated phenotype via enhanced proliferation and proinflammatory signaling (i.e., damaged associated molecular patterns, DAMPs), interleukins, complimentary and growth factors from nearby damaged hepatocytes, endothelial cells and immune cells, resulting in promotion of fibrogenesis (Friedman, 2000) (**Figure 1**). Extracellular matrix (ECM) molecules (ex, type I and III collagen) are secreted by activated HSCs and accumulate to form scar tissue in the space of Disse (Friedman, 2008a) (**Figure 1**). This scar tissue functions to protect the liver from further damage; however, sustained activation of HSCs leads to chronic fibrosis and cirrhosis (Lee and Friedman, 2011; Friedman et al., 2013).

Activated HSCs play an essential role in fibrosis and hepatocarcinogenesis (Tsuchida and Friedman, 2017). Mediators in the activation of HSCs and the hepatic tumor microenvironment (TME) consist of transforming growth factor beta (TGFB), platelet derived growth factor (PDGF), connective tissue growth factor (CCN2, previously CTGF), vascular endothelial growth factor (VEGF), viral infection, focal adhesion kinase-matrix metalloproteinase 9 (FAK-MMP9) signaling, p53/21, phosphatidylinositol 3-kinase/protein kinase B (AKT) (PI3K/AKT), mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK), and interleukin 6/signal transducer and activator of transcription 3 (IL-6/STAT3) signaling pathways (Han et al., 2014; Fabregat et al., 2016; Makino et al., 2018). Furthermore, previous studies have shown HSC activation to be regulated by the tumor suppressor phosphatase and tensin homolog deleted on chromosome 10 (PTEN) (Takashima et al., 2009; Bian et al., 2012). These characteristics of activated HSCs and the role of the TME lay the foundation for exploring the relationship between the hepatic TME, activated HSCs, and hepatocarcinogenesis. This review will focus on the mechanisms regulating HSC activation and how they contribute to fibrosis, cirrhosis and HCC development (Wright et al., 2014; Coll et al., 2015; Das et al., 2020) and promising clinical therapies associated with HSCs (Dong et al., 2018; Li et al., 2020).

THE ROLE OF ACTIVATED HEPATIC STELLATE CELLS IN HEPATIC FIBROSIS, CIRRHOSIS, AND PROGRESSION TO HCC


Hepatic fibrosis is a major risk factor for HCC development. Furthermore, activation of HSCs is a driver of hepatic fibrosis, cirrhosis and HCC (Tsuchida and Friedman, 2017). Due to the lack of effective liver fibrosis treatments, there is a need to better understand the molecular pathways and mediators of HSC activation to develop advantageous, targeted therapies for liver disease. Hepatic fibrosis is a result of chronic inflammation (post liver injury) characterized by secretion of excess ECM components, resulting in a wound healing response that will produce a "scar" in the liver (Friedman, 2008b; Lee and Friedman, 2011). Chronic inflammation can be induced by HBV/HCV infection, alcohol abuse, FLD (including non-alcoholic fatty liver disease and non-alcoholic steatohepatitis) or other metabolic disorders of the normal liver (Higashi et al., 2017). During post chronic liver inflammation, HSC activation supports the development of fibrosis and later cirrhosis, characterized by liver function impairment, portal vein hypertension and jaundice (Gines et al., 1987; de Franchis, 2000; D'Amico et al., 2006). Therefore, preventing the activation and proliferation of HSCs in cases of hepatic fibrosis has the potential to delay progression to HCC.

The Role of DNA Methylation in HSC Activation

Recently, studies have begun focusing on the epigenetic regulation of HSCs to uncover the complexity of HSC activation (Tsuchida and Friedman, 2017). The tumor suppressor gene PTEN is an important negative regulator of HSC activation, which is silenced through promoter hypermethylation in tumors (Cairns et al., 1997; Salvesen et al., 2001; Soria et al., 2002; Furuta et al., 2004; Roman-Gomez et al., 2004; Mirmohammadsadegh et al., 2006; Wiencke et al., 2007; Tao et al., 2011; van Eggermond et al., 2011; Bian et al., 2012). Furthermore, the inhibition of PTEN activity leads to a constitutive activation of HSCs, which can perpetuate hepatic fibrosis (Bian et al., 2012). In addition, activated HSCs have been shown to possess altered DNA methylation and hydroxymethylation marks. For example, Page et al. (2016) showed that activated HSCs lose their fibrogenic phenotype when DNA methyltransferase 3a (DNMT3a) is suppressed in a carbon tetra chloride (CCL₄) rat model of liver fibrosis *in vivo*. The continued discovery of novel mediators of HSC activation, including epigenetic regulators, will help better understand the role of activated HSCs in hepatic fibrosis.

Pro-fibrogenic Cytokines in HSC Activation During Hepatic Fibrosis

Transforming growth factor beta is one of the key fibrogenic cytokines that drive hepatic fibrosis and regulate HSC activation (Fabregat et al., 2016; Tsuchida and Friedman, 2017; Dewidar et al., 2019). The bona-fide TGFB signaling pathway requires latent TGFB to be cleaved and activated by thrombospondin



1 (TSP1). Activated TGFB then binds to TGFB receptor 2 (TGFB_{R2}), inducing phosphorylation of mothers against decapentaplegic homolog 2 and 3 (SMAD2, SMAD3) which translocate into the nucleus to regulate downstream gene expression of pro-fibrotic genes (Breitkopf et al., 2006;

Friedman, 2008a; Liu et al., 2013; Meng et al., 2016; Tsuchida and Friedman, 2017; Murphy-Ullrich and Suto, 2018). TGFB can also activate the MAPK p38, ERK, and c-jun N-terminal kinase (JNK) pathways to regulate HSC activation (Engel et al., 1999; Hanafusa et al., 1999). Central to liver injury and HSC

activation is a pathological increase in expression and activation of TGF β in the ECM (Tsuchida and Friedman, 2017). During HSC activation, TGF β targets and binds to HSCs, inducing phosphorylation of SMAD3 to promote production of type I and II collagen (Breitkopf et al., 2006; Friedman, 2008a). Activated HSCs also promote increased production of TGF β 1 and TSP1 further enhancing the profibrogenic activities of TGF β (Breitkopf et al., 2005). Moreover, attenuation of TSP1 activated TGF β can be achieved with a TSP1 antagonist peptide and has been shown to decrease liver fibrosis in a dimethyl-nitrosamine liver fibrosis model (Li et al., 2017; Murphy-Ullrich and Suto, 2018), demonstrating the importance of this relationship. Along with TGF β , the DNA demethylase tet methylcytosine dioxygenase 3 (TET3) is also upregulated in mouse and human fibrotic livers (Xu et al., 2020). Xu et al. (2020) established TGF β 1 stimulation could increase TET3 levels along with increased profibrotic gene expression in LX-2 cells, a human HSC cell line. Furthermore, TET3 was shown to upregulate the TGF β pathway genes TSP1 and TGF β 2, suggesting a positive feedback loop between TET3 and TGF β 1 to promote HSC activation and hepatic fibrosis (Xu et al., 2020). In addition to TGF β signaling influencing HSC activation, hepatocyte produced extracellular matrix protein 1 (ECM1) has been shown to suppress TGF β levels and prevent HSC activation in an *in vivo* ECM1 knockout mouse model (Fan et al., 2019). Although TGF β remains a strong mediator of HSC activation and fibrogenesis, additional profibrogenic cytokines contribute to this process as well.

Another essential cytokine involved in HSC activation is PDGF (Tsuchida and Friedman, 2017). PDGF levels are increased in human cirrhotic livers compared to normal, healthy livers (Pinzani et al., 1996; Ikura et al., 1997; Stock et al., 2007; Tsuchida and Friedman, 2017). Upon liver injury in both humans and rodents, PDGF receptor beta (PDGFR β) expression increases in HSCs to drive HSC activation, proliferation and migration (Wong et al., 1994; Borkham-Kamphorst et al., 2004). Pdgf-c, a member of Pdgf family, is highly expressed on membrane receptors of hepatocytes in a transgenic Pdgf-c mouse model that resulted in dynamic liver fibrosis (Wright et al., 2014), suggesting that HSC activation may include Pdgf-c signaling. Pdgf is further supported as an effective activator of HSCs through the fibrotic role of Agrin (Agrn), a secreted proteoglycan induced by Pdgf-induced HSC activation in HCC in Diethyl nitrosamine (DEN)-induced HCC Sprague Dawley rat model (Lv et al., 2017). In this study the authors showed that Pdgf acts as an activator of the HSCs, which was inhibited by blocking the binding of Pdgf to its receptor. The authors also demonstrated that Agrin from activated HSC supernatant increased proliferation, metastasis, and invasion of SMMC-7721 (a human HCC cell line) and promoted epithelial to mesenchymal transition (EMT) (Lv et al., 2017). Overall, this study supports the role of PDGF-induced HSC activation resulting in fibrosis and HCC.

The matricellular protein CCN2, known for mediating fibrosis in various organs including the liver (Hall-Glenn and Lyons, 2011; Jun and Lau, 2011; Kodama et al., 2011; Lipson et al., 2012), has also been shown to activate HSCs and promote tumor progression via HSC secretion of the IL-6 and STAT3 *in vitro* (Makino et al., 2018). A potential clinical player in

the cellular crosstalk between HSCs and HCC cells is stroma-derived fibroblast growth factor 9 (FGF9). Interestingly, a 2020 study conducted by Seitz et al. found that only activated HSCs expressed FGF9 compared to HCC cells. In HCC tissues, activated HSC overexpression of FGF9 reduced sensitivity to therapeutic agents and was associated with poor prognosis (Seitz et al., 2020), suggesting FGF9 as a potential therapeutic target and prognostic tool for HCC. Altogether, these findings support the notion of the growth factors PDGF-C and CCN2 as activators of HSCs and FGF9 as a potential clinical target for HCC.

Association of Resident Liver Lymphocytes With Activated HSCs During Hepatic Fibrosis and Cirrhosis

Pathogenesis of liver fibrosis also involves resident liver lymphocytes including Type I and Type II Natural Killer T (NKT), Natural Killer (NK) cells and innate lymphoid cells (ILCs) (Wang and Zhang, 2019). Specifically, the interaction between activated HSCs and these hepatic lymphocytes is important (Wang and Zhang, 2019). The innate role of NKT cells is to defend against pathogens by recruiting circulating lymphocytes (Racanelli and Rehermann, 2006). Once activated, NKT cells can induce HSC activation via production of pro-inflammatory cytokines and release osteopontin and Hedgehog (Hh) ligands (Syn et al., 2012; Wehr et al., 2013) to aid in fibrosis development (Wang and Yin, 2015; Bandyopadhyay et al., 2016). However, it has been established that NK cells, along with NKT cells, protect the liver by preventing infection, tumor formation (Racanelli and Rehermann, 2006) and fibrogenesis (Melhem et al., 2006; Radaeva et al., 2006) in liver fibrosis mouse models (Muhanna et al., 2011; Gur et al., 2012; Hou et al., 2012) and HCV patients in various clinical studies (Glassner et al., 2012; Gur et al., 2012; Kramer et al., 2012). Furthermore, NK cells have an anti-fibrotic effect on HSCs by inducing activated HSC apoptosis (Radaeva et al., 2006). However, this is a temporary effect, and could result in apoptosis-resistant activated HSCs (Radaeva et al., 2007). Additionally, type 3 innate lymphoid cells (ILC3s) function as pro-fibrotic effectors in the liver (Muhanna et al., 2007; Bjorklund et al., 2016). Through co-culturing experiments with LX-2 cells, ILC3s promoted fibrogenesis via Interleukin-17A (IL-17A) and Interleukin-22 (IL-22), resulting in IL-22 inhibition of interferon gamma (IFN γ) to indirectly enhance fibrogenesis (Wang et al., 2018). These data suggest an important role for resident liver immune cells in liver fibrosis through interactions with activated HSCs (Schon and Weiskirchen, 2014). Furthermore, they suggest the involvement of the innate immune system in relation to HSC activation in fibrosis; thus, warranting further study of the roles of activated HSCs in enhancing and suppressing fibrosis.

Even though stimulation of innate immunity has been shown to have a pivotal role in anti-viral and anti-tumor defenses in addition to fibrosis suppression, the regulation of innate immunity during chronic liver injury still needs to be elucidated. In a CCl $_4$ mouse model, Infg induced NK cell activation was decreased in late liver fibrosis (advanced scarring) compared to early fibrosis (minimal scarring) (Jeong et al., 2011). The authors further demonstrated that the anti-fibrotic roles of NK cells

are suppressed during advanced liver injury through increased expression of suppressor of cytokine signaling 1 (Socs1) and Tgfb (Jeong et al., 2011). This study also showed *in vitro* evidence of early activated HSC (4 days co-culture of HSCs and liver NK cells) induced NK cell activation via natural killer group 2 member D (NKG2D), whereas this was abolished in intermediately activated HSCs (8 days co culture of HSCs and liver NK cells) due to increased levels of TGFB1 and downregulation of NKG2D (Jeong et al., 2011). These results establish that although NK cells interact with activated HSCs to mitigate liver fibrosis during chronic liver injury, this process can be suppressed through increased TGFB and SOCS1 produced by activated HSCs *in vitro*.

In a 2017 study, Shi et al. (2017) showed that in liver cirrhosis patients, activated HSCs interact with purified NK cells through HSC-derived TGFB regulation of emperipolesis (the presence of an intact cell within the cytoplasm of another cell). This process was mediated through TGFB and evidenced by significantly reduced NK cell emperipolesis when activated HSCs were treated with an anti-TGFB antibody. The NK cells inside activated HSCs were also apoptotic as observed through positive terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining which indicates DNA fragmentation, a trait of cellular apoptosis (Krylkova et al., 2012). This suggests that activated HSCs reduce the anti-fibrotic roles of NK cells *in vitro* through activated HSC-derived TGFB and programmed death of NK cells to promote fibrosis (Shi et al., 2017). This study demonstrates that activated HSC-derived TGFB and NK cells work together to diminish NK cell anti-fibrotic capabilities and promote fibrosis in liver cirrhosis patients.

Cellular Senescence of Activated HSCs During Hepatic Fibrosis

Senescence of activated HSCs has been shown to suppress liver fibrosis (Hoare et al., 2010; Jin et al., 2016; Nishizawa et al., 2016). In cancer, cellular senescence is a known tumor suppressive mechanism (Sager, 1991) activated by oncogenic stress (Campisi and d'Adda di Fagagna, 2007) and may also be essential for regulating liver fibrosis and cirrhosis (Tsuchida and Friedman, 2017). Cancer cell senescence is maintained by the tumor suppressor proteins p53/p21, p16^{Ink4a} and retinoblastoma (Rb) (Campisi and d'Adda di Fagagna, 2007; Collado et al., 2007). Furthermore, transgenic p53^{-/-} mice, with p53 knocked out specifically in HSCs, that were treated with CCl₄ showed increased fibrosis compared to WT p53 mice. This phenotype indicates that activated HSCs have the ability to undergo senescence resulting in decreased liver fibrosis *in vivo* (Krizhanovsky et al., 2008). This study also revealed retained fibrotic lesions in p53^{-/-}; Cdkn2a/Arf^{-/-} mice 20 days post CCl₄ treatment, along with increased smooth muscle actin (Acta2), Tgfb, and Ki67 (marker of proliferative cells) expression. These findings suggest that activated HSCs during liver fibrosis can also evade cellular senescence to proliferate and secrete ECM components to begin developing the hepatic TME (Krizhanovsky et al., 2008).

HSC activation is one of the first responses to LSEC (liver sinusoidal endothelial cell) injury in the liver. LSECs serve as a

permeable barrier between hepatocytes and the bloodstream and are characterized by fenestrations and a disorganized basement membrane, making them one of the most permeable types of endothelial cells (DeLeve and Maretta-Mira, 2017; Poisson et al., 2017). This permeability allows for efficient transport of solutes and metabolites throughout the liver. In addition, LSECs are an active contributor to the production of excess ECM proteins during liver fibrosis (Natarajan et al., 2017). Recent data shows that removal of senescent LSECs promotes liver fibrosis (Grosse et al., 2020). Through genetic lineage tracing mouse models, Grosse, et al. demonstrated that the majority of senescent cells were vascular endothelial cells, mostly LSECs in liver sinusoids, in addition to macrophages and adipocytes to a reduced extent. The authors also showed that both continuous and acute elimination of senescent cells disrupted blood-tissue barriers resulting in liver and perivascular tissue fibrosis. Overall, this study establishes that senescent cells involved in preventing liver fibrosis are primarily LSECs, rather than hepatic stellate cells (Grosse et al., 2020). While senescent LSECs serve as a barrier against fibrosis, the activation, proliferation and transformation of HSCs caused by liver injury, in addition to their ability to evade senescence once activated, are important developmental processes that provide a suitable microenvironment required for fibrosis, cirrhosis and later HCC development (Lashen et al., 2020).

ACTIVATED HEPATIC STELLATE CELLS WITHIN THE HEPATIC TUMOR MICROENVIRONMENT

The interplay between liver tumor cells and the hepatic TME is crucial to the initiation and progression of HCC (Hernandez-Gea et al., 2013; Zhou et al., 2019). The TME is defined as a peritumoral space (Alfarouk et al., 2011; Joyce and Fearon, 2015; Spill et al., 2016) contributing to the acquisition of various hallmark traits of cancer, including sustained proliferative signaling and activation of invasion, metastasis, and angiogenesis (Hanahan and Weinberg, 2011). Furthermore, the TME can be divided into two major components: (1) cellular and (2) non-cellular. Activated HSCs are a part of the cellular component and exhibit essential biological functions such as promotion of fibrogenesis and ECM remodeling to positively influence HCC tumorigenesis (Friedman, 2008a; Amann et al., 2009; Coulouarn and Clement, 2014).

In addition to HSCs, cellular components of the hepatic TME include stromal hepatocytes, immune cells such as myeloid-derived suppressor cells (MDSCs) (Fu et al., 2007; Hoechst et al., 2008), tumor associated macrophages (TAMs) (Budhu and Wang, 2006; Budhu et al., 2006; Jeong et al., 2011), and cancer associated fibroblasts (CAFs) (Yin et al., 2019). Non-cellular components include cytokines such as Interleukin-6 (IL-6) (Budhu and Wang, 2006; Budhu et al., 2006) and Interleukin-22 (IL-22) (Jiang et al., 2011), growth factors such as VEGF (Coulouarn et al., 2012), TGFB (Thompson et al., 2015), PDGF (Tsuchida and Friedman, 2017), and CCN2 (Makino et al., 2018). Additional non-cellular components include matrix metalloproteinases (MMPs), their inhibitors

(Novikova et al., 2017) and proteoglycans (Theocharis et al., 2010) (Table 1). The following studies will cover the different roles activated HSCs play in HCC progression through their interaction with the other cellular and non-cellular components of the hepatic TME.

Cellular Crosstalk Between Activated HSCs and Cellular Components of the Hepatic TME

The hepatic TME consists of various immune cells to create an immunosuppressed environment in order to maintain HCC tumor growth (Lu et al., 2019). Activated HSCs contribute to this immunosuppressed environment by secreting cytokines which induce MDSC expansion (Maher, 2001; Hui et al., 2004; Yu et al., 2004; Pan et al., 2008; Gabrilovich and Nagaraj, 2009; Hsieh et al., 2013). In an orthotopic liver tumor mouse model, activated HSCs significantly increased regulatory T cell (Treg) and MDSC expression to benefit HCC growth in the spleen, bone marrow, and tumor tissues (Zhao et al., 2014). Furthermore, activated HSCs secrete angiogenic growth factors to form new vasculature within the TME (Coulouarn et al., 2012; Heindryckx and Gerwins, 2015). These functions of activated HSCs create a link to the circulatory system for supplying nutrients to the tumor. Immune cells may also regulate activation of HSCs *in vitro*, demonstrated by Interleukin 20 (IL-20) activation of HSCs, resulting in upregulation of TGF β 1 and type I collagen, and increased proliferation and migration of activated HSCs (Chiu et al., 2014). The same study further indicated that these fibrogenic phenotypes could be attenuated with an anti-IL-20 receptor (IL-20R1) monoclonal antibody, proposing IL-20 as a significant activator of HSCs and fibrogenesis. Taken together, activated HSCs may have an important role in promoting an immunosuppressed and angiogenic hepatic TME to support aggressive HCC cell growth.

Crosstalk Between Activated HSCs and Non-cellular Components of the Hepatic TME

In addition to interacting with other hepatic TME cellular components, HSCs also respond to the non-cellular components of the liver TME (Hall-Glenn and Lyons, 2011; Jun and Lau, 2011; Kodama et al., 2011; Lipson et al., 2012). An example of such is the response to CCN2 produced from hepatic tumor cells (Makino et al., 2018). Makino et al. (2018) demonstrated that elevated CCN2 expression positively correlated with activated HSCs, indicated by smooth muscle actin (ACTA2) expression, in both mouse and human liver tumors. Furthermore, the authors showed that anti-CCN2 reduced IL-6 production in LX-2 cells and inhibited STAT3 activation in HepG2 (human HCC cell line) cells (Makino et al., 2018). This study was the first to establish HCC-cell-derived CCN2 activates HSCs in the TME, thus, accelerating the progression of HCC through cytokine production. These results also support the need for further exploration of CCN2 and other ECM proteins involved in the activation of HSCs.

While HSCs respond to growth factors such as CCN2; once activated, HSCs can also modulate the ECM through secretion and upregulation of proteins such as MMPs (Lachowski et al., 2019), which are needed for HCC tumor migration (Scheau et al., 2019). Studies have shown MMP2 and MMP9 (the most commonly studied MMPs in HCC EMT) to be important for the invasive potential of HCC tumors through degradation and remodeling of collagen in the ECM (Wang et al., 2014; Sun et al., 2018). Moreover, signaling between FAK and MMP9 is considered to be one of the main pathways that promotes HCC cell invasion and metastasis (Chen et al., 2010; Jia et al., 2011). Thus, it is plausible to speculate whether this signaling pathway is promoted through HSC activation. This hypothesis was explored by Han et al. (2014), who investigated whether activated HSCs promote FAK-MMP9 signaling *in vitro*. First, elevated numbers activated HSCs were shown to associate with tumor invasion of the portal vein, advanced tumor node metastasis staging, and lesser tumor differentiation. Thereafter, the number of activated HSCs, quantified by cytoplasmic ACTA2 expression, were positively correlated with the expression levels of phosphorylated FAK (p-FAK) and MMP9 in HCC. Furthermore, the authors used a co-culture experiment to demonstrate the activation of FAK-MMP9 signaling in HCC cells in the presence of activated HSC conditioned medium and with co-culture of activated HSCs. Additionally, inhibition of FAK-MMP9 signaling via small interfering RNA (siRNA) for FAK (siFAK) abrogated the migratory and invasive effects of activated HSCs on HCC cells (Han et al., 2014). These data show that FAK-MMP9 signaling is promoted by activated HSCs and plays a role in modulating metastasis of HCC following activation of HSCs; thus, highlighting the crosstalk between tumor cells and activated HSCs in the hepatic TME.

Micro-RNA Involvement in HSC Activation in the Hepatic TME

In addition to ECM components such as MMPs and growth factors, recent advances have emphasized the significant roles played by micro-RNAs (miRNAs) in the TME (Cheng et al., 2015). This is demonstrated through the ability of miRNAs in tumor cells to transform the microenvironment by sustaining cancer hallmark traits and non-cell-autonomous signaling pathways (Suzuki et al., 2015). miRNAs are small non-coding RNAs (20-25 nucleotides in length) that regulate gene expression by binding to target mRNA transcripts through a seed sequence at the 5' end of the miRNA (Bartel, 2004). In cancer cells, miRNAs are aberrantly expressed compared to normal cells, with expression patterns varying between different types of cancer cells (Karube et al., 2005; Lu et al., 2005; Merritt et al., 2008; Garzon et al., 2009). Additionally, activation and inactivation of HSCs can be controlled by profibrogenic and antifibrogenic miRNAs (Tsuchida and Friedman, 2017). Interestingly, miRNAs have been shown to possess dual roles as oncogenes and tumor suppressors in cancer cells (Zhang et al., 2007). The following studies investigate these dual roles in relation to the activation of HSCs in the hepatic TME.

TABLE 1 | Components of the hepatic TME.

Cellular	References	Non-cellular	References
Stromal hepatocytes: connective tissue cells that provide support to the epithelial cells of the liver	Tahmasebi Birgani and Carloni, 2017	ECM Proteins (Matrix metalloproteinases, collagens, proteoglycans, lamins): key players in the invasive potential of HCC tumors by modulating the ECM	Novikova et al., 2017
Hepatic stellate cells: mesenchymal cells found in the liver that contribute to the hepatic TME by proliferating and promoting fibrosis when activated	Friedman, 2008a; Coulouarn and Clement, 2014	Growth factors (TGFB, PDGF, CCN2, VEGF, HGF): signaling proteins that stimulate the expression pathways of pro-fibrotic genes; stimulate HSC activation, proliferation and migration	Coulouarn et al., 2012; Thompson et al., 2015; Makino et al., 2018
Immune cells (Ex. Tumor Associated Macrophages and MDSCs): components of the hepatic TME that interact with activated HSCs by creating an immunosuppressed environment promoting HCC tumor growth and maintenance	Budhu and Wang, 2006; Budhu et al., 2006; Fu et al., 2007; Hoechst et al., 2008; Jeong et al., 2011	Cytokines (IL-6, IL-8, IL-22): small proteins involved in a range of cell signaling that help drive fibrosis, HSC activation, and contribute to angiogenesis	Chiu et al., 2014; Sevic et al., 2019
Cancer associated fibroblasts: type of cancer stromal cell critical to tumorigenesis regulation by possessing the ability to remodel the ECM and secrete proteins such as cytokines and VEGF	Yin et al., 2019		

Both cellular and non-cellular components cooperate in the hepatic TME to aid in hepatocarcinogenesis.

Oncogenic miRNAs (oncomiRs) derived from the extracellular vesicles (EV) of HCC cells mediate communication between HCC cells and activated HSCs (Daugaard et al., 2017). Interestingly, crosstalk between miRNAs and TME components is partly mediated by exosomes, a type of EV produced in the endosome of eukaryotic cells that can transfer DNA, RNA and proteins to other cells (Zhang et al., 2015; Kosaka, 2016). Results from Li J. et al. (2019) found that EVs released by HepG2 and Huh7 (human HCC cell line) cells contained elevated oncomiRs. As a result, activated HSCs released their own EVs which stimulated HCC invasion, epithelial to mesenchymal transition (EMT) and activation of the AKT/ERK signaling pathway (Li J. et al., 2019). This suggests a positive feedback loop between exosomal oncomiRs of activated HSCs and HCC cells to promote hepatocarcinogenesis. Moreover, the authors observed upregulation of three specific oncomiRs: miR-21, miR-221, and miR-151. These results are further examples of the crosstalk between HCC cells and activated HSCs creating a pro-metastatic phenotype (Li J. et al., 2019).

Likewise, miR-1426 is an oncomiR shown to promote tumorigenesis, metastasis, and migration in multiple cancer types (Xu et al., 2019). Recently, a miRNA expression microarray study revealed a robust increase in miR-1246 in HCC cell lines when co-cultured with activated HSCs (Huang et al., 2020). These results reflect *in vitro* evidence that activated HSCs induce miR-1246 expression in HCC cell lines to promote metastasis. Moreover, miR-1246 and its target RAR related orphan receptor alpha (RORA), promoted EMT *in vitro* and *in vivo* in nude mice indicated by enhanced HCC cell migration, decreased *E*-cadherin, and increased vimentin protein expression. As a result of miR-1246 overexpression in PLC cells (human liver hepatoma cell line), the binding of RORA and beta-catenin (CTNNB1) in the cytoplasm was increased. This process was reversed with RORA knockdown, suggesting that the binding of RORA to beta-catenin prevents beta-catenin

nuclear translocation and activation of the Wnt/beta-catenin signaling pathway. Furthermore, both miR-1246 and RORA were effectively used as independent prognostic markers in HCC tissue (Huang et al., 2020). This data suggests that miR-1246:RORA is a key component in the tumorigenic influence of activated HSCs on HCC cells, and that targeting the miR-1246:RORA axis may slow HCC progression (Huang et al., 2020).

However, miRNAs can also function as tumor suppressors in cancer cells (Zhang et al., 2007). For example, miRNA-212-3p has been shown to suppress cancer cell growth in other forms of cancer such as renal cell carcinoma (Gu et al., 2017) non-small-cell lung cancer (Tang et al., 2017) and glioblastoma (Tang et al., 2017). However, the effects, if any, in HCC remained unclear. Thus, Chen et al. (2019) examined the relationship between miRNA-212-3p and CCN2 in the hepatic TME. The authors showed that microRNA-212-3p inhibited proliferation of HCC cell lines through suppression of CCN2. Additionally, miR-212-3p was downregulated in HCC cell lines and tissues, and negatively correlated with vascular invasion and the absence of a fibrous tumor capsule. This fibrous capsule is formed by host liver mesenchymal cells, instead of HCC cells, and prevents possible invasion of HCC to the host liver (Ishizaki et al., 2001). These findings demonstrate a tumor suppressor role of miRNA-212-3p through its interaction with CCN2, a significant ECM component of the hepatic TME. Moreover, results from this study raise the question of whether microRNA-212-3p also inhibits the CCN2 mediated cytokine production in activated HSCs, given that the experiments from this study were carried out in HCC cell lines and tissues only.

Activated HSC Regulation of Angiogenesis Within the Hepatic TME

The effects of activated HSCs on angiogenesis in HCC have also been investigated over the past decade. Angiogenesis within the

TME is essential for tumor progression, metastasis and invasion (Mittal et al., 2014). Zhu et al. (2015) identified Interleukin-8 (IL-8) as a contributing factor to angiogenesis in HCC. Interestingly, IL-8 was highly expressed in HCC stroma and was mainly derived from activated HSCs rather than from HCC cells. Furthermore, an IL-8 neutralizing antibody was demonstrated to suppress tumor angiogenesis in Hep3B cells (a human HCC cell line) treated with conditioned media from activated HSCs. The authors also demonstrated similar results *in vivo* through a chick embryo chorioallantoic membrane (CAM) assay. Most recently, Lin et al. observed that activated HSCs are the primary source of secreted angiopoietin-1 (Ang-1) in human HCC cells *in vitro*. This not only describes the promotion of HCC angiogenesis through activated HSCs and Ang-1 expression, but also opens the potential of Ang-1 as an anti-angiogenic therapeutic target in HCC (Lin et al., 2020). These findings identify angiogenic factors produced by activated HSCs in the hepatic TME to promote hepatocarcinogenesis.

Intriguingly, a 2019 investigation found that activated HSCs induced angiogenesis in HCC via upregulation of glioma associated oncogene 1 (Gli-1), a member of the Hh signaling pathway (Yan et al., 2017). Furthermore, this study established that 3,5,4'-trihydroxy-*trans*-stilbene (trade name: Resveratrol), a polyphenol compound found in red-wine, grapes, berries and peanuts and believed to act as an antioxidant, also hindered HCC progression driven by HSCs through targeting Gli-1. Specifically, activated HSC induced angiogenesis in HCC via upregulation of Gli-1 expression, stimulated reactive oxygen species (ROS) production and increased HCC cell invasiveness. Resveratrol further abolished activated HSC-stimulated angiogenesis and suppressed ROS production and IL-6 and C-X-C chemokine receptor type 4 (CXCR4) expression in HepG2 cells by downregulating Gli-1 expression (Yan et al., 2017). In a separate study, Han et al. (2019) also described Resveratrol to possess tumor-suppressive effects through tumor microenvironment modulation across several types of cancers including HCC. This suggests the possibility of Gli-1 as a potential target for angiogenesis prevention in HCC.

QUIESCENT HEPATIC STELLATE CELLS

While activated HSCs play a major role in the formation of fibrosis and the hepatic TME, recent studies have also delved into the role of quiescent HSCs (qHSCs), as they are needed to maintain a healthy liver (Coll et al., 2015; Das et al., 2020). Uncovering the mechanisms preserving this phenotype could increase the scope of HSC targeted therapies.

Coll et al. (2015) conducted a study which examined these mechanisms and included a miRNA microarray analysis of isolated human qHSCs. The microarray revealed that HSCs express 259 miRNAs. In contrast, when HSCs were activated *in vitro*, 212 of these miRNAs were upregulated and the other 47 miRNAs were downregulated (Coll et al., 2015) suggesting a role for miRNAs in maintenance of qHSCs. Furthermore, the interactions between the miRNA target genes in qHSCs were also associated with HSC activation. Specifically, miRNA-192 was

chosen for further *in vivo* study due to having 28 target genes in qHSCs and demonstrating decreased expression in cirrhotic liver samples compared to healthy samples. To elucidate the role of miRNA-192 in qHSCs *in vivo*, HSCs were isolated from two liver fibrosis mouse models and showed decreased miRNA-192 expression compared to healthy mouse HSCs. Furthermore, miRNA-192 overexpression resulted in inhibited Tgfb1 signaling and Pdgf-induced HSC migration in primary mouse HSC cells. This data supports miRNA-192 as a regulator of qHSCs through suppressing target genes needed for HSC activation (Coll et al., 2015). Thus, these findings support additional study into the miRNA-192 targeted genes involved in HSC activation to increase the scope of HSC targeted therapies.

Further supporting the significance of qHSCs, Das et al. demonstrated *in vitro* qHSC induction of cancer cell apoptosis via a caspase-independent mechanism (Das et al., 2020). This mechanism was established in rat hepatoma cells treated with qHSC conditioned media. The study determined that qHSC induction of apoptosis required increased apoptosis-inducing factor (AIF) expression, nuclear localization and DNA fragmentation, and resulted in eventual cell death (Das et al., 2020). This data illustrates the ability of qHSCs to increase toxicity and sensitivity to established chemotherapeutic agents, such as doxorubicin (Das et al., 2020), and could lead to augmentation of therapeutic strategies already in existence to increase their success.

CLINICAL IMPLICATIONS AND RELEVANCE

Incidence of HCC in the U.S. has increased substantially in the past two decades (Singal et al., 2019; Das et al., 2020; Wu et al., 2020), and the American Cancer Society estimates 32,107 new cases of HCC will be diagnosed in 2020 along with 22,620 deaths (Society, 2020). Additionally, HCC incidence is three times higher in men than in women and the highest incidence is observed amongst patients greater than 70 years old with a steep mortality observed in patients ages 55–69 and older (Beal et al., 2017; Society, 2020). Unfortunately, over 80% of HCC patients present in advanced stages are not amenable to potentially curative surgical therapy, which combined with a paucity of effective systemic therapies, leads to a high morbidity and mortality rate (Zhou et al., 2018; Li D. et al., 2019). Ongoing investigation has focused on the process of tumor progression and potential therapeutic targets to create clinically relevant treatments for patients with HCC. Implicated in this research is the critical role of activated HSCs, from which 80–90% of HCC cells develop (Shiraha et al., 2020).

Metformin

As mentioned previously, TGFB and PDGF induce HSC activation to contribute to the cellular crosstalk between tumor and stromal cells (Zhang and Friedman, 2012; Tsuchida and Friedman, 2017). GDF15, a member of the TGFB superfamily, is a biomarker for stress responses as a result of cancer treatment damage such as hypoxia and chemotherapy




FIGURE 2 | The relationship amongst HCC tumor cells, tumor angiogenesis and HSCs in the hepatic TME. Quiescent HSCs are activated by numerous factors such as liver damage, inflammation, PDGF and TGFB signaling, hepatitis and viral infection. Additionally, HSC activation can be epigenetically regulated by demethylation of the tumor suppressor PTEN and IL-20 activation of HSCs via upregulation of TGFB1. Activated HSCs go on to produce ECM proteins and alter the ECM accompanied by proteoglycans and VEGF. Cytokines such as IL-6 are released which promote HCC tumor cell proliferation. CCN2 produced from HCC cells can activate HSCs, in contrast, CCN2 is inhibited through the oncogenic miRNA (oncomir) miR-212-3p which decreases HCC cell invasion *in vitro*. Other oncomirs secreted by HCC extracellular vesicles can regulate signaling between HSCs and HCC cells. In addition, IL-8, MMP-9, and Ang-1 contribute to tumor-associated angiogenesis. These factors allow for HCC tumor proliferation, invasion, and metastasis.

(Kelly et al., 2009; Corre et al., 2013) and may be a clinically relevant target for HCC. Common liver cancer therapy utilizes transarterial chemoembolization (TACE) that involves

chemotherapy embolization-induced hypoxia to damage HCC cells; however, this therapy also induces stress in the surrounding HCC tissue (Dong et al., 2018). As a result of the intense

cellular stress, GDF15 is secreted by the treatment-damaged HCC cells, and may have the ability to promote fibrosis through activated HSCs (Dong et al., 2018). To elucidate this process, Dong et al. (2018) demonstrated that HCC cells under “TACE-like” conditions showed increased levels of GDF15 via activation of the p38 MAPK, ERK1/2 and JNK pathways. Metformin, a common FDA approved drug, is known to target the JNK/p38MAPK pathway (Wu et al., 2011). Dong et al. (2018) further illustrated that in activated HSC cells, GDF15 promoted HSC proliferation and collagen production indicated by increased type I collagen protein levels cellular 5-Ethynyl-2'-deoxyuridine (Edu) incorporation, a thymidine analog which incorporates into the DNA of dividing cells (Salic and Mitchison, 2008). Conversely, Metformin was able to target the JNK pathway and suppress GDF15 expression resulting in decreased collagen synthesis and proliferation of activated HSCs *in vitro* and *in vivo* (Dong et al., 2018). These results suggest an opportunity to slow the progression of HCC by targeting activated HSCs with Metformin, a widely available drug.

Sorafenib

Another FDA approved drug, Sorafenib, is a multi-target anti-angiogenic tyrosine kinase inhibitor (Hasskarl, 2014). Sorafenib was the first systemic therapy approved for treatment of HCC after it was shown to increase mean survival time by 2–3 months (Llovet et al., 2008; Cheng et al., 2009). A recently published study suggested that a novel system of biodegradable dendritic polymeric nanoparticles loaded with Sorafenib enhanced HCC therapy (Li et al., 2020). Through the MTT assay, this system induced higher cytotoxicity of HCC cells than a PEG-conjugated nanoparticle system containing Sorafenib and free-Sorafenib *in vitro*. In addition, tumor growth was significantly subdued in mice with HepG2 xenografts, with minimal side effects (Li et al., 2020). Moreover, a previous nanoparticle related study published in 2018 demonstrated that combined delivery of Sorafenib with a mitogen activated protein kinase (MEK) inhibitor using C-X-C motif chemokine receptor 4 (CXCR4)-targeted nanoparticles reduced hepatic fibrosis and prevented tumor development (Sung et al., 2018). This study further assessed the effects of Sorafenib on activated HSCs and established that combined delivery of Sorafenib and a MEK inhibitor through CXCR4-targeted nanoparticles prevented ERK activation in activated HSCs and had anti-fibrotic effects in the CCl₄-mouse model (Sung et al., 2018).

Bevacizumab

Continued interest in the critical role of activated HSCs in the regulation of angiogenesis will likely increase in light of recently released preliminary results of the IMbrave150 trial which challenges the longstanding paradigm of first line Sorafenib for advanced unresectable HCC. This study was a phase III randomized control trial of 336 patients who received either atezolizumab (PD-L1 inhibitor) plus Bevacizumab (VEGF-A inhibitor) or Sorafenib. Preliminary results from the trial demonstrated a significant overall survival benefit, with patients in the combination arm not reaching a median survival versus a median survival of 13.2 months with Sorafenib ($p < 0.0001$)

(Cheng et al., 2019). Bevacizumab has been shown *in vivo* to decrease expression of profibrogenic genes TGFB and ACTA2, as well as decreasing overall HSC activation, altogether attenuating hepatic fibrosis in a CCl₄-rat model (Huang et al., 2013). Although final results of the study have not yet been published, it is widely expected to be practice changing and highlights the importance of both the immune and angiogenic crosstalk in the hepatic TME and HCC progression.

Potential Alternative Therapies

Finally, there is an interest in therapeutically targeting the cytokine-HSC interaction. An IL-8-neutralizing antibody (Zhu et al., 2015) and an anti-CCN2 neutralizing antibody which lead to reduced IL-6 production (Makino et al., 2018) have both shown therapeutic potential in suppressing tumor progression of HCC *in vitro* and *in vivo* with a xenograft murine model. These neutralizing antibodies, which both target interleukin cytokines, are potential clinical therapies focusing on the relationship between activated HSCs and TME. Additionally, a 2015 study established the angiogenin inhibitor neomycin as a potential HCC therapy. This study demonstrated that neomycin decreased HSC activation with conditioned media or recombinant angiogenin *in vitro* (Barcena et al., 2015). Furthermore, neomycin administration reduced tumor growth of HepG2-LX2 cells co-injected into mice, suggesting that angiogenin secretion by HCC cells favors tumor development via induction of HSC activation and ECM remodeling. These findings not only suggest that targeting angiogenin signaling may be of potential relevance in HCC management, but also establishes neomycin as a potential clinical treatment for HCC.

DISCUSSION

Hepatic stellate cells activation is the central event of hepatic fibrosis and the development of cirrhosis and HCC. A fundamental gap in knowledge is the crosstalk between activated HSCs, the hepatic ECM and HCC tumor cells. Recent studies have focused on targeted molecular therapeutic strategies for liver fibrosis and cirrhosis (van der Heide et al., 2019). Thus, utilizing multiple biomarkers may lead to optimized early detection of HCC (Ismail and Pinzani, 2011; Tuohetahuntala et al., 2017). Data published in 2017 suggested a new therapeutic option to target and increase NK activity in patients with chronic hepatitis infection preceding hepatic fibrosis (Shi et al., 2017). Other recent studies discussing potential targeted therapies such as PDGF-C and TGFB are under exploration. Ultimately, elucidating the mechanistic links between activated HSCs through all stages of fibrosis and cirrhosis will lead to a better understanding of HCC tumorigenesis.

Due to the intricate relationship between the hepatic TME, tumor development and HCC progression, recent studies have begun focusing on the role of activated HSCs, one of the prominent factors involved in the hepatic TME. The hepatic TME provides a niche which includes both cellular components, such as HSCs and non-cellular components, being the ECM and

ECM proteins. As activated HSCs become ECM producing cells during liver fibrosis, secreted chemokines, cytokines, and growth factors prime the overall TME for supporting HCC proliferation.

Owing to the complexity of the TME, it is difficult to therapeutically target one pathway as they all have functional redundancies; however, targeting a pivotal cellular component of the hepatic TME, such as activated HSCs, may be more feasible. Pertinent to this review, studies focusing on the role of activated HSCs in the TME could lead to activated HSC targeted therapies that may affect activated HSC related factors such as TGFB, PDGF, MMP-9, CCN2, and oncogenic miRNAs. Other targets that warrant further study and serve as promising areas for therapeutic exploration include IL-8, Ang-1, and Gli-1. A proposed schematic illustrating the relationship between activated HSCs in the hepatic TME and HCC is shown in **Figure 2**.

Collectively the literature covered in this review outline the significance of activated HSCs in the hepatic TME. In addition to cellular crosstalk within the TME, activated HSCs play a crucial role in HCC progression through the TME's non-cellular components. These recent studies provide examples of cytokines, growth factors, ECM components and microRNAs, which are all crucial non-cellular components of the hepatic TME involved in the relationship between activated HSCs and HCC development.

This review also concentrated on clinical implications that highlight potential therapies for HCC through targeting activated HSCs and their relationship with HCC cells. In addition to enhancing the efficacy of current therapeutic agents

such as Metformin, Sorafenib, and Atezolizumab/Bevacizumab, potential alternative therapies include neomycin and neutralizing antibodies against IL-22 and CCN2.

A deeper understanding of how the hepatic TME, most notably activated HSCs, interacts with the primary tumor and non-tumor cells will propel advances in effective diagnostic and prognostic tools. Ongoing investigations are imperative in order to develop more effective treatments for HCC and augment current therapies to increase their success.

AUTHOR CONTRIBUTIONS

AB and RB equally drafted the review, compiled the authors' contributions and references, wrote the manuscript, and produced the table and figures. RL, KP, KZ, DD, RB, AS, and KK contributed by writing and editing various sections within the manuscript. HD provided the overall support, intellectual input, critical evaluation, and all rounds of editing. All the authors read and approved the manuscript.

FUNDING

This work was supported by grants from the American Liver Foundation (A20-0085), American Gastroenterological Association (AGA2020-21-04), and the American Cancer Society (130042-IRG-16-244-10-IRG).

REFERENCES

- Alfarouk, K. O., Muddathir, A. K., and Shayoub, M. E. (2011). Tumor acidity as evolutionary spite. *Cancers* 3, 408–414. doi: 10.3390/cancers3010408
- Amann, T., Bataille, F., Spruss, T., Muhlbauer, M., Gabele, E., Scholmerich, J., et al. (2009). Activated hepatic stellate cells promote tumorigenicity of hepatocellular carcinoma. *Cancer Sci.* 100, 646–653. doi: 10.1111/j.1349-7006.2009.01087.x
- Ananthakrishnan, A., Gogineni, V., and Saeian, K. (2006). Epidemiology of primary and secondary liver cancers. *Semin. Intervent. Radiol.* 23, 47–63. doi: 10.1055/s-2006-939841
- Armstrong, G. L., Wasley, A., Simard, E. P., McQuillan, G. M., Kuhnert, W. L., and Alter, M. J. (2006). The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann. Intern. Med.* 144, 705–714. doi: 10.7326/0003-4819-144-10-200605160-00004
- Bandyopadhyay, K., Marrero, I., and Kumar, V. (2016). NKT cell subsets as key participants in liver physiology and pathology. *Cell. Mol. Immunol.* 13, 337–346. doi: 10.1038/cmi.2015.115
- Barazani, Y., Hiatt, J. R., Tong, M. J., and Busuttill, R. W. (2007). Chronic viral hepatitis and hepatocellular carcinoma. *World J. Surg.* 31, 1243–1248. doi: 10.1007/s00268-007-9041-3
- Barcana, C., Stefanovic, M., Tutusaus, A., Martinez-Nieto, G. A., Martinez, L., Garcia-Ruiz, C., et al. (2015). Angiogenin secretion from hepatoma cells activates hepatic stellate cells to amplify a self-sustained cycle promoting liver cancer. *Sci. Rep.* 5:7916. doi: 10.1038/srep07916
- Bartel, D. P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297. doi: 10.1016/s0092-8674(04)00045-5
- Beal, E. W., Tumin, D., Kabir, A., Moris, D., Zhang, X. F., Chakedis, J., et al. (2017). Trends in the mortality of hepatocellular carcinoma in the United States. *J. Gastrointest. Surg.* 21, 2033–2038. doi: 10.1007/s11605-017-3526-7
- Bian, E. B., Huang, C., Ma, T. T., Tao, H., Zhang, H., Cheng, C., et al. (2012). DNMT1-mediated PTEN hypermethylation confers hepatic stellate cell activation and liver fibrogenesis in rats. *Toxicol. Appl. Pharmacol.* 264, 13–22. doi: 10.1016/j.taap.2012.06.022
- Bjorklund, A. K., Forkel, M., Picelli, S., Konya, V., Theorell, J., Friberg, D., et al. (2016). The heterogeneity of human CD127(+) innate lymphoid cells revealed by single-cell RNA sequencing. *Nat. Immunol.* 17, 451–460. doi: 10.1038/ni.3368
- Blaner, W. S., O'Byrne, S. M., Wongsiriroj, N., Kluwe, J., D'Ambrosio, D. M., Jiang, H., et al. (2009). Hepatic stellate cell lipid droplets: a specialized lipid droplet for retinoid storage. *Biochim. Biophys. Acta* 1791, 467–473. doi: 10.1016/j.bbali.2008.11.001
- Borkham-Kamphorst, E., Herrmann, J., Stoll, D., Treptau, J., Gressner, A. M., and Weiskirchen, R. (2004). Dominant-negative soluble PDGF-beta receptor inhibits hepatic stellate cell activation and attenuates liver fibrosis. *Lab. Invest.* 84, 766–777. doi: 10.1038/labinvest.3700094
- Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R. L., Torre, L. A., and Jemal, A. (2018). Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* 68, 394–424. doi: 10.3322/caac.21492
- Breitkopf, K., Godoy, P., Ciucian, L., Singer, M. V., and Dooley, S. (2006). TGF-beta/Smad signaling in the injured liver. *Z Gastroenterol.* 44, 57–66. doi: 10.1055/s-2005-858989
- Breitkopf, K., Sawitza, I., Westhoff, J. H., Wickert, L., Dooley, S., and Gressner, A. M. (2005). Thrombospondin 1 acts as a strong promoter of transforming growth factor beta effects via two distinct mechanisms in hepatic stellate cells. *Gut* 54, 673–681. doi: 10.1136/gut.2004.042911
- Budhu, A., Forgues, M., Ye, Q. H., Jia, H. L., He, P., Zanetti, K. A., et al. (2006). Prediction of venous metastases, recurrence, and prognosis in hepatocellular carcinoma based on a unique immune response signature of the liver microenvironment. *Cancer Cell* 10, 99–111. doi: 10.1016/j.ccr.2006.06.016
- Budhu, A., and Wang, X. W. (2006). The role of cytokines in hepatocellular carcinoma. *J. Leukoc. Biol.* 80, 1197–1213. doi: 10.1189/jlb.0506297

- Cairns, P., Okami, K., Halachmi, S., Halachmi, N., Esteller, M., Herman, J. G., et al. (1997). Frequent inactivation of PTEN/MMAC1 in primary prostate cancer. *Cancer Res.* 57, 4997–5000.
- Campisi, J., and d'Adda di Fagnaga, F. (2007). Cellular senescence: when bad things happen to good cells. *Nat. Rev. Mol. Cell Biol.* 8, 729–740. doi: 10.1038/nrm2233
- Carloni, V., Luong, T. V., and Rombouts, K. (2014). Hepatic stellate cells and extracellular matrix in hepatocellular carcinoma: more complicated than ever. *Liver Int.* 34, 834–843. doi: 10.1111/liv.12465
- Chen, J. Q., Ou, Y. L., Huang, Z. P., Hong, Y. G., Tao, Y. P., Wang, Z. G., et al. (2019). MicroRNA-212-3p inhibits the Proliferation and Invasion of Human Hepatocellular Carcinoma cells by suppressing CCN2 expression. *Sci. Rep.* 9:9820. doi: 10.1038/s41598-019-46088-w
- Chen, J. S., Huang, X. H., Wang, Q., Chen, X. L., Fu, X. H., Tan, H. X., et al. (2010). FAK is involved in invasion and metastasis of hepatocellular carcinoma. *Clin. Exp. Metastasis* 27, 71–82. doi: 10.1007/s10585-010-9306-3
- Cheng, A. L., Kang, Y. K., Chen, Z., Tsao, C. J., Qin, S., Kim, J. S., et al. (2009). Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol.* 10, 25–34. doi: 10.1016/S1470-2045(08)70285-7
- Cheng, A. L., Qin, S., Ikeda, M., Galle, P., Ducreux, M., Zhu, A., et al. (2019). IMbrave150: efficacy and safety results from a phase III study evaluating atezolizumab (atezo) + bevacizumab (bev) vs sorafenib (Sor) as first treatment (tx) for patients (pts) with unresectable hepatocellular carcinoma (HCC). *Ann. Oncol.* 30, ix186–ix187. doi: 10.1093/annonc/mdz446.002
- Cheng, C. J., Bahal, R., Babar, I. A., Pincus, Z., Barrera, F., Liu, C., et al. (2015). MicroRNA silencing for cancer therapy targeted to the tumour microenvironment. *Nature* 518, 107–110. doi: 10.1038/nature13905
- Chiu, Y. S., Wei, C. C., Lin, Y. J., Hsu, Y. H., and Chang, M. S. (2014). IL-20 and IL-20R1 antibodies protect against liver fibrosis. *Hepatology* 60, 1003–1014. doi: 10.1002/hep.27189
- Choo, S. P., Tan, W. L., Goh, B. K. P., Tai, W. M., and Zhu, A. X. (2016). Comparison of hepatocellular carcinoma in Eastern versus Western populations. *Cancer* 122, 3430–3446. doi: 10.1002/ncr.30237
- Coll, M., El Taghdouini, A., Perea, L., Mannaerts, I., Vila-Casadesus, M., Blaya, D., et al. (2015). Integrative miRNA and gene expression profiling analysis of human quiescent hepatic stellate cells. *Sci. Rep.* 5:11549. doi: 10.1038/srep11549
- Collado, M., Blasco, M. A., and Serrano, M. (2007). Cellular senescence in cancer and aging. *Cell* 130, 223–233. doi: 10.1016/j.cell.2007.07.003
- Corre, J., Hebraud, B., and Bourin, P. (2013). Concise review: growth differentiation factor 15 in pathology: a clinical role? *Stem Cells Transl. Med.* 2, 946–952. doi: 10.5966/sctm.2013-0055
- Coulouarn, C., and Clement, B. (2014). Stellate cells and the development of liver cancer: therapeutic potential of targeting the stroma. *J. Hepatol.* 60, 1306–1309. doi: 10.1016/j.jhep.2014.02.003
- Coulouarn, C., Corlu, A., Glaise, D., Guenon, I., Thorgerirsson, S. S., and Clement, B. (2012). Hepatocyte-stellate cell cross-talk in the liver engenders a permissive inflammatory microenvironment that drives progression in hepatocellular carcinoma. *Cancer Res.* 72, 2533–2542. doi: 10.1158/0008-5472.CAN-11-3317
- D'Amico, G., Garcia-Tsao, G., and Pagliaro, L. (2006). Natural history and prognostic indicators of survival in cirrhosis: a systematic review of 118 studies. *J. Hepatol.* 44, 217–231. doi: 10.1016/j.jhep.2005.10.013
- Das, D., Fayazzadeh, E., Li, X., Koirala, N., Wadera, A., Lang, M., et al. (2020). Quiescent hepatic stellate cells induce toxicity and sensitivity to doxorubicin in cancer cells through a caspase-independent cell death pathway: central role of apoptosis-inducing factor. *J. Cell Physiol.* 235, 6167–6182. doi: 10.1002/jcp.29545
- Daugaard, I., Sanders, K. J., Idica, A., Vittayarukskul, K., Hamdorf, M., Krog, J. D., et al. (2017). miR-151a induces partial EMT by regulating E-cadherin in NSCLC cells. *Oncogenesis* 6:e366. doi: 10.1038/oncsis.2017.66
- de Franchis, R. (2000). Updating consensus in portal hypertension: report of the Baveno III consensus workshop on definitions, methodology and therapeutic strategies in portal hypertension. *J. Hepatol.* 33, 846–852. doi: 10.1016/s0168-8278(00)80320-7
- DeLeve, L. D., and Maretta-Mira, A. C. (2017). Liver sinusoidal endothelial cell: an update. *Semin. Liver Dis.* 37, 377–387. doi: 10.1055/s-0037-1617455
- Dewidar, B., Meyer, C., Dooley, S., and Meindl-Beinker, A. N. (2019). TGF-beta in hepatic stellate cell activation and liver fibrogenesis—updated 2019. *Cells* 8:1419. doi: 10.3390/cells8111419
- Dong, G., Ma, M., Lin, X., Liu, H., Gao, D., Cui, J., et al. (2018). Treatment-damaged hepatocellular carcinoma promotes activities of hepatic stellate cells and fibrosis through GDF15. *Exp. Cell Res.* 370, 468–477. doi: 10.1016/j.yexcr.2018.07.011
- El-Serag, H. B. (2012). Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology* 142, 1264–1273.e1. doi: 10.1053/j.gastro.2011.12.061
- Engel, M. E., McDonnell, M. A., Law, B. K., and Moses, H. L. (1999). Interdependent SMAD and JNK signaling in transforming growth factor-beta-mediated transcription. *J. Biol. Chem.* 274, 37413–37420. doi: 10.1074/jbc.274.52.37413
- Fabregat, I., Moreno-Caceres, J., Sanchez, A., Dooley, S., Dewidar, B., Giannelli, G., et al. (2016). TGF-beta signalling and liver disease. *FEBS J.* 283, 2219–2232. doi: 10.1111/febs.13665
- Fan, W., Liu, T., Chen, W., Hammad, S., Longerich, T., Hausser, I., et al. (2019). ECM1 prevents activation of transforming growth factor beta, hepatic stellate cells, and fibrogenesis in mice. *Gastroenterology* 157, 1352–1367.e13. doi: 10.1053/j.gastro.2019.07.036
- Friedman, S. L. (2000). Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *J. Biol. Chem.* 275, 2247–2250. doi: 10.1074/jbc.275.4.2247
- Friedman, S. L. (2008a). Hepatic stellate cells: protean, multifunctional, and enigmatic cells of the liver. *Physiol. Rev.* 88, 125–172. doi: 10.1152/physrev.00013.2007
- Friedman, S. L. (2008b). Mechanisms of hepatic fibrogenesis. *Gastroenterology* 134, 1655–1669. doi: 10.1053/j.gastro.2008.03.003
- Friedman, S. L., Sheppard, D., Duffield, J. S., and Violette, S. (2013). Therapy for fibrotic diseases: nearing the starting line. *Sci. Transl. Med.* 5:167sr161. doi: 10.1126/scitranslmed.3004700
- Fu, J., Xu, D., Liu, Z., Shi, M., Zhao, P., Fu, B., et al. (2007). Increased regulatory T cells correlate with CD8 T-cell impairment and poor survival in hepatocellular carcinoma patients. *Gastroenterology* 132, 2328–2339. doi: 10.1053/j.gastro.2007.03.102
- Furuta, J., Umabayashi, Y., Miyamoto, K., Kikuchi, K., Otsuka, F., Sugimura, T., et al. (2004). Promoter methylation profiling of 30 genes in human malignant melanoma. *Cancer Sci.* 95, 962–968. doi: 10.1111/j.1349-7006.2004.tb03184.x
- Gabrilovich, D. I., and Nagaraj, S. (2009). Myeloid-derived suppressor cells as regulators of the immune system. *Nat. Rev. Immunol.* 9, 162–174. doi: 10.1038/nri2506
- Garzon, R., Calin, G. A., and Croce, C. M. (2009). MicroRNAs in cancer. *Annu. Rev. Med.* 60, 167–179. doi: 10.1146/annurev.med.59.053006.104707
- Gines, P., Quintero, E., Arroyo, V., Teres, J., Bruguera, M., Rimola, A., et al. (1987). Compensated cirrhosis: natural history and prognostic factors. *Hepatology* 7, 122–128. doi: 10.1002/hep.1840070124
- Glassner, A., Eisenhardt, M., Kramer, B., Korner, C., Coenen, M., Sauerbruch, T., et al. (2012). NK cells from HCV-infected patients effectively induce apoptosis of activated primary human hepatic stellate cells in a TRAIL-, FasL- and NKG2D-dependent manner. *Lab. Invest.* 92, 967–977. doi: 10.1038/labinvest.2012.54
- Grosse, L., Wagner, N., Emelyanov, A., Molina, C., Lacas-Gervais, S., Wagner, K. D., et al. (2020). Defined p16^{High} senescent cell types are indispensable for mouse healthspan. *Cell Metab.* 32, 87–99.e6. doi: 10.1016/j.cmet.2020.05.002
- Gu, C., Wang, Z., Jin, Z., Li, G., Kou, Y., Jia, Z., et al. (2017). MicroRNA-212 inhibits the proliferation, migration and invasion of renal cell carcinoma by targeting X-linked inhibitor of apoptosis protein (XIAP). *Oncotarget* 8, 92119–92133. doi: 10.18632/oncotarget.20786
- Gur, C., Doron, S., Kfir-Erenfeld, S., Horwitz, E., Abu-Tair, L., Safadi, R., et al. (2012). NKp46-mediated killing of human and mouse hepatic stellate cells attenuates liver fibrosis. *Gut* 61, 885–893. doi: 10.1136/gutjnl-2011-301400
- Hall-Glenn, F., and Lyons, K. M. (2011). Roles for CCN2 in normal physiological processes. *Cell Mol. Life Sci.* 68, 3209–3217. doi: 10.1007/s00018-011-0782-7
- Han, S., Han, L., Yao, Y., Sun, H., Zan, X., and Liu, Q. (2014). Activated hepatic stellate cells promote hepatocellular carcinoma cell migration and invasion via

- the activation of FAK-MMP9 signaling. *Oncol. Rep.* 31, 641–648. doi: 10.3892/or.2013.2872
- Han, Y., Jo, H., Cho, J. H., Dhanasekaran, D. N., and Song, Y. S. (2019). Resveratrol as a tumor-suppressive nutraceutical modulating tumor microenvironment and malignant behaviors of cancer. *Int. J. Mol. Sci.* 20:925. doi: 10.3390/ijms20040925
- Hanafusa, H., Ninomiya-Tsuji, J., Masuyama, N., Nishita, M., Fujisawa, J., Shibuya, H., et al. (1999). Involvement of the p38 mitogen-activated protein kinase pathway in transforming growth factor-beta-induced gene expression. *J. Biol. Chem.* 274, 27161–27167. doi: 10.1074/jbc.274.38.27161
- Hanahan, D., and Weinberg, R. A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646–674. doi: 10.1016/j.cell.2011.02.013
- Hasskarl, J. (2014). Sorafenib: targeting multiple tyrosine kinases in cancer. *Recent Expressions Cancer Res.* 201, 145–164. doi: 10.1007/978-3-642-54490-3_8
- Heindryckx, F., and Gerwens, P. (2015). Targeting the tumor stroma in hepatocellular carcinoma. *World J. Hepatol.* 7, 165–176. doi: 10.4254/wjh.v7.i2.165
- Hernandez-Gea, V., Toffanin, S., Friedman, S. L., and Llovet, J. M. (2013). Role of the microenvironment in the pathogenesis and treatment of hepatocellular carcinoma. *Gastroenterology* 144, 512–527. doi: 10.1053/j.gastro.2013.01.002
- Higashi, T., Friedman, S. L., and Hoshida, Y. (2017). Hepatic stellate cells as key target in liver fibrosis. *Adv. Drug Deliv. Rev.* 121, 27–42. doi: 10.1016/j.addr.2017.05.007
- Hoare, M., Das, T., and Alexander, G. (2010). Ageing, telomeres, senescence, and liver injury. *J. Hepatol.* 53, 950–961. doi: 10.1016/j.jhep.2010.06.009
- Hoechst, B., Ormandy, L. A., Ballmaier, M., Lehner, F., Kruger, C., Manns, M. P., et al. (2008). A new population of myeloid-derived suppressor cells in hepatocellular carcinoma patients induces CD4(+)CD25(+)Foxp3(+) T cells. *Gastroenterology* 135, 234–243. doi: 10.1053/j.gastro.2008.03.020
- Hou, X., Yu, F., Man, S., Huang, D., Zhang, Y., Liu, M., et al. (2012). Negative regulation of *Schistosoma japonicum* egg-induced liver fibrosis by natural killer cells. *PLoS Negl. Trop. Dis.* 6:e1456. doi: 10.1371/journal.pntd.0001456
- Hsieh, C. C., Chou, H. S., Yang, H. R., Lin, F., Bhatt, S., Qin, J., et al. (2013). The role of complement component 3 (C3) in differentiation of myeloid-derived suppressor cells. *Blood* 121, 1760–1768. doi: 10.1182/blood-2012-06-440214
- Huang, J. L., Fu, Y. P., Gan, W., Liu, G., Zhou, P. Y., Zhou, C., et al. (2020). Hepatic stellate cells promote the progression of hepatocellular carcinoma through microRNA-1246-RORalpha-Wnt/beta-Catenin axis. *Cancer Lett.* 476, 140–151. doi: 10.1016/j.canlet.2020.02.012
- Huang, Y., Feng, H., Kan, T., Huang, B., Zhang, M., Li, Y., et al. (2013). Bevacizumab attenuates hepatic fibrosis in rats by inhibiting activation of hepatic stellate cells. *PLoS One* 8:e73492. doi: 10.1371/journal.pone.0073492
- Hui, A. Y., Dannenberg, A. J., Sung, J. J., Subbaramaiah, K., Du, B., Olinga, P., et al. (2004). Prostaglandin E2 inhibits transforming growth factor beta 1-mediated induction of collagen alpha 1(I) in hepatic stellate cells. *J. Hepatol.* 41, 251–258. doi: 10.1016/j.jhep.2004.04.033
- Ikura, Y., Morimoto, H., Ogami, M., Jomura, H., Ikeoka, N., and Sakurai, M. (1997). Expression of platelet-derived growth factor and its receptor in livers of patients with chronic liver disease. *J. Gastroenterol.* 32, 496–501. doi: 10.1007/bf02934089
- Ishizaki, M., Ashida, K., Higashi, T., Nakatsukasa, H., Kaneyoshi, T., Fujiwara, K., et al. (2001). The formation of capsule and septum in human hepatocellular carcinoma. *Virchows Arch.* 438, 574–580. doi: 10.1007/s004280000391
- Ismail, M. H., and Pinzani, M. (2011). Reversal of hepatic fibrosis: pathophysiological basis of antifibrotic therapies. *Hepat. Med.* 3, 69–80. doi: 10.2147/HMER.S9051
- Jeong, W. I., Park, O., Suh, Y. G., Byun, J. S., Park, S. Y., Choi, E., et al. (2011). Suppression of innate immunity (natural killer cell/interferon-gamma) in the advanced stages of liver fibrosis in mice. *Hepatology* 53, 1342–1351. doi: 10.1002/hep.24190
- Jia, Y. L., Shi, L., Zhou, J. N., Fu, C. J., Chen, L., Yuan, H. F., et al. (2011). Epimorphin promotes human hepatocellular carcinoma invasion and metastasis through activation of focal adhesion kinase/extracellular signal-regulated kinase/matrix metalloproteinase-9 axis. *Hepatology* 54, 1808–1818. doi: 10.1002/hep.24562
- Jiang, R., Tan, Z., Deng, L., Chen, Y., Xia, Y., Gao, Y., et al. (2011). Interleukin-22 promotes human hepatocellular carcinoma by activation of STAT3. *Hepatology* 54, 900–909. doi: 10.1002/hep.24486
- Jin, H., Lian, N., Zhang, F., Chen, L., Chen, Q., Lu, C., et al. (2016). Activation of PPARgamma/P53 signaling is required for curcumin to induce hepatic stellate cell senescence. *Cell Death Dis.* 7:e2189. doi: 10.1038/cddis.2016.92
- Joyce, J. A., and Fearon, D. T. (2015). T cell exclusion, immune privilege, and the tumor microenvironment. *Science* 348, 74–80. doi: 10.1126/science.aaa6204
- Jun, J. I., and Lau, L. F. (2011). Taking aim at the extracellular matrix: CCN proteins as emerging therapeutic targets. *Nat. Rev. Drug Discov.* 10, 945–963. doi: 10.1038/nrd3599
- Kanwal, F., Hoang, T., Kramer, J. R., Asch, S. M., Goetz, M. B., Zeringue, A., et al. (2011). Increasing prevalence of HCC and cirrhosis in patients with chronic hepatitis C virus infection. *Gastroenterology* 140, 1182–1188.e1. doi: 10.1053/j.gastro.2010.12.032
- Karube, Y., Tanaka, H., Osada, H., Tomida, S., Tatematsu, Y., Yanagisawa, K., et al. (2005). Reduced expression of Dicer associated with poor prognosis in lung cancer patients. *Cancer Sci.* 96, 111–115. doi: 10.1111/j.1349-7006.2005.00015.x
- Kelly, J. A., Lucia, M. S., and Lambert, J. R. (2009). p53 controls prostate-derived factor/macrophage inhibitory cytokine/NSAID-activated gene expression in response to cell density, DNA damage and hypoxia through diverse mechanisms. *Cancer Lett.* 277, 38–47. doi: 10.1016/j.canlet.2008.11.013
- Kodama, T., Takehara, T., Hikita, H., Shimizu, S., Shigekawa, M., Tsunematsu, H., et al. (2011). Increases in p53 expression induce CCN2 synthesis by mouse and human hepatocytes and result in liver fibrosis in mice. *J. Clin. Invest.* 121, 3343–3356. doi: 10.1172/JCI44957
- Kosaka, N. (2016). Decoding the secret of cancer by means of extracellular vesicles. *J. Clin. Med.* 5:22. doi: 10.3390/jcm5020022
- Kramer, B., Korner, C., Kebschull, M., Glassner, A., Eisenhardt, M., Nischalke, H. D., et al. (2012). Natural killer p46High expression defines a natural killer cell subset that is potentially involved in control of hepatitis C virus replication and modulation of liver fibrosis. *Hepatology* 56, 1201–1213. doi: 10.1002/hep.25804
- Krizhanovskiy, V., Yon, M., Dickins, R. A., Hearn, S., Simon, J., Miething, C., et al. (2008). Senescence of activated stellate cells limits liver fibrosis. *Cell* 134, 657–667. doi: 10.1016/j.cell.2008.06.049
- Kyrylkova, K., Kyryachenko, S., Leid, M., and Kioussi, C. (2012). Detection of apoptosis by TUNEL assay. *Methods Mol. Biol.* 887, 41–47. doi: 10.1007/978-1-61779-860-3_5
- Lachowski, D., Cortes, E., Rice, A., Pinato, D., Rombouts, K., and Del Rio Hernandez, A. (2019). Matrix stiffness modulates the activity of MMP-9 and TIMP-1 in hepatic stellate cells to perpetuate fibrosis. *Sci. Rep.* 9:7299. doi: 10.1038/s41598-019-43759-6
- Lashen, S. A., Elshafei, M. M., Hablass, F. H., Alsayed, E. A., and Hassan, A. A. (2020). Liver stiffness as a predictor of hepatocellular carcinoma behavior in patients with hepatitis C related liver cirrhosis. *Hepatobiliary Pancreat Dis. Int.* 19, 22–28. doi: 10.1016/j.hbpd.2019.11.004
- Lee, U. E., and Friedman, S. L. (2011). Mechanisms of hepatic fibrogenesis. *Best Pract. Res. Clin. Gastroenterol.* 25, 195–206. doi: 10.1016/j.bpg.2011.02.005
- Li, D., Sedano, S., Allen, R., Gong, J., Cho, M., and Sharma, S. (2019). Current treatment landscape for advanced hepatocellular carcinoma: patient outcomes and the impact on quality of life. *Cancers* 11:841. doi: 10.3390/cancers11060841
- Li, J., Yan, Y., Ang, L., Li, X., Liu, C., Sun, B., et al. (2019). Extracellular vesicles-derived OncomiRs mediate communication between cancer cells and cancer-associated hepatic stellate cells in hepatocellular carcinoma microenvironment. *Carcinogenesis* 41, 223–234. doi: 10.1093/carcin/bgz096
- Li, Y., Turpin, C. P., and Wang, S. (2017). Role of thrombospondin 1 in liver diseases. *Hepatol. Res.* 47, 186–193. doi: 10.1111/hepr.12787
- Li, Z., Ye, L., Liu, J., Lian, D., and Li, X. (2020). Sorafenib-loaded nanoparticles based on biodegradable dendritic polymers for enhanced therapy of hepatocellular carcinoma. *Int. J. Nanomed.* 15, 1469–1480. doi: 10.2147/IJN.S237335
- Lin, N., Meng, L., Lin, J., Chen, S., Zhang, P., Chen, Q., et al. (2020). Activated hepatic stellate cells promote angiogenesis in hepatocellular carcinoma by secreting angiopoietin-1. *J. Cell Biochem.* 121, 1441–1451. doi: 10.1002/jcb.29380
- Lipson, K. E., Wong, C., Teng, Y., and Spong, S. (2012). CCN2 is a central mediator of tissue remodeling and fibrosis and its inhibition can reverse the process of fibrosis. *Fibrogenesis Tissue Repair* 5(Suppl. 1):S24. doi: 10.1186/1755-1536-5-S1-S24

- Liu, Y., Liu, H., Meyer, C., Li, J., Nadalin, S., Konigsrainer, A., et al. (2013). Transforming growth factor-beta (TGF-beta)-mediated connective tissue growth factor (CCN2) expression in hepatic stellate cells requires Stat3 signaling activation. *J. Biol. Chem.* 288, 30708–30719. doi: 10.1074/jbc.M113.478685
- Lovet, J. M., Ricci, S., Mazzaferro, V., Hilgard, P., Gane, E., Blanc, J. F., et al. (2008). Sorafenib in advanced hepatocellular carcinoma. *N. Engl. J. Med.* 359, 378–390. doi: 10.1056/NEJMoa0708857
- Lozano, R., Naghavi, M., Foreman, K., Lim, S., Shibuya, K., Aboyans, V., et al. (2012). Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 380, 2095–2128. doi: 10.1016/S0140-6736(12)61728-0
- Lu, C., Rong, D., Zhang, B., Zheng, W., Wang, X., Chen, Z., et al. (2019). Current perspectives on the immunosuppressive tumor microenvironment in hepatocellular carcinoma: challenges and opportunities. *Mol. Cancer* 18:130. doi: 10.1186/s12943-019-1047-6
- Lu, J., Getz, G., Miska, E. A., Alvarez-Saavedra, E., Lamb, J., Peck, D., et al. (2005). MicroRNA expression profiles classify human cancers. *Nature* 435, 834–838. doi: 10.1038/nature03702
- Lv, X., Fang, C., Yin, R., Qiao, B., Shang, R., Wang, J., et al. (2017). Agrin para-secreted by PDGF-activated human hepatic stellate cells promotes hepatocarcinogenesis in vitro and in vivo. *Oncotarget* 8, 105340–105355. doi: 10.18632/oncotarget.22186
- Maher, J. J. (2001). Interactions between hepatic stellate cells and the immune system. *Semin. Liver Dis.* 21, 417–426. doi: 10.1055/s-2001-17555
- Makino, Y., Hikita, H., Kodama, T., Shigekawa, M., Yamada, R., Sakamori, R., et al. (2018). CCN2 Mediates tumor-stroma interactions between hepatoma cells and hepatic stellate cells to accelerate HCC progression. *Cancer Res.* 78, 4902–4914. doi: 10.1158/0008-5472.CAN-17-3844
- Melhem, A., Muhanna, N., Bishara, A., Alvarez, C. E., Ilan, Y., Bishara, T., et al. (2006). Anti-fibrotic activity of NK cells in experimental liver injury through killing of activated HSC. *J. Hepatol.* 45, 60–71. doi: 10.1016/j.jhep.2005.12.025
- Meng, X. M., Nikolic-Paterson, D. J., and Lan, H. Y. (2016). TGF-beta: the master regulator of fibrosis. *Nat. Rev. Nephrol.* 12, 325–338. doi: 10.1038/nrneph.2016.48
- Merritt, W. M., Lin, Y. G., Han, L. Y., Kamat, A. A., Spannuth, W. A., Schmandt, R., et al. (2008). Dicer, Drosha, and outcomes in patients with ovarian cancer. *N. Engl. J. Med.* 359, 2641–2650. doi: 10.1056/NEJMoa0803785
- Mirmohammadsadegh, A., Marini, A., Nambiar, S., Hassan, M., Tannapfel, A., Ruzicka, T., et al. (2006). Epigenetic silencing of the PTEN gene in melanoma. *Cancer Res.* 66, 6546–6552. doi: 10.1158/0008-5472.CAN-06-0384
- Mittal, K., Ebos, J., and Rini, B. (2014). Angiogenesis and the tumor microenvironment: vascular endothelial growth factor and beyond. *Semin. Oncol.* 41, 235–251. doi: 10.1053/j.seminoncol.2014.02.007
- Muhanna, N., Abu Tair, L., Doron, S., Amer, J., Azzeh, M., Mahamid, M., et al. (2011). Amelioration of hepatic fibrosis by NK cell activation. *Gut* 60, 90–98. doi: 10.1136/gut.2010.211136
- Muhanna, N., Horani, A., Doron, S., and Safadi, R. (2007). Lymphocyte-hepatic stellate cell proximity suggests a direct interaction. *Clin. Exp. Immunol.* 148, 338–347. doi: 10.1111/j.1365-2249.2007.03353.x
- Murphy-Ullrich, J. E., and Suto, M. J. (2018). Thrombospondin-1 regulation of latent TGF-beta activation: a therapeutic target for fibrotic disease. *Matrix Biol.* 6, 28–43. doi: 10.1016/j.matbio.2017.12.009
- Natarajan, V., Harris, E. N., and Kidambi, S. (2017). SECs (Sinusoidal Endothelial Cells), liver microenvironment, and fibrosis. *BioMed. Res. Int.* 2017:4097205. doi: 10.1155/2017/4097205
- Nishizawa, H., Iguchi, G., Fukuoka, H., Takahashi, M., Suda, K., Bando, H., et al. (2016). IGF-I induces senescence of hepatic stellate cells and limits fibrosis in a p53-dependent manner. *Sci. Rep.* 6:34605. doi: 10.1038/srep34605
- Nordenstedt, H., White, D. L., and El-Serag, H. B. (2010). The changing pattern of epidemiology in hepatocellular carcinoma. *Dig. Liver Dis.* 42(Suppl. 3), S206–S214. doi: 10.1016/S1590-8658(10)60507-5
- Novikova, M. V., Khromova, N. V., and Kopnin, P. B. (2017). Components of the hepatocellular carcinoma microenvironment and their role in tumor progression. *Biochemistry (Mosc)* 82, 861–873. doi: 10.1134/S0006297917080016
- Page, A., Paoli, P., Moran Salvador, E., White, S., French, J., and Mann, J. (2016). Hepatic stellate cell transdifferentiation involves genome-wide remodeling of the DNA methylation landscape. *J. Hepatol.* 64, 661–673. doi: 10.1016/j.jhep.2015.11.024
- Pan, P. Y., Wang, G. X., Yin, B., Ozao, J., Ku, T., Divino, C. M., et al. (2008). Reversion of immune tolerance in advanced malignancy: modulation of myeloid-derived suppressor cell development by blockade of stem-cell factor function. *Blood* 111, 219–228. doi: 10.1182/blood-2007-04-086835
- Parkin, D. M., Bray, F., Ferlay, J., and Pisani, P. (2005). Global cancer statistics, 2002. *CA Cancer J. Clin.* 55, 74–108. doi: 10.3322/canjclin.55.2.74
- Pinzani, M., Milani, S., Herbst, H., DeFranco, R., Grappone, C., Gentilini, A., et al. (1996). Expression of platelet-derived growth factor and its receptors in normal human liver and during active hepatic fibrogenesis. *Am. J. Pathol.* 148, 785–800.
- Poisson, J., Lemoine, S., Boulanger, C., Durand, F., Moreau, R., Valla, D., et al. (2017). Liver sinusoidal endothelial cells: physiology and role in liver diseases. *J. Hepatol.* 66, 212–227. doi: 10.1016/j.jhep.2016.07.009
- Racanelli, V., and Rehermann, B. (2006). The liver as an immunological organ. *Hepatology* 43(2 Suppl. 1), S54–S62. doi: 10.1002/hep.21060
- Radaeva, S., Sun, R., Jaruga, B., Nguyen, V. T., Tian, Z., and Gao, B. (2006). Natural killer cells ameliorate liver fibrosis by killing activated stellate cells in NKG2D-dependent and tumor necrosis factor-related apoptosis-inducing ligand-dependent manners. *Gastroenterology* 130, 435–452. doi: 10.1053/j.gastro.2005.10.055
- Radaeva, S., Wang, L., Radaev, S., Jeong, W. I., Park, O., and Gao, B. (2007). Retinoic acid signaling sensitizes hepatic stellate cells to NK cell killing via upregulation of NK cell activating ligand RAE1. *Am. J. Physiol. Gastrointest. Liver Physiol.* 293, G809–G816. doi: 10.1152/ajpgi.00212.2007
- Rawla, P., Sunkara, T., Muralidharan, P., and Raj, J. P. (2018). Update in global trends and aetiology of hepatocellular carcinoma. *Contemp. Oncol. (Pozn)* 22, 141–150. doi: 10.5114/wo.2018.78941
- Roman-Gomez, J., Jimenez-Velasco, A., Castillejo, J. A., Agirre, X., Barrios, M., Navarro, G., et al. (2004). Promoter hypermethylation of cancer-related genes: a strong independent prognostic factor in acute lymphoblastic leukemia. *Blood* 104, 2492–2498. doi: 10.1182/blood-2004-03-0954
- Sager, R. (1991). Senescence as a mode of tumor suppression. *Environ. Health Perspect.* 93, 59–62. doi: 10.1289/ehp.919359
- Salic, A., and Mitchison, T. J. (2008). A chemical method for fast and sensitive detection of DNA synthesis in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 105, 2415–2420. doi: 10.1073/pnas.0712168105
- Salvesen, H. B., MacDonald, N., Ryan, A., Jacobs, I. J., Lynch, E. D., Akslen, L. A., et al. (2001). PTEN methylation is associated with advanced stage and microsatellite instability in endometrial carcinoma. *Int. J. Cancer* 91, 22–26. doi: 10.1002/1097-0215(20010101)91:1<22::aid-ijc1002<3.0.co;2-s
- Scheau, C., Badarau, I. A., Costache, R., Caruntu, C., Mihai, G. L., Didilescu, A. C., et al. (2019). The Role of matrix metalloproteinases in the epithelial-mesenchymal transition of hepatocellular carcinoma. *Anal. Cell Pathol. (Amst)* 2019:9423907. doi: 10.1155/2019/9423907
- Schon, H. T., and Weiskirchen, R. (2014). Immunomodulatory effects of transforming growth factor-beta in the liver. *Hepatobiliary Surg. Nutr.* 3, 386–406. doi: 10.3978/j.issn.2304-3881.2014.11.06
- Seitz, T., Freese, K., Dietrich, P., Thasler, W. E., Bosserhoff, A., and Hellerbrand, C. (2020). Fibroblast Growth Factor 9 is expressed by activated hepatic stellate cells and promotes progression of hepatocellular carcinoma. *Sci. Rep.* 10:4546. doi: 10.1038/s41598-020-61510-4
- Sevic, I., Spinelli, F. M., Cantero, M. J., Reszegi, A., Kovalszky, I., Garcia, M. G., et al. (2019). “The role of the tumor microenvironment in the development and progression of hepatocellular carcinoma,” in *Hepatocellular Carcinoma*, ed. J. E. E. Tirnitz-Parker (Brisbane: Codon Publications).
- Shi, J., Zhao, J., Zhang, X., Cheng, Y., Hu, J., Li, Y., et al. (2017). Activated hepatic stellate cells impair NK cell anti-fibrosis capacity through a TGF-beta-dependent emperipolesis in HBV cirrhotic patients. *Sci. Rep.* 7:44544. doi: 10.1038/srep44544
- Shiraha, H., Iwamura, M., and Okada, H. (2020). Hepatic stellate cells in liver tumor. *Adv. Exp. Med. Biol.* 1234, 43–56. doi: 10.1007/978-3-030-37184-5_4
- Singal, A. G., Parikh, N. D., Rich, N. E., John, B. V., and Pillai, A. (2019). “Hepatocellular carcinoma surveillance and staging,” in *Hepatocellular Carcinoma: Translational Precision Medicine Approaches*, ed. Y. Hoshida (Cham: Humana Press), 27–51. doi: 10.1007/978-3-030-21540-8_2
- Society, A. C. (2020). *Cancer Facts & Figures 2020*. Atlanta: American Cancer Society.

- Sokolovic, A., Sokolovic, M., Boers, W., Elferink, R. P., and Bosma, P. J. (2010). Insulin-like growth factor binding protein 5 enhances survival of LX2 human hepatic stellate cells. *Fibrogenesis Tissue Repair* 3:3. doi: 10.1186/1755-1536-3-3
- Soria, J. C., Lee, H. Y., Lee, J. I., Wang, L., Issa, J. P., Kemp, B. L., et al. (2002). Lack of PTEN expression in non-small cell lung cancer could be related to promoter methylation. *Clin. Cancer Res.* 8, 1178–1184.
- Spill, F., Reynolds, D. S., Kamm, R. D., and Zaman, M. H. (2016). Impact of the physical microenvironment on tumor progression and metastasis. *Curr. Opin. Biotechnol.* 40, 41–48. doi: 10.1016/j.copbio.2016.02.007
- Stock, P., Monga, D., Tan, X., Micsenyi, A., Loizos, N., and Monga, S. P. (2007). Platelet-derived growth factor receptor- α : a novel therapeutic target in human hepatocellular cancer. *Mol. Cancer Ther.* 6, 1932–1941. doi: 10.1158/1535-7163.MCT-06-0720
- Sun, S. J., Wang, N., Sun, Z. W., Chen, J., and Cui, H. W. (2018). MiR-5692a promotes the invasion and metastasis of hepatocellular carcinoma via MMP9. *Eur. Rev. Med. Pharmacol. Sci.* 22, 4869–4878. doi: 10.26355/eurrev_201808_15623
- Sung, Y. C., Liu, Y. C., Chao, P. H., Chang, C. C., Jin, P. R., Lin, T. T., et al. (2018). Combined delivery of sorafenib and a MEK inhibitor using CXCR4-targeted nanoparticles reduces hepatic fibrosis and prevents tumor development. *Theranostics* 8, 894–905. doi: 10.7150/thno.21168
- Suzuki, H. I., Katsura, A., Matsuyama, H., and Miyazono, K. (2015). MicroRNA regulons in tumor microenvironment. *Oncogene* 34, 3085–3094. doi: 10.1038/onc.2014.254
- Syn, W. K., Agboola, K. M., Swiderska, M., Michelotti, G. A., Liaskou, E., Pang, H., et al. (2012). NKT-associated hedgehog and osteopontin drive fibrogenesis in non-alcoholic fatty liver disease. *Gut* 61, 1323–1329. doi: 10.1136/gutjnl-2011-301857
- Tahmasebi Birgani, M., and Carloni, V. (2017). Tumor microenvironment, a paradigm in hepatocellular carcinoma progression and therapy. *Int. J. Mol. Sci.* 18:405. doi: 10.3390/ijms18020405
- Takashima, M., Parsons, C. J., Ikejima, K., Watanabe, S., White, E. S., and Rippe, R. A. (2009). The tumor suppressor protein PTEN inhibits rat hepatic stellate cell activation. *J. Gastroenterol.* 44, 847–855. doi: 10.1007/s00535-009-0073-3
- Tang, T., Huan, L., Zhang, S., Zhou, H., Gu, L., Chen, X., et al. (2017). MicroRNA-212 functions as a tumor-suppressor in human non-small cell lung cancer by targeting SOX4. *Oncol. Rep.* 38, 2243–2250. doi: 10.3892/or.2017.5885
- Tao, H., Huang, C., Yang, J. J., Ma, T. T., Bian, E. B., Zhang, L., et al. (2011). MeCP2 controls the expression of RASAL1 in the hepatic fibrosis in rats. *Toxicology* 290, 327–333. doi: 10.1016/j.tox.2011.10.011
- Theocharis, A. D., Skandalis, S. S., Tzanakakis, G. N., and Karamanos, N. K. (2010). Proteoglycans in health and disease: novel roles for proteoglycans in malignancy and their pharmacological targeting. *FEBS J.* 277, 3904–3923. doi: 10.1111/j.1742-4658.2010.07800.x
- Thompson, A. I., Conroy, K. P., and Henderson, N. C. (2015). Hepatic stellate cells: central modulators of hepatic carcinogenesis. *BMC Gastroenterol.* 15:63. doi: 10.1186/s12876-015-0291-5
- Tsushima, T., and Friedman, S. L. (2017). Mechanisms of hepatic stellate cell activation. *Nat. Rev. Gastroenterol. Hepatol.* 14, 397–411. doi: 10.1038/nrgastro.2017.38
- Tuohetahuntala, M., Molenaar, M. R., Spee, B., Brouwers, J. F., Wubbolts, R., Houweling, M., et al. (2017). Lysosome-mediated degradation of a distinct pool of lipid droplets during hepatic stellate cell activation. *J. Biol. Chem.* 292, 12436–12448. doi: 10.1074/jbc.M117.778472
- van der Heide, D., Weiskirchen, R., and Bansal, R. (2019). Therapeutic targeting of hepatic macrophages for the treatment of liver diseases. *Front. Immunol.* 10:2852. doi: 10.3389/fimmu.2019.02852
- van Eggermond, M. C., Boom, D. R., Klous, P., Schooten, E., Marquez, V. E., Wierda, R. J., et al. (2011). Epigenetic regulation of CIITA expression in human T-cells. *Biochem. Pharmacol.* 82, 1430–1437. doi: 10.1016/j.bcp.2011.05.026
- Wang, B., Ding, Y. M., Fan, P., Wang, B., Xu, J. H., and Wang, W. X. (2014). Expression and significance of MMP2 and HIF-1 α in hepatocellular carcinoma. *Oncol. Lett.* 8, 539–546. doi: 10.3892/ol.2014.2189
- Wang, H., and Yin, S. (2015). Natural killer T cells in liver injury, inflammation and cancer. *Expert Rev. Gastroenterol. Hepatol.* 9, 1077–1085. doi: 10.1586/17474124.2015.1056738
- Wang, S., Li, J., Wu, S., Cheng, L., Shen, Y., Ma, W., et al. (2018). Type 3 innate lymphoid cell: a new player in liver fibrosis progression. *Clin. Sci. (Lond.)* 132, 2565–2582. doi: 10.1042/CS20180482
- Wang, Y., and Zhang, C. (2019). The roles of liver-resident lymphocytes in liver diseases. *Front. Immunol.* 10:1582. doi: 10.3389/fimmu.2019.01582
- Wasley, A., Grytdal, S., Gallagher, K., and Centers for Disease Control and Prevention (2008). Surveillance for acute viral hepatitis—United States, 2006. *MMWR Surveill. Summ.* 57, 1–24.
- Wehr, A., Baeck, C., Heymann, F., Niemietz, P. M., Hammerich, L., Martin, C., et al. (2013). Chemokine receptor CXCR6-dependent hepatic NK T Cell accumulation promotes inflammation and liver fibrosis. *J. Immunol.* 190, 5226–5236. doi: 10.4049/jimmunol.1202909
- Wiencke, J. K., Zheng, S., Jelluma, N., Tihan, T., Vandenberg, S., Tamguney, T., et al. (2007). Methylation of the PTEN promoter defines low-grade gliomas and secondary glioblastoma. *Neuro Oncol.* 9, 271–279. doi: 10.1215/15228517-2007-003
- Wong, L., Yamasaki, G., Johnson, R. J., and Friedman, S. L. (1994). Induction of beta-platelet-derived growth factor receptor in rat hepatic lipocytes during cellular activation in vivo and in culture. *J. Clin. Invest.* 94, 1563–1569. doi: 10.1172/JCI117497
- Wright, J. H., Johnson, M. M., Shimizu-Albergine, M., Bauer, R. L., Hayes, B. J., Surapitschat, J., et al. (2014). Paracrine activation of hepatic stellate cells in platelet-derived growth factor C transgenic mice: evidence for stromal induction of hepatocellular carcinoma. *Int. J. Cancer* 134, 778–788. doi: 10.1002/ijc.28421
- Wu, M., Miao, H., Fu, R., Zhang, J., and Zheng, W. (2020). Hepatic stellate cell: a potential target for hepatocellular carcinoma. *Curr. Mol. Pharmacol.* doi: 10.2174/1874467213666200224102820 [Epub ahead of print].
- Wu, N., Gu, C., Gu, H., Hu, H., Han, Y., and Li, Q. (2011). Metformin induces apoptosis of lung cancer cells through activating JNK/p38 MAPK pathway and GADD153. *Neoplasma* 58, 482–490. doi: 10.4149/neo_2011_06_482
- Wynn, T. A. (2008). Cellular and molecular mechanisms of fibrosis. *J. Pathol.* 214, 199–210. doi: 10.1002/path.2277
- Xu, Y., Sun, X., Zhang, R., Cao, T., Cai, S. Y., Boyer, J. L., et al. (2020). A positive feedback loop of TET3 and TGF- β 1 promotes liver fibrosis. *Cell Rep.* 30, 1310–1318.e5. doi: 10.1016/j.celrep.2019.12.092
- Xu, Y. F., Hannafon, B. N., Khatri, U., Gin, A., and Ding, W. Q. (2019). The origin of exosomal miR-1246 in human cancer cells. *RNA Biol.* 16, 770–784. doi: 10.1080/15476286.2019.1585738
- Yan, Y., Zhou, C., Li, J., Chen, K., Wang, G., Wei, G., et al. (2017). Resveratrol inhibits hepatocellular carcinoma progression driven by hepatic stellate cells by targeting Gli-1. *Mol. Cell. Biochem.* 434, 17–24. doi: 10.1007/s11010-017-3031-z
- Yin, C., Evason, K. J., Asahina, K., and Stainier, D. Y. (2013). Hepatic stellate cells in liver development, regeneration, and cancer. *J. Clin. Invest.* 123, 1902–1910. doi: 10.1172/JCI66369
- Yin, Z., Dong, C., Jiang, K., Xu, Z., Li, R., Guo, K., et al. (2019). Heterogeneity of cancer-associated fibroblasts and roles in the progression, prognosis, and therapy of hepatocellular carcinoma. *J. Hematol. Oncol.* 12:101. doi: 10.1186/s13045-019-0782-x
- Yu, M. C., Chen, C. H., Liang, X., Wang, L., Gandhi, C. R., Fung, J. J., et al. (2004). Inhibition of T-cell responses by hepatic stellate cells via B7-H1-mediated T-cell apoptosis in mice. *Hepatology* 40, 1312–1321. doi: 10.1002/hep.20488
- Zhang, B., Pan, X., Cobb, G. P., and Anderson, T. A. (2007). microRNAs as oncogenes and tumor suppressors. *Dev. Biol.* 302, 1–12. doi: 10.1016/j.ydbio.2006.08.028
- Zhang, D. Y., and Friedman, S. L. (2012). Fibrosis-dependent mechanisms of hepatocarcinogenesis. *Hepatology* 56, 769–775. doi: 10.1002/hep.25670
- Zhang, L., Valencia, C. A., Dong, B., Chen, M., Guan, P. J., and Pan, L. (2015). Transfer of microRNAs by extracellular membrane microvesicles: a nascent crossstalk model in tumor pathogenesis, especially tumor cell-microenvironment interactions. *J. Hematol. Oncol.* 8:14. doi: 10.1186/s13045-015-0111-y
- Zhao, W., Zhang, L., Xu, Y., Zhang, Z., Ren, G., Tang, K., et al. (2014). Hepatic stellate cells promote tumor progression by enhancement of immunosuppressive cells in an orthotopic liver tumor mouse model. *Lab. Invest.* 94, 182–191. doi: 10.1038/labinvest.2013.139

- Zhou, B. Y., Gong, J. H., Cai, X. Y., Wang, J. X., Luo, F., Jiang, N., et al. (2019). An imbalance between stellate cells and gammadeltaT cells contributes to hepatocellular carcinoma aggressiveness and recurrence. *Hepatol. Int.* 13, 631–640. doi: 10.1007/s12072-019-09969-w
- Zhou, Y., Ren, H., Dai, B., Li, J., Shang, L., Huang, J., et al. (2018). Hepatocellular carcinoma-derived exosomal miRNA-21 contributes to tumor progression by converting hepatocyte stellate cells to cancer-associated fibroblasts. *J. Exp. Clin. Cancer Res.* 37:324. doi: 10.1186/s13046-018-0965-2
- Zhu, B., Lin, N., Zhang, M., Zhu, Y., Cheng, H., Chen, S., et al. (2015). Activated hepatic stellate cells promote angiogenesis via interleukin-8 in hepatocellular carcinoma. *J. Transl. Med.* 13:365. doi: 10.1186/s12967-015-0730-7

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Barry, Baldeosingh, Lamm, Patel, Zhang, Dominguez, Kirton, Shah and Dang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.