



# Stimulation of Non-canonical NF- $\kappa$ B Through Lymphotoxin- $\beta$ -Receptor Impairs Myogenic Differentiation and Regeneration of Skeletal Muscle

Manuel Schmidt<sup>1</sup>, Anja Weidemann<sup>1</sup>, Christine Poser<sup>1</sup>, Anne Bigot<sup>2</sup> and Julia von Maltzahn<sup>1\*</sup>

<sup>1</sup> Leibniz Institute on Aging, Fritz Lipmann Institute, Jena, Germany, <sup>2</sup> Center of Research in Myology-UMRS 974, Institute of Myology, INSERM, Sorbonne Université, Paris, France

## OPEN ACCESS

### Edited by:

D. Cornelison,  
University of Missouri, United States

### Reviewed by:

Alessandra Cecchini,  
University of Missouri, United States  
Eusebio Perdiguerro,  
Pompeu Fabra University, Spain

### \*Correspondence:

Julia von Maltzahn  
julia.vonmaltzahn@leibniz-fli.de

### Specialty section:

This article was submitted to  
Stem Cell Research,  
a section of the journal  
Frontiers in Cell and Developmental  
Biology

**Received:** 07 June 2021

**Accepted:** 16 September 2021

**Published:** 05 October 2021

### Citation:

Schmidt M, Weidemann A,  
Poser C, Bigot A and von Maltzahn J  
(2021) Stimulation of Non-canonical  
NF- $\kappa$ B Through  
Lymphotoxin- $\beta$ -Receptor Impairs  
Myogenic Differentiation  
and Regeneration of Skeletal Muscle.  
*Front. Cell Dev. Biol.* 9:721543.  
doi: 10.3389/fcell.2021.721543

Myogenic differentiation, muscle stem cell functionality, and regeneration of skeletal muscle are cellular processes under tight control of various signaling pathways. Here, we investigated the role of non-canonical NF- $\kappa$ B signaling in myogenic differentiation, muscle stem cell functionality, and regeneration of skeletal muscle. We stimulated non-canonical NF- $\kappa$ B signaling with an agonistically acting antibody of the lymphotoxin beta receptor (LT $\beta$ R). Interestingly, we found that stimulation of non-canonical NF- $\kappa$ B signaling through the LT $\beta$ R agonist impairs myogenic differentiation, muscle stem cell function, and regeneration of skeletal muscle. Furthermore, we show that stimulation of non-canonical NF- $\kappa$ B signaling by the LT $\beta$ R agonist coincides with activation of canonical NF- $\kappa$ B signaling. We suggest a direct crosstalk between canonical and non-canonical NF- $\kappa$ B signaling during myogenic differentiation which is required for proper myogenic differentiation and thereby regeneration of skeletal muscle.

**Keywords:** NF-kappa B, NF- $\kappa$ B, LT $\beta$ R, lymphotoxin- $\beta$ -receptor, muscle stem cell, satellite cell, regeneration, myogenic differentiation

## INTRODUCTION

Skeletal muscle makes up approximately 40% of the total body weight of an adult, it contributes to several bodily functions such as voluntary movements and breathing. Furthermore, skeletal muscle can adapt to physiological demands such as growth, training, or injury (Frontera and Ochala, 2015). It also has a remarkable ability to regenerate, a process which depends on muscle stem cells (MuSCs), also termed satellite cells (Lepper et al., 2011; Murphy et al., 2011; Brack and Rando, 2012; Wang and Rudnicki, 2012; Wang et al., 2014). MuSCs are characterized by the expression of the transcription factor Pax7 which is essential for their functionality (Lepper et al., 2009; von Maltzahn et al., 2013). Under resting conditions, they are quiescent and located between the plasmalemma of the myofiber and the basement membrane (Wang et al., 2014; Frontera and Ochala, 2015; Ancel et al., 2021). Upon injury, MuSCs become activated, proliferate, differentiate, and replace damaged muscle tissue (Schmidt et al., 2019). One remarkable characteristic of MuSCs is their ability to self-renew thereby sustaining the stem cell pool over multiple cycles of growth and regeneration (Brack and Rando, 2012; Wang and Rudnicki, 2012; Wang et al., 2014). The functionality of MuSCs is a tightly controlled process, intrinsic as well as extrinsic cues affect MuSC proliferation,

differentiation, and self-renewal. The extrinsic signals can derive from the stem cell niche or from the blood stream (systemic factors) (Bentzinger et al., 2010; Chang and Rudnicki, 2014; Dinulovic et al., 2017). Intrinsic changes in MuSCs or in their direct niche can dramatically compromise MuSC functionality resulting in impaired regeneration of skeletal muscle as observed during aging (Price et al., 2014; Sousa-Victor et al., 2014; Brack and Munoz-Canoves, 2016; Schwörer et al., 2016).

Regeneration of skeletal muscle resembles embryonic myogenesis, differentiation of MuSCs and myogenic progenitor cells is accompanied by a transcription factor cascade comprising key myogenic regulatory factors including MyoD, myf5, myogenin, and myogenic regulatory factor 4 (MRF-4) (Chal and Pourquie, 2017; Hernandez-Hernandez et al., 2017). Activation of MuSCs coincides with the upregulation of MyoD, a transcription factor promoting myogenic differentiation (Hernandez-Hernandez et al., 2017; Ancel et al., 2021). Activated/committed MuSCs then progress in the myogenic program to become myogenic progenitor cells which are characterized by the expression of MyoD while expression of Pax7 is lost. Upon induction of myogenin expression, the cells further differentiate to become myocytes which then fuse to form multinucleated myotubes. In the final step of myogenesis, those myotubes mature into myofibers, which are the functional units of skeletal muscle (Forcina et al., 2019; Schmidt et al., 2019).

Nuclear factor- $\kappa$ B (NF- $\kappa$ B) signaling plays an important role during differentiation in multiple tissues including skeletal muscle (Ghosh and Karin, 2002; Hayden and Ghosh, 2004; Kucharczak et al., 2004). The NF- $\kappa$ B family comprises five members: p65/RelA, RelB, c-Rel, p100/p52, and p105/p50, which can form homo- or heterodimers. Depending on the NF- $\kappa$ B signaling molecules, one can divide NF- $\kappa$ B signaling into the canonical (classical) and non-canonical (alternative) pathway (Oeckinghaus and Ghosh, 2009; Zhang et al., 2017). Activation of the canonical pathway by TNF- $\alpha$  leads to the phosphorylation of inhibitor of kappa B (I $\kappa$ B), which causes the sequestration of p65/RelA and p50 into the cytoplasm. Hence, NF- $\kappa$ B transcription factors (like p65/RelA, p50, and c-Rel) shuttle into the nucleus and initiate expression of target genes which then drive proliferation, differentiation, or survival pathways (Shih et al., 2011). While the canonical NF- $\kappa$ B pathway is activated by TNF- $\alpha$ , the non-canonical pathway can be stimulated by binding of lymphotoxin  $\alpha$  and  $\beta$  to lymphotoxin- $\beta$ -receptor (LT $\beta$ R). After activation of the non-canonical NF- $\kappa$ B pathway, the precursor protein p100 is phosphorylated by a I $\kappa$ B kinase  $\alpha$  (IKK $\alpha$ ) homodimer, which mediates the partial proteolysis of p100 thereby generating the active form p52. p52 then forms a heterodimer with RelB and translocates into the nucleus (Oeckinghaus and Ghosh, 2009; Zhang et al., 2017). Of note, non-canonical NF- $\kappa$ B signaling was demonstrated to be independent of IKK $\beta$  (Mackay et al., 1996; Muller and Siebenlist, 2003; Shih et al., 2011).

In recent years the role of NF- $\kappa$ B signaling during myogenesis has been investigated focusing on the impact of the canonical pathway. Canonical NF- $\kappa$ B signaling is thought to control myogenesis by limiting myogenic differentiation. For instance, deletion of p65/RelA was shown to enhance

the expression of myogenic transcription factors thereby accelerating myogenic differentiation (Bakkar et al., 2008). While myogenic differentiation was thought to be mainly controlled by canonical NF- $\kappa$ B signaling, non-canonical NF- $\kappa$ B signaling was demonstrated to be important for myotube maintenance and metabolic regulation. For instance, MyoD in cooperation with RelB regulates the oxidative metabolism through binding to enhancer sites of the PGC-1 $\beta$  gene thereby regulating mitochondrial biogenesis (Bakkar et al., 2012; Shintaku et al., 2016).

Here, we investigated how stimulation of non-canonical NF- $\kappa$ B signaling affects myogenic differentiation, muscle stem cell function, and regeneration of skeletal muscle. We demonstrate that stimulation of the non-canonical NF- $\kappa$ B signaling pathway via the LT $\beta$ R impairs myogenic differentiation and limits muscle stem cell functionality. This then leads to impaired regeneration of skeletal muscle. Interestingly, we observe impaired regeneration and myogenic differentiation following stimulation of non-canonical NF- $\kappa$ B signaling in wt and in mdx mice. Furthermore, we noticed a compensatory upregulation of members of the canonical NF- $\kappa$ B pathway following stimulation of the LT $\beta$ R. This suggests that a tight regulation and crosstalk of canonical and non-canonical NF- $\kappa$ B signaling is essential for proper myogenic differentiation and muscle stem cell functionality.

## MATERIALS AND METHODS

### Mice

Adult (12 weeks of age) male C57/BL6Jrj mice obtained from Janvier Laboratories or male mdx mice (12 weeks of age) from Charles River Laboratories were used in this study. All mice were maintained inside a SPF facility. All animal procedures were in accordance with the European Union (EU) directive 2010/63/EU and approved by the Animal Welfare Committee of the Thüringer Landesamt für Lebensmittelsicherheit und Verbraucherschutz (TLV; 03-011/14; Bad Langensalza, Germany).

For regeneration experiments, TA muscles of WT or mdx mice were injured by intramuscular injection of 50  $\mu$ l CTX (10  $\mu$ M) followed by injection of solvent (0.9% NaCl), LT $\beta$ R agonist (10  $\mu$ g for WT and 30  $\mu$ g for mdx mice) or LT $\beta$ R antagonist (30  $\mu$ g) at 3 days post injury (dpi) under isoflurane anesthesia. Muscles were isolated 10 or 14 dpi, embedded in cryoprotective medium (Tissue-Tek O.C.T. with 10% sucrose) and frozen in liquid nitrogen.

### Cell Culture, Transfection and Modulation of NF- $\kappa$ B Signaling

Muscle stem cells were isolated via FACS from hind-limb muscles of 12-week-old male C57/BL6Jrj with a positive selection for  $\alpha$ -7-Integrin as described before (Pasut et al., 2012), cultured and plated on collagen-coated dishes (Corning) in culture medium containing Ham's F10 medium (Thermo Fisher Scientific), 20% fetal bovine serum (FBS) (Thermo Fisher Scientific), 2% Penicillin/Streptomycin (Thermo Fisher Scientific), and 5 ng/ml

basic fibroblast growth factor (bFGF) (Thermo Fisher Scientific). Differentiation of primary murine myoblasts was initiated by a switch of culture media to differentiation medium which contains DMEM (Thermo Fisher Scientific), 10% HS (horse serum, Thermo Fisher Scientific), and 2% Penicillin/Streptomycin (Thermo Fisher Scientific). Human myoblasts from DMD patients and healthy controls [AB1190 Clone 1 – healthy, AB1023DMD11Q Clone 1 and AB1071DMD13PV Clone 5 – DMD, were cultured in skeletal muscle cell growth media (PromoCell, C-23060) with 15% FBS (Thermo Fisher Scientific), 1% GlutaMAX (Thermo Fisher Scientific), 1% Gentamicine (Thermo Fisher Scientific), and 1% Penicillin/Streptomycin (Thermo Fisher Scientific)]. Differentiation of human myoblasts was initiated with skeletal muscle differentiation medium (PromoCell, C-23061) and 1% Penicillin/Streptomycin (Thermo Fisher Scientific). Murine or human myoblasts and MuSCs on floating myofibers were transfected with either non-targeting siRNA (Dharmacon) or si-Rela (Dharmacon) with Lipofectamine RNAiMAX (Thermo Fisher Scientific) according to the protocol provided by the manufacturer. Stimulation or inhibition of non-canonical NF- $\kappa$ B signaling in murine cells was performed by addition of antibodies either acting agonistically (LT $\beta$ R agonist; Aldevron; concentration: 1  $\mu$ g/ $\mu$ l) or acting antagonistically (LT $\beta$ R antagonist; concentration: 0.6  $\mu$ g/ $\mu$ l, obtained from T. Hehlhans, Regensburg, Germany) to the differentiation medium (Weidemann et al., 2016). As a control a mouse IgG antibody (Sigma-Aldrich, 12-370) or equal amount of the solvent (PBS) were added to the medium. Stimulation of non-canonical NF- $\kappa$ B signaling in human myoblasts was performed by addition of recombinant mouse anti-human LT $\beta$ R antibody (CBE11) (1  $\mu$ g/ $\mu$ l, Creative Biolabs) known to act agonistically. Canonical NF $\kappa$ B signaling was inhibited by addition of either TPCA-1 or IMD0454 (Abcam; 1  $\mu$ M final concentration) dissolved in DMSO to the medium, as a control equal amounts of DMSO were used.

## Myofiber Isolation and Culture

Preparation and cultivation of *extensor digitorum longus* (EDL) fibers was performed as described previously (Schmidt et al., 2020). In brief, EDL muscles were dissected under sterile conditions and incubated in DMEM with 0.2% collagenase (from *Clostridium histolyticum*, Sigma-Aldrich) at 37°C in a water bath for 1 h. Dissociation of single myofibers was accomplished by trituration with sterile glass pipettes and single myofibers were cultured in myofiber culture medium (DMEM, 20% FBS, 1% chicken embryo extract; United States Biological) at 37°C, 5% CO<sub>2</sub> and 95% humidity for 72 h. The LT $\beta$ R agonist (concentration: 1  $\mu$ g/ $\mu$ l) or LT $\beta$ R antagonist (concentration: 0.6  $\mu$ g/ $\mu$ l) were added to the culture medium directly after isolation. Myofibers with their adjacent muscle stem cells were fixed with 2% paraformaldehyde (PFA) for 5 min, washed twice with PBS (pH 7.4) and permeabilized with PBS, 0.1% Triton X-100, 0.1 M glycine (pH 7.4) for 10 min. Myofibers were then incubated with blocking solution [5% HS in PBS (pH 7.4)] for 1 h followed by incubation with the primary antibodies directed against Pax7 (undiluted, PAX7; DSHB) and MyoD (1:100, Invitrogen, #PA5-23078) at 4°C overnight. After three washing steps with PBS (pH 7.4) the myofibers were incubated with the

secondary antibodies for 45 min at room temperature (RT). Afterwards, myofibers were washed three times with PBS (pH 7.4), incubated with DAPI (50  $\mu$ g/ml) for 5 min, washed twice with PBS (pH 7.4) and then mounted in PermaFluor mounting medium (Thermo Fisher Scientific) on glass microscope slides. Analysis was carried out using an Axio Observer D1 (Carl Zeiss).

## Histochemistry

Muscles frozen in liquid nitrogen were cut in 14  $\mu$ m cross-sections. Hematoxylin-Eosin (H&E) staining was performed as described earlier (von Maltzahn et al., 2013) with the Leica stainer XL. Images were taken with the Axio Observer D1 (Carl Zeiss).

## Immunostaining and Antibodies

Muscles frozen in liquid nitrogen were cut in 14  $\mu$ m cross-sections. Cross-sections were fixed with 2% PFA for 10 min, washed twice with PBS (pH 7.4), permeabilized with 0.1% Triton X-100, 0.1 M glycine in PBS (pH 7.4) for 10 min, incubated first with blocking solution [5% HS in PBS (pH 7.4)] and then with primary antibodies in blocking solution at 4°C overnight. Afterwards, samples were washed three times with PBS (pH 7.4) and incubated with secondary antibodies in blocking solution for 45 min at RT. After washing with PBS (pH 7.4) three times, nuclei were counterstained with DAPI (50  $\mu$ g/ml) and washed twice with PBS (pH 7.4). Cross-sections were mounted in PermaFluor (Thermo Fisher Scientific). The following antibodies were used: mouse anti-myosin (undiluted, MF20; DSHB), mouse anti-myogenin (undiluted, F5D, DSHB), mouse anti-Pax7 (undiluted, PAX7, DSHB), rabbit anti-MyoD (1:250, Thermo Fisher Scientific, #PA5-23078), chicken anti-Laminin (1:500, LC Bio, LS-C96142), rabbit anti-Laminin (1:1000, MERCK, L9393), and mouse anti-embryonic MHC (undiluted, F1.652, DSHB). Analyses were carried out using the Axio Observer D1 or the Axiovert 200 Apotome (Carl Zeiss). The number of Pax7<sup>+</sup> MuSCs was enumerated by counting Pax7<sup>+</sup> cells per muscle cross-section divided by the total area of the cross-section.

## Immunoblot Analyses

Mouse and human myoblasts were lysed in RIPA buffer for 20 min on ice followed by sonication (10 cycles, 30 s on and 5 s off). Between 10 and 20  $\mu$ g of protein were separated on a 8% Bis-Tris protein gel. Then, a transfer to a nitrocellulose membrane (VWR) was carried out followed by incubation of the membrane in blocking solution (3% BSA in TBST) for 1 h at RT and then with primary antibodies in blocking solution at 4°C overnight. Afterward, membranes were washed three times with TBST, incubated with secondary antibodies in blocking solution for 45 min at RT and washed three times with TBST. Membranes were incubated with Pierce<sup>TM</sup> ECL Western Blotting Substrate (Thermo Fisher Scientific) and imaged using the Bio-Rad MyImager. Cytoplasmatic and nuclear fractions were generated as described previously (Weidemann et al., 2016). The following primary antibodies were used: rabbit anti-RelA (1:200, Santa Cruz, sc-372), rabbit anti-phospho-RelA (1:1000, Cell Signaling, #3033), rabbit anti- NF- $\kappa$ B2 p100/p52 (1:1000, Cell Signaling, #4882), rabbit anti-RelB (1:1000, Cell Signaling, #4882), rabbit anti-phospho RelB (Ser552) (1:1000,

Cell Signaling, #4999S), rabbit anti-IKB $\alpha$  (1:1000, Cell Signaling, #2859), rabbit anti-phospho-IKK $\alpha$ / $\beta$  (1:1000, Cell Signaling, #2697), mouse anti-myosin (undiluted, MF20, DSHB), mouse anti-myogenin (undiluted, F5D, DSHB), rabbit anti-MyoD (1:100, Thermo Fisher Scientific, #PA5-23078), rabbit anti-LT $\beta$ R (1:500, Abcam, ab70063), mouse anti-GAPDH (1:200, Santa Cruz, sc-365062), mouse anti-PARP-1 (C2-10) (1:4000, Santa Cruz, sc-53643). The following secondary antibodies were used: goat anti-rabbit Immunoglobulins/HRP (1:1000, Dako, P0448), goat anti-mouse Immunoglobulins/HRP (1:1500, Dako, P0447).

## Real-Time PCR

Total RNA was isolated with TriFast (PeqLab) reagent followed by additional purification using the Qiagen RNA isolation kit. Reverse transcription was performed with the iScript cDNA synthesis kit (Bio-Rad) according to the protocol provided by the manufacturer. SYBR Green real-time PCR (IQ SYBR Supermix, Bio-Rad) was performed on a real-time thermocycler (Bio-Rad). Primers used are listed in **Table 1**.

## Statistical Analysis

A minimum of three and a maximum of five replicates were analyzed for each experiment presented. Data are shown as SEM. Statistical significance was assessed by Student's *t* test (unpaired, two-tailed) or one-way or two-way ANOVA, using Microsoft Excel or Prism after testing for normal distribution using the Shapiro–Wilk test. In cases where the Shapiro–Wilk test failed, a Kruskal–Wallis test was used (indicated in the respective figure legend). A *p* value < 0.05 was considered significant.

## RESULTS


### Activation of Non-canonical NF- $\kappa$ B Signaling by Stimulating the LT $\beta$ R Impairs Myogenic Differentiation

So far, mainly the role of canonical NF- $\kappa$ B signaling during myogenic differentiation and MuSC function has been investigated in detail (Bakkar et al., 2008; Bakkar and Guttridge, 2010; He et al., 2013). The role of non-canonical NF- $\kappa$ B signaling during myogenic differentiation, especially its activation via the LT $\beta$ R remained elusive. To determine

the activation status of non-canonical NF- $\kappa$ B signaling during myogenic differentiation, we first investigated expression and phosphorylation (corresponding to the activation status) of members of the canonical and non-canonical NF- $\kappa$ B signaling pathway during myogenic differentiation in primary murine myoblasts. Interestingly, we found that members of both, the canonical and the non-canonical signaling pathway are activated during early myogenic differentiation. For instance, we detected an increase in the phosphorylated/activated forms of RelA and RelB during myogenic differentiation suggesting that also non-canonical NF- $\kappa$ B signaling plays an important role during myogenic differentiation (**Supplementary Figures 1A,B**). To investigate the role of non-canonical NF- $\kappa$ B signaling in myogenic differentiation, we differentiated primary murine myoblasts for 5 days in the presence of an antibody directed against the LT $\beta$ R (LT $\beta$ R agonist) known to stimulate non-canonical NF- $\kappa$ B signaling through binding to the LT $\beta$ R (**Figure 1A** and **Supplementary Figures 1C,D**; Lukashchuk et al., 2006; Weidemann et al., 2016). Stimulation of non-canonical NF- $\kappa$ B signaling at induction of differentiation caused a significant decrease in myotube size compared to myotubes which were differentiated in the presence of the solvent or an IgG control antibody (**Figures 1B,C**). Next, we asked whether stimulation of non-canonical NF- $\kappa$ B signaling exclusively affects early myogenic differentiation or if myotube maintenance is also affected. Therefore, we differentiated myoblasts for 4 days to obtain multinucleated myotubes and then added the LT $\beta$ R agonist followed by an additional day of culture in differentiation medium (**Figure 1D**). Interestingly, also this stimulation for 24 h of already formed myotubes resulted in a reduction of myotube diameter suggesting that also myotube maintenance is affected or that activation of non-canonical NF- $\kappa$ B signaling can cause myotube atrophy. However, the effect size was reduced compared to activation of non-canonical NF- $\kappa$ B signaling at the induction of myogenic differentiation (**Figures 1E,F**). We further confirmed that addition of the LT $\beta$ R agonist activates non-canonical NF- $\kappa$ B signaling in myogenic cells. Stimulation with the LT $\beta$ R agonist results in the cleavage of the precursor p100 into the active form p52 in myotubes which were differentiated in the presence of the LT $\beta$ R agonist (**Supplementary Figures 1C,D**). Surprisingly, we also detected an increased phosphorylation of RelA, a member of the canonical NF- $\kappa$ B pathway in both conditions (**Supplementary Figures 1C,D** and **Figure 1O**).

**TABLE 1** | Primers used for quantitative RT-PCR.

Name	Forward sequence	Reverse sequence
MyoD	CTA CAG TGG CGA CTC AGA T	CAC TGT AGT AGG CGG TGT C
Myogenin	CAG TAC ATT GAG CGC CTA C	AAG GCA ACA GAC ATA TCC TC
Mef2a	GAT GCT GAC GAT TAC TTT GA	TCG AAT CTG CTA ATG TTG AG
PGC1 $\alpha$	GAG TCT GAA AGG GCC AAA CA	ACG GTG CAT TCC TCA ATT TC
LT $\beta$ R	TTA TCG CAT AGA AAA CCA GAC TTG C	TCA AAG CCC AGC ACA ATG TC
NF- $\kappa$ B2 p100	GCT AAT GTG AAT GCC CGG AC	CTT TGG GTA TCC CTC TCA GGC
Relb	GTG ACC TCT CTT CCC TGT CAC TAA C	CAG CGT GAC GCT GCT CAG T
Rela	CCT CTC ACA TCC GAT TTT TGA T	GTC CCG TGA AAT ACA CCT CAA T
GAPDH	ATG CCA GTG AGC TTC CCG TC	CAT CAC CAT CTT CCA GGA GC



**FIGURE 1** | Activation of the LT $\beta$ R pathway impairs myogenic differentiation. **(A)** Experimental scheme of myoblast differentiation in the presence of the LT $\beta$ R agonist. **(B)** Representative images of myotubes incubated with the control or the LT $\beta$ R agonist with antibodies directed to myosin heavy chain (MHC, green). Nuclei are counterstained with DAPI (blue). Scale bar = 50  $\mu$ m. **(C)** 5 days culture of myoblasts in the presence of the LT $\beta$ R agonist markedly reduces myotube diameter compared to control treatments.  $n = 3$ . **(D)** Experimental scheme of myoblast differentiation with incubation of myotubes at day 4 of differentiation for 24 h with the LT $\beta$ R agonist or a control. **(E)** Representative images of myotubes incubated with the control or the LT $\beta$ R agonist with antibodies directed to myosin heavy chain (MHC, green). Nuclei are counterstained with DAPI (blue). Scale bar = 50  $\mu$ m. **(F)** Incubation at day 4 of differentiation of 24 h with the LT $\beta$ R agonist results in reduced myotube diameter.  $n = 3$ . **(G)** qRT-PCR for *pgc1alpha*. qRT-PCR of myotubes after incubation with the LT $\beta$ R demonstrates significant upregulation of *myoD* **(H)** and *myogenin* **(I)**, but not *mef2a* **(J)**. qRT-PCR analysis of LT $\beta$ R pathway target genes shows no markedly influence on LT $\beta$ R expression **(K)**, but a significantly increase of p100 **(L)**, *relb* **(M)** and classical NF- $\kappa$ B target gene *rela* mRNA **(N)**, especially after incubation at day 4 for 24 h with the LT $\beta$ R agonist.  $n = 3$ . **(O)** Activation of the canonical NF- $\kappa$ B pathway by the LT $\beta$ R agonist shown by increased p-RelA and p-IKK. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ . Error bars represent SEM. In **(A)** a Kruskal–Wallis test was used.

Hence, we conclude that stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist impairs myogenic differentiation and myotube maintenance while also resulting in the activation of the canonical NF- $\kappa$ B pathway.

Non-canonical NF- $\kappa$ B was reported to be involved in myotube maintenance and metabolic regulation involving *pgc1 $\alpha$*  (Bakkar et al., 2012; Shintaku et al., 2016). Therefore, we investigated the expression of the mitochondrial biogenesis marker peroxisome proliferator-activated receptor gamma coactivator 1-alpha (*pgc1 $\alpha$* ) after stimulating the non-canonical NF- $\kappa$ B pathway with the LT $\beta$ R agonist. We detected a reduced expression of *pgc1 $\alpha$*  in myotubes stimulated with the LT $\beta$ R agonist (Figure 1G). Furthermore, we determined the expression of *myod* and *myogenin*, two markers of early myogenic differentiation and *mef2a*, known to control myogenic fusion. We found increased expression levels of the three markers after stimulation of the non-canonical NF- $\kappa$ B pathway with the LT $\beta$ R agonist (Figures 1H–J). This further strengthens our finding that activation of non-canonical NF- $\kappa$ B signaling impairs myogenic differentiation since terminally differentiated myotubes do not express *myod* and only very few amounts of *myogenin* (Schmidt et al., 2019). *Myod* and *myogenin* are typical markers for myoblasts and myocytes, cell types which are expressed during early myogenic differentiation.

Next, we asked whether stimulation with the LT $\beta$ R agonist affects mRNA expression of members of the non-canonical and canonical NF- $\kappa$ B signaling pathways. We found that stimulation with the LT $\beta$ R agonist does not affect mRNA levels of *lt $\beta$ r* (Figure 1K). However, we detected a significant increase in mRNA levels of *relb* and *p100/nf $\kappa$ b1* – as expected (Figures 1L,M). We then investigated, if also the mRNA levels of members of the canonical NF- $\kappa$ B signaling pathway are affected. Indeed, we found mRNA levels of *rela* to be increased upon stimulation with the LT $\beta$ R agonist (Figure 1N). The increased mRNA levels of *rela* coincide with increased phosphorylation of RelA and increased phosphorylation of the kinase IKK $\alpha$ , which inactivates the p65/RelA inhibitor IKB $\alpha$  (Figure 1O). Therefore, we conclude, that stimulating LT $\beta$ R with the LT $\beta$ R agonist results in impaired myogenesis via activation of the non-canonical NF- $\kappa$ B pathway concomitant with – a potentially compensatory – activation of canonical NF- $\kappa$ B pathway.

### Inhibiting Canonical NF- $\kappa$ B Signaling Counteracts Impaired Myogenic Differentiation Following Stimulation of Non-canonical NF- $\kappa$ B Signaling


Next, we asked whether active canonical NF- $\kappa$ B signaling is a prerequisite for impaired myogenic differentiation after stimulating the non-canonical NF- $\kappa$ B pathway with the LT $\beta$ R agonist. Therefore, we differentiated primary murine myoblasts for 3 days in the presence of the LT $\beta$ R agonist and a siRNA targeting *rela* to silence canonical NF- $\kappa$ B signaling (Figure 2A). As an additional control we used a LT $\beta$ R antagonistic antibody to inhibit non-canonical NF- $\kappa$ B signaling. Stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist resulted in impaired myogenic differentiation as evidenced by a reduced

myotube diameter (Figures 2B,C). This suggests that stimulation of non-canonical NF- $\kappa$ B signaling affects early myogenic differentiation (Figures 2B,C) as well as growth/maintenance of myotubes (Supplementary Figures 1A–F). As expected, we did not observe any effect on myogenic differentiation after inhibiting non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R antagonist (Figures 2B,C). However, knockdown of *rela* resulted in enhanced myogenic differentiation independent of stimulation or inhibition of non-canonical NF- $\kappa$ B signaling (Figures 2B,C). To demonstrate that indeed early myogenic differentiation is impaired through activation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist, we analyzed the myotube diameter after 2 days of differentiation. Thereby we found the myotube size (Supplementary Figures 1E,F) to be significantly reduced when differentiation was induced in the presence of the LT $\beta$ R agonist. Also inhibition of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R antagonist in already formed myotubes did not affect their size. However, inhibition of non-canonical NF- $\kappa$ B signaling with TPCA-1, an inhibitor of IKK $\beta$  or through silencing RelA resulted in larger myotubes independent of activation or inhibition of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist or antagonist (Supplementary Figures 1E,F).

Next, we evaluated the activation status of members of the canonical and the non-canonical NF- $\kappa$ B pathway after knockdown of *rela* or stimulation of the LT $\beta$ R in 3 days differentiated myotubes (Figures 2D,E). Interestingly, knockdown of *rela per se* did not alter the abundance or phosphorylation status of members of the non-canonical NF- $\kappa$ B signaling pathway (p100, p52, RelB, p-RelB). Stimulation of non-canonical NF- $\kappa$ B signaling by the LT $\beta$ R agonist resulted in increased p52 levels by cleavage of p100. However – in line with the results obtained earlier – we observed an increased phosphorylation of RelA after stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist. We conclude that stimulation of the LT $\beta$ R impairs myogenic differentiation through activating non-canonical NF- $\kappa$ B signaling concomitant with activation of the canonical pathway.


### Stimulation of Non-canonical NF- $\kappa$ B Signaling Impairs Muscle Regeneration and Muscle Stem Cell Functionality in wt Mice

Regeneration of skeletal muscle depends on MuSCs and their full functionality including the ability to undergo myogenic differentiation and self-renewal. To determine, if stimulation or inhibition of non-canonical NF- $\kappa$ B signaling affects regeneration of skeletal muscle, we performed cardiotoxin (CTX) mediated injury of the tibialis anterior muscle of adult C57BL/6 mice. This was followed by a single injection of the solvent (control), the LT $\beta$ R agonist or the LT $\beta$ R antagonist into the injured muscle at day 3 after injury (Figure 3A). We found newly formed myofibers to be reduced in size [measured as minimal fiber feret, identified by staining for embryonic MHC (eMHC)] when muscles were injected with the LT $\beta$ R agonist (Figures 3B,C and Supplementary Figure 2). Furthermore, all myofibers undergoing regeneration in muscles injected



with the LT $\beta$ R agonist displayed a smaller myofiber diameter compared to myofibers of muscles which were injected with the solvent (control) or the LT $\beta$ R antagonist (Figures 3B,D,E). However, inhibition of non-canonical NF- $\kappa$ B signaling *per se* only moderately enhanced regeneration of skeletal muscle

(Figures 3B–E and Supplementary Figure 2). In summary, these data suggest that stimulation of non-canonical NF- $\kappa$ B signaling via the LT $\beta$ R agonist results in impaired regeneration of skeletal muscle in wt mice while inhibition of non-canonical NF- $\kappa$ B signaling does not affect the regeneration process.



**FIGURE 3** | Activation of the non-canonical NF- $\kappa$ B signaling by the LT $\beta$ R agonist impairs regeneration in wt mice. **(A)** Experimental scheme depicting CTX injury and injection of the LT $\beta$ R agonist, LT $\beta$ R antagonist or solvent control at day 3 after injury. **(B)** Representative images of HE staining and immunostaining with antibodies directed to laminin (green) and embryonic MHC (red). Nuclei are counterstained with DAPI (blue). Scale bar = 200  $\mu$ m. **(C)** Injection of the LT $\beta$ R agonist results in decreased myofiber size of eMHC-positive myofibers compared to control and antagonist injected muscles.  $n = 5$ . **(E)** Size distribution of myofibers undergoing regeneration as in **(D)**  $n = 5$ . **(F)** Number of Pax7-positive cells in regenerating muscles following injection of LT $\beta$ R agonist or antagonist.  $n = 5$ . **(G)** Injection of the LT $\beta$ R agonist reduces the number of committed MuSCs (Pax7<sup>+</sup>/MyoD<sup>+</sup>),  $n = 5$ . **(H)** Injection of the LT $\beta$ R agonist reduces the number of myoblasts (Pax7<sup>-</sup>/MyoD<sup>+</sup>).  $n = 5$ . \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ . Error bars represent SEM.

Interestingly though, numbers of self-renewing MuSCs (Pax7<sup>+</sup>/MyoD<sup>-</sup>) were not altered in both conditions while the number of activated MuSCs (Pax7<sup>+</sup>/MyoD<sup>+</sup>) and

further differentiated cells (Pax7<sup>-</sup>/MyoD<sup>+</sup>) was reduced after stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist (**Figures 3F–H**). This further supports our notion




that stimulation of non-canonical NF- $\kappa$ B signaling impairs myogenic differentiation and interferes with MuSC functionality.

Next, we asked whether stimulation of non-canonical NF- $\kappa$ B signaling is directly affecting MuSCs or if the effect on regeneration of skeletal muscle is mediated by other cells. Therefore, we performed experiments using the floating myofiber culture system. Here, MuSCs are cultured on their adjacent myofibers, which allows their direct manipulation while being cultured in their endogenous niche (Hüttner et al., 2019). After 72 h of culture single MuSCs form clusters of cells on the myofiber which can then be examined for their differentiation status, e.g., by immunostaining for Pax7 and MyoD (a marker for early differentiation). Stimulation of non-canonical NF- $\kappa$ B signaling by the LT $\beta$ R agonist led to an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells, which are non-differentiated, self-renewing cells. Interestingly, inhibition of the NF- $\kappa$ B signaling by the LT $\beta$ R antagonist also did not affect the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells (Figure 4A). However, stimulation of NF- $\kappa$ B signaling by the LT $\beta$ R agonist did not cause a significant change in the number of clusters per myofiber (Figure 4B) while the number of cells per cluster and the amount of differentiating MuSCs (Pax7<sup>-</sup>/MyoD<sup>+</sup>) were reduced (Figures 4C,D). Of note, also here inhibition of NF- $\kappa$ B signaling via the LT $\beta$ R antagonist

had no effect. To conclude that indeed differentiation of MuSCs is impaired by activation of NF- $\kappa$ B signaling with the LT $\beta$ R agonist, we investigated different time points (Supplementary Figure 3). Thereby we show that the percentage of non-differentiated MuSCs (Pax7<sup>+</sup>/MyoD<sup>-</sup>) was already increased after 24 h and 48 h of culture while the total number of MuSCs per myofiber was not affected (Supplementary Figures 3A–D). Also at these early time points, inhibition of NF- $\kappa$ B signaling by the LT $\beta$ R antagonist did not affect MuSC behavior. A closer look at the clusters formed after 72 h of culture in the presence of the LT $\beta$ R agonist confirmed our observation that activation of NF- $\kappa$ B signaling impairs myogenic differentiation, e.g., demonstrated by the reduced percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> MuSCs being further differentiated myogenic cells (Supplementary Figures 3E,F).

To get a better understanding of the interplay between canonical and non-canonical NF- $\kappa$ B signaling in MuSCs, we cultured MuSCs on floating myofibers in the presence of the LT $\beta$ R agonist and inhibitors of canonical NF- $\kappa$ B signaling (Figures 4E–H and Supplementary Figure 4). Interestingly, we observed very similar effects on myogenic differentiation of MuSCs when they were cultured in the presence of the LT $\beta$ R agonist alone or if canonical NF- $\kappa$ B signaling was inhibited in addition. Of note, we observed similar effects when using



**FIGURE 4 |** Activation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist impairs differentiation of MuSCs. **(A)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist results in an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells per cluster while addition of the LT $\beta$ R antagonist has no effect. **(B)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist or the antagonist does not affect cluster number. **(C)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist leads to a reduction in the number of cells per cluster while culture in the presence of the antagonist increases the number of cells per cluster. **(D)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist results in a decrease in the percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells per cluster while addition of the LT $\beta$ R antagonist has no effect. **(E)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist and/or the IKK inhibitor results in an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells per cluster. **(F)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist and/or the IKK inhibitor results in a reduction in the percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells per cluster. **(G)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist results in an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells per cluster independent of knockdown of *relA*. **(H)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist results in a reduction in the percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells per cluster independent of knockdown of *RelA*. *n* = 4 C57BL/6, 3 months of age. \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001. Error bars represent SEM.

TPCA-1, IMD0354 and siRNA targeting RelA in addition to the LT $\beta$ R agonist. However, the effects on MuSC differentiation we observed with TPCA-1 might be caused by inhibition of JAK/STAT signaling since this inhibitor was described to also affect signaling via STAT3, a pathway known to be important for MuSC functionality (Nan et al., 2014; Price et al., 2014; Tierney et al., 2014). Also IMD0354 was shown to affect STAT signaling, namely signaling via STAT1 (Tanaka et al., 2005). In all cases we observed an impaired myogenic differentiation suggesting that inhibition of canonical NF- $\kappa$ B signaling cannot compensate for activation of non-canonical NF- $\kappa$ B signaling in MuSCs.


### Inhibition of Non-canonical NF- $\kappa$ B Signaling Improves Regeneration in mdx Mice

Next, we asked if stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist impairs MuSC function and thereby regeneration of skeletal muscle in muscles which are continuously regenerating. Therefore, we investigated how stimulation or inhibition of non-canonical NF- $\kappa$ B signaling affects MuSCs from mdx mice, a mouse model for Duchenne muscular dystrophy, often used as a model for continuous muscle regeneration (Dangain and Vrbova, 1984; Anderson et al., 1987; **Figure 5A**). However, it is known that MuSC divisions in mdx mice are reduced after 72 h of culture compared to wt mice (Dumont et al., 2015). When we stimulated non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist in MuSCs from mdx mice, we observed a slight reduction in the cluster number per myofiber (**Figures 5B,E**). This is different from our observation in wt mice, probably due to the different activation status of MuSCs from mdx mice compared to wt mice and their impaired ability to undergo asymmetric divisions (Dumont et al., 2015). Interestingly, inhibition of non-canonical NF- $\kappa$ B signaling did not affect cluster number or composition of clusters in MuSCs from mdx mice (**Figures 5B–E**). However, the number of Pax7<sup>+</sup>/MyoD<sup>+</sup> early differentiating cells was reduced after stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist (**Figures 5C,E**) similar to the effect we observed in wt mice. This reduction in committed cells coincided with a reduction in the number of Pax7<sup>-</sup>/MyoD<sup>+</sup> cells, which represent further differentiated cells (**Figures 5D,E**). Also here, the results are in line with the results obtained in wt mice (**Figures 4A–D**). Of note, neither the number of cells per cluster, the number of self-renewing cells (Pax7<sup>+</sup>/MyoD<sup>-</sup> cells) nor the number of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells were significantly changed following manipulation of non-canonical NF- $\kappa$ B signaling in MuSCs from mdx mice (**Figure 5**). In summary, these data suggest that stimulation of non-canonical NF- $\kappa$ B signaling results in impaired myogenic commitment and early differentiation after acute activation in wt cells (**Figure 4**) as well as in already activated MuSCs from mice which undergo continuous regeneration (**Figure 5**). Furthermore, we show that stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist can directly affect MuSCs.


Next, we asked whether interfering with the activity of non-canonical NF- $\kappa$ B signaling in muscles undergoing continuous regeneration affects MuSC function and regeneration after acute injury. Therefore, we performed an acute muscle injury of mdx mice followed by intramuscular injection of the LT $\beta$ R agonist or LT $\beta$ R antagonist (**Figure 6A**). We found that stimulation of non-canonical NF- $\kappa$ B signaling only slightly impaired regeneration in mdx mice similar to the effect observed in wt mice (**Figures 6B–D**). However, inhibition of non-canonical NF- $\kappa$ B signaling resulted in a slight improvement in the regeneration of mdx muscles after acute injury (**Figures 6B–D**). These results were surprising to us since inhibition of non-canonical NF- $\kappa$ B signaling in wt mice did not affect regeneration (**Figure 3**). However, dystrophic muscles are characterized by an increased/constant inflammatory response (Manning and O'Malley, 2015). Here, I $\kappa$ B kinase/NF- $\kappa$ B (IKK/NF- $\kappa$ B) signaling was shown to be persistently elevated in immune cells and regenerating myofibers (Acharyya et al., 2007). One might therefore speculate that the improved regeneration in mdx mice after inhibition of non-canonical NF- $\kappa$ B signaling is not due to a direct effect on MuSCs but rather an indirect one mediated by the immune cells. Next, we quantified MuSC numbers in mdx mice after stimulation or inhibition of non-canonical NF- $\kappa$ B signaling. Here, stimulation of non-canonical NF- $\kappa$ B signaling did not affect MuSC numbers while its inhibition resulted in a slight increase, reminiscent of the effect observed in wt mice (**Figures 6E,F**). In summary, these results suggest that stimulation of non-canonical NF- $\kappa$ B signaling via the LT $\beta$ R agonist impairs myogenic differentiation and regeneration of muscles after acute injury or undergoing continuous regeneration. However, inhibition of non-canonical NF- $\kappa$ B signaling improves regeneration of muscles of mdx mice. We hypothesize that inhibition of non-canonical NF- $\kappa$ B signaling primarily affects non-myogenic cells in mdx muscles and that improvements of regeneration are rather secondary effects in this specific mouse model.

### Stimulation of Non-canonical NF- $\kappa$ B Signaling Impairs Myogenic Differentiation Independent of Dystrophin Status in Human Myoblasts

We next wondered whether loss of dystrophin expression affects the response to stimulation or inhibition of non-canonical NF- $\kappa$ B signaling also in myogenic cells from humans. Therefore, we investigated myogenic differentiation of human myoblasts from DMD patients and healthy control in response to stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist (**Figure 7A**). Stimulation of non-canonical NF- $\kappa$ B signaling resulted in impaired myogenic differentiation, reminiscent of the phenotype observed in murine myoblasts (**Figures 7B,C**). Interestingly, we found a very similar response in DMD patient derived myoblasts to stimulation of non-canonical NF- $\kappa$ B signaling (**Figures 7B,C**). However, we observed enhanced myogenic differentiation after inhibition of canonical NF- $\kappa$ B signaling with a siRNA targeting *RELA*. Of note, knockdown of *RELA* enhanced myogenic differentiation independent of




**FIGURE 5 |** Manipulating the activation of the non-canonical NF- $\kappa$ B signaling pathway affects myogenic differentiation of MuSCs from mdx mice. **(A)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist causes an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> independent of knockdown of *relA*. **(B)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist leads to a reduction in the number of clusters per myofiber independent of knockdown of *relA*. Incubation with the LT $\beta$ R antagonist does not affect the number of clusters per myofiber. **(C)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist causes a reduction in the number of cells per cluster while incubation with the LT $\beta$ R antagonist does not affect the number of cells per cluster. **(D)** Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R agonist causes a reduction in the percentage of Pax7<sup>-</sup>/MyoD<sup>+</sup> independent of knockdown of *relA*. **(E)** Representative images of clusters of MuSCs after culture for 72 h in the presence of the LT $\beta$ R agonist, LT $\beta$ R antagonist, or a solvent control. MyoD in green, Pax7 in red. Nuclei are counterstained with DAPI (blue). Scale bar = 50  $\mu$ m.  $n = 4$ , 3 months of age, \* $p < 0.05$ . Error bars represent SEM.



stimulation of non-canonical NF- $\kappa$ B signaling in myoblasts from healthy control and DMD patients (**Figures 7B,C** and **Supplementary Figure 5**). In accordance with the data we obtained in murine myotubes we found a slight activation of

the canonical NF- $\kappa$ B pathway following stimulation with the LT $\beta$ R agonist independent of the dystrophin status (**Figures 7D,E** and **Supplementary Figure 5**). We conclude that stimulation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist inhibits



myogenic differentiation in murine and human myoblasts independent of the dystrophin status.

In summary, we demonstrate that stimulation of non-canonical NF- $\kappa$ B signaling impairs myogenic differentiation. We show that the LT $\beta$ R agonist directly affects MuSCs and their progeny causing impaired regeneration of skeletal muscle.

## DISCUSSION

Myogenesis and regeneration of skeletal muscle are tightly controlled processes requiring the precise activation and

inhibition of signaling pathways to allow proper differentiation and self-renewal (Schmidt et al., 2019; Henze et al., 2020). Canonical and non-canonical NF- $\kappa$ B signaling are important signaling pathways whose tight regulation is required for cellular processes such as differentiation, cell adhesion, migration or inflammation, and survival. Of note, non-canonical NF- $\kappa$ B signaling was shown to be important for the regulation of differentiation processes (Shih et al., 2011; Zhang et al., 2017). However, a crosstalk between canonical and non-canonical NF- $\kappa$ B signaling is thought to occur during differentiation, e.g., during osteogenic differentiation. While Caron et al. (2012) showed that RelA/p65 mediated signaling facilitates

chondrogenic differentiation, a study by Franzoso et al. (1997) demonstrated that the simultaneous knockout of the RelA and RelB binding partners p50 and p52 resulted in impaired osteogenic differentiation causing osteopetrosis in mice. However, deletion of p100 – the precursor of p52 – results in endochondral ossification in mice, a phenotype which can be partially rescued by additional deletion of RelB. This further supports the notion that a regulated crosstalk between canonical and non-canonical NF- $\kappa$ B signaling is important for proper differentiation (Jimi et al., 2019). Regarding the role of NF- $\kappa$ B signaling in myogenesis and regeneration of skeletal muscle, research mainly focused on members of the canonical NF- $\kappa$ B signaling axis. For instance, the importance of canonical NF- $\kappa$ B signaling for the control of myogenic differentiation was demonstrated using p65/RelA deficient myoblasts (Bakkar et al., 2008; Bakkar and Guttridge, 2010).

Here, we focused on the role of non-canonical NF- $\kappa$ B signaling in myogenic differentiation, *in vitro* and *in vivo*. Thereby, we demonstrate that stimulation of non-canonical NF- $\kappa$ B signaling through an agonistically acting antibody directed against LT $\beta$ R (LT $\beta$ R agonist) impairs myogenic differentiation of myoblasts and MuSCs. We suggest that this is one of the reasons for the impaired regeneration of skeletal muscle following activation of the non-canonical NF- $\kappa$ B signaling pathway (**Supplementary Figure 6**). Through experiments using the floating myofiber culture system, we demonstrate that activation of the non-canonical NF- $\kappa$ B signaling pathway can also directly affect myogenic differentiation. We show that non-canonical NF- $\kappa$ B signaling regulates myogenic differentiation and that overly active non-canonical NF- $\kappa$ B signaling is detrimental for progression of myogenic differentiation and thereby regeneration of skeletal muscle. While stimulation of non-canonical NF- $\kappa$ B signaling impairs myogenic differentiation in wt mice, inhibition of non-canonical NF- $\kappa$ B signaling with an antagonistically working antibody directed against the LT $\beta$ R (LT $\beta$ R antagonist) does not affect myogenic differentiation or MuSC function, neither *in vivo* nor *in vitro*. However, inhibition of non-canonical NF- $\kappa$ B signaling in mdx mice during acute regeneration improves the regenerative outcome (**Figure 6**). We suggest that the enhanced regeneration in mdx mice is caused by secondary effects of the LT $\beta$ R antagonist, e.g., by blocking the elevated inflammatory response of invading immune cells observed in dystrophic muscles and not by direct effects on MuSCs (Manning and O'Malley, 2015). This hypothesis is supported by the finding that differentiation of MuSCs on floating myofibers from mdx mice is not improved by manipulating the activity of non-canonical NF- $\kappa$ B signaling. Furthermore, differentiation of myoblasts derived from healthy controls or DMD patients is similarly impaired after stimulation of non-canonical NF- $\kappa$ B signaling suggesting that overly active non-canonical NF- $\kappa$ B signaling inhibits myogenic differentiation in a similar manner in mice and men. Furthermore, these data support our notion that stimulation of non-canonical NF- $\kappa$ B signaling directly targets myogenic cells while inhibition of non-canonical NF- $\kappa$ B signaling rather affects non-myogenic cells.

Stimulation of non-canonical NF- $\kappa$ B signaling in myoblasts might lead to impaired activation of the myogenic transcription

factor cascade required for proper myogenic differentiation (Hernandez-Hernandez et al., 2017). Increased levels of RelB in the nucleus might interfere with binding of MyoD or myogenin to specific promoter regions thereby impairing myogenic differentiation. However, changes in mRNA stabilization of *myod* or *myogenin* could also cause impaired differentiation after stimulation with the LT $\beta$ R agonist, similar to RelA interfering with *myoD* stability (Guttridge et al., 2000). In addition to changes in *myoD* or *myogenin* stability, modulation or interaction with TWEAK might contribute to the impaired regeneration/MuSC function following stimulation of non-canonical NF- $\kappa$ B signaling. TWEAK (TNF $\alpha$ -like weak inducer of apoptosis), a member of the TNF family, was shown to regulate myoblast fusion through non-canonical NF- $\kappa$ B signaling in skeletal muscle (Enwere et al., 2012, 2014). Deregulation of Pax7 expression or function might also contribute to the impaired differentiation of MuSCs following stimulation of non-canonical NF- $\kappa$ B signaling. He et al. (2013) demonstrated that cancer cachexia leads to stimulation of canonical NF- $\kappa$ B signaling in MuSCs which causes increased Pax7 levels. Pax7 was shown to be required for myogenic differentiation and MuSC functionality, e.g., Pax7 deficient mice display severe malformations of skeletal muscle and die a couple of weeks after birth (Seale et al., 2000). In contrast, Straughn et al. (2018) demonstrated that inhibition of specific components of canonical NF- $\kappa$ B signaling causes precocious differentiation of MuSCs. However, we observed impaired myogenic differentiation of MuSCs and myoblasts after stimulation of non-canonical NF- $\kappa$ B signaling suggesting that non-canonical NF- $\kappa$ B signaling limits myogenic differentiation, potentially through increasing Pax7 levels or *myoD* mRNA stability.

Of note, we observed a slight upregulation of members of the canonical NF- $\kappa$ B signaling pathway when we stimulated non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist (**Supplementary Figures 1, 5**). This suggests an immediate crosstalk of canonical and non-canonical NF- $\kappa$ B signaling during myogenic differentiation. These findings are in line with data which revealed crosstalk of canonical and non-canonical NF- $\kappa$ B signaling and activation of first RelA/p50 and then RelB/p52 after induction of LT $\beta$ R in mouse embryonic fibroblasts (Muller and Siebenlist, 2003; Basak et al., 2007). The idea of a direct crosstalk between the NF- $\kappa$ B signaling pathways is further supported by another report, in which a site-activation of the canonical NF- $\kappa$ B pathway during adipocyte differentiation was observed when non-canonical NF- $\kappa$ B signaling was stimulated by the agonistically acting antibody directed against LT $\beta$ R (Weidemann et al., 2016). Activation of non-canonical NF- $\kappa$ B signaling is generally thought to be related to developmental cues (Shih et al., 2011). However, developmental LT $\beta$ R signaling causes the disruption of the I $\kappa$ B $\beta$  inhibitory complex; this then allows the translocation of RelA/p50 into the nucleus (Basak et al., 2007). This mechanism shown in fibroblasts might also occur in myogenic differentiation since we observed an increase in nuclear RelA protein upon stimulation of the LT $\beta$ R. This might mediate the crosstalk between canonical and non-canonical NF- $\kappa$ B signaling during myogenesis.

The importance of a fine regulation of canonical and non-canonical NF- $\kappa$ B signaling for proper myogenesis is further underscored by studies in dystrophic mice. Here, canonical NF- $\kappa$ B signaling was shown to be overly active (Acharyya et al., 2007). The phenotype of mdx mice was improved, when the aberrant activation of canonical NF- $\kappa$ B signaling in dystrophic conditions was controlled by orally available inhibitors of p65/RelA (Hammers et al., 2016). Interestingly, the authors also observed a slight decrease in activation of non-canonical NF- $\kappa$ B signaling in mdx muscles. We suggest that crosstalk between canonical and non-canonical NF- $\kappa$ B is occurring during myogenic differentiation and that is important for keeping an adequate balance of myogenic progression and self-renewal.

In summary, our study demonstrates that stimulation of non-canonical NF- $\kappa$ B signaling with an agonistically acting LT $\beta$ R antibody impairs myogenic differentiation, MuSC function and thereby regeneration of skeletal muscle. Furthermore, we provide evidence that a fine balance in the activation of canonical and non-canonical signaling is required for proper myogenic differentiation and MuSC function.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors upon request, without undue reservation.

## ETHICS STATEMENT

The animal study was reviewed and approved by the Thüringer Landesamt für Lebensmittelsicherheit und Verbraucherschutz.

## AUTHOR CONTRIBUTIONS

MS, CP, and JM designed and performed the most experiments, analyzed the data, and interpreted the results. AB generated patient derived myoblasts. MS, AW, and JM analyzed the data, interpreted the results, and wrote the manuscript. All authors contributed to the article and approved the submitted version.

## FUNDING

This work was supported by grants from the Deutsche Forschungsgemeinschaft (MA-3975/2-1) and the Deutsche Krebshilfe (DKH-JvM-861005) to JM. The FLI is a member of the Leibniz Association and is financially supported by the Federal Government of Germany and the State of Thuringia.

## ACKNOWLEDGMENTS

The authors thank Thomas Hehlhans for providing the LT $\beta$ R blocking antibody and the Myoline platform from the Institute

of Myology in Paris for providing the human myoblasts. The authors gratefully acknowledge support from the FLI Core Facilities Imaging, Mouse as well as the FLI Core Services Histology.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcell.2021.721543/full#supplementary-material>

**Supplementary Figure 1** | Canonical and non-canonical NF- $\kappa$ B pathway during myogenesis. Separation of cytoplasmatic (A) and nuclear fraction (B) from primary murine myoblasts undergoing differentiation shows activation of canonical (p-RelA) and non-canonical (p52) signaling during myogenic differentiation. (C,D) Culture in the presence of the LT $\beta$ R agonist for the respective times causes cleavage of p100 to active p52 and increased phosphorylation of RelA. (E) Differentiation of primary myoblasts in the presence of the LT $\beta$ R agonist results in a reduced myotube diameter while addition of the LT $\beta$ R antagonist does not affect myotube size. Inhibition of the IKKB with TPCA-1 increases the myotube diameter. (F) Differentiation of primary myoblasts in the presence of the LT $\beta$ R agonist results in a reduced myotube diameter while addition of the LT $\beta$ R antagonist does not affect myotube size. Knockdown of RelA results in an increase in myotube diameter. Scale bar = 50  $\mu$ m.  $n = 4$ , 3 months of age, \* $p < 0.05$ ; \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . Error bars represent SEM.

**Supplementary Figure 2** | CTX mediated injury of TA muscles from wt mice followed by injection of the LT $\beta$ R agonist or LT $\beta$ R antagonist. (A) TA muscles from mice injected with the LT $\beta$ R antagonist are slightly heavier than muscles injected with the LT $\beta$ R agonist. (B) Myofiber size distribution of eMHC-positive myofibers,  $n = 5$ . Error bars represent SEM.

**Supplementary Figure 3** | Activation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist impairs early myogenic differentiation of MuSCs. (A) Culture of isolated myofibers for 24 h in the presence of the LT $\beta$ R agonist results in an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells per cluster while addition of the LT $\beta$ R antagonist has no effect. (B) Culture of isolated myofibers for 24 h in the presence of the LT $\beta$ R agonist or antagonist does not affect the number of cells per myofiber. (C) Culture of isolated myofibers for 48 h in the presence of the LT $\beta$ R agonist results in an increase in the percentage of Pax7<sup>+</sup>/MyoD<sup>-</sup> cells per cluster while addition of the LT $\beta$ R antagonist has no effect. (D) Culture of isolated myofibers for 48 h in the presence of the LT $\beta$ R agonist or antagonist does not affect the number of cells per myofiber. (E) Culture of isolated myofibers for 72 h in the presence of the LT $\beta$ R results in an increase in the number of single cells per myofiber while the percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells per cluster is reduced. (F) The LT $\beta$ R antagonist does not affect the number of single cells per myofiber nor the percentage of Pax7<sup>+</sup>/MyoD<sup>+</sup> cells per cluster.  $n = 4$ , 3 months of age, \* $p < 0.05$ , \*\* $p < 0.01$ . Error bars represent SEM.

**Supplementary Figure 4** | MuSC differentiation is impaired after activation of non-canonical NF- $\kappa$ B signaling with the LT $\beta$ R agonist independent of inhibition of the canonical NF- $\kappa$ B pathway. (A) Culture of MuSCs on their adjacent myofibers for 72 h in the presence of the LT $\beta$ R agonist and an inhibitor of IKKB results in a reduction in the number of cells per cluster. (B) The percentage of Pax7<sup>-</sup>/MyoD<sup>+</sup> cells cell per cluster after 72 h in culture under the respective conditions.  $n = 4$ , 3 months of age, \* $p < 0.05$ ; \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . Error bars represent SEM.

**Supplementary Figure 5** | Activation of canonical and non-canonical NF- $\kappa$ B signaling in differentiating human myoblasts. Investigation of activation of canonical (RelA, p-RelA) and non-canonical (p100, p52, and LT $\beta$ R) NF- $\kappa$ B total protein levels by immunoblot analyses. Incubation with the LT $\beta$ R agonist results in increased phosphorylation of RelA and also cleavage of p100 to p52. Knockdown of RELA leads to the activation of the non-canonical pathway after addition of the LT $\beta$ R agonist, while the phosphorylation status of RelA is not affected.

**Supplementary Figure 6** | Overly active non-canonical NF- $\kappa$ B signaling impairs myogenic differentiation, muscle stem cell function, and regeneration of skeletal muscle.

## REFERENCES

- Acharyya, S., Villalta, S. A., Bakkar, N., Bupha-Intr, T., Janssen, P. M., Carathers, M., et al. (2007). Interplay of IKK/NF-kappaB signaling in macrophages and myofibers promotes muscle degeneration in Duchenne muscular dystrophy. *J. Clin. Invest.* 117, 889–901. doi: 10.1172/jci30556
- Ancel, S., Stuelsatz, P., and Feige, J. N. (2021). Muscle stem cell quiescence: controlling stemness by staying asleep. *Trends Cell Biol.* 31, 556–568. doi: 10.1016/j.tcb.2021.02.006
- Anderson, J. E., Ovalle, W. K., and Bressler, B. H. (1987). Electron microscopic and autoradiographic characterization of hindlimb muscle regeneration in the mdx mouse. *Anat. Rec.* 219, 243–257. doi: 10.1002/ar.1092190305
- Bakkar, N., and Guttridge, D. C. (2010). NF-kappaB signaling: a tale of two pathways in skeletal myogenesis. *Physiol. Rev.* 90, 495–511. doi: 10.1152/physrev.00040.2009
- Bakkar, N., Ladner, K., Canan, B. D., Liyanarachchi, S., Bal, N. C., Pant, M., et al. (2012). IKK $\alpha$  and alternative NF-kappaB regulate PGC-1 $\beta$  to promote oxidative muscle metabolism. *J. Cell Biol.* 196, 497–511. doi: 10.1083/jcb.201108118
- Bakkar, N., Wang, J., Ladner, K. J., Wang, H., Dahlman, J. M., Carathers, M., et al. (2008). IKK/NF-kappaB regulates skeletal myogenesis via a signaling switch to inhibit differentiation and promote mitochondrial biogenesis. *J. Cell Biol.* 180, 787–802. doi: 10.1083/jcb.200707179
- Basak, S., Kim, H., Kearns, J. D., Tergaonkar, V., O'Dea, E., Werner, S. L., et al. (2007). A fourth IkkappaB protein within the NF-kappaB signaling module. *Cell* 128, 369–381.
- Bentzinger, C. F., von Maltzahn, J., and Rudnicki, M. A. (2010). Extrinsic regulation of satellite cell specification. *Stem Cell Res Ther.* 1:27. doi: 10.1186/scrt27
- Brack, A. S., and Munoz-Canoves, P. (2016). The ins and outs of muscle stem cell aging. *Skelet Muscle* 6:1. doi: 10.1016/j.csb.2016.06.001
- Brack, A. S., and Rando, T. A. (2012). Tissue-specific stem cells: lessons from the skeletal muscle satellite cell. *Cell Stem Cell* 10, 504–514. doi: 10.1016/j.stem.2012.04.001
- Caron, M. M., Emans, P. J., Surtel, D. A., Cremers, A., Voncken, J. W., Welting, T. J., et al. (2012). Activation of NF-kappaB/p65 facilitates early chondrogenic differentiation during endochondral ossification. *PLoS One* 7:e33467. doi: 10.1371/journal.pone.0033467
- Chal, J., and Pourquie, O. (2017). Making muscle: skeletal myogenesis in vivo and in vitro. *Development* 144, 2104–2122. doi: 10.1242/dev.151035
- Chang, N. C., and Rudnicki, M. A. (2014). Satellite cells: the architects of skeletal muscle. *Curr. Top. Dev. Biol.* 107, 161–181.
- Dangain, J., and Vrbova, G. (1984). Muscle development in mdx mutant mice. *Muscle Nerve* 7, 700–704. doi: 10.1002/mus.880070903
- Dinulovic, I., Furrer, R., and Handschin, C. (2017). Plasticity of the muscle stem cell microenvironment. *Adv. Exp. Med. Biol.* 1041, 141–169. doi: 10.1007/978-3-319-69194-7\_8
- Dumont, N. A., Wang, Y. X., von Maltzahn, J., Pasut, A., Bentzinger, C. F., Brun, C. E., et al. (2015). Dystrophin expression in muscle stem cells regulates their polarity and asymmetric division. *Nat. Med.* 21, 1455–1463. doi: 10.1038/nm.3990
- Enwere, E. K., Holbrook, J., Lejmi-Mrad, R., Vineham, J., Timusk, K., Sivaraj, B., et al. (2012). TWEAK and cIAP1 regulate myoblast fusion through the noncanonical NF-kappaB signaling pathway. *Sci. Signal.* 5:ra75. doi: 10.1126/scisignal.2003086
- Enwere, E. K., Lacasse, E. C., Adam, N. J., and Korneluk, R. G. (2014). Role of the TWEAK-Fn14-cIAP1-NF-kappaB signaling axis in the regulation of myogenesis and muscle homeostasis. *Front. Immunol.* 5:34. doi: 10.3389/fimmu.2014.00034
- Forcina, L., Miano, C., Pelosi, L., and Musaro, A. (2019). An overview about the biology of skeletal muscle satellite cells. *Curr. Genomics* 20, 24–37. doi: 10.2174/1389202920666190116094736
- Franzoso, G., Carlson, L., Xing, L., Poljak, L., Shores, E. W., Brown, K. D., et al. (1997). Requirement for NF-kB in osteoclast and B-cell development. *Genes Dev.* 11, 3482–3496. doi: 10.1101/gad.11.24.3482
- Frontera, W. R., and Ochala, J. (2015). Skeletal muscle: a brief review of structure and function. *Calcif. Tissue Int.* 96, 183–195. doi: 10.1007/s00223-014-9915-y
- Ghosh, S., and Karin, M. (2002). Missing pieces in the NF-kB puzzle. *Cell* 109, S81–S96.
- Guttridge, D. C., Mayo, M. W., Madrid, L. V., Wang, C.-Y., et al. (2000). NF-kB-Induced loss of MyoD messenger RNA: possible role in muscle decay and cachexia. *Science* 289, 2363–2366. doi: 10.1126/science.289.5488.2363
- Hammers, D. W., Sleeper, M. M., Forbes, S. C., Coker, C. C., Jirousek, M. R., Zimmer, M., et al. (2016). Disease-modifying effects of orally bioavailable NF-kappaB inhibitors in dystrophin-deficient muscle. *JCI Insight* 1:e90341.
- Hayden, M. S., and Ghosh, S. (2004). Signaling to NF-kB. *Genes Dev.* 18, 2195–2224.
- He, W. A., Berardi, E., Cardillo, V. M., Acharyya, S., Aulino, P., Thomas-Ahner, J., et al. (2013). NF-kappaB-mediated Pax7 dysregulation in the muscle microenvironment promotes cancer cachexia. *J. Clin. Invest.* 123, 4821–4835. doi: 10.1172/jci68523
- Henze, H., Jung, M. J., Ahrens, H. E., Steiner, S., and von Maltzahn, J. (2020). Skeletal muscle aging - Stem cells in the spotlight. *Mech. Ageing Dev.* 189:111283. doi: 10.1016/j.mad.2020.111283
- Hernandez-Hernandez, J. M., Garcia-Gonzalez, E. G., Brun, C. E., and Rudnicki, M. A. (2017). The myogenic regulatory factors, determinants of muscle development, cell identity and regeneration. *Semin. Cell Dev. Biol.* 72, 10–18. doi: 10.1016/j.semcdb.2017.11.010
- Hüttner, S. S., Ahrens, H. E., Schmidt, M., Henze, H., Jung, M. J., Schüler, S. C., et al. (2019). “Isolation and culture of individual myofibers and their adjacent muscle stem cells from aged and adult skeletal muscle,” in *Stem Cells and Aging: Methods and Protocols*, ed K. Turksen (New York, NY: Springer New York), 25–36. doi: 10.1007/978-1-4939-9209-209
- Jimi, E., Huang, F., and Nakatomi, C. (2019). NF-kB signaling regulates physiological and pathological chondrogenesis. *Int. J. Mol. Sci.* 20:6275. doi: 10.3390/ijms20246275
- Kucharczak, J., Simmons, M. J., Fan, Y., and Gélinas, C. (2004). To be, or not to be: NF-kB is the answer – role of Rel/NF-kB in the regulation of apoptosis. *Oncogene* 23:8858. doi: 10.1038/sj.onc.1207744
- Lepper, C., Conway, S. J., and Fan, C. M. (2009). Adult satellite cells and embryonic muscle progenitors have distinct genetic requirements. *Nature* 460, 627–631. doi: 10.1038/nature08209
- Lepper, C., Partridge, T. A., and Fan, C.-M. (2011). An absolute requirement for Pax7-positive satellite cells in acute injury-induced skeletal muscle regeneration. *Development* 138:3639.
- Lukashev, M., LePage, D., Wilson, C., Bailly, V., Garber, E., Lukashin, A., et al. (2006). Targeting the lymphotoxin-beta receptor with agonist antibodies as a potential cancer therapy. *Cancer Res.* 66, 9617–9624.
- Mackay, F., Majeau, G. R., Hochman, P. S., and Browning, J. L. (1996). Lymphotoxin  $\beta$  receptor triggering induces activation of the nuclear factor  $\kappa$ B transcription factor in some cell types. *J. Biol. Chem.* 271, 24934–24938. doi: 10.1074/jbc.271.40.24934
- Manning, J., and O'Malley, D. (2015). What has the mdx mouse model of duchenne muscular dystrophy contributed to our understanding of this disease? *J. Muscle Res. Cell Motil.* 36, 155–167. doi: 10.1007/s10974-015-9406-4
- Muller, J. R., and Siebenlist, U. (2003). Lymphotoxin beta receptor induces sequential activation of distinct NF-kappa B factors via separate signaling pathways. *J. Biol. Chem.* 278, 12006–12012. doi: 10.1074/jbc.m210768200
- Murphy, M. M., Lawson, J. A., Mathew, S. J., Hutcheson, D. A., and Kardon, G. (2011). Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration. *Development* 138, 3625–3637. doi: 10.1242/dev.064162
- Nan, J., Du, Y., Chen, X., Bai, Q., Wang, Y., Zhang, X., et al. (2014). TPCA-1 is a direct dual inhibitor of STAT3 and NF-kappaB and regresses mutant EGFR-associated human non-small cell lung cancers. *Mol. Cancer Ther.* 13, 617–629. doi: 10.1158/1535-7163.mct-13-0464
- Oeckinghaus, A., and Ghosh, S. (2009). The NF-kappaB family of transcription factors and its regulation. *Cold Spring Harb. Perspect. Biol.* 1:a000034.
- Pasut, A., Olynyk, P., and Rudnicki, M. A. (2012). Isolation of muscle stem cells by fluorescence activated cell sorting cytometry. *Methods Mol. Biol.* 798, 53–64. doi: 10.1007/978-1-61779-343-1\_3
- Price, F. D., von Maltzahn, J., Bentzinger, C. F., Dumont, N. A., Yin, H., Chang, N. C., et al. (2014). Inhibition of JAK-STAT signaling stimulates adult satellite cell function. *Nat. Med.* 20, 1174–1181. doi: 10.1038/nm.3655
- Schmidt, M., Poser, C., and von Maltzahn, J. (2020). Wnt7a counteracts cancer cachexia. *Mol. Therapy - Oncolyt.* 16, 134–146. doi: 10.1016/j.omto.2019.12.011
- Schmidt, M., Schuler, S. C., Hüttner, S. S., von Eyss, B., and von Maltzahn, J. (2019). Adult stem cells at work: regenerating skeletal muscle. *Cell Mol. Life Sci.* 76, 2559–2570. doi: 10.1007/s00018-019-03093-6



- Schwörer, S., Becker, F., Feller, C., Baig, A. H., Köber, U., Henze, H., et al. (2016). Epigenetic stress responses induce muscle stem-cell ageing by Hoxa9 developmental signals. *Nature* 540, 428–432. doi: 10.1038/nature20603
- Seale, P., Sabourin, L. A., Girgis-Gabardo, A., Mansouri, A., Gruss, P., and Rudnicki, M. A. (2000). Pax7 is required for the specification of myogenic satellite cells. *Cell* 102, 777–786. doi: 10.1016/s0092-8674(00)00066-0
- Shih, V. F., Tsui, R., Caldwell, A., and Hoffmann, A. (2011). A single NF-kappaB system for both canonical and non-canonical signaling. *Cell Res.* 21, 86–102. doi: 10.1038/cr.2010.161
- Shintaku, J., Peterson, J. M., Talbert, E. E., Gu, J. M., Ladner, K. J., Williams, D. R., et al. (2016). MyoD regulates skeletal muscle oxidative metabolism cooperatively with alternative NF-kappaB. *Cell Rep.* 17, 514–526. doi: 10.1016/j.celrep.2016.09.010
- Sousa-Victor, P., Gutarra, S., Garcia-Prat, L., Rodriguez-Ubreva, J., Ortet, L., Ruiz-Bonilla, V., et al. (2014). Geriatric muscle stem cells switch reversible quiescence into senescence. *Nature* 506, 316–321. doi: 10.1038/nature13013
- Straughn, A. R., Hindi, S. M., Xiong, G., and Kumar, A. (2018). Canonical NF- $\kappa$ B signaling regulates satellite stem cell homeostasis and function during regenerative myogenesis. *J. Mol. Cell Biol.* 11, 53–66. doi: 10.1093/jmcb/mjy053
- Tanaka, A., Konno, M., Muto, S., Kambe, N., Morii, E., Nakahata, T., et al. (2005). A novel NF-kappaB inhibitor, IMD-0354, suppresses neoplastic proliferation of human mast cells with constitutively activated c-kit receptors. *Blood* 105, 2324–2331. doi: 10.1182/blood-2004-08-3247
- Tierney, M. T., Aydogdu, T., Sala, D., Malecova, B., Gatto, S., Puri, P. L., et al. (2014). STAT3 signaling controls satellite cell expansion and skeletal muscle repair. *Nat. Med.* 20, 1182–1186. doi: 10.1038/nm.3656
- von Maltzahn, J., Jones, A. E., Parks, R. J., and Rudnicki, M. A. (2013). Pax7 is critical for the normal function of satellite cells in adult skeletal muscle. *Proc. Natl. Acad. Sci. U S A.* 110, 16474–16479. doi: 10.1073/pnas.1307680110
- Wang, Y. X., Dumont, N. A., and Rudnicki, M. A. (2014). Muscle stem cells at a glance. *J. Cell Sci.* 127, 4543–4548.
- Wang, Y. X., and Rudnicki, M. A. (2012). Satellite cells, the engines of muscle repair. *Nat. Rev. Mol. Cell Biol.* 13:127. doi: 10.1038/nrm3265
- Weidemann, A., Lovas, A., Rauch, A., Andreas, N., von Maltzahn, J., Riemann, M., et al. (2016). Classical and alternative NF-kappaB signaling cooperate in regulating adipocyte differentiation and function. *Int. J. Obes. (Lond)* 40, 452–459. doi: 10.1038/ijo.2015.198
- Zhang, Q., Lenardo, M. J., and Baltimore, D. (2017). 30 years of NF-kappaB: a blossoming of relevance to human pathobiology. *Cell* 168, 37–57. doi: 10.1016/j.cell.2016.12.012

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Schmidt, Weidemann, Poser, Bigot and von Maltzahn. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.