



# *In Vitro* Differentiation of Preosteoblast-Like Cells, MC3T3-E1, to Adipocytes Is Enhanced by 1,25(OH)<sub>2</sub> Vitamin D<sub>3</sub>

Elisha Pendleton and Nalini Chandar\*

Department of Biochemistry, Chicago College of Osteopathic Medicine, Midwestern University, Downers Grove, IL, United States

## OPEN ACCESS

### Edited by:

Paula H. Stern,  
Northwestern University,  
United States

### Reviewed by:

Melissa Orlandin Premaor,  
Universidade Federal de Santa  
Maria, Brazil  
David M. Findlay,  
University of Adelaide, Australia

### \*Correspondence:

Nalini Chandar  
nchand@midwestern.edu

### Specialty section:

This article was submitted  
to Bone Research,  
a section of the journal  
Frontiers in Endocrinology

**Received:** 20 December 2016

**Accepted:** 26 May 2017

**Published:** 16 June 2017

### Citation:

Pendleton E and Chandar N (2017)  
*In Vitro* Differentiation of  
Preosteoblast-Like Cells, MC3T3-E1,  
to Adipocytes Is Enhanced by  
1,25(OH)<sub>2</sub> Vitamin D<sub>3</sub>.  
*Front. Endocrinol.* 8:128.  
doi: 10.3389/fendo.2017.00128

Osteoblasts and adipocytes originate from common mesenchymal progenitor cells and are controlled by specific transcription factors. While 1,25-dihydroxyvitamin D<sub>3</sub> (vitamin D) is known to be an important factor for osteoblast differentiation, there are conflicting reports regarding its effect on adipogenesis. In this study, we attempted to understand the effect of exposure of preosteoblasts (MC3T3-E1) to adipogenic media with and without vitamin D and determined the expression of adipogenic genes during this process. Our studies show that while transdifferentiation of preosteoblasts occurred on exposure to adipogenic media, the effect of vitamin D treatment was synergistic resulting in several hundred fold increase in adipocyte transcription factors C/EBP $\alpha$  and peroxisome proliferator-activated receptor-gamma ( $P < 0.001$ ) along with an increase in markers of adipogenesis and accumulation of lipid droplets in cells. Vitamin D treatment was also accompanied by 100-fold to 700-fold increases in vitamin D receptor expression during the treatment period ( $P < 0.001$ ). To determine how the effect of vitamin D might compare to other genetic manipulations that promote adipogenic differentiation, we stably knocked down retinoblastoma expression in MC3T3-E1 cells. Recent studies have suggested retinoblastoma (Rb1) tumor suppressor gene function to be critical to maintain osteoblasts function and inhibit adipocyte differentiation. We exposed MC3T3-E1 cells with reduced Rb1 expression to adipogenic media and found an increase in adipogenic differentiation when compared to cells with a full complement of Rb dosage. However, the extent of the change was not as dramatic as seen with vitamin D. These studies show that preosteoblasts are sensitive and respond to these manipulations that favor the adipocytic phenotype. While vitamin D is not known to directly affect targets in adipogenesis, our observations may have resulted from the malleability of preosteoblast genome in MC3T3-E1 cells, which allowed adipocyte specific gene expression under appropriate stimuli. Why this pathway is influenced and subverted by an anabolic bone factor such as vitamin D remains to be determined.

**Keywords:** vitamin D, osteoblasts, MC3T3, Rb1, adipogenic differentiation, transdifferentiation

## INTRODUCTION

Osteoblasts and adipocytes are derived from mesenchymal stem cells, and many studies have suggested the presence of plasticity between these two cell types (1, 2). Aging and estrogen deficiency are also known to produce an increase in the bone marrow adiposity as well as decrease in the ability of mesenchymal precursor to progress toward osteogenesis (2, 3). Several studies have also suggested the

occurrence of both osteoblastic and adipocytic phenotype in the same cell population (4). In fact, a preadipocyte cell line has been derived from osteoblastic cell population from mouse calvaria (5).

One of the factors that regulate adipogenesis is peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ), which belongs to the class of nuclear hormone receptors that are affected by specific ligands such as fatty acids as well as by cross talk with other molecules. PPAR $\gamma$  is both sufficient and necessary for conversion of preadipocytes into adipocytes and is a master regulator of adipogenesis (6).

Vitamin D is an important bone anabolic agent, and several bone specific genes are directly regulated by this hormone. 1,25-Dihydroxyvitamin D<sub>3</sub>, the active form of vitamin D functions through its receptor, vitamin D receptor (VDR) a ligand dependent transcription factor, belonging to the same class of nuclear receptor family of transcription factors as PPARs. The similarity between these two factors is that they mediate their action by heterodimerizing with retinoic acid receptor in the regulation of gene expression. Studies on vitamin D effects on osteoblasts are mixed. While some studies show vitamin D to inhibit adipocyte differentiation, other studies have suggested adipocytic conversion of bone marrow osteoblasts by vitamin D (6–8). While vitamin D has been shown to inhibit adipocytic differentiation of preadipocytes, there are studies that demonstrate a stimulatory effect on other mesenchymal derived cells (7, 8).

There is also cross talk between PPAR $\gamma$  and VDR as VDR elements have been discovered in the PPAR genes demonstrating an influence of vitamin D over PPAR system (9). PPAR $\gamma$  also appears to bind to VDR and inhibit vitamin D dependent transactivation (10).

Retinoblastoma tumor suppressor gene (Rb1) function is important for controlling the cell cycle progression and differentiation of several cell types (11). Loss of Rb1 gene function predisposes individuals to retinoblastoma and osteosarcomas which are commonly seen as secondary tumors in the pediatric population. Recent animal studies have suggested a role for pRb in the regulation of mesenchymal cell fate towards osteoblasts, by potentiating the effect of Runx2 a master regulator for osteoblast differentiation (12). pRb1 is also known to be important for adipocytic differentiation as it is required for activating several of the transcription factors involved in adipocyte differentiation (13). However, complete loss of Rb1 expression from mesenchymal stem cells predisposes them to an adipocytic cell fate (14). While the role is complex and yet to be clearly elucidated, Rb1 is generally considered to regulate cell fate and in the absence of Rb1 cells default to an adipogenic phenotype (14, 15).

In this study, we attempted to determine the effect exposure to adipogenesis promoting (AP) media on MC3T3-E1 preosteoblasts. We also determined how the presence of vitamin D might affect the expression of adipocytic phenotype. The effect of Rb1 deficiency in osteoblasts and their transdifferentiation ability to adipocytes was also studied.

## MATERIALS AND METHODS

### Cell Culture

The MC3T3-E1 subclone 4 (ATCC CRL-2593) is a preosteoblast cell line isolated from mouse calvaria and is a well-established

model to study osteoblast differentiation. Cell lines were grown in alpha-minimum essential media ( $\alpha$ MEM) (Life Technologies) with 10% fetal bovine serum (FBS) (Atlanta Pharmaceuticals) under 5% CO<sub>2</sub> at 37°C.

### Stable Rb1 Line

MC3T3-E1 cells were used to create Rb1 deficient line using shRNA technology. A stable clone of cells containing less than 40% of wild-type Rb1 expression was used for these studies. A control MC3T3-E1 line containing a stably expressing scrambled shRNA control was also used.

### Adipogenic Media and Vitamin D Treatment

The AP media used consisted of 2% FBS and  $\alpha$ MEM containing 0.5 mM 3-isobutyl-1-methylxanthine (Sigma-Aldrich, St. Louis, MO, USA), 1 $\mu$ M dexamethasone (Sigma), 10  $\mu$ g/ml Insulin (Sigma). The AP media were changed every 48 h after the initial treatment, and the cells were maintained in an incubator at 37°C and 5% carbon dioxide.

Vitamin D (1,25-dihydroxyvitamin D<sub>3</sub>) was purchased from Sigma-Aldrich, St. Louis, MO, USA and utilized at a final concentration of 20 nM dissolved in ethanol in AP media. All experiments were carried out in triplicates and repeated thrice.

To analyze expression of adipocyte markers after exposure to AP media, we plated equal numbers of cells and grew them to 70% confluency in regular media (10% FBS in DMEM) and switched them to serum free for 48 h before exposing them to AP media described above. Media were changed every 48 h, and a plate was harvested each day for 6 days. A similar protocol was followed for vitamin D treatment. Control cells received the same volume of the solvent used for vitamin D (ethanol).

For analyses of adipogenic expression in Rb1 knockdown cells, we used shRNA technology to stably reduce Rb1 expression. Several clones were first characterized and analyzed, and in this study, we used a clone with a 50% stable reduction of Rb1. Control MC3T3-E1 cells carried a stable scrambled shRNA sequence. These cells were treated as described above, except the vitamin D treatment was not performed on these cells. Validation of knockdown was carried out at the level of RNA and protein.

### SyberGreen Real-time PCR

RNA was isolated using TRI reagent (Sigma). A Two-step RT-PCR was performed using Bullseye EvaGreen qPCR master Mix (Midsci) and RT-PCR Reagents Kit with Step One Plus Real Time Machine (Applied Biosystems, Roche, IN). RNA sample were converted into cDNA, using a high-capacity cDNA RT kit. Analysis of the results was performed with the 7300 RT-PCR System RQ Software version 1.4 (Applied Biosystems). Efficiency of the control and treatment samples were checked using the slope of the graph of  $\Delta$ CT values versus log of the total RNA sample. If the calculated slope was less than 0.1, then comparative CT method was used to compare the gene expression between control and treatment cells. These were normalized according to a housekeeping gene,  $\beta$ -actin, serving as an internal control.

## Primer Pairs Used in Real-time PCR

Primers were designed using Integrated DNA Technologies, Coralville, IA, USA PrimerQuest Tool. The Genbank number was used to import the sequence, and the parameters of  $T_m$  of 60° and amplicon size of 75–125 bp were used. Sequence sets that resulted with amplicon ~100 bp were chosen. Primer pair sequences used are shown in **Table 1**.

## Protein Isolation and Western Blot Assay

Cells were lysed with buffer, containing 50 mM DTT, 2% SDS, 10% glycerol, 65 mM Tris-HCl pH 6.8, with protease inhibitor cocktail (cComplete ULTRA Tablets Mini Roche protease inhibitor). The lysate was collected and sonicated with a Bioruptor 300 (Diagenode, NJ, USA). Protein concentrations were measured, and a standard curve was generated using a Pierce BCA Protein Assay Kit (Thermo Scientific). Rabbit VDR (Proteintech, Chicago, IL, USA) and Rabbit C/EBP $\alpha$  (Proteintech) were used. SuperSignal® West Pico Chemiluminescent Substrate (Thermo Scientific) was used to visualize the membrane using BioRad ChemiDoc MP Imaging System (BIORAD, CA, USA).

## Luciferase Assay

Luciferase assay for PPAR $\gamma$  activity was conducted with PPAR $\gamma$  luciferase construct (a kind gift of Dr. B. Spiegelman, Addgene #1015) using BrightGlow (Promega, WI, USA) and measured with a GLOMAX 20/20 Luminometer. Transfections were carried out on MC3T3 E1 cells (control) or cells with stable Rb1 knockdown using Superfect (Qiagen) according to manufacturer recommendations. Forty-eight hours after transfection, cells were trypsinized, pelleted, and then resuspended in serum free

media. An aliquot was used to count cells for normalization, the rest was assayed using Promega's Bright-Glo™ Luciferase Assay Kit (Promega, cat. no. E2610). Equal volumes of the resuspended cells and Bright-Glo™ were mixed and incubated for 2 min. Samples were read for 10 s, each after a 2 s delay. Luciferase readings for each of the samples were then normalized according to the number of cells per milliliters. All measurements were carried out on triplicate samples, and experiments were repeated at least thrice.

## Statistical Analyses

Data were analyzed by one or two way ANOVA for effects of vitamin D, media type (regular media, adipogenic media) and/or time point (as appropriate) using GraphPad Prism 7 statistical software (GraphPad Software Inc., La Jolla, CA, USA) A *post hoc* Bonferroni's test was used when appropriate. Data are expressed as Mean  $\pm$  SE. A *P* value of <0.05 was considered significant.

## RESULTS



To elucidate if exposure of AP media to the preosteoblast cell line MC3T3-E1 produced transdifferentiation to an adipocytic phenotype, we exposed cells for 6 days to the media and analyzed RNA expression for several transcription factors that regulate adipogenic differentiation. The transcriptional program regulating adipogenesis is well understood and proceeds through sequential expression of different transcription factors (16). Our data on MC3T3-E1 cells followed the expected changes that occur during adipogenesis where C/EBP $\beta$  and C/EBP $\delta$  expression precedes expression of PPAR $\gamma$  and C/EBP $\alpha$ . As seen in **Figure 1A**, the two transcription factors (C/EBP $\beta$  and C/EBP $\delta$ ) were increased within a day of treatment to about fourfold and a steady increase that continued through the treatment ( $P < 0.01$  for C/EBP $\beta$ ,  $P < 0.05$  for C/EBP $\delta$ ). C/EBP $\beta$  showed a larger increase (15-fold) than C/EBP $\delta$  (5-fold). This was followed by an increase in PPAR $\gamma$  after 3 and 6 days of treatment ( $P < 0.01$ ). An increase in C/EBP $\alpha$  was observed on Day 6 on AP media. The changes in these transcription factors were accompanied by an increase in expression of several adipocyte markers such as FABP4, UCP1, and SREBF1 (**Figure 1B**). Positive Oil Red O staining confirmed the presence of lipid droplets, which were observed in MC3T3-E1 cells exposed to AP media but not regular media (**Figure 1C**).

To determine the effect of vitamin D on osteoblasts in the presence of adipogenic differentiation factors, we exposed cells to 20 nM vitamin D for a week with media change every other day. Real-time PCR analyses of adipocyte transcription factors revealed a large upregulation of expression with vitamin D treatment. **Figure 2** shows changes seen with and without vitamin D treatment of MC3T3-E1 cells. The overall sequence of expression of the transcription factors followed what we saw in cells without vitamin D. However, there was a dramatic statistically significant ( $P < 0.001$ ) interaction between media and vitamin D in the upregulation of several factors. The only transcription factor that did not appear to be affected by vitamin D was C/EBP $\beta$ , which showed a change similar to that of cells treated only with adipogenic media. C/EBP $\delta$  showed about a 50-fold increase, PPAR $\gamma$

**TABLE 1** | Mouse specific primer sequences used for quantitative real-time PCR analysis.

Primer	Sequence
VDR forward	GGCTTCCACTTCAACGCTATG
VDR reverse	TGCTCCGCCTGAAGAAACC
FABP4 forward	CCGCAGACGACAGGAAGGT
FABP4 reverse	AGGCCCCCGCCATCT
Srebf1 forward	GCGGTTGGCACAGAGCTT
Srebf1 reverse	CTGTGGCCTCATGTAGGAATACC
AdipoQ forward	GACACAAAAGGGCTCAGGAT
AdipoQ reverse	TGGGCAGGATTAAGAGGAACA
Ucp1 forward	ACTGCCACACCTCCAGTCATT
Ucp1 reverse	CTTTGCCTCACTCAGGATTGG
C/EBP $\alpha$ forward	CAAGAACAGCAACGAGTACCG
C/EBP $\alpha$ reverse	GTCACTGGTCAACTCCAGCAC
C/EBP $\beta$ forward	GGGTTGTTGATGTTTTGGTTT
C/EBP $\beta$ reverse	GAAACGGAAAAGGTTCTCAAAA
C/EBP $\delta$ forward	CGACTTCAGCGCTACATTGA
C/EBP $\delta$ reverse	GAAGAGGTCGGCGAAGAGTT
Peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) forward	TGAAAGAAGCGGTGAACCACTG
PPAR $\gamma$ reverse	TGGCATCTCTGTGTCAACCATG
$\beta$ -Actin forward	TGTCCACCTTCCAGCAGATGT
$\beta$ -Actin reverse	AGCTCAGTA ACAGTCCGCCTAG

Primer sequences were designed using PrimerQuest tool (Integrated DNA Technologies) as described in Section "Materials and Methods."



a 200-fold increase, and C/EBP $\alpha$  a 300-fold increase in activity when compared to cells treated only with adipogenic media.

To confirm that this was a result of vitamin D receptor mediated action, we measured expression of vitamin D receptor

(VDR) during vitamin D treatment. We treated MC3T3-E1 cells with adipogenic media with 20 nM 1,25 dihydroxyvitamin D with frequent changes of media as described in Section "Materials and Methods." Control cells received the same amount of vehicle

(EtOH). Addition of vitamin D produced a maximal increase of 600-fold 4 days ( $P < 0.001$ ) after treatment with media with other days showing several fold elevation of expression (Figures 3A,B), demonstrating that the action of vitamin D did indeed result in an increase in its cognate receptor levels. Western blots for PPAR $\gamma$  (not shown) and C/EBP $\alpha$  (Figure 3C) also confirmed the changes seen in RNA expression.

## Expression of Adipogenic Marker Genes


We tested the expression of FABP4 and SREBF1 after vitamin D treatment. When compared to untreated cells, treatment with vitamin D produced dramatic increases in FABP4 and SREBF1 expression (Figure 4A). Transcripts for both these markers were increased by vitamin D treatment in the range of 100-fold and higher. Consistent with these results, we observed an increase in lipid droplet formation when cells were grown in the presence of vitamin D. Figure 4B showing that vitamin D was able to stimulate the transdifferentiation of osteoblasts to adipocytes when cultured in the presence of adipogenic differentiation medium.


## Effect of pRb1 Deficiency on Adipocytic Differentiation of Osteoblasts

While our results with vitamin D suggested that the manipulation of preosteoblast MC3T3-E1 cells to express an adipocytic program is influenced dramatically at the level of transcription even though adipogenic genes are not known to be under hormone regulation by vitamin D. Therefore, we sought to determine if perturbation of key genes known to regulate adipogenesis might also produce a similar outcome. Recent studies have suggested that the presence of Rb1 is essential for osteoblast cell fate (14, 15).

Other studies have demonstrated a role for pRb in the regulation of C/EBPs (13). pRb is also known to suppress PPAR $\gamma$  activity and activate Runx2 a osteoblast master regulator during mesenchymal cell differentiation (14). In this study, we attempted to determine how exposure to adipogenic media might affect osteoblasts with a deficiency in Rb1 expression. Figure 5 inset shows the generation and validation of Rb knockdown in MC3T3-E1. Other than a slight increase in growth rate (1.18-fold higher than control cells receiving scrambled control) that was not statistically significant (not shown), the cells did not appear to have any discernable changes to morphology. Several clones were analyzed and characterized, and we used one such clone to determine the effect of Rb deficiency on adipogenic program. As shown in Figure 5A, PPAR $\gamma$  expression trended toward being generally higher in Rb KD cells when compared to control cells but the data appeared highly variable. C/EBP $\alpha$  and C/EBP $\beta$  were also generally higher in Rb1 deficient cells and showed a clear increase with Rb loss ( $P < 0.001$ ) during exposure to adipogenic media. On the other hand C/EBP $\delta$  appeared to be unaffected by Rb1 loss (Figure 5A). The increase in adipocyte differentiation of the Rb1 knockdown cells was also confirmed with Oil Red O staining (Figure 5B) and the expression of adipocyte makers FABP4, UCP1, and SREBF1 (Figure 5C). We included UCP1 a marker of brown adipose tissue in our analyses as pRb has been shown to play a role in white and brown adipogenesis (13). While the extent of elevation of these markers was moderate when compared to what we observed with vitamin D treatment, nevertheless deficiency of Rb1 expression produced 20-fold to 100-fold increase in markers such as FAB4, UCP1, and SREBF1 (Figure 5C).

To confirm that these manipulations actually produced changes in PPAR $\gamma$  function, we conducted a luciferase assay





**FIGURE 4** | Adipocyte marker expression levels during vitamin D treatment in the presence of adipogenesis promoting (AP) media. **(A)** Real-time PCR of adipocyte markers after vitamin D treatment. Other details are similar to that shown in **Figure 2**. All statistics were calculated relative to control with no vitamin D treatment with O time set at 1. **(B)** Oil Red O staining of cells after vitamin D exposure demonstrates increased fatty vacuoles (left). Control cells received AP media and vehicle with no vitamin D (right) 20x.

with cells used in the above experiments. A construct containing six copies of PPAR $\gamma$  response elements were transiently transfected into cells followed by exposure to adipogenic media for 3 days in the presence and absence of vitamin D. As shown in **Figure 6**, maximal activity was seen after vitamin D treatment but just exposure to AP media was sufficient to produce a statistically significant ( $P < 0.001$ ) increase in activity in control MC3T3-E1 cells. Loss of Rb1 expression also resulted in a statistically significant increase in PPAR $\gamma$  activity ( $P < 0.001$ ) (**Figure 6**).

## DISCUSSION

Adipocytes and osteoblasts share a common precursor in the bone marrow, and it is known that there is a high degree of plasticity between the two cell lineages (1). MC3T3-E1 is a clonal preosteoblast cell line that was originally isolated from new born mouse calvaria (17). To demonstrate plasticity in this line, investigators have generally employed genetic manipulations that have included overexpression of adipogenic transcription factors (18), treatment with drugs (19), etc. In this study, we attempted to understand the influence of adipogenic media with and without vitamin D or the effect of reduction in Rb1 expression on MC3T3-E1 cells. Our studies show that exposure of these cells to media generally utilized to promote adipogenesis in preadipocyte cell lines such as 3T3-L1 can produce gene expression characteristic of adipocytes in determined preosteoblasts. However, we were surprised to find a synergistic enhancement of adipocyte differentiation factors and

associated markers on treatment with vitamin D. While the role of vitamin D receptor and vitamin D on adipogenesis has not been clearly elucidated, it is well known that vitamin D receptors are necessary and present in adipocytes (20, 21). In fact, studies have demonstrated an inhibition by vitamin D of adipocyte specific expression of PPAR $\gamma$  and other factors in preadipocytes (22). None of the other transcription factors involved in adipocyte differentiation are also known to be regulated by vitamin D. Since VDR is expressed in response to vitamin D treatment, it appears that some of the effect could be mediated at the level of ligand receptor interaction. However, it is also known that vitamin D might have non-genomic actions (23) that may not involve the classical steroid hormone mediated stimulation of gene expression. Irrespective of the mechanism, treatment of preosteoblasts with vitamin D did not inhibit but promoted adipocyte differentiation. We cannot exclude the fact that adipogenic media components 3-isobutyl-1-methylxanthine and insulin may have played a role in influencing the adipocytic phenotype of these cells (20, 21).

While the final outcome of MC3T3-E1 cells exposed to adipogenic media was the same with and without vitamin D, there were important differences in expression of C/EBP factors. In the case exposure to just the media, the changes in gene expression agreed with previously published observations during preadipocyte differentiation to adipocytes (24). With vitamin D, our results show a dramatic increase in the case of C/EBP $\alpha$ , which went up several hundred fold and C/EBP $\delta$  which was increased 20-fold to 40-fold over just the media alone. While the extent of changes seen in C/EBP $\beta$  after exposure to AP media

was the highest, with addition of vitamin the effect remained the same without any enhancement we saw in CEBP $\alpha$  and C/EBP $\delta$  (Figure 2). In a preadipocyte cell line 3T3-L1, liganded vitamin D appears to inhibit CEBP $\beta$  while unliganded VDR is seen necessary for adipogenesis (21). Our results may represent an inhibition of CEBP $\beta$  expression by vitamin D treatment, but the fact that upregulation provided by the media alone was still retained, might explain the overall enhancement in adipogenesis.

Previous work has suggested that the tumor suppressor gene Rb1 might be critical for cell fate determination and it has been shown that loss of pRb1 in mesenchymal stem cells cause them to default to an adipocytic cell lineage (14). We therefore used a cell line that has stable reduction in Rb1 expression to demonstrate

that a similar phenotype of enhanced adipogenesis was seen. Interestingly, in these cells, there was a steady state increase in PPAR $\gamma$  expression and agrees with previous work that suggests that pRb inhibits PPAR $\gamma$  expression in mesenchymal stem cells (25). C/EBP $\alpha$  and C/EBP $\beta$  expression was generally increased with Rb1 loss, but C/EBP $\delta$  expression was unaffected. This was unlike our observations with vitamin D and might relate to independent roles of Rb1 in regulating these transcription factors. It is also known the Rb1 regulation of adipogenesis might depend on several factors, since pRb regulates not only the CEBPs but also cell cycle control through E2Fs. In addition, loss of Rb1 promotes brown adipose tissue conversion as evidenced by markers such as UCP1 by these cells (Figure 5C).






FIGURE 5 | Continued



Our studies generally reveal that uninduced preosteoblasts maintain their plasticity toward adipocytes and factors in the adipogenic media appear to be sufficient to induce MC3T3-E1 cells to express the adipocyte phenotype. While this is not surprising, what is surprising is the fact that treatment with vitamin D in the presence of adipogenic factors dramatically enhances gene expression in preosteoblasts toward adipocytes. While this result was unexpected, studies that have looked at genomic binding of transcription factors in differentiated and undifferentiated cells demonstrate that this might relate to the plasticity of the epigenome (26). Both histone modifications and transcription factor binding appear to be different in mature versus immature osteoblasts (26, 27). While there may be a default pathway or preference for a particular cell fate, it appears to be easily perturbed at the level of transcription factor binding (26). In fact, these investigators show that transdifferentiation of preosteoblasts to adipocytes is associated with rapid changes to chromatin in regions containing the adipocyte specific genes in MC3T3-E1 cells. It is also known that unliganded VDR remains bound to a number of target genes with the genome and undergo modifications with vitamin D treatment. It is possible that vitamin D treatment might not have directly contributed to the activation of adipocyte specific gene expression but may have altered the transcription factor binding sites of VDR and essentially mitigated the inhibition of VDR targets. It is known that vitamin D exerts both positive and negative effects on osteoblast differentiation (25) and is dependent on timing. In immature osteoblasts, vitamin D



downregulated the master regulator of osteoblast differentiation Runx2 while increasing PPAR activity and adipocyte differentiation (28). While our studies have not investigated in parallel how osteoblast differentiation may be affected in the presence of adipogenic media and vitamin D, it is likely that competition of adipogenic and osteogenic transcription factors for binding on DNA might have influenced the response seen. There is much to be learned about vitamin D targets as several of the sites where VDR bound on chromatin is not known targets of vitamin D action (27). Our results suggest the need for further functional studies to uncover vitamin D-dependent pathways that govern adipocyte–osteoblast transdifferentiation.

## REFERENCES

- Berendsen AD, Olsen BR. Osteoblast-adipocyte lineage plasticity in tissue development, maintenance and pathology. *Cell Mol Life Sci* (2014) 71(3):493–7. doi:10.1007/s00018-013-1440-z
- Savopoulos Ch, Dokos Ch, Kaiafa G, Hatzitolios A. Adipogenesis and osteoblastogenesis: trans-differentiation in the pathophysiology of bone disorders. *Hippokratia* (2011) 15(1):18–21.
- Guo L, Li X, Tang QQ. Transcriptional regulation of adipocyte differentiation: a central role for CCAAT/enhancer-binding protein (C/EBP)  $\beta$ . *J Biol Chem* (2015) 290(2):755–61. doi:10.1074/jbc.R114.619957
- Bellows CG, Heersche JN. The frequency of common progenitors for adipocytes and osteoblasts and of committed and restricted adipocyte and osteoblast progenitors in fetal rat calvaria cell populations. *J Bone Miner Res* (2001) 16(11):1983–93. doi:10.1359/jbmr.2001.16.11.1983
- Kodama HA, Amagai Y, Koyama H, Kasai S. Hormonal responsiveness of a preadipose cell line derived from newborn mouse calvaria. *J Cell Physiol* (1982) 112(1):83–8. doi:10.1002/jcp.1041120113
- Poulos SP, Dodson MV, Culver MF, Hausman GJ. The increasingly complex regulation of adipocyte differentiation. *Exp Biol Med (Maywood)* (2016) 241(5):449–56. doi:10.1177/1535370215619041
- Ji S, Doumit ME, Hill RA. Regulation of adipogenesis and key adipogenic gene expression by 1, 25-dihydroxyvitamin D in 3T3-L1 cells. *PLoS One* (2015) 10(6):e0126142. doi:10.1371/journal.pone.0126142
- Narvaez CJ, Simmons KM, Brunton J, Salinero A, Chittur SV, Welsh JE. Induction of STEAP4 correlates with 1,25-dihydroxyvitamin D3 stimulation of adipogenesis in mesenchymal progenitor cells derived from human adipose tissue. *J Cell Physiol* (2013) 228(10):2024–36. doi:10.1002/jcp.24371
- Dunlop TW, Väisänen S, Frank C, Molnár F, Sinkkonen L, Carlberg C. The human peroxisome proliferator-activated receptor delta gene is a primary target of 1 $\alpha$ ,25-dihydroxyvitamin D3 and its nuclear receptor. *J Mol Biol* (2005) 349(2):248–60. doi:10.1016/j.jmb.2005.03.060
- Alimirah F, Peng X, Yuan L, Mehta RR, von Knethen A, Choubey D, et al. Crosstalk between the peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) and the vitamin D receptor (VDR) in human breast cancer cells: PPAR $\gamma$  binds to VDR and inhibits 1 $\alpha$ ,25-dihydroxyvitamin D3 mediated transactivation. *Exp Cell Res* (2012) 318(19):2490–7. doi:10.1016/j.yexcr.2012.07.020
- Dyson NJ. RB1: a prototype tumor suppressor and an enigma. *Genes Dev* (2016) 30(13):1492–502. doi:10.1101/gad.282145.116
- Thomas D, Kansara M. Epigenetic modifications in osteogenic differentiation and transformation. *J Cell Biochem* (2006) 98(4):757–69. doi:10.1002/jcb.20850
- Hallenborg P, Feddersen S, Madsen L, Kristiansen K. The tumor suppressors pRB and p53 as regulators of adipocyte differentiation and function. *Expert Opin Ther Targets* (2009) 13(2):235–46. doi:10.1517/14712590802680141
- Calo E, Quintero-Estades JA, Danielian PS, Nedelcu S, Berman SD, Lees JA. Rb regulates fate choice and lineage commitment in vivo. *Nature* (2010) 466(7310):1110–4. doi:10.1038/nature09264
- Gündüz V, Kong E, Bryan CD, Hinds PW. Loss of the retinoblastoma tumor suppressor protein in murine calvaria facilitates immortalization of osteoblast-adipocyte bipotent progenitor cells characterized by low expression of N-cadherin. *Mol Cell Biol* (2012) 32(13):2561–9. doi:10.1128/MCB.06453-11
- Siersbaek R, Mandrup S. Transcriptional networks controlling adipocyte differentiation. *Cold Spring Harb Symp Quant Biol* (2011) 76:247–55. doi:10.1101/sqb.2011.76.010512
- Sudo H, Kodama HA, Amagai Y, Yamamoto S, Kasai S. In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *J Cell Biol* (1983) 96(1):191–8. doi:10.1083/jcb.96.1.191
- Kim SW, Her SJ, Kim SY, Shin CS. Ectopic overexpression of adipogenic transcription factors induces transdifferentiation of MC3T3-E1 osteoblasts. *Biochem Biophys Res Commun* (2005) 327(3):811–9. doi:10.1016/j.bbrc.2004.12.076
- Weivoda MM, Hohl RJ. Geranylgeranyl pyrophosphate stimulates PPAR $\gamma$  expression and adipogenesis through the inhibition of osteoblast differentiation. *Bone* (2012) 50(2):467–76. doi:10.1016/j.bone.2011.09.056
- Wood RJ. Vitamin D and adipogenesis: new molecular insights. *Nutr Rev* (2008) 66(1):40–6. doi:10.1111/j.1753-4887.2007.00004.x
- Blumberg JM, Tzamelis I, Astapova I, Lam FS, Flier JS, Hollenberg AN. Complex role of the vitamin D receptor and its ligand in adipogenesis in 3T3-L1 cells. *J Biol Chem* (2006) 281(16):11205–13. doi:10.1074/jbc.M510343200
- Kong J, Li YC. Molecular mechanism of 1,25-dihydroxyvitamin D3 inhibition of adipogenesis in 3T3-L1 cells. *Am J Physiol Endocrinol Metab* (2006) 290(5):E916–24. doi:10.1152/ajpendo.00410.2005
- Farach-Carson MC, Ridall AL. Dual 1,25-dihydroxyvitamin D3 signal response pathways in osteoblasts: cross-talk between genomic and membrane-initiated pathways. *Am J Kidney Dis* (1998) 31(4):729–42. doi:10.1053/ajkd.1998.v31.pm9531195
- Ntambi JM, Young-Cheul K. Adipocyte differentiation and gene expression. *J Nutr* (2000) 130(12):3122S–6S.
- Fajas L, Egler V, Reiter R, Hansen J, Kristiansen K, Debril MB, et al. The retinoblastoma-histone deacetylase 3 complex inhibits PPAR $\gamma$  and adipocyte differentiation. *Dev Cell* (2002) 3(6):903–10. doi:10.1016/S1534-5807(02)00360-X
- Meyer MB, Benkusky NA, Sen B, Rubin J, Pike JW. Epigenetic plasticity drives adipogenic and osteogenic differentiation of marrow-derived mesenchymal stem cells. *J Biol Chem* (2016) 291(34):17829–47. doi:10.1074/jbc.M116.736538
- Meyer MB, Benkusky NA, Lee CH, Pike JW. Genomic determinants of gene regulation by 1,25-dihydroxyvitamin D3 during osteoblast-lineage cell differentiation. *J Biol Chem* (2014) 289(28):19539–54. doi:10.1074/jbc.M114.578104
- Kim JH, Seong S, Kim K, Kim I, Jeong BC, Kim N. Downregulation of Runx2 by 1,25-dihydroxyvitamin D3 induces the transdifferentiation of osteoblasts to adipocytes. *Int J Mol Sci* (2016) 17(5):770–81. doi:10.3390/ijms17050770

## AUTHOR CONTRIBUTIONS

EP designed and performed experiments, analyzed data, and helped write the paper; NC designed experiments, analyzed data, and wrote the paper.

## ACKNOWLEDGMENTS

This work was supported by intramural funds from MWU. No experiments were performed on animals. Midwestern University's policies and national guidelines do not require ethics approval for this study.



**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Pendleton and Chandar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.