



Conservation of a Critically Endangered Endemic Halophyte of West Portugal: A Microcosm Assay to Assess the Potential of Soil Technology for Species Reintroduction

Ana Cortinhas, Teresa Cardoso Ferreira, Maria Manuela Abreu and Ana D. Caperta*

Linking Landscape, Environment, Agriculture and Food (LEAF), Instituto Superior de Agronomia (ISA), Universidade de Lisboa, Lisbon, Portugal

OPEN ACCESS

Edited by:

Terry D. Macfarlane,
Conservation and Attractions (DBCA),
Australia

Reviewed by:

Pavol Elias,
Slovak University of Agriculture,
Slovakia

Ana Maria Rodrigues Cassiolato,
São Paulo State University, Brazil

*Correspondence:

Ana D. Caperta
anadelaunay@isa.ulisboa.pt

Specialty section:

This article was submitted to
Conservation and Restoration
Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 09 September 2020

Accepted: 13 April 2021

Published: 04 May 2021

Citation:

Cortinhas A, Ferreira TC,
Abreu MM and Caperta AD (2021)
Conservation of a Critically
Endangered Endemic Halophyte
of West Portugal: A Microcosm Assay
to Assess the Potential of Soil
Technology for Species
Reintroduction.
Front. Ecol. Evol. 9:604509.
doi: 10.3389/fevo.2021.604509

The soil system has been frequently overlooked during plant reintroduction planning and practice since working with soils and plant roots can be difficult, particularly in saline environments. Coastal saline environments are major contributors to regional and global biodiversity and an important source of endemic species. However, various species are in decline or considered threatened, particularly halophytes (salt tolerant) due to negative anthropic impacts. The Lusitanian endemic halophyte *Limonium daveaui* formerly had a large distribution range along the west coast of Portugal but currently it shows a restricted distribution in the Tagus estuary. Field surveys revealed that this critically endangered species forms few local populations with small size invaded by exotic species. In this study, we investigated the potential utilization of Technosols, an innovative sustainable, ecological engineering method combined with brackish water irrigation for potential *L. daveaui* reintroduction in native habitats. Seed germination percentages were evaluated in different environmental conditions. Through a microcosm assay, a Technosol was constructed using a saline Fluvisol with a mixture of low value inorganic and organic wastes, which were chemically characterized. Plants were cultivated in the Fluvisol and Technosol and irrigated with brackish water collected in the nearby area. To assess plant growth, morphometric parameters and the plants' physiological status were assessed and the fresh and dry biomass determined. Results showed that seed germination was higher on moist filter paper with distilled water than in Fluvisol or Technosol. Plants grown in Technosol had a greater development, with higher values of photosynthetic indexes and biomass production than in Fluvisol. Our findings provide a basis for future *in situ* conservation studies and support the idea that eco-friendly soil technology approaches are beneficial to conserve rare halophyte species.

Keywords: estuarine water, *Limonium daveaui*, fluvisol, tailored soils, plant restoration, soil technology, wastes

INTRODUCTION

Reintroduction of native species is an important issue in conservation and has been increasingly acknowledged in international treaties and legislation, particularly endemic species (Maunder, 1992; Godefroid et al., 2011; Maschinski and Albrecht, 2017). Conservation of coastal endemics from rocky cliffs, sand dunes and salt marshes is challenging due to the high degree of habitat specialization of these species and their vulnerability to disturbance by several natural factors, invasive species and human activities (Baumberger et al., 2012; Martins et al., 2013; Caperta et al., 2014; Lechuga-Lago et al., 2017; Olsson et al., 2019).

The Portuguese coast is rich in endemic flora due to a singular biogeographic position (Braun-Blanquet et al., 1972; Asensi et al., 1993; van der Maarel and van der Maarel-Versluys, 1996; Costa et al., 2012) and about 35% of all Portuguese Natura 2000 habitats consist of coastal habitats (Costa et al., 2007; Duarte et al., 2016). The critically endangered endemic species *Limonium daveaui* Erben had a large distribution in the marshes of the Tagus Estuary (SW Portugal, SW Europe) in the past (Erben, 1978, 1993; Franco, 1984; Espírito-Santo et al., 2012; Caperta and Carapeto, 2020) but currently it is narrowly distributed in this area (Caperta and Carapeto, 2020). Among the reasons invoked for the continuous decline of the species in this estuary are the deterioration of its habitat quality due to invasive species (e.g., *Carpobrotus edulis*) and anthropic pressures (Caperta and Carapeto, 2020) like industry activities (petrochemical and steelwork) with huge requirements for space and suitable terrain for industrial complexes in the Lisbon Metropolitan Area (Costa, 2013; Fernandes et al., 2020). Moreover, the Tagus estuary is also affected by other anthropic activities (chemicals, ship construction and repair and cement manufacture), agriculture-fertilizers and pesticides (Duarte et al., 2014). Due to these activities, this estuary was considered as contaminated since 1985 (Figueroes et al., 1985), with high levels of anthropic Hg in Fluvisols (Cesário et al., 2016).

A detailed understanding of the biology and microenvironmental requirements including soil is needed for successful reintroduction of rare species (Holl and Hayes, 2006; Dunwiddie and Martin, 2016). Conservation practitioners are increasingly using diverse soil-based technologies to propagate and restore rare plant species (Haskins and Pence, 2012). One of the most challenging aspects of rare species reintroduction is the low success obtained when seeds rather than seedlings are used to start new populations *in situ* (Davy, 2002; Godefroid et al., 2011; Albrecht and Maschinski, 2012). Physically adverse environments and soil disturbance have also been shown to limit the possible success of species reintroduction using directly sown seeds (Wang et al., 2014; Menges et al., 2016; Cao et al., 2018; van Regteren et al., 2020), as observed in salt marshes where soil disturbance reduces successful seedling establishment (Balke et al., 2011). In the surface layer of sediments/soils, poor drainage may result in the build-up of potentially toxic dissolved species, high salinity, and anoxia that make seed germination difficult (Spencer et al., 2017). An

alternative environmentally friendly management of Fluvisols is via elaboration of Technosols, man-made tailored soils, whose properties and pedogenesis are dominated by their technical origin (IUSS Working Group, 2015). These soils could be elaborated from wastes and employed in the subsequent recovery of degraded and/or polluted soils, improving their physicochemical and biological properties (Macia et al., 2014; Cortinhas et al., 2020), while contributing to the circular economy (Breure et al., 2018).

This study aims to assess the potential of tailored soils to improve (i) soil properties, (ii) seed germination, (iii) seedling establishment, and (iv) plant growth of an endemic, critically endangered species, through a microcosm assay using estuarine water irrigation. The study also hopes to develop awareness of eco-friendly soil technologies for conservation of rare species.

MATERIALS AND METHODS

Study Area and Species

The Fluvial beach “Praia do Sobralinho,” Sobralinho, in the Tagus Estuary (38°54'16.1"N, 9°01'09.0"W), in the Municipal Council of Vila Franca de Xira¹, is integrated in the Lisbon metropolitan area, on the west coast of Portugal (Instituto da Conservação da Natureza e das Florestas [ICNF], 2012; **Figure 1**). The saline tide reaches about 50 km upstream from the river mouth, near Vila Franca de Xira (Guerreiro et al., 2015).

The upper saltmarsh area encompasses plant communities with diverse halophytes found in NATURA 2000: Habitat 1420 Mediterranean and thermo-Atlantic halophilous scrubs (*Sarcocornetea fruticosi*) and in the priority Habitat 2130 Mediterraneo-Atlantic fixed gray dunes (The Environmental Protection Agency of Portugal [APA], 2011), where the Lusitanian endemic *Limonium daveaui* (syn. *Limonium auriculae-ursifolium* (Poir.) Druce subsp. *lusitanicum* (Pignatti) Pignatti; *Limonium ramosissimum* (Poir.) Maire subsp. *confusum* (Gren. et Godr.) Pignatti (Pignatti, 1971) thrives (Espírito-Santo et al., 2012). This species comprises perennial plants with a basal rosette, and oblanceolate-spatulate glabrous, three parallel veined leaves. The inflorescences (scapes) are branched in the lower third, with a few sterile branches, and present bluish violaceous flowers blooming between June and August (Erben, 1978, 1993; Franco, 1984).

Microcosm Assay

A microcosm assay was set up with a Salic Fluvisol (Eutric) (FLU; IUSS Working Group, 2015) collected in Sobralinho salt marsh area, which was dried and sieved (<2 mm) and a Technosol (TEC). The TEC was produced with 85% of FLU and 15% of organic/inorganic wastes mixture that were added and mixed manually. The wastes mixture used as amendments was constituted by sludge and waste kieselguhr from breweries, medium sand (0.25–0.5 mm), gravel limestone (2 mm < Ø < 5 mm) and residual biomass obtained from

¹<https://www.cm-vfxira.pt/>




FIGURE 1 | The study area located in Sobralinho salt marsh area, on the right bank of the river Tagus, in the Municipal Council of Vila Franca de Xira, west Portugal.

pruning, in the proportion 1.5:0.5:3:2:3, respectively. The FLU and TEC were potted and incubated at 70% of the maximum water-holding capacity, in the dark, for 28 days. After the incubation, and before seedling transplantation, composite samples from each pot (0–15 cm of depth) were collected and analyzed. This process was repeated at the end of the assay. For the microcosm assay, the seedlings were obtained from seeds germinated on wet filter paper with deionized water and after 2.5 months (~10 cm) they were transplanted to the substrata, in five replicates, irrigated with brackish water collected from a channel located in the Tagus estuary, keeping the substrata at 70% of the maximum water-holding capacity (400 ml).

Fluvisol, Amendments, and Brackish Water Characterization

The FLU, amendments and TEC were characterized following methods in Póvoas and Barral (1992) for: pH and electric conductivity (EC) in a water suspension (1:2.5 *m/V*); extractable K ($K_{extract}$; Egner–Riehm method) and P ($P_{extract}$; Olsen method); total N (N_{total} ; Kjeldahl method); organic C (C_{org}) by wet combustion; micronutrients (Lakanen and Erviö, 1971). For the FLU, the cation exchange capacity (CEC) by 1 mol/dm³ ammonium acetate, particle size distribution analysis by sieving and sedimentation, and multielemental concentration by ICP-MS after acid digestion (Activation Laboratories, 2020) were determined.

The brackish water was analyzed for: pH (potentiometry), EC (conductivity), Cl[−] (Mohr, 1945), HCO₃[−] (titration method using HCl solution methyl orange as indicator), Na, Ca, Mg, K, P, Fe, Zn, Mn, and Cu (atomic absorption spectrometry). The Na adsorption ratio (SAR) was calculated by

$$\left(\frac{[Na]}{([Ca] + [Mg])^{1/2}} \right)$$
 (Ayers and Westcot, 1985; Lesch and Suarez, 2009).

Germination Tests

The *L. daveaui* seeds were collected in Samouco salterns complex and germinated in transparent boxes containing filter paper, TEC and FLU, respectively. Each box contained 36 seeds equally distant from each other and 170 ml of deionized water. The boxes were exposed at the temperature 20–25 °C with a 16 h light and 8 h dark photoperiod, in a controlled chamber (Rumed) as described in Róis et al. (2012).

Biomass, Morphometric, and Physiological Parameters

To understand and compare the evolution of plant vegetative and reproductive growth in the substrata, morphometric and photosynthetic parameters were measured every 2.5 months, one measure in each season of the year. The measurements were taken at transplantation, in winter (T0); in spring (T1), in summer (T2) and at the end of the microcosm assay, in autumn (T3). The number of live and dead leaves, leaf length and width, scape number and length were determined. For leaf size measurements about three leaves/individual were labeled (15 leaves/treatment) at the beginning of the assay (T0). The Photochemical Reflectance Index (PRI) was measured with a PlantPen model PRI 200, which determined the photosynthetic light use efficiency in these leaves. The Normalized Difference Vegetation Index (NDVI) that is an indicator of chlorophyll content in plants and directly related to their photosynthetic capacity was also evaluated using the same leaves, with a PlantPen model NDVI 300. These measures were non-destructive, they were obtained through an optical window in the instruments. At the end of the experiment, the

fresh and dried biomass of both aerial parts and roots were determined in a digital analytical balance (Sartorius ENTRIS124-1S, 120 × 0.0001 g, Sartorius Stedim Biotech North America Inc., New York, United States). The fresh biomass was oven-dried at 60°C, until a constant weight was reached.

Statistical Analysis

The germination and biomass data were analyzed by one-way ANOVA (substrata) and the remaining data by two-way ANOVA (Time (Substrata). When the data were statistically significant ($p < 0.05$), all pairwise-comparisons were carried out by Tukey HSD post hoc test. A Correlation Matrix Principal Component Analysis (PCA using normalised normalized data) was performed and the Pearson correlation coefficient ($r > 0.7$). was used to correlate the nutrient concentrations of substrata in T0 and the morphometric and photosynthetic indexes in T3. The tests were made using the statistical software R studio version 1.1.423 for Windows.

RESULTS AND DISCUSSION

Characterization of Soils, Amendments, and Irrigation Water

The FLU had a silty clay loam texture, was slightly alkaline, with a high EC and low values of C_{org} , N_{total} , extractable P and K (Table 1, Figure 2A, and Supplementary Figure 1A). These data indicate that this soil is saline-sodic with a 25% of $Na_{exchangeable}$ and 41.48 of SAR. The multielemental composition of the Fluvisol showed that concentrations of the potentially hazardous elements were below the maximum allowed values for agricultural use (The Environmental Protection Agency of Portugal [APA], 2019), except for As concentration (15.6 mg/kg), which was slightly higher than the allowed concentration limit

(11 mg/kg; The Environmental Protection Agency of Portugal [APA], 2019). This can be attributed to a widespread industrial waste such as wastes resulting from chemical and metallurgical plants located close to the river Tagus, deposited in the estuary in the past (De Bettencourt, 1988).

The FLU had a high $Na_{exchangeable}$ percentage and SAR contributing to soil aggregate degradation and colloid dispersion, promoting clogging of soil pores and making plants root penetration difficult. This leads to a lack of drainage, water puddle formation and oxygen deficiency (Brady and Weil, 2008). Further, the ratio $[Ca]/[Mg] < 1.5$ indicates very unfavorable conditions for soil physical properties (INIA-LQARS, 2000). For the saline-sodic soil reclamation it is necessary to reduce the $Na_{exchangeable}$ percentage replacing the Na^+ ion to the Ca^{2+} ion, in the exchange complex (Brady and Weil, 2008). While some studies favor gypsum use (Ca_2SO_4) to form Na_2SO_4 that can be leached (Mozheiko, 1969; Carter et al., 1977), other works support the application of organic compounds in addition to the gypsum to improve soil nutrition, aggregation, water-holding capacity, infiltration rate and decrease in EC (Hanay et al., 2004; Diacono and Montemurro, 2015). Tailored soils can be utilized as an alternative to improve salt-affected soil properties (Macía et al., 2014; Cortinhas et al., 2020). Moreover, the cultivation of species well adapted to the saline environments (halophytes) contribute to soil recovery (Manousaki and Kalogerakis, 2011; Devi et al., 2016; Liang et al., 2017). These plants can survive in saline environments due to protection mechanisms such as the Na^+ and Cl^- compartmentation, Ca^{2+} and K^+ uptake, succulence, synthesis of particular compounds and the presence of salt glands (Flowers et al., 2010; Song and Wang, 2014; Caperta et al., 2020).

In this study, the TEC soil fertility was increased due to high content of N_{total} , C_{org} , extractable P and K present in sludge and waste kieselguhr (Table 1). The residual biomass applied as a source of C and other nutrients in the long run by slow mineralization also contributed to improve soil texture/aggregation. The addition of sand and limestone gravel enhanced the texture and permeability of the Fluvisol used in the TEC construction (Supplementary Figures 1B,D) as also shown previously (Macía et al., 2014; Cortinhas et al., 2020). No significant differences among TEC and FLU were found in terms of pH and CE. Nonetheless, the TEC presented a significant increase in the concentration of all determined nutrients, mainly in P ($p = 2.16 \times 10^{-5}$), Fe ($p = 2 \times 10^{-6}$) and Cu ($p = 2 \times 10^{-6}$), except for Mn ($p = 1.07 \times 10^{-7}$) (Figure 2A). The soil improvement in these nutrients is principally related to the wastewater sludge that was rich in P, Fe and Cu (Table 1). The mixture of amendments was suitable to improve soil fertility, texture, and structure (Supplementary Figure 1B) as well as soil permeability and drainage avoiding waterlogging. These latter two soil properties are extremely important since the remaining populations of *L. daveaui* in the Tagus estuary thrive in Fluvisols and saline sands, mainly inundated with salty water by the equinoctial tides (Caperta and Carapeto, 2020).

The brackish water was strongly saline, had a neutral pH and high concentrations of Cl^- (207 mmol/L), HCO_3^-

TABLE 1 | Chemical characteristics of the Fluvisol and amendments (mean ± SD, $n = 3$) collected in Sobralinho, in the Municipal Council of Vila Franca de Xira, west Portugal.

Parameters	Fluvisol	Wastewater sludge	Waste kieselguhr
pH	8.03 (0.23)	8.47 (0.15)	7.17 (1.12)
EC	5.60 (1.28)	3.18 (0.41)	0.77 (0.45)
C_{org}	19.96 (2.07)	223.13 (19.55)	31.33 (16.37)
N_{total}	1.69 (0.17)	42.19 (4.06)	10.95 (4.33)
$Na_{exchangeable}$	6.54 (0.31)	–	–
$K_{exchangeable}$	2.11 (0.07)	–	–
$K_{extractable}$	0.86 (0.03)	1.92 (2.05)	0.13 (0.02)
$Mg_{exchangeable}$	10.27 (0.31)	–	–
$Ca_{exchangeable}$	10.26 (0.26)	–	–
$P_{extractable}$	0.01 (7.40×10^{-4})	1.38 (0.31)	0.24 (0.08)
CEC	26.07 (0.50)	–	–
ESP	25.10 (1.33)	–	–

Electrical conductivity (EC) is given in mS/cm; C_{org} , N_{total} , $K_{extractable}$ and $P_{extractable}$, in g/kg; $Na_{exchangeable}$, $K_{exchangeable}$, $Mg_{exchangeable}$, $Ca_{exchangeable}$ and Cation Exchangeable Capacity (CEC) in cmol_c/kg; ESP – Exchangeable sodium percentage [$((Na_{exchangeable}/CEC) \times 100)$; Ayers and Westcot, 1985].




TABLE 2 | Chemical characteristics of irrigation water (mean \pm SD, $n = 3$) collected in Sobralinho, in the Municipal Council of Vila Franca de Xira, west Portugal.

Chemical characteristics of brackish estuarine water	
pH	7.78 (0.41)
EC	22.03 (7.09)
Cl ⁻	207 (75.05)
HCO ₃ ⁻	4.37 (0.89)
Na ⁺	187 (58.16)
Ca ²⁺	5.24 (1.49)
Mg ²⁺	22.14 (7.59)
K ⁺	4.02 (1.34)
P	2.59×10^{-2} (1.55×10^{-2})
Fe	4.86×10^{-4} (6.38×10^{-4})
Zn	8.58×10^{-5} (4.53×10^{-5})
Mn	1.77×10^{-3} (1.98×10^{-3})
Cu	1.31×10^{-4} (1.20×10^{-4})
SAR	35.43 (5.81)

The electrical conductivity (EC) is given in mS/cm and anions and cations concentrations in mmol/L; SAR, sodium adsorption ratio $[(\text{Na})/((\text{Ca}) + (\text{Mg}))^{1/2}]$; Ayers and Westcot, 1985; Lesch and Suarez, 2009).


(4.37 mmol/L), Na⁺ (187 mmol/L), Ca²⁺ (5.24 mmol/L), and Mg²⁺ (22.14 mmol/L) (Table 2) as was to be expected in estuarine water (Santos et al., 2017; Cortinhas et al., 2020).

The calculated SAR of water was high (35.4) but tolerated by halophyte species (Haro et al., 1993; Flowers and Colmer, 2008; Table 2). The maritime fog mostly observed from November to February each year in the Lisbon area (Belo-Pereira and Santos, 2016), is deposited as occult precipitation by condensing or sublimating directly onto plant surfaces (Karavoltzos et al., 2017), probably contributing to the water balance of plants in these saline habitats.

After plant growth (T3), the soils irrigated with brackish water show no change in pH values, however the EC values of FLU (5.6 dS/m) and TEC (5.9 dS/m) had a significant ($p < 1 \times 10^{-7}$) increase (34.03; 32.48 dS/m, respectively) due to the high salinity of estuarine water. In T3 (Figure 2B), the organic C, N_{total} and extractable P concentrations did not show significant variations compared to T0 (Figure 2A). At the end of the assay (Figure 2B), the Zn content of TEC increased significantly as well as the extractable K in both soils ($p < 5 \times 10^{-6}$). In contrast, the Fe and Cu values decreased significantly ($p < 9 \times 10^{-4}$) from T0 (Figure 2A) to T3 (Figure 2B).

Seed Germination and Plant Growth

The findings of a significantly different ($p = 5.95 \times 10^{-8}$) higher seed germination percentage in distilled water and in FLU compared to TEC (Figure 3) can be due to the effect of high salinity (i.e., lowest osmotic potential) on *Limonium* seed



germination (Boorman, 1967; Luque et al., 2013; Al Hassan et al., 2017) as well as of other halophyte species (Debez et al., 2018; Lombardi et al., 2019).

Due to FLU low permeability, a water film is retained on the surface of the substrate allowing the seeds to float (Supplementary Figure 1C), whereas at the surface of the relatively porous TEC this water film does not occur (Supplementary Figure 1). Although Fluvisols and rocky deposits seem to be suitable for halophyte seed germination, for endemic *Limonium* species seed germination by directly sowing them in the field did not prove to be successful (Caperta and Vasco Silva, unpublished research).

Transplanting and even translocating plants to start new populations of rare and endemic species have shown better results than using seeds *in situ* (Davy, 2002; Godefroid et al., 2011; Albrecht and Maschinski, 2012). A comparison of plants grown in FLU and in TEC irrigated with estuarine water (Figure 4) revealed that there were no significant differences for the growth parameters measured in plants growing in both soils. However, there was a significant increase in the leaf number in T2 ($p \leq 6.51 \times 10^{-4}$) followed by a small decrease in T3 (Figure 4). Plants grown in TEC presented smaller leaves than plants cultivated in FLU. Nonetheless, the former had a higher number of leaves, which can contribute to a higher photosynthetic rate (Figure 4). Between T2 and T3, the plants started to bloom, which may have influenced a reduction in leaf number. The plants

cultivated in TEC had a higher vegetative and reproductive growth and produced flowering stems that emerged earlier than in FLU. Although only one individual per treatment presented a flowering stem, the plant grown in TEC produced more and larger scapes than that in FLU (Figure 4). The TEC substrate, rich in essential nutrients and with better texture and structure for plants allowed better plant development. These findings suggest that plant cultivation in Technosols contributes to the reproductive success of *L. daveau* as also found previously in *Limonium algarvense* (Cortinhas et al., 2020).



In terms of the physiological status there were no significant differences in NDVI and PRI between plants raised in different substrata (Figure 5). However, the plants cultivated in FLU presented lower values in all the reference periods than individuals grown in TEC, supporting that this soil was a more stressful environment for plants than TEC (Figure 5). The NDVI and PRI values of plants raised in both substrata dropped significantly ($p < 2.4 \times 10^{-6}$) after the spring (T1). While the PRI value only reduced significantly from T1 to T2 ($p = 5 \times 10^{-7}$), the NDVI value decreased significantly between T1-T2 ($p < 1 \times 10^{-7}$) and T2-T3 ($p < 1 \times 10^{-7}$). Stress factors such as leaves senescence, salinity, light excess, or high temperature or in combination can lead to PRI and NDVI reduction (Bieto and Talón, 2008; Yudina et al., 2020). In this study, reduction of these indexes at least could be associated to leaves senescence that occurred when plants start flowering.

The dry mass of roots, aerial part and flowers represented ~40% of their weight. The aerial (leaves and scapes) and roots biomass did not show significant differences among substrata (Figure 6B). However, plants from TEC showed high flowers production than FLU (Figure 6B). The roots of plants cultivated in FLU were concentrated in the bottom of pots (Figure 6A), while in plants grown in TEC roots spread uniformly all over the substrate (Figure 6A). The deficient FLU structure, due to colloid dispersion, is an obstacle to roots penetration and oxygen circulation. The poor structure and the compaction due to high $\text{Na}_{\text{exchangeable}}$ percentage and SAR can lead to plant death as found previously in Cortinhas et al. (2020).

Principal Component Analysis (PCA)

The PCA correlated nutrient concentrations determined in substrata in the beginning of the assay (T0) with the morphometric parameters and the photosynthetic indexes measured in plants at the end of the assay (T3). The first two components explain 71.14% of data variation. The third principal component only corresponded to 12.78% of variation and did not show correlation with any of the evaluated variables ($r < 0.60$); as such, it was not represented in this analysis.

The first principal component represented 53.12% of data variation with the pH and nutrients being the variables having the highest correlation ($r > 0.70$) with that component (Figure 7 and Supplementary Table 1). This contributes





to clearly separate the individuals cultivated in TEC from plants cultivated in FLU (Figure 7). The PCA revealed that TEC had higher nutrient concentrations (except for Mn) but lower pH value than FLU. All the nutrients showed correlation ($r > 0.70$) among them (Supplementary Table 2) and the pH was negatively correlated with C, Fe, and Cu ($-0.77 < r < -0.70$), but positively with Mn ($r = 0.78$).

In the second principal component, which represented 18.02% of data variability (Figure 7), leaf and scape number were the parameters that mostly contributed to the observed variability ($r \geq 0.70$; Supplementary Table 1). All the growth parameters and the values of the photosynthetic indexes were also higher in the plants cultivated in the TEC than plants grown in the FLU. The NDVI was strongly correlated ($0.95 < r < 0.96$) with leaf size (Supplementary Table 2).

CONCLUSION

This work contributes to the knowledge of *L. daveaui* microenvironmental requirements particularly soil properties. The findings demonstrated that eco-friendly soil technologies have an enormous potential to improve the reintroduction success of this species. As such, plants will be pre-grown in the Technosol and then will be planted on the sites in modified soil. The amendments that have shown suitable to improve the physical and chemical properties of the saline-sodic Fluvisol will be added and mixed with this soil. The plants will be subjected to irrigation with brackish water collected directly from a channel located in the Tagus estuary. From a circular economy perspective, this approach that uses cost-effective subproduct/wastes contributes to enhance soil fertility and structure, valuing underused resources such as saline soils, not



used in conventional reintroduction schemes, but appropriate for halophyte species.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Materials**, further inquiries can be directed to the corresponding author/s.

AUTHOR CONTRIBUTIONS

MA and ACa designed and coordinated the study. ACo, TF, MA, and ACa performed the experimental assay. ACo and TF processed the raw data and realized the statistical analyses. ACo, MA, and ACa drafted the manuscript. All authors read and approved the manuscript.

FUNDING

This research was funded by Fundação para a Ciência e a Tecnologia (FCT, www.fct.pt/) by project PTDC/AGRPRO/4285/2014 and grant SFRH/BD/130256/2017 attributed to ACo. Publication costs were funded by FCT through project UID/AGR/04129/2020 (LEAF). The Open Access was funded by LEAF Project UID/AGR/04129/2020.

ACKNOWLEDGMENTS

We want to thank Arq^{td} Vitória Cochicho (Municipal Council Vila Franca de Xira, Lisbon, Portugal; <https://www.cm-vfxira>).

REFERENCES

- Activation Laboratories (2020). *Metals in Soil*. Available at: <https://actlabs.com/environmental/services/metals-in-soil/> (accessed June 1, 2020).
- Al Hassan, M., Estrelles, E., Soriano, P., López-Gresa, M. P., Bellés, J. M., Boscaiu, M., et al. (2017). Unraveling salt tolerance mechanisms in halophytes: a comparative study on four Mediterranean *Limonium* species with different geographic distribution patterns. *Front. Plant Sci.* 8:1438. doi: 10.3389/fpls.2017.01438
- Albrecht, M. A., and Maschinski, J. (2012). "Influence of founder population size, propagule stages, and life history on the survival of reintroduced plant populations," in *Plant Reintroduction in a Changing Climate*, eds J. Maschinski and K. E. Haskins (Washington, DC: Island Press), 171–188. doi: 10.5822/978-1-61091-183-2_10
- Asensi, A., Diez Garretas, B., and van der Maarel, E. (1993). "Dry coastal ecosystems of Portugal," in *Ecosystems of the World, 2A: Dry Coastal Ecosystems: Polar Regions and Europe*, ed. E. van der Maarel (Amsterdam: Elsevier), 342–347.
- Ayers, R. S., and Westcot, D. W. (1985). *Water Quality for Agriculture*. Rome: Food and Agriculture Organization of the United Nations. FAO Irrigation and Drainage Paper, 29 Rev. I.
- Balke, T., Bouma, T. J., Horstman, E. M., Webb, E. L., Erfteimeijer, P. L., and Herman, P. M. (2011). Windows of opportunity: thresholds to mangrove seedling establishment on tidal flats. *Mar. Ecol. Prog. Ser.* 440, 1–9. doi: 10.3354/meps09364
- Baumberger, T., Affre, L., Croze, T., and Mesléard, F. (2012). Habitat requirements and population structure of the rare endangered *Limonium girardianum* in Mediterranean salt marshes. *Flora* 207, 283–293. doi: 10.1016/j.flora.2011.11.008
- Belo-Pereira, M., and Santos, J. A. (2016). A persistent wintertime fog episode at Lisbon airport (Portugal): performance of ECMWF and AROME models. *Meteorol. Appl.* 23, 353–370. doi: 10.1002/met.1560
- Bieto, J. A., and Talón, M. (2008). *Fundamentos de Fisiología Vegetal*. Madrid: McGraw-Hill Interamericana.
- Boorman, L. A. (1967). *Limonium vulgare* Miller and *L. humile* Miller. *J. Ecol.* 55, 221–232. doi: 10.2307/2257727
- Brady, N. C., and Weil, R. R. (2008). *The Nature and Properties of Soils*. New Jersey: Pearson Prentice Hall.
- Braun-Blanquet, J., Rozeira, A., and Pinto da Silva, A. R. (1972). Résultats de deux excursions géobotaniques à travers le Portugal septentrional et moyen. IV. Esquisse sur la végétation dunale. *Agron. Lusit.* 33, 217–234.
- Breure, A. M., Lijzen, J. P. A., and Maring, L. (2018). Soil and land management in a circular economy. *Sci. Total Environ.* 624, 1125–1130. doi: 10.1016/j.scitotenv.2017.12.137
- Cao, H., Zhu, Z., Balke, T., Zhang, L., and Bouma, T. J. (2018). Effects of sediment disturbance regimes on *Spartina* seedling establishment: implications for salt marsh creation and restoration. *Limnol. Oceanogr.* 63, 647–659. doi: 10.1002/lno.10657
- Caperta, A., and Carapeto, A. (2020). "Limonium daveaui: ficha de avaliação do risco de extinção," in *Lista Vermelha da Flora Vasculosa de Portugal Continental*, (Lisboa: Sociedade Portuguesa de Botânica Associação Portuguesa de Ciência da Vegetação).
- Caperta, A. D., Espírito-Santo, M. D., Silva, V., Ferreira, A., Paes, A. P., Róis, A. S., et al. (2014). Habitat specificity of a threatened and endemic,

pt/) for providing permission for Fluvisol and estuarine water collection, and to Dra. Sandra Borges (Sociedade Central de Cervejas e Bebidas, S.A., <http://www.centralcervejas.pt/>) for providing the sludge and waste kieselguhr.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fevo.2021.604509/full#supplementary-material>

Supplementary Figure 1 | Physical appearance of Fluvisol (A) and Technosol (B). The Fluvisol was collected in Sobralinho salt marsh area in the Municipal Council of Vila Franca de Xira, west Portugal. The Technosol was produced with Fluvisol and organic/inorganic wastes' as amendments. Seed germination (white arrows) on a water film retained in the Fluvisol surface (C) and in the Technosol surface, where the water film was not present (D).

Supplementary Table 1 | Correlation matrix between the first two principal components (PC1 and PC2) on the normalized data and the original variables. This matrix includes the pH, CE, organic carbon (C) and nutrient concentration (N total (N), extractable P (P), extractable K (K), Fe, Mn, Zn, Cu) values determined in the substrata in the beginning of the assay (T0) and morphometric parameters (leaf number, leaf length, leaf width, scape number, scape length) and photosynthetic indexes (Normalised Difference Vegetation Index (NDVI) and Photochemical Reflectance Index (PRI)) obtained in plants at the end of the assay (T3).

Supplementary Table 2 | Correlation matrix between variables. This matrix includes the pH, CE, organic carbon (C) and nutrient concentration (N total (N), extractable P (P), extractable K (K), Fe, Mn, Zn, Cu) values determined in the substrata in the beginning of the assay (T0) and morphometric parameters (leaf number, leaf length, leaf width, scape number, scape length) and photosynthetic indexes (Normalised Difference Vegetation Index (NDVI) and Photochemical Reflectance Index (PRI)) obtained in plants at the end of the assay (T3).

- cliff-dwelling halophyte. *AoB Plants* 6:lu032. doi: 10.1093/aobpla/plu032
- Caperta, A. D., Róis, A. S., Teixeira, G., Garcia-Caparros, P., and Flowers, T. J. (2020). Secretory structures in plants: lessons from the *Plumbaginaceae* on their origin, evolution and roles in stress tolerance. *Plant. Cell. Environ.* 43, 2912–2931. doi: 10.1111/pce.13825
- Carter, M. R., Webster, G., and Cairns, R. (1977). Amelioration of a brown solodized solonchets soil by a surface application of gypsum plus ammonium nitrate. *Can. J. Soil. Sci.* 57, 139–145. doi: 10.4141/cjss77-018
- Cesário, R., Monteiro, C. E., Nogueira, M., O'Driscoll, N. J., Caetano, M., Hintelmann, H., et al. (2016). Mercury and methylmercury dynamics in sediments on a protected area of Tagus Estuary (Portugal). *Water Air Soil Pollut.* 227:475. doi: 10.1007/s11270-016-3179-2
- Cortinhas, A., Caperta, A. D., Teixeira, G., Carvalho, L., and Abreu, M. M. (2020). Harnessing sediments of coastal aquaculture ponds through technosols construction for halophyte cultivation using saline water irrigation. *J. Environ. Manag.* 261:109907. doi: 10.1016/j.jenvman.2019.109907
- Costa, J. (2013). *Urbanismo e Adaptação às Alterações Climáticas: As Frentes de Água*. Lisboa: Livros Horizonte.
- Costa, J. C., Monteiro-Henriques, T., Neto, C., Arsénio, P., and Aguiar, C. (2007). The application of the habitats directive in Portugal. *Fitosociologia* 44, 23–28.
- Costa, J. C., Neto, C., Aguiar, C., Capelo, J., Espírito-Santo, M. D., Honrado, J., et al. (2012). Vascular plant communities in Portugal (Continental, the Azores and Madeira). *Glob. Geobot.* 2, 1–180.
- Davy, A. (2002). "Establishment and manipulation of plant populations and communities in terrestrial systems," in *Handbook of Ecological Restoration. Principles of Restoration*, eds M. R. Perrow and A. J. Davy (Cambridge: Cambridge University Press), 223–241.
- De Bettencourt, A. M. M. (1988). On arsenic speciation in the Tagus estuary. *Neth. J. Sea Res.* 22, 205–212. doi: 10.1016/0077-7579(88)90024-5
- Debez, A., Belghith, I., Pich, A., Taamalli, W., Abdely, C., and Braun, H. P. (2018). High salinity impacts germination of the halophyte *Cakile maritima* but primes seeds for rapid germination upon stress release. *Physiol. Plant.* 164, 134–144. doi: 10.1111/ppl.12679
- Devi, S., Nandwal, A. S., Angrish, R., Arya, S. S., Kumar, N., and Sharma, S. K. (2016). Phytoremediation potential of some halophytic species for soil salinity. *Int. J. Phytoremediation* 18, 693–696. doi: 10.1080/15226514.2015.1131229
- Diacono, M., and Montemurro, F. (2015). Effectiveness of organic wastes as fertilizers and amendments in salt-affected soils. *Agriculture* 5, 221–230. doi: 10.3390/agriculture5020221
- Duarte, B., Silva, G., Costa, J. L., Medeiros, J. P., Azeda, C., Sá, E., et al. (2014). Heavy metal distribution and partitioning in the vicinity of the discharge areas of Lisbon drainage basins (Tagus Estuary, Portugal). *J. Sea Res.* 93, 101–111. doi: 10.1016/j.seares.2014.01.003
- Duarte, I., Rego, F. C., Casquilho, J. P., and Arsénio, P. (2016). A relevance index for the habitat areas of Natura 2000 network based on their rarity and representativeness. *Ecol. Indic.* 61, 202–213. doi: 10.1016/j.ecolind.2015.09.015
- Dunwiddie, P. W., and Martin, R. A. (2016). Microsites matter: improving the success of rare species reintroductions. *PLoS One* 11:e0150417. doi: 10.1371/journal.pone.0150417
- Erben, M. (1978). Die Gattung *Limonium* im südwestmediterranen Raum. *Mitt. Bot. Staatssamml. München.* 14, 361–626.
- Erben, M. (1993). "*Limonium* Mill," in *Flora Iberica*, eds S. Castroviejo, C. Aedo, S. Cirujano, M. Lainz, P. Montserrat, R. Morales, et al. (Madrid: Real Jardín Botánico CSIC), 2–143.
- Espírito-Santo, M. D., Alves, H. N., Caperta, A. D., and Moreira, I. (2012). "Plantas endêmicas do litoral de Portugal Continental in Monteiro," in *Gestão e Conservação da Flora e da Vegetação de Portugal e da África Lusófona*, eds A. Gomes da Silva and R. Jorge (Lisboa: ISA Press), 267–302.
- Fernandes, A., Figueira de Sousa, J., Costa, J. P., and Neves, B. (2020). Mapping stakeholder perception on the challenges of brownfield sites' redevelopment in waterfronts: the Tagus Estuary. *Eur. Plan. Stud.* 28, 2247–2464. doi: 10.1080/09654313.2020.1722985
- Figueres, G., Martin, J. M., Meybeck, M., and Seyler, P. (1985). A comparative study of mercury contamination in the Tagus Estuary (Portugal) and major French Estuaries (Gironde, Loire, Rhône). *Estuar. Coast. Shelf Sci.* 20, 183–203. doi: 10.1016/0272-7714(85)90037-X
- Flowers, T. J., and Colmer, T. D. (2008). Salinity tolerance in halophytes. *New Phytol.* 179, 945–963. doi: 10.1111/j.1469-8137.2008.02531.x
- Flowers, T. J., Galal, H. K., and Bromham, L. (2010). Evolution of halophytes: multiple origins of salt tolerance in land plants. *Funct. Plant Biol.* 37, 604–612. doi: 10.1071/FP09269
- Franco, J. D. A. (1984). *Nova Flora de Portugal (Continente e Açores) Vol II Clethraceae – Compositae*. Lisboa: Author's Edition.
- Godefroid, S., Piazza, C., Rossi, G., Buord, S., Stevens, A. D., Aguruijua, R., et al. (2011). How successful are plant species reintroductions? *Biol. Conserv.* 144, 672–682. doi: 10.1016/j.biocon.2010.10.003
- Guerreiro, M., Fortunato, A. B., Freire, P., Rilo, A., Taborada, R., Freitas, M. C., et al. (2015). Evolution of the hydrodynamics of the Tagus estuary (Portugal) in the 21st century. *J. Integr. Coast. Zone Manag.* 15, 65–80. doi: 10.5894/rgci515
- Hanay, A., Büyüksönmez, F., Kiziloglu, F. M., and Canbolat, M. Y. (2004). Reclamation of saline-sodic soils with gypsum and MSW compost. *Compost Sci. Util.* 12, 175–179. doi: 10.1080/1065657X.2004.10702177
- Haro, R., Bañuelos, M. A., Quintero, F. J., Rubio, F., and Rodríguez-Navarro, A. (1993). Genetic basis of sodium exclusion and sodium tolerance in yeast. A model for plants. *Physiol. Plant* 89, 868–874. doi: 10.1111/j.1399-3054.1993.tb05298.x
- Haskins, K. E., and Pence, V. (2012). *Transitioning Plants to New Environments: Beneficial Applications of Soil Microbes in Plant Reintroduction in a Changing Climate*. Washington, DC: Island Press.
- Holl, K. D., and Hayes, G. F. (2006). Challenges to introducing and managing disturbance regimes for *Holocarpha macradenia*, an endangered annual grassland forb. *Biol. Conserv.* 20, 1121–1131. doi: 10.1111/j.1523-1739.2006.00416.x
- INIA-LQARS (2000). *Manual de Fertilização das Culturas*. Lisboa: Laboratório Químico Agrícola Rebelo da Silva.
- Instituto da Conservação da Natureza e das Florestas [ICNF]. (2012). *Reserva Natural do Estuário do Tejo*. Available at: <http://www2.icnf.pt/portal/ap/r-nat/rnet> (accessed May 25, 2020).
- IUSS Working Group (2015). *World Reference Base for Soil Resources 2014 (update 2015) International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*. Rome: FAO. World Soil Resources Report No. 106.
- Karavoltsov, S., Fotiadis, F., Michopoulos, P., Sakellari, A., Plavšić, M., Bourletsikas, A., et al. (2017). Organic complexation of copper in throughfall and open field bulk deposition: influence of the tree canopy of Mediterranean forest ecosystems. *Chemosphere* 167, 28–35. doi: 10.1016/j.chemosphere.2016.09.131
- Lakanen, E., and Erviö, R. (1971). A comparison of eight extractants for the determination of plant available micronutrients in soils. *Acta Agr. Fenn.* 123, 223–232.
- Lechuga-Lago, Y., Novoa, A., Le Roux, J. J., and González, L. (2017). Understanding the influence of urbanization on invasibility: *Carpobrotus edulis* as an exemplar. *Biol. Invasions.* 19, 3601–3611. doi: 10.1007/s10530-017-1593-z
- Lesch, S. M., and Suarez, D. L. (2009). A short note on calculating the adjusted SAR Index. *Am. Soc. Agric. Biol. Eng.* 52, 493–496. doi: 10.13031/2013.26842
- Liang, L., Liu, W., Sun, Y., Huo, X., Li, S., and Zhou, Q. (2017). Phytoremediation of heavy metal contaminated saline soils using halophytes: current progress and future perspectives. *Environ. Rev.* 25, 269–281. doi: 10.1139/er-2016-0063
- Lombardi, T., Bedini, S., and Bertacchi, A. (2019). Germination ecology of the aromatic halophyte *Artemisia caerulescens* L.: influence of abiotic factors and seed after-ripening time. *Folia Geobot.* 54, 115–124. doi: 10.1007/s12224-019-09345-4
- Luque, E. G., Fernández, I. C. D., and Mercado, F. G. (2013). Effect of salinity and temperature on seed germination in *Limonium cossonianum*. *Botany* 91, 12–16. doi: 10.1139/cjb-2012-0157
- Macía, P., Fernández-Costas, C., Rodríguez, E., Sieiro, P., Pazos, M., and Sanromán, M. A. (2014). Technosols as a novel valorisation strategy for an ecological management of dredged marine sediments. *Ecol. Eng.* 67, 182–189. doi: 10.1016/j.ecoleng.2014.03.020
- Manousaki, E., and Kalogerakis, N. (2011). Halophytes present new opportunities in phytoremediation of heavy metals and saline soils. *Ind. Eng. Chem. Res.* 50, 656–660. doi: 10.1021/ie100270x
- Martins, M. C., Neto, C. S., and Costa, J. C. (2013). The meaning of mainland Portugal beaches and dunes' psammophilic plant communities: a contribution to tourism management and nature conservation. *J. Coast Conserv.* 17, 279–299. doi: 10.1007/s11852-013-0232-9
- Maschinski, J., and Albrecht, M. A. (2017). Center for plant conservation's best practice guidelines for the reintroduction of rare plants. *Plant Divers.* 39, 390–395. doi: 10.1016/j.pld.2017.09.006

- Maunder, M. (1992). Plant reintroductions: an overview. *Biodivers. Conserv.* 1, 51–61. doi: 10.1007/BF00700250
- Menges, E. S., Smith, S. A., and Weekley, C. W. (2016). Adaptive introductions: how multiple experiments and comparisons to wild populations provide insights into requirements for long-term introduction success of an endangered shrub. *Plant Divers.* 38, 238–246. doi: 10.1016/j.pld.2016.09.004
- Mohr, A. C. (1945). *U.S. Patent No. 2*, Vol. 391. Washington, DC: Patent and Trademark Office, 328.
- Mozheiko, A. M. (1969). Chemical Reclamation of Sodic Solonchets in the Southern Part of the Middle Dnieper Region by the Application of Gypsum and Calcium Chloride. *Agrokémia és talajtan* 18, 310–314.
- Olsson, L., Barbosa, H., Bhadwal, S., Cowie, A., Delusca, K., Flores-Renteria, D., et al. (2019). “Land Degradation,” in *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*, eds P. R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, et al. (Geneva: IPCC).
- Pignatti, S. (1971). “Studi sui *Limonium* VIII,” in *Notulae Systematicae ad Floram Europaeam Spectantes*, Vol. 64, ed. V. H. Flora Europaea (Heywood: Botanical Journal of the Linnean Society), 353–381.
- Póvoas, I., and Barral, M. F. (1992). *Métodos de Análise de Solos. Comunicações do Instituto de Investigação Científica Tropical*. Lisbon: Instituto de Investigação Científica Tropical. Série de Ciências Agrárias N.º 10.
- Róis, A. S., Teixeira, G., Sharbel, T. F., Fuchs, J., Martins, S., Espírito-Santo, D., et al. (2012). Male fertility versus sterility, cytotype, and DNA quantitative variation in seed production in diploid and tetraploid sea lavenders (*Limonium* sp., *Plumbaginaceae*) reveal diversity in reproduction modes. *Plant Reprod.* 25, 305–318. doi: 10.1007/s00497-012-0199-y
- Santos, E. S., Abreu, M. M., Peres, S., Magalhães, M. C. F., Leitão, S., Pereira, A. S., et al. (2017). Potential of *Tamarix africana* and other halophyte species for phytostabilisation of contaminated salt marsh soils. *J. Soils Sediments* 17, 1459–1473. doi: 10.1007/s11368-015-1333-x
- Song, J., and Wang, B. (2014). Using euhalophytes to understand salt tolerance and to develop saline agriculture: *Suaeda salsa* as a promising model. *Ann. Bot.* 115, 541–553. doi: 10.1093/aob/mcu194
- Spencer, K. L., Carr, S. J., Diggins, L. M., Tempest, J. A., Morris, M. A., and Harvey, G. L. (2017). The impact of pre-restoration land-use and disturbance on sediment structure, hydrology and the sediment geochemical environment in restored saltmarshes. *Sci. Total Environ.* 587, 47–58. doi: 10.1016/j.scitotenv.2016.11.032
- The Environmental Protection Agency of Portugal [APA] (2019). *Solos Contaminados – Valores de Referência Para o Solo*. Amadora: The Ministry of Agriculture, the Sea.
- The Environmental Protection Agency of Portugal [APA] (2011). *Guide for the Assessment of Imminent Threats and Environmental Damages*. Lisboa: The Ministry of Agriculture, the Sea.
- van der Maarel, E., and van der Maarel-Versluys, M. (1996). Distribution and conservation status of littoral vascular plant species along the European coasts. *J. Coast Conserv.* 2, 73–92. doi: 10.1007/bf02743039
- van Regeteren, M., Meesters, E. H., Baptist, M. J., De Groot, A. V., Bouma, T. J., and Elschot, K. (2020). Multiple environmental variables affect germination and mortality of an annual salt marsh pioneer: *Salicornia procumbens*. *Estuar. Coast.* 43, 1489–1501.
- Wang, N., Jiao, J. Y., Lei, D., Chen, Y., and Wang, D. L. (2014). Effect of rainfall erosion: seedling damage and establishment problems. *Land Degrad. Dev.* 25, 565–572. doi: 10.1002/ldr.2183
- Yudina, L., Sukhova, E., Gromova, E., Nerush, V., Vodeneev, V., and Sukhov, V. (2020). A light-induced decrease in the photochemical reflectance index (PRI) can be used to estimate the energy-dependent component of non-photochemical quenching under heat stress and soil drought in pea, wheat, and pumpkin. *Photosynth. Res.* 146, 175–187.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Cortinhas, Ferreira, Abreu and Caperta. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.