

# Exosomal lincRNA-p21 levels may help to distinguish prostate cancer from benign disease

Mustafa Işın<sup>1</sup>, Ege Uysaler<sup>1</sup>, Emre Özgür<sup>1</sup>, Hikmet Köseoğlu<sup>2</sup>, Öner Şanlı<sup>3</sup>, Ömer B. Yücel<sup>2</sup>, Uğur Gezer<sup>1</sup> and Nejat Dalay<sup>1\*</sup>

<sup>1</sup> Department of Basic Oncology, Oncology Institute, Istanbul University, Istanbul, Turkey, <sup>2</sup> Department of Urology, School of Medicine, Istanbul Hospital, Başkent University, Istanbul, Turkey, <sup>3</sup> Department of Urology, Istanbul Medical Faculty, Istanbul University, Istanbul, Turkey

## OPEN ACCESS

### Edited by:

Mohammad Faghihi,  
University of Miami Miller School  
of Medicine, USA

### Reviewed by:

Georges St. Laurent,  
St. Laurent Institute, USA  
Gregory C. Sartor,  
University of Miami Miller School  
of Medicine, USA  
Mohammadreza Hajjari,  
Shahid Chamran University of Ahvaz,  
Iran

### \*Correspondence:

Nejat Dalay,  
Department of Basic Oncology,  
Oncology Institute, Istanbul University,  
Millet Cad., Istanbul 34093, Turkey  
ndalay@yahoo.com

### Specialty section:

This article was submitted to  
RNA,  
a section of the journal  
Frontiers in Genetics

Received: 28 February 2015

Paper pending published:

24 March 2015

Accepted: 14 April 2015

Published: 06 May 2015

### Citation:

Işın M, Uysaler E, Özgür E,  
Köseoğlu H, Şanlı Ö, Yücel ÖB,  
Gezer U and Dalay N (2015)  
Exosomal lincRNA-p21 levels may  
help to distinguish prostate cancer  
from benign disease.  
Front. Genet. 6:168.  
doi: 10.3389/fgene.2015.00168

Exosomes are membranous vesicles containing various biomolecules including lincRNAs which are involved in cellular communication and are secreted from many cells including cancer cells. In our study, we investigated the exosomal GAS5 and lincRNA-p21 lincRNA levels in urine samples from 30 patients with prostate cancer (PCa) and 49 patients with benign prostatic hyperplasia. Quantification of lincRNA molecules was performed by real-time PCR. We observed a significant difference in the exosomal lincRNA-p21 levels between PCa and BPH patients whereas the GAS5 levels did not reveal a difference. Our data suggest that the discriminative potential of exosomal lincRNA-p21 levels may help to improve the diagnostic prediction of the malignant state for patients with PCa.

**Keywords:** prostate cancer, exosome, lincRNA, benign prostatic hyperplasia, non-invasive diagnosis

## Introduction

Prostate cancer (PCa) is the second most common malignancy and accounts for 15% of the cancers in men. It is the fifth leading cause of death and almost 70% of the cases occur in the developed countries (Ferlay et al., 2015). The diagnosis of PCa is performed via histopathological evaluation of biopsy samples which has several known disadvantages like bleeding and infection (Loeb et al., 2013). Although use of the prostate specific antigen (PSA) as a diagnostic marker has improved the detection and management of PCa (Lippi et al., 2009) its low specificity and lack of other predictive parameters for the progression of the disease makes the stratification of the patients with high risk or indolent PCa difficult (Roddam et al., 2005; Nogueira et al., 2010).

Recently, a prostate specific lincRNA, the PCa Gene 3 (PCA3), has been approved as an additional test to determine the need for biopsies in PCa. Unfortunately, negative PCA3 results in indolent cancer carriers and high grade prostatic intraepithelial neoplasia (HGPIN; Morote et al., 2010; Auprich et al., 2011a) renders the biomarker insufficient. The level of TMPRSS2-ERG fusion transcripts, in cells collected from urine after digital examination, have also been evaluated together with the PCA3 levels for a more accurate diagnosis (Leyten et al., 2014). However, none of these approaches can satisfactorily distinguish high-risk from indolent cancer. Therefore identification of new biomarkers to correctly identify patients needing more aggressive treatment would help to prevent individuals with localized tumors from getting unnecessary biopsies and from the side effects of overtreatment.

Exosomes are small membranous vesicles originating from the endosomal compartment which function as messengers in intercellular communication (Simpson et al., 2008; Bang and Thum, 2012; Gezer et al., 2014). They are secreted and released by cells and bind to the receptors on recipient cells, thereby transferring the signal. Exosome secretion is an evolutionary conserved cellular

mechanism dating back to Archaea but their role as cellular messengers has been described only recently (Taylor et al., 2006; Deatherage and Cookson, 2012). Exosomes contain proteins and various types of RNA molecules including lncRNAs. Non-exonic transcripts as a whole constitute the majority of non-ribosomal RNA molecules in the cell and lncRNAs constitute a significant portion of this fraction (Kapranov et al., 2010). Exosomes secreted from prostate can be detected in semen (Ronquist and Brody, 1985) and urine (Pisitkun et al., 2004). In recent years it has been shown that PCA3 and several other microRNAs are present in exosomes isolated from PCa patients (Dijkstra et al., 2014; Huang et al., 2015).

LincRNA-p21 and GAS5 lncRNA act as tumor suppressor molecules in the cellular machinery (Schneider et al., 1988; Huarte et al., 2010). Expression of lincRNA-p21 is stimulated by the p53 tumor suppressor protein and upon transcription it suppresses expression of the genes transcriptionally regulated by p53 by binding to the hnRNP-K complex (Huarte et al., 2010). GAS5 plays a role in the induction of apoptosis. It suppresses several antiapoptotic genes by binding to the glucocorticoid receptor (GR) and hence prevents GR from binding to the glucocorticoid response elements on the target DNA molecule (Kino et al., 2010). Recent data indicate that lncRNA molecules may exhibit tissue- and disease-specific expression which can provide important potential biomarkers specific to the particular cancer types (Yaman Agaoglu et al., 2011; Gezer et al., 2014). However, it should be noted that lncRNA molecules are associated both with cancer and pluripotency which can be a confounding factor (St. Laurent et al., 2013). In this study, we aimed to evaluate the diagnostic utility of exosomal lincRNA-p21 and GAS5 levels in individuals with benign prostatic hyperplasia (BPH) and PCa.

## Materials and Methods

### Exosomal RNA Isolation

Thirty patients with PCa (median age  $64 \pm 6$ ) and 49 patients with BPH (median age  $68 \pm 9$ ) were enrolled in the study. The study was approved by the Ethics Committee of the Istanbul Faculty of Medicine and informed consent was obtained from the participants. All patients were Caucasian and the disease scores of the patients are given in **Table 1**. Urine samples from patients were collected after digital rectal examination and were centrifuged for 10 min at 1000 rpm in a conical tube to remove cell and debris. Supernatant was transferred to another conical tube and centrifuged at 2000 rpm for 10 min to remove remaining debris and bacteria. 10 ml of cell-free urine was then transferred to a new 15 ml tube and stored at  $-80^{\circ}\text{C}$  until use. Exosomal RNA was extracted according to the manufacturer's protocol using the "Urine Exosome RNA Isolation Kit" (Norgen Biotek, Thorold, ON, Canada).

### Quantification of lncRNAs

Exosomal RNA isolated from urine samples were used for cDNA synthesis using the First Strand cDNA Synthesis kit (Thermo Scientific, West Palm Beach, FL, USA) according to the manufacturers' instructions. The real-time amplification of lncRNA molecules was performed using the LightCycler 480 system

**TABLE 1 | Disease scores of the patients.**

Benign prostatic hyperplasia patients		<i>n</i> = 49
IPSS Score (Barry et al., 1992)		
0–7		0
8–19		24
20–35		17
Unknown		8
Prostate cancer patients		<i>n</i> = 30
Gleason score (Epstein, 2010)		
6 (3+3)		19
7 (3+4)		3
7 (4+3)		4
Unknown		4

**TABLE 2 | The primer sequences used in the study.**

Gene	Primer sequence	Reference
LincRNA-p21	F 5' - GGGTGGCTCACTCTTCTGGC -3'	Huarte et al. (2010)
	R 5' - TGGCCTTGCCCGGGCTTGTC -3'	
GAS5	F 5' - CTTCTGGGCTCAAGTGATCCT -3'	Khalil et al. (2009)
	R 5' - TTGTGCCATGAGACTCCATCAG -3'	
GAPDH	F 5' - GCTCTGCTCCTCCTGTTTC -3'	Khalil et al. (2009)
	R 5' - ACGACCAAATCCGTTGACTC -3'	

(Roche, Germany). SYBR Green (Roche) was used as the fluorescent molecule. The primer sequences are shown in **Table 2**.

The PCR reaction included an initial "hot start" for 10 min., followed by 45 cycles of amplification. Each cycle consisted of a denaturation step at  $95^{\circ}\text{C}$  for 10 s, annealing starting at  $60^{\circ}\text{C}$  for 20 s and decreasing by  $2^{\circ}\text{C}$  every two cycles down to  $55^{\circ}\text{C}$ , and amplification at  $72^{\circ}\text{C}$  for 30 s. For quantification of lncRNAs, the  $\Delta\Delta\text{Ct}$  method was used. The GAPDH gene was used as the reference. All experiments were performed twice and the mean values were calculated.


### Statistical Analyses

SPSS® ver.21.0 statistical program was used for statistical analysis. Kruskal–Wallis or Mann–Whitney U tests were used when appropriate to compare the parameters (lncRNA and PSA levels). A *p*-value  $\leq 0.05$  was considered statistically significant.

## Results

Exosomal levels of GAS5 and lincRNA-p21 lncRNAs were evaluated in the urine samples of 49 patients diagnosed with BPH and 30 patients with PCa. The distribution of exosomal lncRNA levels are shown in **Figure 1**. The lincRNA-p21 levels were significantly higher in PCa than in BPH (median; 0.163 vs 0.071; *p* = 0.016, AUC: 0.663). Exosomal GAS5 levels were found to be similar in the two disease groups (median; 1.197 vs 1.235 and *p* = 0.127). The data are shown in **Table 3**.

The PSA levels were higher in the PCa group than in the patients with BPH (*p* < 0.001, median values; 7.7 and 2.16 respectively). In the BPH group no correlation was observed between the



**TABLE 3 | Median values of exosomal lncRNAs and their statistical significances.**

Median	BPH	PCa	Significance (p-value)
GAS5	1.197	1.235	0.127
LincRNA-p21	0.071	0.163	0.016


AUC value of LincRNA-p21: 0.663, CI:95%

IPSS (International Prostate Symptom Score) score and GAS5 or lincRNA-p21 levels. Likewise, no correlation between the clinical stage (Gleason score) and exosomal lncRNA levels was observed in the PCa group. There was no correlation between the PSA levels and the IPSS score in patients with BPH but a correlation was observed between the PSA values and the Gleason score in the PCa group ( $p = 0.123$  and  $0.049$ , respectively).

The sensitivity and specificity of lincRNA-p21 and lincRNA-p21 in combination with PSA were calculated using a cut off value of 2.5 ng/ml for PSA and 0.181 for exosomal lincRNA-p21 expression (Figure 2). The specificity for predicting PCa increased from 63 to 94% when the two parameters were combined while the specificity did not change (Table 4).

## Discussion

Our study is the first report, revealing presence of lincRNA-p21 and GAS5 lncRNA molecules in the exosomes derived from urine samples. Circulating GAS5 and lincRNA-p21 have been previously detected in B-cell malignancies (İşin et al., 2014) but exosomal GAS5 and lincRNA-p21 molecules have previously been only reported in exosomes secreted from HeLa and MCF-7 cell lines but not in human tumors (Gezer et al., 2014). Cellular lincRNA-p21 expression has been suggested to affect global gene expression in different cancers by modulating mRNA translation and suppressing the p53 and Wnt/ $\beta$ -catenin signaling pathways (Dimitrova et al., 2014; Wang et al., 2014). Cellular GAS5



**TABLE 4 | Sensitivity and specificity of exosomal lincRNA-p21 and of the lincRNA-p21/PSA combination.**

	LincRNA-p21	LincRNA-p21 and PSA
Sensitivity	67%	52%
Specificity	63%	94%

expression suggest a role of GAS5 in the regulation of apoptosis in breast cancer cell lines and tumors (Pickard and Williams, 2014) and an inverse association with the mTOR expression in PCa cell lines (Yacqub-Usman et al., 2015). In the present study, we observed significantly higher levels of exosomal lincRNA-p21 levels in the patients with PCa.

It has been shown that one in six of the prostatectomy specimens may contain indolent cancers which usually do not progress to clinical detection during the lifetime of the patient (Epstein et al., 1994). On the other hand, several studies suggest that more than 50% of the cancers which are initially diagnosed as localized tumors are actually advanced at the time of treatment (Chun et al., 2008; Jeldres et al., 2008).

The PSA test is not sensitive enough to predict the presence, extent and risk of recurrence of PCa (Albertsen, 2010; Zeliadt et al., 2010; Friedrich, 2011). Therefore, there is a definite need for non-invasive diagnostic biomarkers which can distinguish the low- and high-risk patients in the clinical decision-making. Even though non-invasive detection of PCA3 levels in urine after digital rectal examination may provide some useful information on the need of a repeated biopsy, evaluation of PCA3 also fails to be specific (Bradley et al., 2013). Dijkstra et al. (2014) have shown that the diagnostic performance of PCA3 in exosomes was found to be more successful than in urine where the PCA3 score was normalized with PSA mRNA in order to achieve a higher performance. Although their study group is quite small the authors suggested an advantage of exosomal PCA3 evaluation

while indicating that their data should be validated in larger clinical cohorts. The discriminative capacity achieved by the exosomal PCA3 in this study (AUC: 0.524) is lower than lincRNA-p21 (AUC: 0.663, CI: 95%) in the present study. These data indicate that analysis of exosomal lincRNA-p21 in urine performs better than PCA3 in detecting PCa.

Recently, there is an active controversy over decreasing the cut-off level of PCA3 from 35 to 25 (Nakanishi et al., 2008; Auprich et al., 2011b; de la Taille et al., 2011; Ploussard et al., 2011) which is expected to increase its diagnostic sensitivity. A meta-analysis by Luo et al. (2014) reported that the sensitivity of PCA3 for detecting PCa ranges from 46.9 to 82.3% and specificity from 55 to 92%. In our study the specificity of lincRNA-p21 for PCa was 94% when combined with PSA.

Expression of TMPRSS2-ERG has also been analyzed in urine samples (Leyten et al., 2014). It has been reported that combination of PCA3 and TMPRSS2-ERG expression increased the sensitivity of detecting PCa. However, the biomarker pair still failed to detect indolent tumors with a Gleason score of  $\geq 7$ .

## References

- Albertsen, P. C. (2010). Efficacy vs effectiveness in prostate-specific antigen screening. *J. Natl. Cancer Inst.* 102, 288–289. doi: 10.1093/jnci/djq002
- Auprich, M., Bjartell, A., Chun, F. K., de la Taille, A., Freedland, S. J., Haese, A., et al. (2011a). Contemporary role of prostate cancer antigen 3 in the management of prostate cancer. *Eur. Urol.* 60, 1045–1054. doi: 10.1016/j.eururo.2011.08.003
- Auprich, M., Chun, F. K., Ward, J. F., Pummer, K., Babaian, R., Agustin, H., et al. (2011b). Critical assessment of preoperative urinary prostate cancer antigen 3 on the accuracy of prostate cancer staging. *Eur. Urol.* 59, 96–105. doi: 10.1016/j.eururo.2010.10.024
- Bang, C., and Thum, T. (2012). Exosomes: new players in cell-cell communication. *Int. J. Biochem. Cell Biol.* 44, 2060–2064. doi: 10.1016/j.biocel.2012.08.007
- Barry, M. J., Fowler, F. J. Jr, O'Leary, M. P., Bruskewitz, R. C., Holtgrewe, H. L., Mebust, W. K., et al. (1992). The American Urological Association symptom index for benign prostatic hyperplasia. The Measurement Committee of the American Urological Association. *J. Urol.* 148, 1549–1557; discussion 1564.
- Bradley, L. A., Palomaki, G. E., Gutman, S., Samson, D., and Aronson, N. (2013). Comparative effectiveness review: prostate cancer antigen 3 testing for the diagnosis and management of prostate cancer. *J. Urol.* 190, 389–398. doi: 10.1016/j.juro.2013.02.005
- Chun, F. K., Haese, A., Ahyai, S. A., Walz, J., Suardi, N., Capitanio, U., et al. (2008). Critical assessment of tools to predict clinically insignificant prostate cancer at radical prostatectomy in contemporary men. *Cancer* 113, 701–709. doi: 10.1002/cncr.23610
- de la Taille, A., Irani, J., Graefen, M., Chun, F., de Reijke, T., Kil, P., et al. (2011). Clinical evaluation of the PCA3 assay in guiding initial biopsy decisions. *J. Urol.* 185, 2119–2125. doi: 10.1016/j.juro.2011.01.075
- Deatherage, B. L., and Cookson, B. T. (2012). Membrane vesicle release in bacteria, eukaryotes, and archaea: a conserved yet underappreciated aspect of microbial life. *Infect. Immun.* 80, 1948–1957. doi: 10.1128/IAI.06014-11
- Dijkstra, S., Birker, I. L., Smit, F. P., Leyten, G. H., de Reijke, T. M., van Oort, I. M., et al. (2014). Prostate cancer biomarker profiles in urinary sediments and exosomes. *J. Urol.* 191, 1132–1138. doi: 10.1016/j.juro.2013.11.001
- Dimitrova, N., Zamudio, J. R., Jong, R. M., Soukup, D., Resnick, R., Sarma, K., et al. (2014). LincRNA-p21 activates p21 in cis to promote Polycomb target gene expression and to enforce the G1/S checkpoint. *Mol. Cell* 54, 777–790. doi: 10.1016/j.molcel.2014.04.025
- Epstein, J. I. (2010). An update of the Gleason grading system. *J. Urol.* 183, 433–440. doi: 10.1016/j.juro.2009.10.046
- Epstein, J. I., Walsh, P. C., Carmichael, M., and Brendler, C. B. (1994). Pathologic and clinical findings to predict tumor extent of nonpalpable (stage T1c) prostate cancer. *JAMA* 271, 368–374. doi: 10.1001/jama.1994.03510290050036
- Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., et al. (2015). Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int. J. Cancer* 136, E359–E386. doi: 10.1002/ijc.29210
- Friedrich, M. J. (2011). Debate continues on use of PSA testing for early detection of prostate cancer. *JAMA* 305, 2273–2275. doi: 10.1001/jama.2011.777
- Gezer, U., Özgür, E., Cetinkaya, M., Isin, M., and Dalay, N. (2014). Long non-coding RNAs with low expression levels in cells are enriched in secreted exosomes. *Cell Biol. Int.* 38, 1076–1079. doi: 10.1002/cbin.10301
- Huang, X., Yuan, T., Liang, M., Du, M., Xia, S., Dittmar, R., et al. (2015). Exosomal miR-1290 and miR-375 as prognostic markers in castration-resistant prostate cancer. *Eur. Urol.* 67, 33–41. doi: 10.1016/j.eururo.2014.07.035
- Huarte, M., Guttman, M., Feldser, D., Garber, M., Koziol, M. J., Kenzelmann-Broz, D., et al. (2010). A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell* 142, 409–419. doi: 10.1016/j.cell.2010.06.040
- Isin, M., Özgür, E., Cetin, G., Erten, N., Aktan, M., Gezer, U., et al. (2014). Investigation of circulating lncRNAs in B-cell neoplasms. *Clin. Chim. Acta.* 431, 255–259. doi: 10.1016/j.cca.2014.02.010
- Jeldres, C., Suardi, N., Walz, J., Hutterer, G. C., Ahyai, S., Lattouf, J. B., et al. (2008). Validation of the contemporary epstein criteria for insignificant prostate cancer in European men. *Eur. Urol.* 54, 1306–1313. doi: 10.1016/j.eururo.2007.11.057
- Kapranov, P., St Laurent, G., Raz, T., Ozsolak, F., Reynolds, C. P., and Sorensen, P. H. B. (2010). The majority of total nuclear-encoded non-ribosomal RNA in a human cell is 'dark matter' un-annotated RNA. *BMC Biol.* 8:149. doi: 10.1186/1741-7007-8-149
- Khalil, A. M., Guttman, M., Huarte, M., Garber, M., Raj, A., Rivea Morales, D., et al. (2009). Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc. Natl. Acad. Sci. U.S.A.* 106, 11667–11672. doi: 10.1073/pnas.0904715106
- Kino, T., Hurt, D. E., Ichijo, T., Nader, N., and Chrousos, G. P. (2010). Noncoding RNA GAS5 is a growth arrest and starvation-associated repressor of the glucocorticoid receptor. *Sci. Signal.* 3, 8. doi: 10.1126/scisignal.2000568
- Leyten, G. H., Hessels, D., Jannink, S. A., Smit, F. P., de Jong, H., Cornel, E. B., et al. (2014). Prospective multicentre evaluation of PCA3 and TMPRSS2-ERG gene fusions as diagnostic and prognostic urinary biomarkers for prostate cancer. *Eur. Urol.* 65, 534–542. doi: 10.1016/j.eururo.2012.11.014
- Lippi, G., Montagnana, M., Guidi, G. C., and Plebani, M. (2009). Prostate specific antigen-based screening for prostate cancer in the third millennium: useful or hype? *Ann. Med.* 41, 480–489. doi: 10.1080/07853890903156468
- Loeb, S., Vellekoop, A., Ahmed, H. U., Catto, J., Emberton, M., Nam, R., et al. (2013). Systematic review of complications of prostate biopsy. *Eur. Urol.* 64, 876–892. doi: 10.1016/j.eururo.2013.05.049

- Luo, Y., Gou, X., Huang, P., and Mou, C. (2014). Prostate cancer antigen 3 test for prostate biopsy decision: a systematic review and meta analysis. *Chin. Med. J.* 127, 1768–1774.
- Morote, J., Rigau, M., Garcia, M., Mir, C., Ballesteros, C., Planas, J., et al. (2010). Behavior of the PCA3 gene in the urine of men with high grade prostatic intraepithelial neoplasia. *World J. Urol.* 28, 677–680. doi: 10.1007/s00345-010-0580-0
- Nakanishi, H., Groskopf, J., Fritsche, H. A., Bhadkamkar, V., Blase, A., Kumar, S. V., et al. (2008). PCA3 molecular urine assay correlates with prostate cancer tumor volume: implication in selecting candidates for active surveillance. *J. Urol.* 179, 1804–1809; discussion 1809–1810. doi: 10.1016/j.juro.2008.01.013
- Nogueira, L., Corradi, R., and Eastham, J. A. (2010). Other biomarkers for detecting prostate cancer. *BJU Int.* 105, 166–169. doi: 10.1111/j.1464-410X.2009.09088.x
- Pickard, M. R., and Williams, G. T. (2014). Regulation of apoptosis by the long non-coding RNA GAS5 in breast cancer cells: implications for chemotherapy. *Breast Cancer Res. Treat.* 145, 359–370. doi: 10.1007/s10549-014-2974-y
- Pisitkun, T., Shen, R. F., and Knepper, M. A. (2004). Identification and proteomic profiling of exosomes in human urine. *Proc. Natl. Acad. Sci. U.S.A.* 101, 13368–13373. doi: 10.1073/pnas.0403453101
- Ploussard, G., Durand, X., Xylinas, E., Moutereau, S., Radulescu, C., Forque, A., et al. (2011). Prostate cancer antigen 3 score accurately predicts tumour volume and might help in selecting prostate cancer patients for active surveillance. *Eur. Urol.* 59, 422–429. doi: 10.1016/j.eururo.2010.11.044
- Roddam, A. W., Duffy, M. J., Hamdy, F. C., Ward, A. M., Patnick, J., Price, C. P., et al. (2005). NHS Prostate Cancer Risk Management Programme. Use of prostate-specific antigen (PSA) isoforms for the detection of prostate cancer in men with a PSA level of 2–10 ng/ml: systematic review and meta-analysis. *Eur. Urol.* 48, 386–399. doi: 10.1016/j.eururo.2005.04.015
- Ronquist, G., and Brody, I. (1985). The prostasome: its secretion, and function in man. *Biochim. Biophys. Acta Rev. Biomembr.* 822, 203–218. doi: 10.1016/0304-4157(85)90008-5
- Schneider, C., King, R. M., and Philipson, L. (1988). Genes specifically expressed at growth arrest of mammalian cells. *Cell* 54, 787–793. doi: 10.1016/S0092-8674(88)91065-3
- Simpson, R. J., Jensen, S. S., and Lim, J. W. (2008). Proteomic profiling of exosomes: current perspectives. *Proteomics* 8, 4083–4099. doi: 10.1002/pmic.200800109
- St. Laurent, G., Shtokalo, D., Dong, B., Tackett, M. R., Fan, X., Lazorthes, S., et al. (2013). VlnRNAs controlled by retroviral elements are a hallmark of pluripotency and cancer. *Genome Biol.* 14, R73. doi: 10.1186/gb-2013-14-7-r73
- Taylor, D. D., Akyol, S., and Gercel-Taylor, C. (2006). Pregnancy-associated exosomes and their modulation of T cell signaling. *J. Biol. Chem.* 281, 1534–1542. doi: 10.4049/jimmunol.176.3.1534
- Wang, G., Li, Z., Zhao, Q., Zhu, Y., Zhao, C., Li, X., et al. (2014). lincRNA-p21 enhances the sensitivity of radiotherapy for human colorectal cancer by targeting the Wnt/ $\beta$ -catenin signaling pathway. *Oncol. Rep.* 31, 1839–1845. doi: 10.3892/or.2014.3047
- Yacqub-Usman, K., Pickard, M. R., and Williams, G. T. (2015). Reciprocal regulation of GAS5 lncRNA levels and mTOR inhibitor action in prostate cancer cells. *Prostate* 75, 693–705. doi: 10.1002/pros.22952
- Yaman Agaoglu, F., Kovancilar, M., Dizdar, Y., Darendeliler, E., Holdenrieder, S., Dalay, N., et al. (2011). Investigation of miR-21, miR-141, and miR-221 in blood circulation of patients with prostate cancer. *Tumour Biol.* 32, 583–588. doi: 10.1007/s13277-011-0154-9
- Zeliadt, S. B., Hoffman, R. M., Etzioni, R., Ginger, V. A., and Lin, D. W. (2010). What happens after an elevated PSA test: the experience of 13, 591 veterans. *J. Gen. Intern. Med.* 25, 1205–1210. doi: 10.1007/s11606-010-1468-9

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Işın, Uysaler, Özgür, Köseoglu, Şanlı, Yücel, Gezer and Dalay. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.