



Functional Analyses of a Novel Splice Variant in the *CHD7* Gene, Found by Next Generation Sequencing, Confirm Its Pathogenicity in a Spanish Patient and Diagnose Him with CHARGE Syndrome

Olatz Villate^{1,2}, Nekane Ibarluzea¹, Eugenia Fraile-Bethencourt³, Alberto Valenzuela³, Eladio A. Velasco³, Detelina Grozeva⁴, F. L. Raymond⁴, María P. Botella⁵ and María-Isabel Tejada^{1,2,6*}

¹ Biocruces Health Research Institute, Barakaldo, Spain, ² Molecular Genetics Laboratory, Genetics Service, Cruces University Hospital, Barakaldo, Spain, ³ Splicing and Cancer Laboratory, Instituto de Biología y Genética Molecular, Consejo Superior de Investigaciones Científicas, Universidad de Valladolid, Valladolid, Spain, ⁴ Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Cambridge, United Kingdom, ⁵ Department of Pediatrics, Araba University Hospital, Vitoria, Spain, ⁶ Clinical Group, Centro de Investigación Biomédica en Red de Enfermedades Raras, Madrid, Spain

OPEN ACCESS

Edited by:

Enrico Baruffini,
Università degli Studi di Parma, Italy

Reviewed by:

Paolo Ghirri,
University of Pisa, Italy
Theodora Katsila,
University of Patras, Greece

*Correspondence:

María-Isabel Tejada
mariaisabel.tejadaminguez@
osakidetza.eus

Specialty section:

This article was submitted to
Genetic Disorders,
a section of the journal
Frontiers in Genetics

Received: 10 November 2017

Accepted: 08 January 2018

Published: 26 January 2018

Citation:

Villate O, Ibarluzea N, Fraile-Bethencourt E, Valenzuela A, Velasco EA, Grozeva D, Raymond FL, Botella MP and Tejada M-I (2018) Functional Analyses of a Novel Splice Variant in the *CHD7* Gene, Found by Next Generation Sequencing, Confirm Its Pathogenicity in a Spanish Patient and Diagnose Him with CHARGE Syndrome. *Front. Genet.* 9:7. doi: 10.3389/fgene.2018.00007

Mutations in *CHD7* have been shown to be a major cause of CHARGE syndrome, which presents many symptoms and features common to other syndromes making its diagnosis difficult. Next generation sequencing (NGS) of a panel of intellectual disability related genes was performed in an adult patient without molecular diagnosis. A splice donor variant in *CHD7* (c.5665 + 1G > T) was identified. To study its potential pathogenicity, exons and flanking intronic sequences were amplified from patient DNA and cloned into the pSAD[®] splicing vector. HeLa cells were transfected with this construct and a wild-type minigene and functional analysis were performed. The construct with the c.5665 + 1G > T variant produced an aberrant transcript with an insert of 63 nucleotides of intron 28 creating a premature termination codon (TAG) 25 nucleotides downstream. This would lead to the insertion of 8 new amino acids and therefore a truncated 1896 amino acid protein. As a result of this, the patient was diagnosed with CHARGE syndrome. Functional analyses underline their usefulness for studying the pathogenicity of variants found by NGS and therefore its application to accurately diagnose patients.

Keywords: *CHD7*, next generation sequencing, alternative splicing, minigene, CHARGE syndrome

INTRODUCTION

The advent of next generation sequencing (NGS) technology which allows the sequencing either of the whole genome or of the expressed genes (exome) in one analysis, is transforming the process of genetic testing. NGS is being used extensively to diagnose diseases and find novel causative mutations for disease phenotypes. However, detailed analysis conclusively confirming

these variants, as well as the underlying molecular mechanisms explaining the diseases, are often lacking.

Actually, 100s of 1000s of DNA variants are detected in massive sequencing projects of genetic disorders and interestingly, many estimations have shown that an unexpectedly large fraction of genetic diseases are caused by variants that disrupt the splicing process (Wang and Cooper, 2007), ranging from 15 to >60% (Lopez-Bigas et al., 2005).

CHD7 is a gene located on chromosome 8q12.1 encoding the chromodomain helicase DNA-binding (CHD) protein 7, which belongs to a family of nine CHD proteins that can modify chromatin structure (Vissers et al., 2004). Among them *CHD7* is a transcriptional regulator that binds to enhancer elements in the nucleoplasm. CHARGE syndrome is characterized by Coloboma, Heart defects, Atresia of the choanae, Retardation of growth and development, Genital hypoplasia and Ear abnormalities, and approximately 60–70% of the patients have pathogenic mutations in *CHD7*, the major causative gene of this syndrome (Zentner et al., 2010b). While most *CHD7* mutations are nonsense or frameshift and predicted to be loss of function (Zentner et al., 2010a), the incidence of splice mutations is low, around 12% according to the Human Gene Mutation Database (HGMD) (Stenson et al., 2017). Moreover, splice mutations are often based on bioinformatic predictions and functional analyses confirming the pathogenicity of the mutations are lacking. Here, we describe the functional consequences of a novel splice mutation in *CHD7* found by NGS in a patient without molecular diagnosis.

BACKGROUND

Case Report

The male patient was the second child born to non-consanguineous Spanish healthy parents (24 and 30 years old, at the time of birth), with an unremarkable family history. He was born after 40 weeks of an uneventful pregnancy at a local hospital in 1977 and birth parameters were: weight 2,860 g (20th percentile), a length of 49 cm (5th percentile) and head circumference 36 cm (50th percentile). Apgar scores were 9 and 9 at 1 and 5 min, respectively.

At birth, he had an acceptable general appearance with good skin color, good muscle tone and normal active movements, but he showed facial dysmorphic features, including right choanal atresia resulting in a respiratory insufficiency, abnormal placement of the parietals, retrognathia of the lower jaw, narrow palate and glossoptosis, bilateral dysplastic, and low-set ears, protrusion of the right eye with megalocornea and papillar coloboma of the left eye. There were no thorax anomalies, neither in the limbs nor in the genitalia. At the age of 10 days, a systolic murmur was detected and therefore a congenital heart anomaly was suspected. At 21 days he was transferred to a reference hospital in Barcelona (Spain) where they found a cardiomegaly, an interventricular communication and an arteriovenous shunt and he was diagnosed with an atypical Treacher-Collins-Franceschetti syndrome.

During his 1st year of life he was admitted to his local hospital on many occasions due to breathing and swallowing difficulties requiring artificial ventilation and nasogastric tube feeding. Biochemical tests were negative, with no evidence of metabolic disease. At 1 year of age he had surgery due to his heart malformations in a reference center in Navarra (Spain).

It was early on when doctors realized that his psychomotor development was also delayed. He had autistic features and developed no speech. His weight-stature development was normal. Clinical data throughout his life are scarce but he was repeatedly admitted to the hospital because of recurrent respiratory infections, dyspnea, swallowing difficulties, gastrointestinal bleedings and a hyper-excitability which was difficult to control with a severe intellectual disability (ID). His parents had always cared for him at home until he died in 2013, at the age of 36. He had never been seen by an expert in Medical Genetics.

The molecular study of this patient began in 2010 in the context of a research project. This study was approved by the ethics committee for clinical research of Araba University Hospital (Vitoria, Spain). Informed consent was obtained from his parents before the extraction of peripheral blood samples for genetic analyses and they provided written consent to publish the report. Test results for karyotype, fragile X syndrome and arrays-CGH (60k) revealed no abnormalities. In 2011, he was included in a panel sequencing study of 565 ID-related genes within the UK10K project due to the scarcity of clinical data. A novel splice mutation in the *CHD7* gene was observed: c.5665 + 1G > T (Grozeva et al., 2015). This variant was not observed in gnomAD variant frequency database of more than 100,000 sequenced individuals (Lek et al., 2016). The presence of the variant in the patient was validated by Sanger sequencing and was absent in the parents, confirming that it is a *de novo* mutation (Figure 1A). When the parents received the result and the altered diagnosis of CHARGE syndrome, the patient had already died. Further analysis of the variant was performed to inform the recurrence risk for extended family members.

MATERIALS AND METHODS

Functional consequences of the mutation c.5665 + 1G > T in *CHD7* were tested by a minigene assay, as RNA from the patient was not available. First, to evaluate the potential impact of the variant on splicing, Human Splicing Finder¹ and NNSplice² splicing prediction programs were used. This bioinformatic analysis of the c.5665 + 1G > T variant in *CHD7* predicted the disruption of the canonical splice donor of exon 28 and also detected the presence of a cryptic donor site in the intron 28, 64 nucleotides downstream (NNSplice = 0.59) (Figure 1B). The +1G nucleotide is conserved in 100% of the 5' splice site recognized by the major spliceosome.

A minigene was constructed with an insert of 1480 bp corresponding to the exons 26, 27 and 28 and the flanking

¹<http://www.umd.be/HSF/>

²http://www.fruitfly.org/seq_tools/splice.html




FIGURE 2 | Functional analysis of the variant using a minigene strategy. **(A)** Exons 26–28, complete 26 and 27 introns and part of flanking introns 25 and 28 (intron 25, 355 bp, intron 28, 298 bp) of the *CHD7* gene were amplified (total size 1480 bp) and cloned into the splicing vector pSAD[®]. V1 and V2 are constitutive exons of the vector that in eukaryotic cells produce the splicing reactions (marked in the figure) with the cloned exons. **(B)** The RT-PCR functional assay of the wild-type (above) and mutant (below) minigenes was performed with a FAM-labeled first fluorescence (blue peaks). Capillary electrophoresis was performed in triplicate on an automated sequencer with the Genescan LIZ 1200 size standard (orange peaks; Applied Biosystems) and Peak Scanner (Applied Biosystems) analysis. The asterisk indicates a possible aberrant transcript. **(C)** Partial sequence of the RT-PCR product generated by the c.5665 + 1G > T variant. After sequencing, the insertion of 63 nt of intron 28 can be observed. No traces of other aberrant transcripts detected in the fragment analysis were observed. In the upper right the sequence of the border between exons 28 and V2 of the canonical transcript produced by the wild-type minigene can be observed.

have been reported in CHARGE syndrome, 96 of them being splice mutations (Stenson et al., 2017). The c.5665 + 1G > T variant is not reported in HGMD and our patient is the first described with this variant although there are several splice variants in that region.

To confirm the pathogenicity of the novel mutation, functional assays were performed. The feasibility of performing

functional analyses depends on the availability and accessibility of the required samples which can be a major challenge. In this context, a large number of methods, including model systems, can be used for functional interpretation of genome sequence variants. In our case, bioinformatic analyses suggested that the mutation could affect the splicing process. The ideal manner to study it would be to use the patient's RNA but in this case,

it was impossible to obtain because the patient was already deceased. Subsequently, a reliable and straightforward method to assess splicing was required. *Ex vivo* assays of DNA variants with splicing reporter minigenes have emerged to solve this problem (Bottillo et al., 2007; Lu et al., 2007; Jaijo et al., 2011; Acedo et al., 2015). Minigenes allow precise quantification of a single-mutant allele effect without the interference of the wild-type counterpart in patient samples. Another advantage of this approach is the high reproducibility of physiological/pathological splicing patterns by virtue of keeping the genomic context of each exon. Our functional assay showed that the novel splice donor variant c.5665 + 1G > T has a complete impact on the splicing of the *CHD7* gene and the effect would be a truncated protein.

The large 2997 amino acid CHD7 protein contains two chromodomains at its N terminus, followed by centrally located SNF2 and helicase domains; three conserved region (CR) domains; a switching-defective protein 3, adaptor 2, nuclear receptor corepressor; transcription factor IIIB (SANT) domain; two Brahma and Kismet (BRK) domains of unknown function; and, at the C terminus a leucine-zipper domain (Kim et al., 2008). It has been previously described that CHD7 can bind to the p53 promoter, thereby negatively regulating p53 expression, and that CHD7 loss in mouse neural crest cells or samples from patients with CHARGE syndrome results in p53 activation (Van Nostrand et al., 2014). The effect of the splice mutation found in this patient on the CHD7 protein will be the loss of the SANT and the BRK domains. The SANT domain may mediate binding to either DNA or modified histones (Schnetz et al., 2009) so the truncated protein will lose the ability to bind to the DNA or histones.

The minigene-construct allows the analysis of multiple variants from different exons. Therefore, the minigene containing exons 26–28 and the flanking intronic sequences of the *CHD7* gene we have constructed, could be used for the analysis of other splice mutations in that region for which there are not functional analysis yet (Bartels et al., 2010).

CONCLUDING REMARKS

Functional analyses are very useful for studying the pathogenicity of variants found by NGS. In the case of novel mutations in

the *CHD7* gene, the analysis of the functional consequences is very useful not only in subjects with typical signs not recognized at birth like our case, but also in subjects with atypical clinical signs. Specifically, in the present study our findings demonstrate that the novel variant c.5665 + 1G > T has a complete impact on the splicing of the *CHD7* gene resulting in a CHARGE syndrome and therefore improves our understanding of the genetic causes of CHARGE syndrome which is useful for accurately diagnosing patients and for providing genetic counseling to families.

AUTHOR CONTRIBUTIONS

MB provided patient clinical data and samples; EF-B, AV, EV, DG, FR, OV, and NI collected data and performed the experiments; OV and M-IT designed the study. All authors revised the manuscript critically, approved the final manuscript as submitted and agreed to be accountable for all aspects of the work.

FUNDING

This work was funded by Jesús de Gangoiti Barrera Foundation (FJGB15/005). The EAV laboratory is funded by projects of the Spanish Ministry of Economy and Competitiveness, National Plan for R & D 2013–2016, ISCIII (FIS: PI13/01749) co-financed by FEDER from Regional Development European Funds (European Union) and the project CSI090U14 of the Regional ministry of Education (ORDER EDU/122/2014) (Castilla y León, Spain). This study made use of data generated by the UK10K Project. Funding for the UK10K Project was provided by the Wellcome Trust under award WT091310.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2018.00007/full#supplementary-material>

REFERENCES

- Acedo, A., Hernandez-Moro, C., Curiel-Garcia, A., Diez-Gomez, B., and Velasco, E. A. (2015). Functional classification of *BRCA2* DNA variants by splicing assays in a large minigene with 9 exons. *Hum. Mutat.* 36, 210–221. doi: 10.1002/humu.22725
- Bartels, C. F., Scacheri, C., White, L., Scacheri, P. C., and Bale, S. (2010). Mutations in the *CHD7* gene: the experience of a commercial laboratory. *Genet. Test. Mol. Biomarkers* 14, 881–891. doi: 10.1089/gtmb.2010.0101
- Bottillo, I., De Luca, A., Schirinzi, A., Guida, V., Torrente, I., Calvieri, S., et al. (2007). Functional analysis of splicing mutations in exon 7 of *NF1* gene. *BMC Med. Genet.* 8:4. doi: 10.1186/1471-2350-8-4
- Fraile-Bethencourt, E., Diez-Gomez, B., Velasquez-Zapata, V., Acedo, A., Sanz, D. J., and Velasco, E. A. (2017). Functional classification of DNA variants by hybrid minigenes: identification of 30 spliceogenic variants of *BRCA2* exons 17 and 18. *PLOS Genet.* 13:e1006691. doi: 10.1371/journal.pgen.1006691
- Grozeva, D., Carss, K., Spasic-Boskovic, O., Tejada, M. I., Gecz, J., Shaw, M., et al. (2015). Targeted next-generation sequencing analysis of 1,000 individuals with intellectual disability. *Hum. Mutat.* 36, 1197–1204. doi: 10.1002/humu.22901
- Jaijo, T., Aller, E., Aparisi, M. J., Garcia-Garcia, G., Hernan, I., Gamundi, M. J., et al. (2011). Functional analysis of splicing mutations in *MYO7A* and *USH2A* genes. *Clin. Genet.* 79, 282–288. doi: 10.1111/j.1399-0004.2010.01454.x
- Janssen, N., Bergman, J. E., Swertz, M. A., Tranebjaerg, L., Lodahl, M., Schoots, J., et al. (2012). Mutation update on the *CHD7* gene involved in CHARGE syndrome. *Hum. Mutat.* 33, 1149–1160. doi: 10.1002/humu.22086
- Kim, H. G., Kurth, I., Lan, F., Melicani, I., Wenzel, W., Eom, S. H., et al. (2008). Mutations in *CHD7*, encoding a chromatin-remodeling protein, cause idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Am. J. Hum. Genet.* 83, 511–519. doi: 10.1016/j.ajhg.2008.09.005

- Kohmoto, T., Shono, M., Naruto, T., Watanabe, M., Suga, K., Nakagawa, R., et al. (2016). A novel frameshift mutation of *CHD7* in a Japanese patient with CHARGE syndrome. *Hum. Genome Var.* 3:16004. doi: 10.1038/hgv.2016.4
- Lalani, S. R., Hefner, M. A., Belmont, J. W., and Davenport, S. L. H. (1993). "CHARGE syndrome" in *GeneReviews*® [Internet], eds M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens et al (Seattle, WA: University of Washington).
- Lee, B., Duz, M. B., Sagong, B., Koparir, A., Lee, K. Y., Choi, J. Y., et al. (2016). Revealing the function of a novel splice-site mutation of *CHD7* in CHARGE syndrome. *Gene* 576, 776–781. doi: 10.1016/j.gene.2015.11.006
- Lek, M., Karczewski, K. J., Minikel, E. V., Samocha, K. E., Banks, E., Fennell, T., et al. (2016). Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 536, 285–291. doi: 10.1038/nature19057
- Lopez-Bigas, N., Audit, B., Ouzounis, C., Parra, G., and Guigo, R. (2005). Are splicing mutations the most frequent cause of hereditary disease? *FEBS Lett.* 579, 1900–1903. doi: 10.1016/j.febslet.2005.02.047
- Lu, Z. X., Peng, J., and Su, B. (2007). A human-specific mutation leads to the origin of a novel splice form of neuropsin (KLK8), a gene involved in learning and memory. *Hum. Mutat.* 28, 978–984. doi: 10.1002/humu.20547
- Schnetz, M. P., Bartels, C. F., Shastri, K., Balasubramanian, D., Zentner, G. E., Balaji, R., et al. (2009). Genomic distribution of *CHD7* on chromatin tracks H3K4 methylation patterns. *Genome Res.* 19, 590–601. doi: 10.1101/gr.086983.108
- Stenson, P. D., Mort, M., Ball, E. V., Evans, K., Hayden, M., Heywood, S., et al. (2017). The Human Gene Mutation Database: towards a comprehensive repository of inherited mutation data for medical research, genetic diagnosis and next-generation sequencing studies. *Hum. Genet.* 136, 665–677. doi: 10.1007/s00439-017-1779-6
- Van Nostrand, J. L., Brady, C. A., Jung, H., Fuentes, D. R., Kozak, M. M., Johnson, T. M., et al. (2014). Inappropriate p53 activation during development induces features of CHARGE syndrome. *Nature* 514, 228–232. doi: 10.1038/nature13585
- Vissers, L. E., van Ravenswaaij, C. M., Admiraal, R., Hurst, J. A., de Vries, B. B., Janssen, I. M., et al. (2004). Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nat. Genet.* 36, 955–957. doi: 10.1038/ng1407
- Wang, G. S., and Cooper, T. A. (2007). Splicing in disease: disruption of the splicing code and the decoding machinery. *Nat. Rev. Genet.* 8, 749–761. doi: 10.1038/nrg2164
- Zentner, G. E., Hurd, E. A., Schnetz, M. P., Handoko, L., Wang, C., Wang, Z., et al. (2010a). *CHD7* functions in the nucleolus as a positive regulator of ribosomal RNA biogenesis. *Hum. Mol. Genet.* 19, 3491–3501. doi: 10.1093/hmg/ddq265
- Zentner, G. E., Layman, W. S., Martin, D. M., and Scacheri, P. C. (2010b). Molecular and phenotypic aspects of *CHD7* mutation in CHARGE syndrome. *Am. J. Med. Genet. A* 152A, 674–686. doi: 10.1002/ajmg.a.33323

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Villate, Ibarluzea, Fraile-Bethencourt, Valenzuela, Velasco, Grozeva, Raymond, Botella and Tejada. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.