



# The X Files: “The Mystery of X Chromosome Instability in Alzheimer’s Disease”

Vladan P. Bajic<sup>1\*†</sup>, Magbubah Essack<sup>2†</sup>, Lada Zivkovic<sup>3</sup>, Alan Stewart<sup>4</sup>, Sonja Zafirovic<sup>1</sup>, Vladimir B. Bajic<sup>2‡</sup>, Takashi Gojobori<sup>2,5</sup>, Esmā Isenovic<sup>1</sup> and Biljana Spremo-Potparevic<sup>3</sup>

<sup>1</sup> Laboratory for Radiobiology and Molecular Genetics, Vinca Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia, <sup>2</sup> Computational Bioscience Research Center (CBRC), Computer, Electrical and Mathematical Sciences and Engineering (CEMSE) Division, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia, <sup>3</sup> Department of Physiology, Faculty of Pharmacy, University of Belgrade, Belgrade, Serbia, <sup>4</sup> School of Medicine, University of St Andrews, St Andrews, United Kingdom, <sup>5</sup> Biological and Environmental Sciences and Engineering Division (BESE), King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia

## OPEN ACCESS

### Edited by:

Ivan Y. Iourov,  
Mental Health Research Center of  
Russian Academy of Medical  
Sciences, Russia

### Reviewed by:

Huntington Potter,  
University of Colorado Denver,  
United States  
Anja Weise,  
University Hospital Jena,  
Germany

### \*Correspondence:

Vladan P. Bajic  
vladanbajic@yahoo.com

<sup>†</sup>These authors have contributed  
equally in this work and share first  
authorship

<sup>‡</sup>Deceased

### Specialty section:

This article was submitted to  
Human Genomics,  
a section of the journal  
Frontiers in Genetics

Received: 25 September 2019

Accepted: 13 December 2019

Published: 28 January 2020

### Citation:

Bajic VP, Essack M, Zivkovic L,  
Stewart A, Zafirovic S, Bajic VB,  
Gojobori T, Isenovic E and  
Spremo-Potparevic B (2020) The X  
Files: “The Mystery of X Chromosome  
Instability in Alzheimer’s Disease”.  
*Front. Genet.* 10:1368.  
doi: 10.3389/fgene.2019.01368

Alzheimer’s disease (AD) is a neurodegenerative disease that affects millions of individuals worldwide and can occur relatively early or later in life. It is well known that genetic components, such as the amyloid precursor protein gene on chromosome 21, are fundamental in early-onset AD (EOAD). To date, however, only the apolipoprotein E4 (ApoE4) gene has been proved to be a genetic risk factor for late-onset AD (LOAD). In recent years, despite the hypothesis that many additional unidentified genes are likely to play a role in AD development, it is surprising that additional gene polymorphisms associated with LOAD have failed to come to light. In this review, we examine the role of X chromosome epigenetics and, based upon GWAS studies, the PCDHX11 gene. Furthermore, we explore other genetic risk factors of AD that involve X-chromosome epigenetics.

**Keywords: X chromosome, Alzheimer’s disease, sex chromosome dosage, protocadherin 11, centromere instability**

## INTRODUCTION

In the first two decades of the 21st century, the proportion of individuals living with Alzheimer’s disease (AD) [AD (MIM: 104300)] has been on the rise with an increasingly aging population. Today, two basic forms of AD exist, early-onset AD (EOAD) and late-onset AD (LOAD). EOAD correlates with the occurrence of mutations on specific genes that have given rise to inherited forms of the disease, whilst LOAD - which occurs later in life - has no specified etiology (Smith, 1998; Selkoe, 2001). Familial studies have identified a point mutation associated with EOAD on chromosome 21. This mutation is located in a gene called amyloid precursor protein (APP), and all members of these families show signs of the Alzheimer’s phenotype at a relatively early stage of

**Abbreviations:** AD, Alzheimer’s disease; EOAD, Early onset AD; GWAS, genome-wide association studies; LOAD, Late onset AD; NRC/MASC, N-methyl-D-aspartate receptor complex/membrane-associated guanylate kinase-associated signaling complex; PCDH11X, protocadherin 11; PSD, postsynaptic density; SCD, Sex Chromosome Dosage; Xist, X-inactive-specific transcript gene; Xi, inactive X chromosome.

life (Wiseman et al., 2018). Novel mutations located on chromosomes 14 and 1 in genes encoding presenilin-1 and presenilin-2 have also been identified in EOAD (Guven et al., 2019). Unfortunately, specific genetic determinants that can explain the high prevalence of LOAD have yet to be identified.

Today we are aware that EOAD comprises only 1-3% of all AD cases (Smith, 1998; Selkoe, 2001; Bekris et al., 2010). In LOAD subjects, disease prevalence changes with age; 5% after 65 years of age, 20% after 75 years of age, 30% after 80 years of age (Bekris et al., 2010). Also women are twice as likely to suffer from AD than men (Pike, 2017). This prevalence is suggested to be due to differences in the life expectancy between males and females and to hormonal status (Vest and Pike, 2013; Pike, 2017). Studies with twins clearly signal that a strong genetic component is present in LOAD cases (Gatz et al., 2006; Seripa et al., 2009). Various genes have been implicated in AD and identified by using genetic approaches, such as Genome-Wide Association Studies (GWAS). However, the only "single gene" risk factor for LOAD without opposition in the research community concerns the gene encoding apolipoprotein E4 (Giri et al., 2017). In LOAD, the percentage of individuals carrying the at-risk allele of the ApoE4 gene was found to be between 20% and 70%, suggesting that there are additional genetic, and perhaps also epigenetic, factors that underlie the development of LOAD (Slooter et al., 1998; Giri et al., 2017). Carrasquillo et al. (Carrasquillo et al., 2009) found that an alteration of a single-nucleotide polymorphism (SNP; rs5984894) on the Xq21.3 in a gene called protocadherin 11 (PCDH11X) in a cohort of women, was significantly associated with LOAD (**Figure 1**). Other GWAS, however, have been unable to confirm the existence of these connections (Beecham et al., 2010; Wu et al., 2010; Miar et al., 2011; Chung et al., 2013). We hypothesize that one of the possible answers to these observed genetic discrepancies is based on the epigenetics of the X chromosome.

We have previously identified centromere impairment or premature centromere separation (PCS) of the X chromosome in neuronal nuclei of the cerebral cortex in AD women (Spremo-Potparevic et al., 2008). In addition, Yurov et al. discovered X chromosome aneuploidy in AD-affected neurons (Yurov et al., 2014), which suggests that premature centromere separation is a


mechanism of X chromosome instability (Spremo-Potparevic et al., 2008). Epigenetically, chromosome X can be affected by skewed X chromosome inactivation, asynchronous replication patterns of the inactive X chromosome (Xi), X-inactivation escape, aneuploidy, and premature centromere separation. All these epigenetic X chromosome changes could potentially affect X chromosome genes through changes in sex chromosome dosage (SCD), and consequently promote AD pathogenesis (Amiel et al., 1998; Gribnau et al., 2005; Ahn and Lee, 2008; Spremo-Potparevic et al., 2008; Hong and Reiss, 2014; Mugford et al., 2014; Yurov et al., 2014; Bajic et al., 2015a; Balaton and Brown, 2016; Le Gall et al., 2017; Graham et al., 2019). Raznahan et al. found recently that sex chromosome dosage not only influenced the adjacent sex chromosomes X and Y, but also autosomal gene expression (Raznahan et al., 2018).

## The X Chromosome Is Unique

In women, there is a systematic demand to compensate for SCD by silencing one of the copies of the X chromosome. With two X-chromosomes, women are more prone to inheriting potentially deleterious mutations in X-encoded genes, which, because of Xi, may all be expressed in different cells. The first finding of inactivation of the X chromosome was reported by Lyon, (1961). It was found that one of the X chromosomes, paternal or maternal, was always inactivated, suggesting that an inactivation mechanism only allows active transcription at one X chromosome (Splinter et al., 2011). This process of X-chromosome inactivation (XCI) evolved as a mechanism to regulate gene dosage. As a compensation mechanism, it does not affect all genes equally, and those genes that are not affected are known to escape XCI [termed escapees; (Pessia et al., 2012)].

Human embryos initially have non-random imprinted XCI, where the X-chromosome from the mother remains active, and XCI applies only to the X-chromosome inherited from the father. The imprint is not constant; XCI resets at the embryonic implantation stage. At this point the XCI reset leaves the maternal and paternal X open to random inactivation (Sun and Lee, 2006). Because XCI at this stage is random it causes most women to be mosaic for two cell lines, one harboring the active chromosome, the paternal X, and the other the maternal X. The randomness of this process causes an XCI ratio of approximately 50%:50% to be associated with the two cell lines in the female population. However, on rare occasions, in approximately 9% of the female population, a bias towards one of the two X chromosomes produces a skewed ratio (> 80%:20%; (Amos-Landgraf et al., 2006). In this regard, Renault et al., analyzed the distribution of X-inactivation patterns (the relative abundance of the two cell populations) in a large cohort study of normal females, and reported that human XCI distribution pattern is more genetically influenced in comparison to the Xi model, which suggests a completely random selection of XCI (Renault et al., 2013).

The genetically influenced selection of XCI may be indicative of mutations in genes (Orstavik, 2009; Shvetsova et al., 2019), suggesting the inactive X chromosome often harbors the mutated allele of an X-linked gene. This would mean that with a 50%:50% XCI ratio, wild type cells generally ameliorate disease



phenotypes. Changes in the XCI ratio towards an increased expression of mutated genes can increase disease phenotype severity, as it is in the case of female hemophilia A (Renault et al., 2007), and sideroblastic anemia (Cazzola et al., 2000), where the majority of cells express the mutated allele. Changes in the XCI ratio where expression of the mutated allele is increased to exhibit the disease phenotype can also occur, as in Rett syndrome. In this case, the hemizygous mutation of the methyl-CpG-binding protein 2 (MeCP2) gene in males causes lethality, while the MeCP2 heterozygous mutation in females weakens such phenotypic consequences. It seems that the loss of MeCP2 function contributes to Rett syndrome, while the gain in MeCP2 dosage does not necessarily ameliorate the disease phenotype but may manifest as a less aggressive form in other neurological diseases. Increased expression of MeCP2 was found to be associated with other neurological diseases, such as AD and Huntington's disease (Amir et al., 1999; Ausio et al., 2014; McFarland et al., 2014; Maphis et al., 2017).

Xi acquires several features of heterochromatin, such as hypermethylation, hypercondensation, altered replication patterns (late vs. early), and depletion of acetylated histones (Chow and Brown, 2003; Ng et al., 2007). Methylation patterns have been extensively used to determine the inactive chromosome (Shvetsova et al., 2019), enabling an analysis of non-random inactivation processes in diseases that are X chromosome-linked (Yuan et al., 2015). In our published study we suggest that changes in the inactivation patterns of the X-chromosome could have an impact on AD pathogenesis (Bajic et al., 2015a).

## BRAIN AND THE X CHROMOSOME

The X chromosome harbors 3-5% of all the genes in a genome (Skuse, 2005). There has been a debate on how many genes reside on the X chromosome and how many genes are expressed in the brain alone, compared to genes that are X-linked and expressed in the placenta, testes, muscles, and ovary. It is estimated that between 1,100 and 1,500 genes are present on the X chromosome (Skuse, 2005; Laumonnier et al., 2007). By using the Mart View software it was found that 1,500 X-linked genes are expressed in the brain, which represent numerous candidate genes that could be responsible for X-linked brain diseases (Laumonnier et al., 2007). Many of the proteins expressed from the genes linked to the X chromosome represent channels, receptors, repair, transcription factors, and DNA/RNA binding proteins. Most of these proteins are located in the postsynaptic cleft and postsynaptic density (PSD) and are regulated through signaling complexes (Nguyen and Distèche, 2006; Laumonnier et al., 2007). It is intriguing that even if the X chromosome harbors 3-5% of all the genes, it is responsible for 10% of all diseases with Mendelian inheritance (Germain, 2006).

Another aspect that makes the X chromosome unique is that it harbors a higher proportion of brain-expressed miRNAs than would be expected (Goncalves et al., 2019), with 20% of these related to autoimmune diseases such as rheumatoid arthritis and

systemic lupus erythematosus (Khalifa et al., 2016). Most of these miRNA are clustered, for example, miR532/188, miR-221/222, miR-98/Let7f, and miR-363/106a/20b/92a (DeMarco et al., 2019; Goncalves et al., 2019). Many of these are also intronic and it is believed that they are co-transcribed and co-expressed with other genes linked to chromosome X (deleted X-linked genes) and may be susceptible to SCD, skewing, and Xi escape processes. It is important to point out that inflammation and altered immunity are features of AD (Forloni et al., 1992; Hauss-Wegrzyniak et al., 1998; Eikelenboom et al., 2000; McGeer and McGeer, 2002; Castellani et al., 2008; Krstic et al., 2012; Bajic et al., 2015c; Regen et al., 2017).

How these X-linked genes interact with genes controlling the immune system in AD is still unknown. For individual genes involved in diseases of the brain, a more complex hypothesis is that interplay occurs in disease genes embedded in multiprotein neuronal complexes. Many of the most important components of neuronal complexes are encoded on the X chromosome (Laumonnier et al., 2007). Such complexes, which are essential for neuronal plasticity, cognitive processes, and cell signaling, are thought to be in the PSD cleft (Muddashetty et al., 2011; Yudowski et al., 2013). Taking N-methyl-D-aspartate receptor/membrane-associated guanylate kinase-associated signaling complex/(NRC/MASC) as an example; combining its 185 proteins and with the other proteins in PSD gives a total of 1100 proteins. The X chromosome plays an essential role, and the percentage of genes related to synaptic plasticity, some 86% of all the genes in NRC/MASC are genes linked to chromosome X (Grant et al., 2005; Laumonnier et al., 2007). It is interesting that these genes are also presented or expressed in human cognitive disorders (Grant et al., 2005; Pocklington et al., 2006; Fernandez et al., 2009; Tam et al., 2009). An analysis of the number of altered proteins in X-linked mental retardation disorders shows that from 69 genes currently known, 19 (or 28%) of these genes belong to postsynaptic proteins (Laumonnier et al., 2007). The same pattern is conserved in the mouse X chromosome, and this suggests a network of multiprotein complexes functioning as integrated entities or complex molecular machines. If one component of this complex machinery is disrupted, the whole complex/network fails thus impairing the overall role of the multiprotein complex in processes of cognition (Grant et al., 2005; Nguyen and Distèche, 2006; Pocklington et al., 2006; Laumonnier et al., 2007; Fernandez et al., 2009; Tam et al., 2009).

## X-Linked miRNA and the Brain

The X chromosome is enriched in ncRNAs and harbors several miRNAs essential to brain function (Goncalves et al., 2019). It is important to note that miRNAs not only affect mRNA through translation repression but also work through other ncRNAs, such as lncRNAs and circRNAs, affecting downstream genes (Khalifa et al., 2016; DeMarco et al., 2019; Goncalves et al., 2019). Bian et al. revealed that a miRNA located on the X chromosome, a miR-374 family member, plays a role in cell growth and differentiation not only in various cancers, but also in AD. This miR-374 member is located at the X chromosome

inactivation center and targets the VEGF, PTEN, Wnt, and Fas signaling pathways (Bian et al., 2019). Importantly, the PTEN pathway is of importance to the progression of AD through a mechanism that includes altered autophagy (Wani et al., 2019), mitophagy (Fang et al., 2019), and apoptosis (Cui et al., 2017). A report by Manzine et al. suggested that miR-374 directly targets the beta-secretase 1 to regulate the progress of AD, as the levels of miR-374 were significantly decreased in comparison to controls (Manzine et al., 2018). In addition to miR-374, several miRNAs have also been found to correlate to X chromosome-linked intellectual disability syndrome, and among them are miR-223-3p, miR-362-5p, miR-504-5p.1, miR-361-5p, miR-505-3p.1 and miR-505-3p.2. All these miRNAs act as key regulators of genes linked to chromosome X but also of many autosomal intellectual disability genes that are connected in a complex network (Goncalves et al., 2019).

In the future, it is hoped that further work will reveal the extent to which genes on the X chromosome and miRNAs expressed in the brain, that together regulate processes including nervous system development, cell proliferation and transcription regulation, are altered by X chromosome skewing and asynchronous replication, which lead to aneuploidy and deregulation of cohesion dynamics in AD. Also, RNA genes that are linked to the X chromosome are prone to escape inactivation of the X chromosome (Peeters et al., 2019). These epigenetic processes may prove to be gender-associated as research shows that expression of an X-linked miRNA in rheumatoid arthritis is more prevalent in women than in men (Khalifa et al., 2016).

## PCDH11X

Carrasquillo et al. previously identified an SNP (rs 5984894) on the X chromosome (Xq21,3) in a gene called PCDH11X (Carrasquillo et al., 2009). This locus is associated with LOAD in women of European origin from the USA. The PCDH11X gene encodes the protein, protocadherin 11. Women who are homozygous for this SNP have a greater risk of developing AD, not only when compared to women without the SNP, but also when compared to women that are heterozygotes, and male hemizygotes (Carrasquillo et al., 2009). Zubenko et al. reported that the DXS1047 genotype is correlated with AD (Zubenko et al., 1999) and that this genotype is associated with the PCDH11X gene (Zubenko et al., 1998). The results from the same authors indicate an association between the variation in the PCDH11X gene and the risk of acquiring AD, but these results have not been confirmed in other GWAS (Beecham et al., 2010; Wu et al., 2010; Miar et al., 2011). Our suggestion is that these discrepancies in GWAS results may well be due to the changes in the epigenetics of the X chromosome.

## Does PCDH11X Escape X Inactivation?

Pseudoautosomal genes and functional Y chromosome orthologues (X-linked genes with Y homology) tend to escape X inactivation (Disteche et al., 2002; Brown and Grealley, 2003). Sudbrak et al. reported that PCDH11X expression might also

escape X inactivation, and this assumption was verified by using an X chromosome-specific cDNA microarray where elevated expression of PCDH11X was identified in cells expressing multiple X chromosomes (Sudbrak et al., 2001). Lopes et al. indirectly found that PCDH11X expression was higher in women than in men by looking at CpG islands and their methylation patterns. By using bisulfite sequencing analysis, the same authors found the absence of CpG island methylation on both the active and the Xi chromosomes and that these processes coincide with possible PCDH11X escape from X inactivation (Lopes et al., 2006). Another study found that PCDH11X can undergo asynchronous replication, and that PCDH11X is also prone to escape the inactivation process (Wilson et al., 2007). Replication asynchrony of the X pseudoautosomal locus has been identified (Vorsanova et al., 2001), and suggests that other genes that replicate asynchronously are also prone to escape inactivation (Anderson and Brown, 2005; Carrel and Willard, 2005; Escamilla-Del-Arenal et al., 2011).

## PCDH11X Asynchronous Replication

Xi is associated with a sequence of epigenetic modifications (Chow and Brown, 2003), and goes through a phase of changes involving DNA methylation and histone modification resulting in Xi condensation in a body called the Barr body. This results in changes in DNA replication – more specifically, the Xi in the S phase replicates later than its active counterpart. Imperfect chromosome replication can be a consequence of “escapees” (genes that escape the inactivation process). Such genes include hypoxanthine-guanine phosphoribosyltransferase and Fragile X-chromosome genes that display asynchronicity. The X-inactive-specific transcript (Xist) gene (important for inactivation) that is expressed from the Xi also replicates asynchronously (Boggs and Chinault, 1994; Aladjem and Fu, 2014). Wilson et al. reported that PCDH11X displays replication asynchrony in both female and male cells (Wilson et al., 2007). The data from these authors, together with those from others (Orstavik, 2009), show that a complex relationship exists between X-inactivation, replication asynchrony, and the status of expression of individual genes on chromosome X (Bajic et al., 2008; Bajic et al., 2009).

It thus appears that synchronous replication occurs more frequently than previously thought, and is found not only through imprinting, but also through randomized monoallelic expression, pathologies, and tandem duplications (Wilson et al., 2007). Clinically, an increase in asynchronous replication increases the risk in women for aneuploidy (Amiel et al., 2000). The relationship between centromere instability, control of replication, and nondisjunction are best exemplified by the fact that young women that have children with Down's syndrome have twice the risk of developing AD (Hardy et al., 1989; Goate et al., 1990; Fidani et al., 1992; Schellenberg et al., 1992; Schupf et al., 1994; Petersen et al., 2000; Schupf et al., 2001; Migliore et al., 2006; Migliore et al., 2009; Iourov et al., 2010; Goate and Hardy, 2012).

Chromosomes 21, X, and 18 were primarily affected, showing repeated non-disjunction and centromere impairment (Potter and Geller, 1996; Geller and Potter, 1999; Petersen et al., 2000;



Migliore et al., 2006; Migliore et al., 2009; Iourov et al., 2010; Potter, 2016). We suggest that X chromosome replication asynchrony is likely to lead to accelerated instability of chromosome X in AD (Bajic et al., 2009).

## SEX CHROMOSOME DOSAGE (SCD): AN ENGINE OF STABILITY

The crosstalk that exists between X chromosomes and autologous genes is a relatively new paradigm that has emerged as a result of the biology of sex differences, and gives rise to the question of how SCD shapes the genome function. To explore this, human sex aneuploidies were analyzed from a genome-wide expression dataset by Raznahan et al. where they found a dosage sensitivity of the X-Y chromosome pair resulting in increased expression of genes that decrease X/Y chromosomal dosage (Raznahan et al., 2018). The most interesting finding was that X-linked genes were found to regulate co-expression of networks of autosomal genes that are SCD-sensitive and, in addition to these findings, suggest that the autosomal genes and their corresponding networks are crucial for cellular functions. This highlights the potential of SCD to affect the occurrence of disease.

The most common aneuploidy in AD is XO mosaicism (Spremo-Potparevic et al., 2004; Spremo-Potparevic et al., 2008; Yurov et al., 2014; Spremo-Potparevic et al., 2015). In respect to SCD and the XO status, Raznahan et al. have demonstrated up-regulation of the protein networks, noncoding RNA metabolism, suppression of the cell cycle, changes in regulation of DNA/chromatin organization, glycolysis, and response to stress (Raznahan et al., 2018). Changes in these collective networks through XO and supernumerary XXY, and XXYY syndromes may enhance the risk of AD (Raznahan et al., 2018; Graham et al., 2019).

There is a small but constant number of neuronal cells that express a different number of chromosomes, such as aneuploidy (Iourov et al., 2006; Yurov et al., 2007; Iourov et al., 2008; Iourov et al., 2009; Yurov et al., 2014), but also copy number variation on chromosome 21, which is crucial in AD (Cai et al., 2014), DNA content variation (Madrigal et al., 2007; Westra et al., 2010), and LINE elements (Evrony et al., 2012).


Mosaic aneuploidy in the brain revealed that not only was chromosome 21 affected in AD, but also that the X chromosome was found to be supernumerary and presumed to be affected through a mechanism that involves altered cohesion/cohesin dynamics (Spremo-Potparevic et al., 2004; Spremo-Potparevic et al., 2008; Bajic et al., 2009; Zivković et al., 2010; Zivkovic et al., 2013; Yurov et al., 2014; Bajic et al., 2015b; Spremo-Potparevic et al., 2015; Yurov et al., 2019).

Yurov et al. (2014) suggested that chromosome 21 might not be the only chromosome to influence changes in genome stability of a neuron, which leads to a cascade of processes that result in neuronal loss. The finding that affected brains show a two-fold increase in X chromosome aneuploidy in the hippocampus and

cerebrum - areas of the brain most affected by AD - suggesting that altered sex chromosome dosage plays a role in the large scale genomic variation in neuronal cells in AD compared to controls. These results have been recently corroborated by the finding that the sex chromosomes were distinct from autosomes in the dorsolateral prefrontal cortex and that X chromosome aneuploidy was associated with a faster rate of cognitive decline which is a hallmark of AD (Graham et al., 2019). Therefore, X chromosome aneuploidy may contribute to aging, but also to processes leading to pathological changes in brains affected by AD.

Previously we proposed the “post-mitotic state-maintained protein hypothesis” where we distinguished aneuploization in the brain as constitutional aneuploidy with non-pathological diversification of the neurons (Bajic et al., 2015b). These aneuploidogenic processes are balanced with cohesin and cohesion-related proteins. Alteration of this balance develops as a link between neuronal development and chromosomal instability, intracellular diversity and human brain diseases including AD (Hong and Reiss, 2014). Looking closely at the overall somatic mosaicism found in the brain, we, together with others, suggest that micro aneuploidy or segmental aneuploidy is a more proper measure of changes in gene dosage leading to AD (Dierssen et al., 2009). These processes are heavily realized when looking at SCD effects on gene expression in humans (Raznahan et al., 2018).

An additional complexity of genome mosaicism in the brain relates to findings concerning DNA and gene copy number variations. Regional variations of DNA content has been identified with higher DNA content found in the frontal cortex and cerebellum compared with other brain regions (Westra et al., 2010). Copy number variations may be considered as an independent genetic factor not related to other genomic changes, suggesting its plays a role in neurodevelopmental disorders in patients with sex chromosome aneuploidies (Haack et al., 2013; Le Gall et al., 2017). It has been reported that 11% of neurons in the brain cortex exhibit a DNA content that is above the diploid level (Fischer et al., 2012), and similar findings have also been reported in the AD brain (Ueberham and Arendt, 2005; Arendt et al., 2010; Yurov, 2017; Barrio-Alonso et al., 2018). These somatic gene variations in neurons appear to be generated by chromosome segregation defects. Some of these cells are expelled by apoptosis, but several cells are introduced as a pool of variability of the neuronal genome. These cell populations are thus vulnerable in the sense that they are more prone to genome instability and thus may contribute to age-related mental disorders, such as AD. Gómez-Ramos et al. presented distinct X-chromosome single nucleotide variants from some sporadic AD samples (Gómez-Ramos et al., 2015). In samples from LOAD patients, a higher number of single nucleotide variants in genes present at the X chromosome were identified using exome sequencing compared to age-matched controls. Two genes that were not previously described as risk factors, UBE2NL and ATXN3L, were found to have variants important for the ubiquitin pathway in LOAD (Gómez-Ramos et al., 2015).



**FIGURE 2 |** X chromosome instability, Sex Chromosome Dosage, Topological changes of Chromosomes, and its possible role in AD.

Maintenance of the interphase state in neurons is an active process. The 3D organization of the genome is correlated to gene expression in the interphase. In the 3D domain, chromosomes occupy preferential positions by self-organizing into topologically-associated domains, which may change due to the cell lineage or stage of the organism (Laskowski et al., 2019). There is a possible exchange between the inactive and active chromosome in gene regulatory information. Cohesin is indirectly associated with the Xi 3D position in the genome. Minajigi et al. reported that a reaction between Xist and cohesins results in the repulsion of the latter from Xi, thus changing its 3D shape (Minajigi et al., 2015). The Xi is much more complex, and it also represents a reservoir of genes that could replace mutated genes from the active X chromosome. At present, this untapped potential known as the X interactome requires further investigation (Minajigi et al., 2015). Progress in understanding the Xist interactome requires more understanding of how it is used and how epigenetically-regulated long ncRNAs potentially influence disease. By utilizing a specific technique named iDRiP, some 200 proteins in the Xist interactome were identified

(Minajigi et al., 2015). Most of the proteins are from several categories, such as cohesins, condensins, topoisomerases, RNA helicase, histone modifiers, methyltransferases, nuclear matrix proteins, and nucleoskeletal factors. Cohesin may play a more important role in the complex relationship between the Xi and active X chromosome (Minajigi et al., 2015). Even though these processes are an important mechanism of diversity, alterations may lead to an increased structural and topological variation of the genome in the brain, enhancing the susceptibility of affected neurons to genome instability that may lead to AD (Bajic et al., 2015b; Graham et al., 2019; Yurov et al., 2019).

A number of publications have reported mislocalization of some critical proteins responsible for chromatin organization and epigenetic modifications in brain diseases including AD (Gill et al., 2007; Lu et al., 2014; Luperchio et al., 2014; Quinodoz and Guttman, 2014; Guo et al., 2015; Mastroeni et al., 2015; Pombo and Dillon, 2015; Sen et al., 2015; Winick-Ng and Rylett, 2018).

All these data suggest, that in AD chromatin, organizers are deregulated and chromatin topology is changed in a manner that alters gene expression leading to synaptic dysfunction, a major

pathological change in AD, and consequently neurodegeneration (Gill et al., 2007; Lu et al., 2014; Luperchio et al., 2014; Quinodoz and Guttman, 2014; Guo et al., 2015; Mastroeni et al., 2015; Pombo and Dillon, 2015; Sen et al., 2015; Winick-Ng and Rylett, 2018).

Xist RNA can act as a scaffold for proteins required to maintain the inactive state of neurons. It has been shown that it can act as a repulsion mechanism that expels architectural factors such as cohesins in order to avoid unwanted chromatin conformation that could increase unfavorable transcription (Raznahan et al., 2018). Minajigi et al. suggest that Xi RNA plays an important role in the organization of how chromosomes are regulated into chromosome territories and that Xi inactivation is fundamentally important in these processes (Minajigi et al., 2015). It could be suggested that X chromosome instability found in AD may result in changes in the Xi pattern, Xi escapees, SCD, and consequently changes in the topological organization, thus altering chromatin organization that may affect already other genes related to AD (Figure 2).

The cohesin-associated protein, shugoshin-1, seems to be fundamental in repressing the accumulation of amyloid- $\beta$  and Tau phosphorylation in shugoshin-1 gene (Sgo1) haploinsufficient mice (Rao et al., 2018).

## SUMMARY

Conflicting results from studies of the PCDH11X gene in AD could be explained by cohort size, ethnicity, and environmental factors *per se* but also by the influence of X chromosome epigenetics. Thus, GWAS of sex chromosomes should take into account any alterations of the epigenetic processes in the X chromosome (Schurz et al., 2019).

## REFERENCES

- Ahn, J., and Lee, J. (2008). X chromosome: X inactivation. *Nature Education* 1, 24.
- Aladjem, M. I., and Fu, H. (2014). A new light on DNA replication from the inactive X chromosome. *Bioessays* 36, 591–597. doi: 10.1002/bies.201400021
- Amiel, A., Avivi, L., Gaber, E., and Fejgin, M. D. (1998). Asynchronous replication of allelic loci in down syndrome. *Eur. J. Hum. Genet.* 6, 359–364. doi: 10.1038/sj.ejhg.5200199
- Amiel, A., Reish, O., Gaber, E., Kedar, I., Diukman, R., and Fejgin, M. (2000). Replication asynchrony increases in women at risk for aneuploid offspring. *Chromosome Res.* 8, 141–150. doi: 10.1023/a:1009246603868
- Amir, R. E., Van den Veyver, I. B., Wan, M., Tran, C. Q., Francke, U., and Zoghbi, H. Y. (1999). Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nat. Genet.* 23, 185–188. doi: 10.1038/13810
- Amos-Landgraf, J. M., Cottle, A., Plenge, R. M., Friez, M., Schwartz, C. E., Longshore, J., et al. (2006). X chromosome-inactivation patterns of 1,005 phenotypically unaffected females. *Am. J. Hum. Genet.* 79, 493–499. doi: 10.1086/507565
- Anderson, C. L., and Brown, C. J. (2005). Epigenetic predisposition to expression of TIMP1 from the human inactive X chromosome. *BMC Genet.* 6, 48. doi: 10.1186/1471-2156-6-48
- Arendt, T., Bruckner, M. K., Mosch, B., and Losche, A. (2010). Selective cell death of hyperplod neurons in Alzheimer's disease. *Am. J. Pathol.* 177, 15–20. doi: 10.2353/ajpath.2010.090955

The findings that chromosome X expresses all of the somatic genomic neuronal variability properties and can *de novo* express several epigenetic mechanisms suggest that the X chromosome instability phenotype may be viewed as an important risk factor in AD pathogenesis.

## AUTHOR CONTRIBUTIONS

All authors (VPB, ME, LZ, AS, SZ, VBB, TG, EI, and BS-P) contributed to the design and writing of the manuscript. VPB and SZ designed the figures.

## FUNDING

This work is part of the collaboration between the Laboratory of Radiobiology and Molecular Genetics, Vinca Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia and King Abdullah University of Science and Technology (KAUST), Computational Bioscience Research Center (CBRC), Thuwal, Saudi Arabia. This work has been supported by grants No. 173033 (EI) and No. 173034 (VPB) from the Ministry of Education, Science and Technological Development, Republic of Serbia and by the KAUST grant OSR#4129 (to EI and VBB), which also supported SZ and VPB. VBB has been supported by the KAUST Base Research Fund (BAS/1/1606-01-01), while VBB and ME have been supported by KAUST Office of Sponsored Research (OSR) grant no. FCC/1/1976-17-01. TG has been supported by the King Abdullah University of Science and Technology (KAUST) Base Research Fund (BAS/1/1059-01-01).

- Ausio, J., Martinez de Paz, A., and Esteller, M. (2014). MeCP2: the long trip from a chromatin protein to neurological disorders. *Trends Mol. Med.* 20, 487–498. doi: 10.1016/j.molmed.2014.03.004
- Bajic, V. P., Spremo-Potparevic, B., Zivkovic, L., Djelic, N., and Smith, M. A. (2008). Is the time dimension of the cell cycle re-entry in AD regulated by centromere cohesion dynamics? *Biosci. Hypotheses* 1, 156–161. doi: 10.1016/j.bihy.2008.03.006
- Bajic, V. P., Spremo-Potparevic, B., Zivkovic, L., Bonda, D. J., Siedlak, S. L., Casadesus, G., et al. (2009). The X-chromosome instability phenotype in Alzheimer's disease: a clinical sign of accelerating aging? *Med. Hypotheses* 73, 917–920. doi: 10.1016/j.mehy.2009.06.046
- Bajic, V., Mandusic, V., Stefanova, E., Bozovic, A., Davidovic, R., Zivkovic, L., et al. (2015a). Skewed X-chromosome inactivation in women affected by Alzheimer's disease. *J. Alzheimers Dis.* 43, 1251–1259. doi: 10.3233/JAD-141674
- Bajic, V., Spremo-Potparevic, B., Zivkovic, L., Isenovic, E. R., and Arendt, T. (2015b). Cohesion and the aneuploid phenotype in Alzheimer's disease: A tale of genome instability. *Neurosci. Biobehav. Rev.* 55, 365–374. doi: 10.1016/j.neubiorev.2015.05.010
- Bajic, V., Stanojevic, B., Zivkovic, L., Cabarkapa, A., Perry, G., Arendt, T., et al. (2015c). Cyclin dependent kinase 11, neuroinflammation and alzheimer's disease: a review. *J. Clin. Cell Immunol.* 6, 305. doi: 10.4172/2155-9899.1000305
- Balaton, B. P., and Brown, C. J. (2016). Escape Artists of the X Chromosome. *Trends Genet.* 32, 348–359. doi: 10.1016/j.tig.2016.03.007
- Barrio-Alonso, E., Hernández-Vivanco, A., Walton, C. C., Perea, G., and Frade, J. M. (2018). Cell cycle reentry triggers hyperploidy and synaptic

- dysfunction followed by delayed cell death in differentiated cortical neurons. *Sci. Rep.* 8, 14316–14316. doi: 10.1038/s41598-018-32708-4
- Beecham, G. W., Naj, A. C., Gilbert, J. R., Haines, J. L., Buxbaum, J. D., and Pericak-Vance, M. A. (2010). PCDH11X variation is not associated with late-onset Alzheimer disease susceptibility. *Psychiatr. Genet.* 20, 321–324. doi: 10.1097/YPG.0b013e32833b635d
- Bekris, L. M., Yu, C. E., Bird, T. D., and Tsuang, D. W. (2010). Genetics of Alzheimer disease. *J. Geriatr. Psychiatry Neurol.* 23, 213–227. doi: 10.1177/0891988710383571
- Bian, H., Zhou, Y., Zhou, D., Zhang, Y., Shang, D., and Qi, J. (2019). The latest progress on miR-374 and its functional implications in physiological and pathological processes. *J. Cell Mol. Med.* 23, 3063–3076. doi: 10.1111/jcmm.14219
- Boggs, B. A., and Chinault, A. C. (1994). Analysis of replication timing properties of human X-chromosomal loci by fluorescence *in situ* hybridization. *Proc. Natl. Acad. Sci. U.S.A.* 91, 6083–6087. doi: 10.1073/pnas.91.13.6083
- Brown, C. J., and Grealley, J. M. (2003). A stain upon the silence: genes escaping X inactivation. *Trends Genet.* 19, 432–438. doi: 10.1016/S0168-9525(03)00177-X
- Cai, X., Evrony, G. D., Lehmann, H. S., Elhosary, P. C., Mehta, B. K., Poduri, A., et al. (2014). Single-cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human brain. *Cell Rep.* 8, 1280–1289. doi: 10.1016/j.celrep.2014.07.043
- Carrasquillo, M. M., Zou, F., Pankratz, V. S., Wilcox, S. L., Ma, L., Walker, L. P., et al. (2009). Genetic variation in PCDH11X is associated with susceptibility to late-onset Alzheimer's disease. *Nat. Genet.* 41, 192–198. doi: 10.1038/ng.305
- Carrel, L., and Willard, H. F. (2005). X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature* 434, 400–404. doi: 10.1038/nature03479
- Castellani, R. J., Lee, H. G., Zhu, X., Perry, G., and Smith, M. A. (2008). Alzheimer disease pathology as a host response. *J. Neuropathol. Exp. Neurol.* 67, 523–531. doi: 10.1097/NEN.0b013e318177eaf4
- Cazzola, M., May, A., Bergamaschi, G., Cerani, P., Rosti, V., and Bishop, D. F. (2000). Familial-skewed X-chromosome inactivation as a predisposing factor for late-onset X-linked sideroblastic anemia in carrier females. *Blood* 96, 4363–4365. doi: 10.1182/blood.V96.13.4363
- Chow, J. C., and Brown, C. J. (2003). Forming facultative heterochromatin: silencing of an X chromosome in mammalian females. *Cell Mol. Life Sci.* 60, 2586–2603. doi: 10.1007/s00018-003-3121-9
- Chung, S. J., Lee, J. H., Kim, S. Y., You, S., Kim, M. J., Lee, J. Y., et al. (2013). Association of GWAS top hits with late-onset Alzheimer disease in Korean population. *Alzheimer Dis. Assoc. Disord.* 27, 250–257. doi: 10.1097/WAD.0b013e31826d7281
- Cui, W., Wang, S., Wang, Z., Wang, Z., Sun, C., and Zhang, Y. (2017). Inhibition of PTEN attenuates endoplasmic reticulum stress and apoptosis via activation of PI3K/AKT pathway in Alzheimer's disease. *Neurochem. Res.* 42, 3052–3060. doi: 10.1007/s11064-017-2338-1
- DeMarco, B., Stefanovic, S., Williams, A., Moss, K. R., Anderson, B. R., Bassell, G. J., et al. (2019). FMRP - G-quadruplex mRNA - miR-125a interactions: implications for miR-125a mediated translation regulation of PSD-95 mRNA. *PLoS One* 14, e0217275. doi: 10.1371/journal.pone.0217275
- Dierssen, M., Hérault, Y., and Estivill, X. (2009). Aneuploidy: from a physiological mechanism of variance to Down syndrome. *Physiol. Rev.* 89, 887–920. doi: 10.1152/physrev.00032.2007
- Disteche, C. M., Filippova, G. N., and Tsuchiya, K. D. (2002). Escape from X inactivation. *Cytogenet. Genome Res.* 99, 36–43. doi: 10.1159/000071572
- Eikelenboom, P., Rozemuller, A. J., Hoozemans, J. J., Veerhuis, R., and van Gool, W. A. (2000). Neuroinflammation and Alzheimer disease: clinical and therapeutic implications. *Alzheimer Dis. Assoc. Disord.* 14 Suppl 1, S54–S61. doi: 10.1097/00002093-200000001-00009
- Escamilla-Del-Arenal, M., da Rocha, S. T., and Heard, E. (2011). Evolutionary diversity and developmental regulation of X-chromosome inactivation. *Hum. Genet.* 130, 307–327. doi: 10.1007/s00439-011-1029-2
- Evrony, G. D., Cai, X., Lee, E., Hills, L. B., Elhosary, P. C., Lehmann, H. S., et al. (2012). Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell* 151, 483–496. doi: 10.1016/j.cell.2012.09.035
- Fang, E. F., Hou, Y., Palikaras, K., Adriaanse, B. A., Kerr, J. S., Yang, B., et al. (2019). Mitophagy inhibits amyloid- $\beta$  and tau pathology and reverses cognitive deficits in models of Alzheimer's disease. *Nat. Neurosci.* 22, 401–412. doi: 10.1038/s41593-018-0332-9
- Fernandez, E., Collins, M. O., Uren, R. T., Kopanitsa, M. V., Komiyama, N. H., Croning, M. D., et al. (2009). Targeted tandem affinity purification of PSD-95 recovers core postsynaptic complexes and schizophrenia susceptibility proteins. *Mol. Syst. Biol.* 5, 269. doi: 10.1038/msb.2009.27
- Fidani, L., Rooke, K., Chartier-Harlin, M. C., Hughes, D., Tanzi, R., Mullan, M., et al. (1992). Screening for mutations in the open reading frame and promoter of the beta-amyloid precursor protein gene in familial Alzheimer's disease: identification of a further family with APP717 Val $\rightarrow$ Ile. *Hum. Mol. Genet.* 1, 165–168. doi: 10.1093/hmg/1.3.165
- Fischer, H. G., Morawski, M., Bruckner, M. K., Mittag, A., Tarnok, A., and Arendt, T. (2012). Changes in neuronal DNA content variation in the human brain during aging. *Aging Cell* 11, 628–633. doi: 10.1111/j.1474-9726.2012.00826.x
- Forloni, G., Demicheli, F., Giorgi, S., Bendotti, C., and Angeretti, N. (1992). Expression of amyloid precursor protein mRNAs in endothelial, neuronal and glial cells: modulation by interleukin-1. *Brain Res. Mol. Brain Res.* 16, 128–134. doi: 10.1016/0169-328x(92)90202-m
- Gómez-Ramos, A., Podlesniy, P., Soriano, E., and Avila, J. (2015). Distinct X-chromosome SNVs from some sporadic AD samples. *Sci. Rep.* 5, 18012. doi: 10.1038/srep18012
- Gatz, M., Reynolds, C. A., Fratiglioni, L., Johansson, B., Mortimer, J. A., Berg, S., et al. (2006). Role of genes and environments for explaining Alzheimer disease. *Arch. Gen. Psychiatry* 63, 168–174. doi: 10.1001/archpsyc.63.2.168
- Geller, L. N., and Potter, H. (1999). Chromosome missegregation and trisomy 21 mosaicism in Alzheimer's disease. *Neurobiol. Dis.* 6, 167–179. doi: 10.1006/nbdi.1999.0236
- Germain, D. P. (2006). "General aspects of X-linked diseases," in *Fabry Disease: Perspectives from 5 Years of FOS*. Eds. A. Mehta, M. Beck and G. Sunder-Plassmann (Oxford: Oxford PharmaGenesis).
- Gill, S. K., Ishak, M., Dobransky, T., Haroutunian, V., Davis, K. L., and Rylett, R. J. (2007). 82-kDa choline acetyltransferase is in nuclei of cholinergic neurons in human CNS and altered in aging and Alzheimer disease. *Neurobiol. Aging* 28, 1028–1040. doi: 10.1016/j.neurobiolaging.2006.05.011
- Giri, M., Shah, A., Upreti, B., and Rai, J. C. (2017). Unraveling the genes implicated in Alzheimer's disease. *Biomed. Rep.* 7, 105–114. doi: 10.3892/br.2017.927
- Goate, A., and Hardy, J. (2012). Twenty years of Alzheimer's disease-causing mutations. *J. Neurochem.* 120 Suppl 1, 3–8. doi: 10.1111/j.1471-4159.2011.07575.x
- Goate, A. M., Hardy, J. A., Owen, M. J., Haynes, A., James, L., Farrall, M., et al. (1990). Genetics of Alzheimer's disease. *Adv. Neurol.* 51, 197–198.
- Goncalves, T. F., Piergiorgio, R. M., Dos Santos, J. M., Gusmao, J., Pimentel, M. M. G., and Santos-Reboucas, C. B. (2019). Network profiling of brain-expressed X-chromosomal microRNA genes implicates shared key microRNAs in intellectual disability. *J. Mol. Neurosci.* 67, 295–304. doi: 10.1007/s12031-018-1235-7
- Graham, E. J., Vermeulen, M., Vardarajan, B., Bennett, D., De Jager, P., Pearce, R. V., 2nd, et al. (2019). Somatic mosaicism of sex chromosomes in the blood and brain. *Brain Res.* 1721, 146345. doi: 10.1016/j.brainres.2019.146345
- Grant, S. G. N., Marshall, M. C., Page, K.-L., Cumiskey, M. A., and Armstrong, J. D. (2005). Synapse proteomics of multiprotein complexes: en route from genes to nervous system diseases. *Hum. Mol. Genet.* 14, R225–R234. doi: 10.1093/hmg/ddi330
- Gribnau, J., Luikenhuis, S., Hochedlinger, K., Monkhorst, K., and Jaenisch, R. (2005). X chromosome choice occurs independently of asynchronous replication timing. *J. Cell Biol.* 168, 365–373. doi: 10.1083/jcb.200405117
- Guo, Y., Xu, Q., Canzio, D., Shou, J., Li, J., Gorkin, D. U., et al. (2015). CRISPR Inversion of CTCF sites alters genome topology and enhancer/promoter function. *Cell* 162, 900–910. doi: 10.1016/j.cell.2015.07.038
- Guyen, G., Erginel-Unaltuna, N., Samanci, B., Gulec, C., Hanagasi, H., and Bilgic, B. (2019). A patient with early-onset Alzheimer's disease with a novel PSEN1 p.Leu424Pro mutation. *Neurobiol. Aging* 84, 238.e1–238.e4. doi: 10.1016/j.neurobiolaging.2019.05.014
- Haack, T. B., Hogarth, P., Gregory, A., Prokisch, H., and Hayflick, S. J. (2013). BPAN: the only X-linked dominant NBIA disorder. *Int. Rev. Neurobiol.* 110, 85–90. doi: 10.1016/b978-0-12-410502-7.00005-3
- Hardy, J., Goate, A., Owen, M., and Rossor, M. (1989). Presenile dementia associated with mosaic trisomy 21 in a patient with a Down syndrome child. *Lancet* 2, 743. doi: 10.1016/s0140-6736(89)90805-2



- Haus-Wegrzyniak, B., Dobrzanski, P., Stoehr, J. D., and Wenk, G. L. (1998). Chronic neuroinflammation in rats reproduces components of the neurobiology of Alzheimer's disease. *Brain Res.* 780, 294–303. doi: 10.1016/s0006-8993(97)01215-8
- Hong, D. S., and Reiss, A. L. (2014). Cognitive and neurological aspects of sex chromosome aneuploidies. *Lancet Neurol.* 13, 306–318. doi: 10.1016/s1474-4422(13)70302-8
- Iourov, I. Y., Vorsanova, S. G., and Yurov, Y. B. (2006). Chromosomal variation in mammalian neuronal cells: known facts and attractive hypotheses. *Int. Rev. Cytol.* 249, 143–191. doi: 10.1016/s0074-7696(06)49003-3
- Iourov, I. Y., Vorsanova, S. G., and Yurov, Y. B. (2008). Molecular cytogenetics and cytogenomics of brain diseases. *Curr. Genomics* 9, 452–465. doi: 10.2174/138920208786241216
- Iourov, I. Y., Vorsanova, S. G., Liehr, T., and Yurov, Y. B. (2009). Aneuploidy in the normal, Alzheimer's disease and ataxia-telangiectasia brain: differential expression and pathological meaning. *Neurobiol. Dis.* 34, 212–220. doi: 10.1016/j.nbd.2009.01.003
- Iourov, I. Y., Vorsanova, S. G., and Yurov, Y. B. (2010). Somatic genome variations in health and disease. *Curr. Genomics* 11, 387–396. doi: 10.2174/138920210793176065
- Khalifa, O., Pers, Y. M., Ferreira, R., Senechal, A., Jorgensen, C., Apparailly, F., et al. (2016). X-Linked miRNAs associated with gender differences in rheumatoid arthritis. *Int. J. Mol. Sci.* 17, 1852–1863. doi: 10.3390/ijms17111852
- Krstic, D., Madhusudan, A., Doehner, J., Vogel, P., Notter, T., Imhof, C., et al. (2012). Systemic immune challenges trigger and drive Alzheimer-like neuropathology in mice. *J. Neuroinflammation* 9, 151. doi: 10.1186/1742-2094-9-151
- Laskowski, A. I., Neems, D. S., Laster, K., Strojny-Okyere, C., Rice, E. L., Konieczna, I. M., et al. (2019). Varying levels of X chromosome coalescence in female somatic cells alters the balance of X-linked dosage compensation and is implicated in female-dominant systemic lupus erythematosus. *Sci. Rep.* 9, 8011. doi: 10.1038/s41598-019-44229-9
- Laumonnier, F., Cuthbert, P. C., and Grant, S. G. (2007). The role of neuronal complexes in human X-linked brain diseases. *Am. J. Hum. Genet.* 80, 205–220. doi: 10.1086/511441
- Le Gall, J., Nizon, M., Pichon, O., Andrieux, J., Audebert-Bellanger, S., Baron, S., et al. (2017). Sex chromosome aneuploidies and copy-number variants: a further explanation for neurodevelopmental prognosis variability? *Eur. J. Hum. Genet.* 25, 930–934. doi: 10.1038/ejhg.2017.93
- Lopes, A. M., Ross, N., Close, J., Dagnall, A., Amorim, A., and Crow, T. J. (2006). Inactivation status of PCDH11X: sexual dimorphisms in gene expression levels in brain. *Hum. Genet.* 119, 267–275. doi: 10.1007/s00439-006-0134-0
- Lu, T., Aron, L., Zullo, J., Pan, Y., Kim, H., Chen, Y., et al. (2014). REST and stress resistance in ageing and Alzheimer's disease. *Nature* 507, 448–454. doi: 10.1038/nature13163
- Luperchio, T. R., Wong, X., and Reddy, K. L. (2014). Genome regulation at the peripheral zone: lamina associated domains in development and disease. *Curr. Opin. Genet. Dev.* 25, 50–61. doi: 10.1016/j.gde.2013.11.021
- Lyon, M. F. (1961). Gene action in the X-chromosome of the mouse (*Mus musculus* L.). *Nature* 190, 372–373. doi: 10.1038/190372a0
- Madrigal, I., Rodríguez-Revilla, L., Armengol, L., González, E., Rodríguez, B., Badenas, C., et al. (2007). X-chromosome tiling path array detection of copy number variants in patients with chromosome X-linked mental retardation. *BMC Genomics* 8, 443. doi: 10.1186/1471-2164-8-443
- Manzine, P. R., Pelucchi, S., Horst, M. A., Vale, F. A. C., Pavarini, S. C. I., Audano, M., et al. (2018). microRNA 221 targets ADAM10 mRNA and is downregulated in Alzheimer's disease. *J. Alzheimers Dis.* 61, 113–123. doi: 10.3233/jad-170592
- Maphis, N. M., Jiang, S., Binder, J., Wright, C., Gopalan, B., Lamb, B. T., et al. (2017). Whole Genome Expression Analysis in a Mouse Model of Tauopathy Identifies MECP2 as a Possible Regulator of Tau Pathology. *Front. Mol. Neurosci.* 10, 69. doi: 10.3389/fnmol.2017.00069
- Mastroeni, D., Delvaux, E., Nolz, J., Tan, Y., Grover, A., Oddo, S., et al. (2015). Aberrant intracellular localization of H3k4me3 demonstrates an early epigenetic phenomenon in Alzheimer's disease. *Neurobiol. Aging* 36, 3121–3129. doi: 10.1016/j.neurobiolaging.2015.08.017
- McFarland, K. N., Huizenga, M. N., Darnell, S. B., Sangrey, G. R., Berezovska, O., Cha, J. H., et al. (2014). MeCP2: a novel Huntingtin interactor. *Hum. Mol. Genet.* 23, 1036–1044. doi: 10.1093/hmg/ddt499
- McGeer, P. L., and McGeer, E. G. (2002). Local neuroinflammation and the progression of Alzheimer's disease. *J. Neurovirol.* 8, 529–538. doi: 10.1080/13550280290100969
- Miar, A., Alvarez, V., Corao, A. I., Alonso, B., Diaz, M., Menendez, M., et al. (2011). Lack of association between protocadherin 11-X/Y (PCDH11X and PCDH11Y) polymorphisms and late onset Alzheimer's disease. *Brain Res.* 1383, 252–256. doi: 10.1016/j.brainres.2011.01.054
- Migliore, L., Boni, G., Bernardini, R., Trippi, F., Colognato, R., Fontana, L., et al. (2006). Susceptibility to chromosome malsegregation in lymphocytes of women who had a down syndrome child in young age. *Neurobiol. Aging* 27, 710–716. doi: 10.1016/j.neurobiolaging.2005.03.025
- Migliore, L., Migheli, F., and Coppede, F. (2009). Susceptibility to aneuploidy in young mothers of down syndrome children. *Sci. World J.* 9, 1052–1060. doi: 10.1100/tsw.2009.122
- Minajigi, A., Froberg, J. E., Wei, C., Sunwoo, H., Kesner, B., Colognori, D., et al. (2015). A comprehensive Xist interactome reveals cohesin repulsion and an RNA-directed chromosome conformation. *Science* 349, aab2276. doi: 10.1126/science.aab2276
- Muddashetty, R. S., Nalavadi, V. C., Gross, C., Yao, X., Xing, L., Laur, O., et al. (2011). Reversible inhibition of PSD-95 mRNA translation by miR-125a, FMRP phosphorylation, and mGluR signaling. *Mol. Cell* 42, 673–688. doi: 10.1016/j.molcel.2011.05.006
- Mugford, J. W., Starmer, J., Williams, R. L., Jr., Calabrese, J. M., Mieczkowski, P., Yee, D., et al. (2014). Evidence for local regulatory control of escape from imprinted X chromosome inactivation. *Genetics* 197, 715–723. doi: 10.1534/genetics.114.162800
- Ng, K., Pullirsch, D., Leeb, M., and Wutz, A. (2007). Xist and the order of silencing. *EMBO Rep.* 8, 34–39. doi: 10.1038/sj.embor.7400871
- Nguyen, D. K., and Disteche, C. M. (2006). Dosage compensation of the active X chromosome in mammals. *Nat. Genet.* 38, 47–53. doi: 10.1038/ng1705
- Orstavik, K. H. (2009). X chromosome inactivation in clinical practice. *Hum. Genet.* 126, 363–373. doi: 10.1007/s00439-009-0670-5
- Peeters, S. B., Korecki, A. J., Baldry, S. E. L., Yang, C., Tosefsky, K., Balaton, B. P., et al. (2019). How do genes that escape from X-chromosome inactivation contribute to Turner syndrome? *Am. J. Med. Genet.* 181, 28–35. doi: 10.1002/ajmg.c.31672
- Pessia, E., Makino, T., Bailly-Bechet, M., McLysaght, A., and Marais, G. A. B. (2012). Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proc. Natl. Acad. Sci.* 109, 5346. doi: 10.1073/pnas.1116763109
- Petersen, M. B., Karadima, G., Samaritaki, M., Avramopoulos, D., Vassilopoulos, D., and Mikkelsen, M. (2000). Association between presenilin-1 polymorphism and maternal meiosis II errors in down syndrome. *Am. J. Med. Genet.* 93, 366–372. doi: 10.1002/1096-8628(20000828)93:5<366::AID-AJMG5>3.0.CO;2-G
- Pike, C. J. (2017). Sex and the development of Alzheimer's disease. *J. Neurosci. Res.* 95, 671–680. doi: 10.1002/jnr.23827
- Pocklington, A. J., Cumiskey, M., Armstrong, J. D., and Grant, S. G. (2006). The proteomes of neurotransmitter receptor complexes form modular networks with distributed functionality underlying plasticity and behaviour. *Mol. Syst. Biol.* 2, 2006 0023. doi: 10.1038/msb4100041
- Pombo, A., and Dillon, N. (2015). Three-dimensional genome architecture: players and mechanisms. *Nat. Rev. Mol. Cell Biol.* 16, 245. doi: 10.1038/nrm3965
- Potter, H., and Geller, L. N. (1996). Alzheimer's disease, Down's syndrome, and chromosome segregation. *Lancet* 348. doi: 10.1016/s0140-6736(05)64399-1
- Potter, H. (2016). Beyond trisomy 21: phenotypic variability in people with down syndrome explained by further chromosome mis-segregation and mosaic aneuploidy. *J. Down Syndr. Chromosom. Abnorm.* 2, 1–6. doi: 10.4172/2472-1115.1000109
- Quinodoz, S., and Guttman, M. (2014). Long noncoding RNAs: an emerging link between gene regulation and nuclear organization. *Trends Cell Biol.* 24, 651–663. doi: 10.1016/j.tcb.2014.08.009
- Rao, C. V., Farooqui, M., Zhang, Y., Asch, A. S., and Yamada, H. Y. (2018). Spontaneous development of Alzheimer's disease-associated brain pathology in

- a Shugoshin-1 mouse cohesinopathy model. *Aging Cell* 17, e12797. doi: 10.1111/ace1.12797
- Raznahan, A., Parikshak, N. N., Chandran, V., Blumenthal, J. D., Clasen, L. S., Alexander-Bloch, A. F., et al. (2018). Sex-chromosome dosage effects on gene expression in humans. *Proc. Natl. Acad. Sci. U.S.A.* 115, 7398–7403. doi: 10.1073/pnas.1802889115
- Regen, F., Hellmann-Regen, J., Costantini, E., and Reale, M. (2017). Neuroinflammation and Alzheimer's disease: implications for microglial activation. *Curr. Alzheimer Res.* 14, 1140–1148. doi: 10.2174/1567205014666170203141717
- Renault, N. K., Dyack, S., Dobson, M. J., Costa, T., Lam, W. L., and Greer, W. L. (2007). Heritable skewed X-chromosome inactivation leads to haemophilia a expression in heterozygous females. *Eur. J. Hum. Genet.* 15, 628–637. doi: 10.1038/sj.ejhg.5201799
- Renault, N. K., Pritchett, S. M., Howell, R. E., Greer, W. L., Sapienza, C., Orstavik, K. H., et al. (2013). Human X-chromosome inactivation pattern distributions fit a model of genetically influenced choice better than models of completely random choice. *Eur. J. Hum. Genet.* 21, 1396–1402. doi: 10.1038/ejhg.2013.84
- Schellenberg, G. D., Bird, T. D., Wijsman, E. M., Orr, H. T., Anderson, L., Nemens, E., et al. (1992). Genetic linkage evidence for a familial Alzheimer's disease locus on chromosome 14. *Science* 258, 668–671. doi: 10.1126/science.1411576
- Schupf, N., Kapell, D., Lee, J. H., Ottman, R., and Mayeux, R. (1994). Increased risk of Alzheimer's disease in mothers of adults with down's syndrome. *Lancet* 344, 353–356. doi: 10.1016/s0140-6736(94)91398-6
- Schupf, N., Kapell, D., Nightingale, B., Lee, J. H., Mohlenhoff, J., Bewley, S., et al. (2001). Specificity of the fivefold increase in AD in mothers of adults with down syndrome. *Neurology* 57, 979–984. doi: 10.1212/wnl.57.6.979
- Schurz, H., Salie, M., Tromp, G., Hoal, E. G., Kinnear, C. J., and Möller, M. (2019). The X chromosome and sex-specific effects in infectious disease susceptibility. *Hum. Genomics* 13, 2. doi: 10.1186/s40246-018-0185-z
- Selkoe, D. J. (2001). Alzheimer's disease results from the cerebral accumulation and cytotoxicity of amyloid beta-protein. *J. Alzheimers Dis.* 3, 75–80. doi: 10.3233/jad-2001-3111
- Sen, A., Nelson, T. J., and Alkon, D. L. (2015). ApoE4 and abeta oligomers reduce BDNF expression via HDAC nuclear translocation. *J. Neurosci.* 35, 7538–7551. doi: 10.1523/jneurosci.0260-15.2015
- Seripa, D., Panza, F., Franceschi, M., D'Onofrio, G., Solfrizzi, V., Dallapiccola, B., et al. (2009). Non-apolipoprotein E and apolipoprotein E genetics of sporadic Alzheimer's disease. *Ageing Res. Rev.* 8, 214–236. doi: 10.1016/j.arr.2008.12.003
- Shvetsova, E., Sofronova, A., Monajemi, R., Gagalova, K., Draisma, H. H. M., White, S. J., et al. (2019). Skewed X-inactivation is common in the general female population. *Eur. J. Hum. Genet.* 27, 455–465. doi: 10.1038/s41431-018-0291-3
- Skuse, D. H. (2005). X-linked genes and mental functioning. *Hum. Mol. Genet.* 14 Spec No 1, R27–R32. doi: 10.1093/hmg/ddi112
- Slooter, A. J., de Knijff, P., Hofman, A., Cruts, M., Breteler, M. M., Van Broeckhoven, C., et al. (1998). Serum apolipoprotein E level is not increased in Alzheimer's disease: the rotterdam study. *Neurosci. Lett.* 248, 21–24. doi: 10.1016/s0304-3940(98)00339-5
- Smith, M. A. (1998). Alzheimer disease. *Int. Rev. Neurobiol.* 42, 1–54. doi: 10.1016/s0074-7742(08)60607-8
- Splinter, E., de Wit, E., Nora, E. P., Klous, P., van de Werken, H. J., Zhu, Y., et al. (2011). The inactive X chromosome adopts a unique three-dimensional conformation that is dependent on Xist RNA. *Genes Dev.* 25, 1371–1383. doi: 10.1101/gad.633311
- Spremo-Potparevic, B., Zivkovic, L., Djelic, N., and Bajic, V. (2004). Analysis of premature centromere division (PCD) of the X chromosome in Alzheimer patients through the cell cycle. *Exp. Gerontol.* 39, 849–854. doi: 10.1016/j.exger.2004.01.012
- Spremo-Potparevic, B., Zivkovic, L., Djelic, N., Plecas-Solarovic, B., Smith, M. A., and Bajic, V. (2008). Premature centromere division of the X chromosome in neurons in Alzheimer's disease. *J. Neurochem.* 106, 2218–2223. doi: 10.1111/j.1471-4159.2008.05555.x
- Spremo-Potparevic, B., Bajic, V., Perry, G., and Zivkovic, L. (2015). Alterations of the X chromosome in lymphocytes of alzheimer's disease patients. *Curr. Alzheimer Res.* 12, 990–996. doi: 10.2174/1567205012666151027124154
- Sudbrak, R., Wiczorek, G., Nuber, U. A., Mann, W., Kirchner, R., Erdogan, F., et al. (2001). X chromosome-specific cDNA arrays: identification of genes that escape from X-inactivation and other applications. *Hum. Mol. Genet.* 10, 77–83. doi: 10.1093/hmg/10.1.77
- Sun, B. K., and Lee, J. T. (2006). “X-Chromosome Inactivation,” in *Encyclopedic Reference of Genomics and Proteomics in Molecular Medicine* (Berlin, Heidelberg: Springer Berlin Heidelberg), 2013–2019.
- Tam, G. W., Redon, R., Carter, N. P., and Grant, S. G. (2009). The role of DNA copy number variation in schizophrenia. *Biol. Psychiatry* 66, 1005–1012. doi: 10.1016/j.biopsych.2009.07.027
- Ueberham, U., and Arendt, T. (2005). The expression of cell cycle proteins in neurons and its relevance for Alzheimer's disease. *Curr. Drug Targets CNS Neurol. Disord.* 4, 293–306. doi: 10.2174/1568007054038175
- Vest, R. S., and Pike, C. J. (2013). Gender, sex steroid hormones, and Alzheimer's disease. *Horm. Behav.* 63, 301–307. doi: 10.1016/j.yhbeh.2012.04.006
- Vorsanova, S. G., Yurov, Y. B., Kolotii, A. D., and Soloviev, I. V. (2001). FISH analysis of replication and transcription of chromosome X loci: new approach for genetic analysis of Rett syndrome. *Brain Dev.* 23 Suppl 1, S191–S195. doi: 10.1016/s0387-7604(01)00364-3
- Wani, A., Gupta, M., Ahmad, M., Shah, A. M., Ahsan, A. U., Qazi, P. H., et al. (2019). Alborixin clears amyloid- $\beta$  by inducing autophagy through PTEN-mediated inhibition of the AKT pathway. *Autophagy* 15, 1810–1828. doi: 10.1080/1548627.2019.1596476
- Westra, J. W., Rivera, R. R., Bushman, D. M., Yung, Y. C., Peterson, S. E., Barral, S., et al. (2010). Neuronal DNA content variation (DCV) with regional and individual differences in the human brain. *J. Comp. Neurol.* 518, 3981–4000. doi: 10.1002/cne.22436
- Wilson, N. D., Ross, L. J. N., Close, J., Mott, R., Crow, T. J., and Volpi, E. V. (2007). Replication profile of PCDH11X and PCDH11Y, a gene pair located in the non-pseudoautosomal homologous region Xq21.3/Yp11.2. *Chromosome Res. Int. J. Mol. Supramol. Evol. Aspects Chromosome Biol.* 15, 485–498. doi: 10.1007/s10577-007-1153-y
- Winick-Ng, W., and Rylett, R. J. (2018). Into the fourth dimension: dysregulation of genome architecture in aging and Alzheimer's disease. *Front. Mol. Neurosci.* 11, 60. doi: 10.3389/fnfmol.2018.00060
- Wiseman, F. K., Pulford, L. J., Barkus, C., Liao, F., Portelius, E., Webb, R., et al. (2018). Trisomy of human chromosome 21 enhances amyloid-beta deposition independently of an extra copy of APP. *Brain* 141, 2457–2474. doi: 10.1093/brain/awy159
- Wu, Z. C., Yu, J. T., Wang, N. D., Yu, N. N., Zhang, Q., Chen, W., et al. (2010). Lack of association between PCDH11X genetic variation and late-onset Alzheimer's disease in a Han Chinese population. *Brain Res.* 1357, 152–156. doi: 10.1016/j.brainres.2010.08.008
- Yuan, D., XiuJuan, W., Yan, Z., JunQin, L., Fang, X., Shirong, Y., et al. (2015). Use of X-chromosome inactivation pattern to analyze the clonality of 14 female cases of kaposi sarcoma. *Med. Sci. Monitor Basic Res.* 21, 116–122. doi: 10.12659/MSMBR.894089
- Yudowski, G. A., Olsen, O., Adesnik, H., Marek, K. W., and Bredt, D. S. (2013). Acute inactivation of PSD-95 destabilizes AMPA receptors at hippocampal synapses. *PLoS One* 8, e53965. doi: 10.1371/journal.pone.0053965
- Yurov, Y. B., Iourov, I. Y., Vorsanova, S. G., Liehr, T., Kolotii, A. D., Kutsev, S. I., et al. (2007). Aneuploidy and confined chromosomal mosaicism in the developing human brain. *PLoS One* 2, e558. doi: 10.1371/journal.pone.0000558
- Yurov, Y. B., Vorsanova, S. G., Liehr, T., Kolotii, A. D., and Iourov, I. Y. (2014). X chromosome aneuploidy in the Alzheimer's disease brain. *Mol. Cytogenet.* 7, 20. doi: 10.1186/1755-8166-7-20
- Yurov, Y. B., Vorsanova, S. G., and Iourov, I. Y. (2019). Chromosome instability in the neurodegenerating brain. *Front. Genet.* 10, 892. doi: 10.3389/fgene.2019.00892
- Yurov, Y. B. (2017). “FISH-Based Assays for Detecting Genomic (Chromosomal) Mosaicism in Human Brain Cells,” in *Genomic Mosaicism in Neurons and Other Cell Types*. Eds. J. M. Frade and F. Gage (London, UK: Springer Nature).
- Zivković, L., Spremo-Potparević, B., Plecas-Solarović, B., Djelić, N., Očić, G., Smiljković, P., et al. (2010). Premature centromere division of metaphase chromosomes in peripheral blood lymphocytes of Alzheimer's disease patients: relation to gender and age. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 65, 1269–1274. doi: 10.1093/gerona/glq148

- Zivkovic, L., Spremo-Potparevic, B., Siedlak, S. L., Perry, G., Plecas-Solarovic, B., Milicevic, Z., et al. (2013). DNA damage in Alzheimer disease lymphocytes and its relation to premature centromere division. *Neurodegener. Dis.* 12, 156–163. doi: 10.1159/000346114
- Zubenko, G. S., Stiffler, J. S., Hughes, H. B., Hurtt, M. R., and Kaplan, B. B. (1998). Initial results of a genome survey for novel Alzheimer's disease risk genes: association with a locus on the X chromosome. *Am. J. Med. Genet.* 81, 196–205. doi: 10.1002/(sici)1096-8628(19980207)81:1<98::aid-ajmg17>3.0.co;2-r
- Zubenko, G. S., Hughes, H. B., and Stiffler, J. S. (1999). Clinical and neurobiological correlates of DXS1047 genotype in Alzheimer's disease. *Biol. Psychiatry* 46, 173–181. doi: 10.1016/s0006-3223(99)00035-9

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Bajic, Essack, Zivkovic, Stewart, Zafirovic, Bajic, Gojobori, Isenovic and Spremo-Potparevic. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.