



## OPEN ACCESS

## EDITED BY

Sundeeep Kumar,  
Indian Council of Agricultural Research  
(ICAR), India

## REVIEWED BY

Balpreet Kaur Dhatt,  
University of Nebraska-Lincoln,  
United States  
Bahram Heidari,  
Shiraz University, Iran

## \*CORRESPONDENCE

Wang Xing,  
✉ xiaoyujiayou\_086@163.com

†These authors share first authorship

RECEIVED 01 April 2023

ACCEPTED 30 June 2023

PUBLISHED 21 July 2023

## CITATION

Li W, Lin M, Li J, Liu D, Tan W, Yin X, Zhai Y,  
Zhou Y and Xing W (2023), Genome-wide  
association study of drought tolerance  
traits in sugar beet germplasms at the  
seedling stage.  
*Front. Genet.* 14:1198600.  
doi: 10.3389/fgene.2023.1198600

## COPYRIGHT

© 2023 Li, Lin, Li, Liu, Tan, Yin, Zhai, Zhou  
and Xing. This is an open-access article  
distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is  
permitted, provided the original author(s)  
and the copyright owner(s) are credited  
and that the original publication in this  
journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Genome-wide association study of drought tolerance traits in sugar beet germplasms at the seedling stage

Wangsheng Li<sup>1,2†</sup>, Ming Lin<sup>3†</sup>, Jiajia Li<sup>1,2</sup>, Dali Liu<sup>1,2</sup>, Wenbo Tan<sup>1,2</sup>,  
Xilong Yin<sup>1,2</sup>, Yan Zhai<sup>1,2</sup>, Yuanhang Zhou<sup>3</sup> and Wang Xing<sup>1,2\*</sup>

<sup>1</sup>National Beet Medium-Term Gene Bank, Heilongjiang University, Harbin, China, <sup>2</sup>Key Laboratory of Sugar Beet Genetics and Breeding, College of Advanced Agriculture and Ecological Environment, Heilongjiang University, Harbin, China, <sup>3</sup>Xinjiang Academy of Agricultural Sciences, Urumqi, China

**Introduction:** Sugar beets are an important crop for global sugar production. Intense drought and the increasing lack of water resources pose a great threat to sugar beet cultivation. It is a priority to investigate favourable germplasms and functional genes to improve the breeding of drought tolerant plants.

**Methods:** Thus, in this study, 328 sugar beet germplasms were used in a genome-wide association study (GWAS) to identify single nucleotide polymorphism (SNP) markers and candidate genes associated with drought tolerance.

**Results:** The results showed that under drought stress (9% PEG-6000), there were 11 significantly associated loci on chromosomes 2, 3, 5, 7, and 9 from the 108946 SNPs filtered using a mixed linear model (MLM). Genome-wide association analysis combined with qRT-PCR identified 13 genes that were significantly differentially expressed in drought-tolerant extreme materials.

**Discussion:** These candidate genes mainly exhibited functions such as regulating sugar metabolism, maintaining internal environmental stability and participating in photosystem repair. This study provides valuable information for exploring the molecular mechanisms of drought tolerance and improvement in sugar beet.

## KEYWORDS

sugar beet germplasms, PEG-6000 drought stress, SNP, GWAS, linkage disequilibrium

## 1 Introduction

Sugar beet is an important sugar crop. It has strong drought tolerance in the middle and late growth period, but weak drought tolerance at seedling stage. Most regions in the world are facing the problem of drought, and the sugar beet plant area of Heilongjiang Province in China is no exception. In spring, there are many southerly winds, large evaporation and small precipitation. The development of drought-tolerant varieties becomes more and more important (Bloch et al., 2006), and the selection and utilization of excellent drought tolerant germplasm resources and drought tolerant genes play an important role in the utilization efficiency and the improvement of agricultural production in water-shortage areas.

Mapping of quantitative trait loci (QTLs) is an effective tool often used to reveal the genetic basis of complex quantitative traits in crops (Huang et al., 2006). However, using traditional molecular markers such as restriction fragment length polymorphism (RFLP)

and simple sequence repeats (SSR), only a few QTLs associated with drought stress can be identified (Jha et al., 2021; Dwiningsih et al., 2022). With the development of functional genomics and transcriptomics in recent years, a large number of genes have been found to be involved in plant drought tolerance, including protein kinases, transcription factors and some structural genes (Wang et al., 2021b). However, only a few genes have been functionally validated. It remains a major challenge to accurately and effectively screen more drought stress-related genes and use them to breed drought-tolerant varieties.

Drought tolerance in plants is a complex quantitative trait that is controlled by multiple genes and involves multiple physiological and biochemical metabolic pathways. Although QTL mapping is a powerful method for detecting genomic regions associated with complex traits, the genetic effects of QTLs may not exist or may simply not be tested in different genetic backgrounds and environments (Shariatipour et al., 2021). Compared with traditional QTL, genome-wide association study (GWAS) can utilize genome-wide single nucleotide polymorphisms (SNPs) as molecular markers to analyze the genetic basis of complex traits, assisted by high-throughput genotyping platforms (Zheng et al., 2021). With the continuous development of high-throughput sequencing and high-resolution metabolic assays, as well as the development of multiple bioinformatics techniques and statistical methods to provide a basis for precise localization of complex trait gene variants, genome-wide association analysis is playing an increasingly important role in mining genetic loci for research (Zheng et al., 2021). Significant markers associated with aboveground biomass, fruit size, lodging score and leaf elongation in beans were identified through a whole genome-wide association analysis of 96 common bean germplasms. Specifically, 7 significant markers were found under irrigated conditions, and 5 significant markers were found under water deficit conditions (Hoyos-Villegas et al., 2017). In 2019, Mathew et al. used genome-wide association analysis to identify 75 significant marker-trait associations for population structure and marker traits of wheat biomass traits under drought and non-stress conditions. A total of 37 presumptive candidate genes were screened through gene annotation on IWGSC RefSeq 1.1 (Mathew et al., 2019). In recent years, the genome-wide association method has been applied in many crop drought tolerance studies and achieved good results. However, the current research on drought tolerance of sugar beet mainly focus on physiological and biochemical responses. Although traditional breeding methods such as analysis of combinatorial ability of lines have been used for selection of superior sugar beet genotypes for sugar related traits (Hassani et al., 2020), GWAS provide more reliable information about markers linked with sugar trait for use in marker-assisted selection. We believe that using genome-wide association analysis in drought tolerance studies of beet germplasm resources can quickly and accurately excavate drought tolerance related genes. Therefore, in this study, 328 beet germplasm resources were simulated by PEG-6000 drought stress. It is hoped to explore significant loci and potential candidate genes related to drought tolerance of beet germplasm resources by genome-wide association analysis, and to provide reference and basis for further research on the isolation of related genes and molecular marker-assisted selection of beet drought tolerance.

## 2 Materials and methods

### 2.1 Test material

The material for this study was provided by the National Beet Medium-Term Gene Bank at Heilongjiang University. The 328 sugar beet germplasm resources used in the trial were from 17 countries (Supplementary Table S1). 200 from China, 9 from Japan, 1 from North Korea, 15 from Russia, 49 from the United States, 1 from Austria, 2 from Belgium, 10 from Poland, 1 from Denmark, 10 from Germany, 3 from France, 12 from the Netherlands, 2 from Romania, 5 from Sweden, 2 from Hungary, 4 from Italy, and 2 from the United Kingdom (Figure 1). This experiment was conducted in a sophisticated climate chamber with a constant environment of 25°C and 60% humidity during the day and 18°C and 60% humidity during the night. There were 4 rows for each genotype, and 6 plants in each row were cultured in Hoagland's solution. After growing one pair of true leaves, two of the four rows of germplasms were selected and treated with PEG-6000 (Livak and Schmittgen) at a concentration of 9% to simulate drought stress (Tan et al., 2023). The remaining two rows of germplasms continued to be cultured in Hoagland's solution as controls (CG). The solution was changed once every 3 days. The materials were sampled after the growth of three pairs of true leaves (Cai et al., 2015).


### 2.2 Phenotype identification and physiological trait assay

The following traits were measured in this test: embryonic axis diameter (stem diameter measured using Vernier callipers) (Yang et al., 2022b), maximum root length (a straightedge was used to measure the maximum length of the root), plant height (measured from the ground to the top of the plant using a straightedge), root fresh weight (the roots were removed, dried on paper towels, and weighed on an electronic balance), leaf fresh weight (all leaves were cut off and weighed on an electronic balance), root dry weight (roots were dried in a ventilated oven at 105°C for 15 min, then transferred to 75°C for drying to a standard weight) (Polash et al., 2018), and leaf dry weight (leaves were dried in a ventilated oven at 105°C for 15 min, then transferred to 75°C for drying to a standard weight) (Yang et al., 2022a). Related tools were Vernier callipers (0.01 mm, Brand: Sheffield, Model: S071012, Origin: Mainland China), electronic balance (0.00001 g, Brand: Mettler-Toledo GmbH, Model: MS105DU/A, Origin: Switzerland), and steel ruler (0.1 cm, Brand: Sheffield, Model: S079020, Origin: China Mainland). The relative leaf water was calculated as follows:

$$\text{Relative leaf water} = (\text{Leaf fresh weight} - \text{Leaf dry weight}) / (\text{Leaf saturated fresh weight} - \text{Leaf dry weight})$$

Superoxide dismutase activity was evaluated by the nitrogen blue tetrazolium method (Spitz and Oberley, 1989). Soluble sugar levels were evaluated by the Anthrone-colorimetric method (Maness, 2010). Proline levels were evaluated by the acidic ninhydrin method (Carillo and Gibon, 2011). Soluble protein levels were evaluated using a BCA Protein Assay Kit (Brand: biosharp, Model: BL521A-3, Origin: Beijing, China).

The drought tolerance coefficient (DTC) was calculated for each trait:



**FIGURE 1**  
328 sugar beet germplasm resources distribution map.

$$DTC_X = DT/NT$$

where  $DTC_X$  is the drought tolerance coefficient of each trait;  $DT$  is the mean value of the traits under drought treatment; and  $NT$  is the mean value of the traits under normal treatment.

## 2.3 Genotype data

**Sequencing:** SNPs were identified by NOVOGENE (Beijing, China) and sequenced using the Illumina HiSeq sequencing platform with double-end (Paired-End) 150 sequencing. BWA (Li and Durbin, 2009) (0.7.17): Interference information was filtered out of the raw data to obtain high-quality clean data, and effective high-quality sequencing data were aligned to the reference genome by BWA software (parameter: mem-t 4-k 32-M) (reference genome download address: [ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/511/025/GCF\\_000511025.2\\_RefBeet-1.2.2/GCF\\_000511025.2\\_RefBeet-1.2.2\\_genomic.fna.gz](ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/511/025/GCF_000511025.2_RefBeet-1.2.2/GCF_000511025.2_RefBeet-1.2.2_genomic.fna.gz)). Samtools (Soskic et al., 2019) (1.9): Samtools was used to transform the format of the sam file and build an index to generate the bai file. GATK (Sun et al., 2019) (4.2.6.1): GATK was used to detect mutations and generate group modification.vcf file. VCFtools (Ratnapriya et al., 2019) (0.1.17): A total of 108946 high-quality loci were screened for subsequent analysis using sequencing depth ( $dp \geq 4$ ), minor allele frequency ( $maf \geq 0.05$ ) and miss rate ( $miss \leq 80\%$ ).

## 2.4 Population structure and LD analysis

VCFtools (0.1.17) was first used to convert the.vcf files ped files and then plink (1.9) software was used to convert ped format to bed format (Cadzow et al., 2014). Population structure analysis of bed files was performed using admixture (1.3.0) software, the population size  $K$  value was preset to 1–10 for classification,

and the minimum value of cross entropy was used to determine the optimal number of classifications (Byrne et al., 2020). Principal component analysis of 108946 high-quality SNP loci was performed using Tassel (5.2.82) software (Brerria et al., 2020), and the PCA results were visualized using R (4.1.0) software. The LD of 328 sugar beet germplasms was analysed by PopLDdecay (3.4.2). The LD intensity ( $r^2$ ) between two SNPs within a certain distance was calculated (Xu et al., 2018).


## 2.5 Genome-wide association study

The kinship analysis was performed using Tassel (5.2.82) software to obtain the kinship matrix (Zeng et al., 2017) and the results were visualized using the pheatmap package in R (4.1.0). The mixed linear model (MLM) (Kang et al., 2015) of Tassel software was used to analyse the PCA and kinship analysis results as covariates. This combination ( $Q + K$ ) is often considered the most powerful, and this model can classify crops into different subgroups based on growth habit or geographical origin. Mixed models corrected with population structure and kinship can effectively reduce false positive results (Wang et al., 2007). Manhattan and Q-Q plots were visualized using the CMplot package of R (4.0.1) (Kumar et al., 2021). The mixed linear model (MLM) equation was as follows:

$$Y = X\alpha + Q\beta + K\mu + e$$

In the equation,  $Y$  is the vector of measured phenotypes,  $X$  is the fixed effect of SNPs,  $P$  is the fixed effect of population structure,  $K$  is the random effect of kinship, and  $e$  is the random error (Dramadri et al., 2021).

We used the powerful annotation function of the internationally recognized UniProt database (Yang and Wang, 2015) to annotate the SNPs. After screening fake SNPs according to genome



**FIGURE 2**  
Comparison of various traits between drought treatment and normal water supply for 328 sugar beet germplasms. Note: NT, normal treatment; DT, drought treatment, 9% PEG-6000 for 7 days.

**TABLE 1 Analysis of variation in drought tolerance coefficient of 328 sugar beet germplasm for each trait.**

Traits	Mean	STDEV	Max	Min	Skewness	Kurtosis	CV(%)
EAD	0.708	0.243	1.676	0.171	0.718	0.848	34.38
PH	0.651	0.154	1.197	0.259	0.506	0.586	23.58
RL	0.914	0.215	1.659	0.475	0.549	0.238	23.55
LFW	0.236	0.188	1.042	0.005	1.411	2.112	79.58
RFW	0.482	0.322	1.879	0.015	1.269	1.87	66.73
RDW	0.634	0.406	2.333	0.022	1.261	1.923	64.16
LDW	0.394	0.278	1.684	0.019	1.341	2.74	70.62
RLW	0.799	0.130	1.226	0.267	-1.062	2.581	16.29
SS	1.611	1.225	6.943	0.152	1.725	3.249	76.03
SOD	1.023	0.263	1.994	0.204	0.532	1.409	25.72
Pro	1.156	0.319	2.185	0.362	0.539	0.446	27.57
SP	1.130	0.415	2.715	0.006	0.831	0.928	36.70

Note: EAD, embryonic axis diameter; PH, plant height; RL, maximum root length; LFW, leaf fresh weight; RFW, root fresh weight; RDW, root dry weight; LDW, leaf dry weight; RLW, relative leaf water; SS, soluble sugar levels; SOD, superoxide dismutase activity; Pro, Proline levels; SP, soluble protein levels.



annotation, we obtained high-quality and significant SNPs with a threshold of  $-\log_{10}p > 6$ .

The SNPs were screened to obtain high-quality and significantly correlated SNPs with a threshold of  $-\log_{10}p > 6$ . The size of the screened SNP interval was estimated based on LD block analysis, and genes in the region near the locus were obtained and functionally annotated. Based on the functional annotations, we could roughly determine which genes might be

associated with the traits of interest and use in subsequent analysis (Bai et al., 2022).


## 2.6 Real-time fluorescence quantitative PCR

Four drought-tolerant (Mono HY53, Neitang578, Gansunongjiazhong, BGRC16137) and four drought-sensitive




germplasms [7501A/BCE, 92017/1-8/1, Tianyanshao (X), 92017/1-4/1] were used as extreme materials to extract total RNA according to the Trizol method. cDNA was synthesized according to the TransScript One-Step gDNA Removal and cDNA Synthesis SuperMix protocol (AT311, Beijing All Style Gold Biotechnology Co., Ltd.). qPCR was performed using the SuperReal PreMix Plus kit (Beijing Tiangen Biochemical Technology Co., Ltd., version FP210831) on a real-time

fluorescent quantitative PCR instrument (Thermo Fisher Scientific Instruments, Shanghai. QuantStudio™ 1 Plus) (Liu et al., 2023). Expression level of 14 genes was analyzed with the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001). *BvGAPDH* (NC\_024800) was used as an internal control to standardize the expression levels of the different samples. All assays were performed in two independent experiments and replicated three times. Primers were summarized in Supplementary Table S5.



**FIGURE 5**  
Kinship heatmap and cluster analysis.




**FIGURE 6**  
Manhattan plot: inner circles indicate single nucleotide polymorphisms (SNPs) associated with the Q-Q plot on the left, outer circles indicate single nucleotide polymorphisms (SNPs) associated with the Q-Q plot on the right, grey dashed lines indicate SNP loci with  $-\log_{10}p = 4.5$ , red dashed lines indicate SNP loci with  $-\log_{10}p = 6$ , and the outermost side indicates SNP density on the chromosome. Q-Q plots: horizontal coordinates indicate expected values, vertical coordinates indicate observed values, and grey areas are 95% confidence intervals.

### 3 Results

#### 3.1 Phenotypic and physiological analysis

The statistical results showed that the measured data were all close to a normal distribution, indicating that the 12 traits were all

typical quantitative traits. Eight traits, including embryonic axis diameter (EAD), plant height (PH), root length (RL), leaf fresh weight (LFW), root fresh weight (RFW), root dry weight (RDW), leaf dry weight (Lander et al., 2001) and leaf relative water content (RLW), were significantly lower in the drought-treated plants than in the control plants. Three indicators, including soluble sugar levels



(SS), soluble protein levels (SP) and free proline levels (Pro), were significantly higher in the drought-treated plants than in the control plants (Figure 2). Drought tolerance coefficients for each trait showed that these traits all exhibited suitable variation, indicating that they were controlled by small effect polygenes with coefficients of variation (CV) ranging from 16.29% to 79.58% (Table 1). The CV is an absolute value reflecting the degree of dispersion of each trait in the population. To better estimate the correlations between markers and traits, it is desirable to use highly dispersed trait data for statistical analysis. The traits in this test exhibited a high degree of dispersion and were suitable for genome-wide association analysis.

## 3.2 Population structure and genetic diversity

Population stratification refers to the existence of subgroups within a population, where the diversity among individuals within the subgroups is greater than the average diversity among individuals within the entire population (Wang et al., 2021a). Different allele frequencies at certain loci between different subpopulations can lead to false positive results when using two subpopulations for association analysis (Zheng et al., 2021). Therefore, before performing association analysis, we used a subset of 108946 SNPs without close linkage for population structure prediction. Using Tassel (5.2.82) software to construct phylogenetic trees by the neighbor-joining (NJ) method, 328 sugar beet germplasms were divided into three subgroups, but the differences between the subgroups were small (Figures 3A, D). The results from a principal component analysis (PCA) of the above three subgroups were consistent with the phylogenetic analysis, and no significant grouping was found (Figure 3E). The

results indicated that the 328 germplasms without significant grouping were suitable for a GWAS. Using admixture software, the number of subgroups (K) was set to 1–10, and each K value was repeated three times to select the optimal K value according to the maximum likelihood method. The cross-validation error (CV error) value was the smallest when  $K = 7$  (Figure 3B), so the Q-matrix with  $K = 7$  was used as a covariate to improve the accuracy of the GWAS (Figure 3C).

A linkage disequilibrium (LD) analysis was carried out, and the LD distance decreased as the physical location of the SNPs on chromosomes gradually increased. The LD distance of 50 kb was equal to half of the maximum value (Figure 4).

The visualization of the kinship matrix showed that the variation between most of the 328 genotypes was low (light blue), and the genetic similarity between individual genotypes was high. The low genetic variation in this study would reduce the occurrence of false positive results (Figure 5).

## 3.3 GWAS of drought tolerance coefficients for each trait



The GWAS results showed (Figure 6; Supplementary Table S2) that a total of 183 SNPs ( $-\log_{10}p > 4.5$ ) were identified in the 12 drought tolerance-related traits, including 11 SNPs with peaks  $>6.0$ , which the study mainly focused on. It was found that 1 SNP on chromosome 7 was associated with DTC\_RLW; 2 SNPs associated with DTC\_LFW were located on chromosomes 2 and 5; 1 SNP on chromosome 3 was associated with DTC\_LDW; 2 SNPs on chromosome 9 were associated with DTC\_RDW; and there were 5 SNPs associated with DTC\_SS on chromosomes 2 and 7. It is worth noting that the two significant loci associated with DTC\_RDW were close to each other with a distance of only 216 bp.

## 3.4 Candidate genes associated with sugar beet drought tolerance

In the five traits ( $-\log_{10}p > 6$ ) of DTC\_RLW, DTC\_LFW, DTC\_LDW, DTC\_RDW, and DTC\_SS, 50 kb upstream and downstream of the significant loci were analyzed by LD decay. A total of 24 genes in sugar beet associated with drought tolerance were identified and annotated (Supplementary Table S3).

Two genes (*BVRB\_7g160020* and *BVRB\_7g160030*) were identified on chromosome 7 at position 4,745,224 may be associated with DTC\_RLW (Figure 7; Supplementary Table S3). *BVRB\_7g160020* encodes a tyrosine-protein phosphatase, RLP2, which is reported to be a new phospho-tyrosine-specific phosphatase belonging to the phosphoprotein phosphatase (PPP) family (Uhrig et al., 2016). *BVRB\_7g160030*, as 1,4-alpha-glucan-branching enzyme 2-2, is mainly involved in amylose synthesis in gluconeogenesis (Han et al., 2022). It was reported that *BVRB\_7g160030* catalytically regulates  $\alpha$  (1–6) glycosidic bond branching synthesis. We believe that *BVRB\_7g160030* is a potential target gene for the drought stress response in sugar beet.


Associated with DTC\_LFW, three genes (*BVRB\_2g033710*, *BVRB\_2g033720* and *BVRB\_2g033730*) were found on chromosome 2. Two genes (*BVRB\_5g114800* and *BVRB\_*



5g114810) were found on chromosome 5. *BVRB\_2g033720*, *BVRB\_2g033730* and *BVRB\_5g114800* were of unknown function (Figure 8; Supplementary Table S3). Through annotation, we found that *BVRB\_2g033710* functions as myosin-15; *BVRB\_*

5g114810 functions as MYB-related protein 306, the MYB structural domain is a peptide segment of approximately 51-52 amino acids containing a series of highly conserved amino acid residues and spacer sequences, and the MYB gene family is






widely involved in plant metabolic regulation (Jiao et al., 2021; Johnson et al., 2022).

Two genes (*BVRB\_3g048910* and *BVRB\_3g048900*) were identified on chromosome 3 position: 998,024, may be associated with DTC\_LDW (Figure 9; Supplementary Table S3). By annotation, we found that *BVRB\_3g048910* functions as the probable methyltransferase PMT3. Methyltransferases use S-adenosylmethionine, betaine and dimethylthetin as methyl donors to generate methionine. *BVRB\_3g048900* functions as serine/threonine-protein kinase tricornered (TOR), which functions as a phosphate donor using ATP to phosphorylate serine and threonine residues on proteins to regulate abscisic acid, growth hormone, glucose and sucrose-mediated signalling (Ahn et al., 2011; Xiong et al., 2013; Deng et al., 2016).

Two close significant loci (position: 1,172,900, position: 1,173,116) on chromosome 9 were found to be associated with DTC\_RDW (Figure 10; Supplementary Table S3). Six gene were found in the region near these two significant loci (*BVRB\_9g203050*, *BVRB\_9g203040*, *BVRB\_9g203030*, *BVRB\_9g203020*, *BVRB\_9g203010* and *BVRB\_9g203000*). *BVRB\_9g203050* functions as photosystem II (PSII) stability/assembly factor HCF136 and is essential for PSII biogenesis. *BVRB\_9g203030* functions as eukaryotic initiation factor 4A-9, which promotes the hydrolysis of ATP and drives RNA deconvolution against abiotic stress; *BVRB\_9g203000* functions as enhancer of AG-4 protein 2, which possesses the dual ability to induce apoptosis and autophagy (Kumar et al., 2015), and is a transcription factor that acts as a flowering repressor by enhancing the expression of genes that delay flowering and suppressing nutritional growth to inhibit reproductive development (Wang et al., 2007).




Two gene regions (*BVRB\_2g036960* and *BVRB\_2g036950*) were found to be associated with DTC\_SS (Figure 11; Supplementary Table S3) in the region near positions 20,553,998 on chromosome 2. *BVRB\_2g036950* is an uncharacterized protein, and *BVRB\_2g036960* functions as a probable membrane-associated kinase regulator 4. Tyrosine phosphorylation controls the activation of brassinosteroid receptors through membrane-released protein kinase inhibitors.


Four SNP loci significantly associated with DTC\_SS (Figure 12; Supplementary Table S3) were found on chromosome 7. No genes were found near position 15,215,427. Two genes (*BVRB\_7g163640* and *BVRB\_7g163650*) were found near position 15,170,171 and they are hypothetical and uncharacterized proteins. Three genes (*BVRB\_7g166470*, *BVRB\_7g166480* and *BVRB\_7g166490*) were found near position 27,948,364, and *BVRB\_7g166470* functions as glycine-rich cell wall structural protein 1, a process that results in the assembly, arrangement of constituent parts, or disassembly of the cell wall. *BVRB\_7g166480* functions as PROTEIN LOW PSII ACCUMULATION 1, which is mainly involved in the efficient assembly of photosynthetic system II. *BVRB\_7g166490* functions as photosystem I reaction centre subunit IV. Two gene (*BVRB\_7g166670* and *BVRB\_7g166680*) were found near positions 28,950 and *BVRB\_7g166670* is an uncharacterized protein, and *BVRB\_7g166680* functions as a ras GTPase-activating protein-binding protein.

### 3.5 qRT-PCR validation of candidate gene expression

By examining the overlap of the identified candidate genes with known drought stress response genes in other crop species, we



**FIGURE 12** Manhattan plot of candidate gene regions associated with DTC\_SS and heatmap of linkage disequilibrium (bottom). The orange vertical line indicates the position of the most significant SNP, and the orange horizontal line indicates  $-\log_{10}p = 6$ .



**FIGURE 13** Expression levels of candidate genes in drought-tolerant extreme materials. Note: (A–D) are drought-sensitive materials. (E–H) are drought-tolerant materials. \*: Indicates that the expression levels of the two groups of materials are different at the 0.05 level. \*\*: Indicates that the expression levels of the two groups of materials are different at the 0.01 level.

identified 14 genes that may be involved in the drought tolerance process in sugar beet. In order to validate the GWAS results, qRT-PCR was performed on these genes (As shown in Figure 13). The results showed that, except for *BVRB\_2g036960*, the other 13 genes were all significantly differently expressed between the drought tolerant and sensitive groups. The expression levels of *BVRB\_7g160020* and *BVRB\_5g114810* were 3.30-fold and 2.57-fold higher in the drought-sensitive group than them in the drought-tolerant group, respectively. *BVRB\_7g160030*, *BVRB\_2g033710*, *BVRB\_3g048910*, *BVRB\_3g048900*, *BVRB\_9g203050*, *BVRB\_9g203030*, *BVRB\_9g203000*, *BVRB\_7g166680*, *BVRB\_7g166470* and *BVRB\_7g166490* in drought-tolerant group were expressed 1.47–4.23-fold of them in the drought-sensitive group. Notably,

the expression level of *BVRB\_7g166480* in the drought-tolerant group was 46.69-fold higher than that of the drought-sensitive group.

## 4 Discussion

### 4.1 Drought tolerance-related traits in sugar beet

Drought stress is the main abiotic stress factor that causes crop yield reduction worldwide. The sugar beet plant is more drought tolerant in the middle and late growth stages but is drought sensitive

in the seedling stage, which causes great taproot loss (Fugate et al., 2018). Enhancing the drought tolerance of seedlings will greatly help to improve the sugar beet crop yield. Sadeghian and Yavari conducted mannitol-simulated drought stress on 9 sugar beet varieties, assessed the dry weight of cotyledons, fresh weight of cotyledons, RFW, RDW, and RL, and concluded that the drought tolerance traits of sugar beet seedlings can be used for selective breeding of the strain with the strongest tolerance (Sadeghian and Yavari, 2004). Islam et al. (2020) screened 7 beet varieties suitable for drought conditions by carrying out drought stress experiments on 11 varieties and measuring related physiological and biochemical traits. Through stepwise regression analysis, Moosavi et al. (2017) found that root and petiole dry weight were suitable as standard traits for drought tolerance of sugar beet, and 7221 drought tolerance genotypes were screened by these traits.

Although breeding by screening drought-tolerant varieties of sugar beet can improve drought tolerance in new varieties, this method has some limitations. The transfer of agronomic traits is easily limited by intermediate reproductive isolation, which is not conducive to genetic improvement of selected crops using genetic resources of closely or distantly related species. This study aims to improve the efficiency and precision of drought-tolerant breeding of sugar beet through research methods at the molecular level.

## 4.2 Genome-wide association analysis of drought tolerance-related traits in sugar beet

With the development of genotyping technology, SNP markers are characterized by high throughput, low cost and high accuracy. The application of genome-wide association analysis is universal, and it can be used to analyze both qualitative and quantitative traits (Alqudah et al., 2020). In 175 rice samples, 13 significant SNPs were screened out and 50 genes were identified, and 10 of these genes were associated with drought and/or other abiotic stress tolerance (Pantalião et al., 2016). It was detected 132 single nucleotide polymorphisms in sesame under drought stress, and 13 potential candidate genes were found to be related with drought tolerance (Li et al., 2018). At present, GWAS has been reported in the drought tolerance studies of many crops, but it is rarely reported in sugar beet. In this study, we explored the functional genes related to drought tolerance of sugar beet through genome-wide association analysis. 328 sugar beet germplasms were analyzed, and phylogenetic tree and PCA analysis showed that there was no obvious population stratification, indicating that the heredity of sugar beet used here was less affected by geographical factors. Considering that MLM model can better control false positive results, we used this model to excavate significant sites (Zhang et al., 2010).

## 4.3 Potential candidate genes associated with sugar beet drought tolerance

A total of 11 SNPs loci were identified by genome-wide association analysis and 25 potential candidate genes were screened out. It was found that 14 genes had been reported to be related with drought tolerance. We conducted qRT-PCR to verify

gene expression levels in the two groups of drought tolerance extreme materials and found that only *BVRB\_2g036960* had no difference in expression between the two groups of materials, and the expression levels of 2 genes in the drought sensitive group were significantly higher than those in the drought tolerant group. The expression levels of 11 genes in drought-tolerant group were significantly higher than those in drought-sensitive group. Therefore, these 13 differentially expressed genes might be key genes which can be used in our next research on drought tolerance in sugar beet.

*BVRB\_7g160020* regulated RLP2 is a tyrosine phosphorylase of the PPP family, and tyrosine phosphorylation is essential for signaling at the cytoplasmic structural domain (Mühlenbeck et al., 2021). We also found that the gene *BVRB\_7g160030*, which is adjacent to the gene *BVRB\_7g160020*, is involved in straight-chain starch synthesis in sugar metabolism.  $\alpha$  (1–6) glycosidic bond branching increases the water solubility of glycogen for storage and increases the number of nonreducing ends, making it easier for biological organisms to aggregate when glycogen is needed for energy supply, and the 1,4- $\alpha$ -glucan-branching enzyme regulates glycogen metabolism and participates in the abiotic stress response (Li et al., 2019).

We found that *BVRB\_7g166470* is involved in the cell wall composition. *BVRB\_3g048910* can regulate methionine, which can mitigate membrane damage by peroxides. It has been shown that the probable methyltransferase PMT3 is involved in the synthesis of phosphatidylcholine (PC), which is an important precursor of lipid signalling (Baldessari and Iglesias, 2012) and a ligand for regulatory proteins (Inatsugi et al., 2009). PC appears to be a major contributor to the adaptive response of plants to environmental stress, and it plays an important role in maintaining cell membrane integrity in complex plant species. The ras GTPase-activating protein-binding protein 2 associated with *BVRB\_7g166680* regulates ascorbic acid, which has strong reducing properties that effectively mitigate peroxide damage (Khazaei et al., 2020). *BVRB\_9g203030* is eukaryotic initiation factor 4A-9, which promotes ATP hydrolysis to provide energy for material transport and signaling in plants, and myosin, which is a component of the cytoskeleton to provide energy for organelle movement, material transport and apical growth (Han et al., 2021). *BVRB\_2g033710* is involved in the components of the cytoskeleton that provide energy for organelle movement, material transport and apex growth (Han et al., 2022). *BVRB\_3g048900* on chromosome 3 was found to regulate abscisic acid, and *BVRB\_5g114810* on chromosome 5 was related to MYB transcription factors. Abscisic acid can regulate MYB, which can promote lateral root growth, enhance root water absorption, and induce old leaf death to reduce water loss under drought conditions (Moosavi et al., 2017). The ENHANCER OF AG-4 protein regulated by *BVRB\_9g203000* can induce apoptosis and prolong nutritional growth (Moosavi et al., 2017; Alqudah et al., 2020).

*BVRB\_9g203050* on chromosome 9 regulates HCF136, which is essential for PSII biogenesis and stability (Meurer et al., 1998); the HCF136 protein is essential for the formation of early intermediates in PSII assembly, including D2 (psbD) and cytochrome b559 (Plücken et al., 2002). D2 can repair PSII and D1 expression dysfunction and maintain the stability of photosynthesis (Kanamaru and Tanaka, 2004). Protein LOW PSII ACCUMULATION (*BVRB\_7g166480*), a required chaperone for

efficient PSII assembly, is a component of the chloroplast cyst-like membrane; it binds to psbA during the biogenesis of PSII and plays an important role in repairing PSII damage to maintain photosystem stability (Peng et al., 2006). Photosystem I (PSI) reaction centre subunit IV (BVRB\_7g166490) is capable of stabilizing the interaction between PsaC and the PSI core, assisting in the docking of iron oxidation reduction protein to PSI and interacting with iron oxidation reduction protein-NADP oxidoreductase (Dutta et al., 2014).

#### 4.4 Potential impact on sugar beet breeding

Although conventional breeding can improve the performance of sugar beet in drought tolerance, artificial populations are limited by population size, insufficient recombination, and often large locus intervals, making selection both time-consuming and inefficient (He et al., 2010). Most of the natural populations used here have been reproduced over long periods of time and multiple generations, with adequate genomic recombination and accurate linkage localization, and these new genomic resources will facilitate genomic evolution in sugar beet and provide a scientific and theoretical basis for relevant research in drought tolerance.

### 5 Conclusion

In this study, 11 loci potentially associated with drought tolerance in sugar beet were screened using a mixed linear model (MLM). Genome-wide association analysis combined with qRT-PCR revealed that 13 genes were significantly differentially expressed between the two extreme groups of materials, with *BVRB\_7g166480* being 46.69-fold more expressed in the drought-tolerant group than in the drought-sensitive group. It prompted us to further investigate the role of these interesting genes in sugar beet drought tolerance improvement in the future.

### Data availability statement

The datasets presented in this study can be found in online repositories. The sequencing data has been uploaded. The link is “<https://www.ncbi.nlm.nih.gov/sra/PRJNA935487>”. The accession no. is PRJNA935487.

### References

- Ahn, C. S., Han, J.-A., Lee, H.-S., Lee, S., and Pai, H.-S. (2011). The PP2A regulatory subunit Tap46, a component of the TOR signaling pathway, modulates growth and metabolism in plants. *Plant Cell*. 23 (1), 185–209. doi:10.1105/tpc.110.074005
- Alqudah, A. M., Sallam, A., Baenziger, P. S., and Börner, A. (2020). Gwas: Fast-forwarding gene identification and characterization in temperate cereals: Lessons from barley—a review. *J. Adv. Res.* 22, 119–135. doi:10.1016/j.jare.2019.10.013
- Bai, X., Wang, X., Wang, Y., Wei, Y., Fu, Y., Rao, J., et al. (2022). Genome-wide association study of six forage traits in ramie (*Boehmeria nivea* L. Gaud). *Plants* 11 (11), 1443. doi:10.3390/plants11111443
- Baldessari, A., and Iglesias, L. (2012). In *Lipases and phospholipases, methods and protocols*. Editor G. Sandoval (New York: Humana Press), 457–470.
- Bloch, D., Hoffmann, C. M., and Märlander, B. (2006). Impact of water supply on photosynthesis, water use and carbon isotope discrimination of sugar beet genotypes. *Eur. J. Agron.* 24 (3), 218–225. doi:10.1016/j.eja.2005.08.004
- Breteria, C. M., Hsieh, C. H., Yen, J.-Y., Nair, R., Lin, C.-Y., Huang, S.-M., et al. (2020). Population structure of the world vegetable center mungbean mini core collection and genome-wide association mapping of loci associated with variation of seed coat luster. *Trop. Plant Biol.* 13 (1), 1–12. doi:10.1007/s12042-019-09236-0
- Byrne, R. P., van Rheenen, W., van den Berg, L. H., Veldink, J. H., and McLaughlin, R. L. (2020). Dutch population structure across space, time and GWAS design. *Nat. Commun.* 11 (1), 4556. doi:10.1038/s41467-020-18418-4
- Cadzow, M., Boocock, J., Nguyen, H. T., Wilcox, P., Merriman, T. R., and Black, M. A. (2014). A bioinformatics workflow for detecting signatures of selection in genomic data. *Front. Genet.* 5, 293. doi:10.3389/fgene.2014.00293

### Author contributions

JL designed the research. WL analyzed the physiology data. DL and WT drafted the article. XY and YAZ performed the experiment. WX revised the article. ML and YUZ were tested for gene expression. All authors contributed to the article and approved the submitted version.

### Funding

The National Sugar Industry Technology System Project (CARS-170102), the Inner Mongolia Autonomous Region “the open competition mechanism to select the best candidates” project entitled “Creation of Elite Beet Germplasm and Breeding of Varieties Suitable for Mechanized Operation” (2022JBG0029), the Accurate Identification of Sugar Beet Germplasm Resources (19230709), the Innovative Training Plan for Young Talents in Heilongjiang Ordinary Undergraduate Colleges and Universities (UNPYSCT-2020) and Tianchi Talent Program.

### Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

### Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2023.1198600/full#supplementary-material>

- Cai, S., Jiang, G., Ye, N., Chu, Z., Xu, X., Zhang, J., et al. (2015). A key ABA catabolic gene, OsABA8ox3, is involved in drought stress resistance in rice. *PLoS One* 10 (2), e0116646. doi:10.1371/journal.pone.0116646
- Carillo, P., and Gibon, Y. (2011). Protocol: Extraction and determination of proline. *PrometheusWiki* 2011, 1–5.
- Deng, K., Yu, L., Zheng, X., Zhang, K., Wang, W., Dong, P., et al. (2016). Target of rapamycin is a key player for auxin signaling transduction in Arabidopsis. *Front. Plant Sci.* 7, 291. doi:10.3389/fpls.2016.00291
- Dramadri, I. O., Nkalubo, S. T., Kramer, D. M., and Kelly, J. D. (2021). Genome-wide association analysis of drought adaptive traits in common bean. *Crop Sci.* 61 (5), 3232–3253. doi:10.1002/csc2.20484
- Dutta, S., Teresinski, H. J., and Smith, M. D. (2014). A split-ubiquitin yeast two-hybrid screen to examine the substrate specificity of atToc159 and atToc132, two Arabidopsis chloroplast preprotein import receptors. *PLoS One* 9 (4), e95026. doi:10.1371/journal.pone.0095026
- Dwiningsih, Y., Thomas, J., Kumar, A., Gupta, C., Gill, N., Ruiz, C., et al. (2022). Identification of QTLs and candidate loci associated with drought-related traits of the K/Z RIL rice population.
- Fugate, K. K., Lafta, A. M., Eide, J. D., Li, G., Lulai, E. C., Olson, L. L., et al. (2018). Methyl jasmonate alleviates drought stress in young sugar beet (*Beta vulgaris* L) plants. *J. Agron. Crop Sci.* 204 (6), 566–576. doi:10.1111/jac.12286
- Han, H., Verstraeten, I., Roosjen, M., Mazur, E., Rýdza, N., Hajný, J., et al. (2021). Rapid auxin-mediated phosphorylation of Myosin regulates trafficking and polarity in Arabidopsis. *BioRxiv*. doi:10.1101/2021.04.13.439603
- Han, M., Xu, M., Su, T., Wang, S., Wu, L., Feng, J., et al. (2022). Transcriptome analysis reveals critical genes and pathways in carbon metabolism and ribosome biogenesis in poplar fertilized with glutamine. *Int. J. Mol. Sci.* 23 (17), 9998. doi:10.3390/ijms23179998
- Hassani, M., Heidari, B., and Stevanato, P. (2020). Combining abilities of sugar beet genotypes for root- and sugar-related traits under multi-environmental trials. *Plant Breed.* 139 (1), 192–206. doi:10.1111/pbr.12755
- He, Z., Xia, X., Bonjean, A. P., Li, F., Wang, B., and Wang, J. (2010). The potential of carcinoembryonic antigen, p53, Ki-67 and glutathion Stransferase- $\pi$  as clinico-histopathological markers for colorectal cancer. *Cereals China* 688 (787), 51–57. doi:10.1016/S1674-8301(10)60008-5
- Hoyos-Villegas, V., Song, Q., and Kelly, J. D. (2017). Genome-wide association analysis for drought tolerance and associated traits in common bean. *Plant Genome* 10 (1), 1–17. doi:10.3835/plantgenome2015.12.0122
- Huang, X., Cloutier, S., Lycar, L., Radovanovic, N., Humphreys, D., Noll, J., et al. (2006). Molecular detection of QTLs for agronomic and quality traits in a doubled haploid population derived from two Canadian wheats (*Triticum aestivum* L). *Theor. Appl. Genet.* 113, 753–766. doi:10.1007/s00122-006-0346-7
- Inatsugi, R., Kawai, H., Yamaoka, Y., Yu, Y., Sekiguchi, A., Nakamura, M., et al. (2009). Isozyme-specific modes of activation of CTP: Phosphorylcholine cytidyltransferase in Arabidopsis thaliana at low temperature. *Plant Cell. Physiology* 50 (10), 1727–1735. doi:10.1093/pcp/pcp115
- Islam, M. J., Kim, J. W., Begum, M. K., Soheli, M. A. T., and Lim, Y.-S. (2020). Physiological and biochemical changes in sugar beet seedlings to confer stress adaptability under drought condition. *Plants* 9 (11), 1511. doi:10.3390/plants9111511
- Jha, U. C., Nayyar, H., Palakurthi, R., Jha, R., Valluri, V., Bajaj, P., et al. (2021). Major QTLs and potential candidate genes for heat stress tolerance identified in chickpea (*Cicer arietinum* L). *Front. Plant Sci.* 12, 655103. doi:10.3389/fpls.2021.655103
- Jiao, Z., Xu, W., Nong, Q., Zhang, M., Jian, S., Lu, H., et al. (2021). An integrative transcriptomic and metabolomic analysis of red pitaya (*Hylocereus polyrhizus*) seedlings in response to heat stress. *Genes* 12 (11), 1714. doi:10.3390/genes12111714
- Johnson, A. A., Venkatesh, B., Jayamma, N., Reddy, B. M., Pandurangiah, M., and Sudhakar, C. (2022). R2R3 MYB transcription factor, AhMYB94 plays a crucial role in stress tolerance of a salt susceptible groundnut cultivar.
- Kanamaru, K., and Tanaka, K. (2004). Roles of chloroplast RNA polymerase sigma factors in chloroplast development and stress response in higher plants. *Biosci. Biotechnol. Biochem.* 68 (11), 2215–2223. doi:10.1271/bbb.68.2215
- Kang, Y., Sakiroglu, M., Krom, N., Stanton-Geddes, J., Wang, M., Lee, Y. C., et al. (2015). Genome-wide association of drought-related and biomass traits with HapMap SNPs in Medicago truncatula. *Plant, Cell. & Environ.* 38 (10), 1997–2011. doi:10.1111/pce.12520
- Khazaei, Z., Esmailpour, B., and Estaji, A. (2020). Ameliorative effects of ascorbic acid on tolerance to drought stress on pepper (*Capsicum annum* L) plants. *Physiology Mol. Biol. Plants* 26 (8), 1649–1662. doi:10.1007/s12298-020-00846-7
- Kumar, D., Das, B., Sen, R., Kundu, P., Manna, A., Sarkar, A., et al. (2015). Andrographolide analogue induces apoptosis and autophagy mediated cell death in U937 cells by inhibition of PI3K/Akt/mTOR pathway. *PLoS One* 10 (10), e0139657. doi:10.1371/journal.pone.0139657
- Kumar, D., Sharma, S., Sharma, R., Pundir, S., Singh, V. K., Chaturvedi, D., et al. (2021). Genome-wide association study in hexaploid wheat identifies novel genomic regions associated with resistance to root lesion nematode (*Pratylenchus thornei*). *Sci. Rep.* 11 (1), 3572. doi:10.1038/s41598-021-80996-0
- Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., et al. (2001). Initial sequencing and analysis of the human genome. *Nature* 409 (6822), 860–921. doi:10.1038/35057062
- Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25 (14), 1754–1760. doi:10.1093/bioinformatics/btp324
- Li, D., Dossa, K., Zhang, Y., Wei, X., Wang, L., Zhang, Y., et al. (2018). GWAS uncovers differential genetic bases for drought and salt tolerances in sesame at the germination stage. *Genes* 9 (2), 87. doi:10.3390/genes9020087
- Li, X., Ding, M., Wang, M., Yan, F., Zhang, Z., Liang, W., et al. (2019). Gene cloning of 1,4- $\alpha$ -glucan branching enzyme (GBE) from nostoc flagelliforme and its differential expression in response to drought stress. *Mol. Plant Breed.* 17 (19), 6305–6313.
- Liu, D., Gao, Z., Li, J., Yao, Q., Tan, W., Xing, W., et al. (2023). Effects of cadmium stress on the morphology, physiology, cellular ultrastructure, and BvHIPP24 gene expression of sugar beet (*Beta vulgaris* L). *Int. J. Phytoremediation* 25 (4), 455–465. doi:10.1080/15226514.2022.2090496
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25 (4), 402–408. doi:10.1006/meth.2001.1262
- Maness, N. (2010). “Extraction and analysis of soluble carbohydrates,” in *Plant stress tolerance* (Springer), 341–370.
- Mathew, I., Shimelis, H., Shayanowako, A. I. T., Laing, M., and Chaplot, V. (2019). Genome-wide association study of drought tolerance and biomass allocation in wheat. *PLoS One* 14 (12), e0225383. doi:10.1371/journal.pone.0225383
- Meurer, J., Plücken, H., Kowallik, K. V., and Westhoff, P. (1998). A nuclear-encoded protein of prokaryotic origin is essential for the stability of photosystem II in Arabidopsis thaliana. *EMBO J.* 17 (18), 5286–5297. doi:10.1093/emboj/17.18.5286
- Moosavi, S. G. R., Ramazani, S. H. R., Hemayati, S. S., and Gholizade, H. (2017). Effect of drought stress on root yield and some morpho-physiological traits in different genotypes of sugar beet (*Beta Vulgaris* L). *J. Crop Sci. Biotechnol.* 20 (3), 167–174. doi:10.1007/s12892-017-0009-0
- Mühlenbeck, H., Bender, K. W., and Zipfel, C. (2021). Importance of tyrosine phosphorylation for transmembrane signaling in plants. *Biochem. J.* 478 (14), 2759–2774. doi:10.1042/BCJ20210202
- Pantalio, G. F., Narciso, M., Guimaraes, C., Castro, A., Colombari, J. M., Bressegello, F., et al. (2016). Genome wide association study (GWAS) for grain yield in rice cultivated under water deficit. *Genetica* 144 (6), 651–664. doi:10.1007/s10709-016-9932-z
- Peng, L., Ma, J., Chi, W., Guo, J., Zhu, S., Lu, Q., et al. (2006). LOW PSII ACCUMULATION1 is involved in efficient assembly of photosystem II in Arabidopsis thaliana. *Plant Cell.* 18 (4), 955–969. doi:10.1105/tpc.105.037689
- Plücken, H., Müller, B., Grohmann, D., Westhoff, P., and Eichacker, L. A. (2002). The HCF136 protein is essential for assembly of the photosystem II reaction center in Arabidopsis thaliana. *FEBS Lett.* 532 (1–2), 85–90. doi:10.1016/S0014-5793(02)03634-7
- Polash, M. A. S., Sakil, M. A., Tahjib-UI-Arif, M., and Hossain, M. A. (2018). Effect of salinity on osmolytes and relative water content of selected rice genotypes. *Trop. Plant Res.* 5 (2), 227–232. doi:10.22271/tpr.2018.v5.i2.029
- Ratnapriya, R., Sosina, O. A., Starostik, M. R., Kwicklis, M., Kapphahn, R. J., Fritsche, L. G., et al. (2019). Retinal transcriptome and eQTL analyses identify genes associated with age-related macular degeneration. *Nat. Genet.* 51 (4), 606–610. doi:10.1038/s41588-019-0351-9
- Sadeghian, S., and Yavari, N. (2004). Effect of water-deficit stress on germination and early seedling growth in sugar beet. *J. Agron. Crop Sci.* 190 (2), 138–144. doi:10.1111/j.1439-037x.2004.00087.x
- Shariatipour, N., Heidari, B., Tahmasebi, A., and Richards, C. (2021). Comparative genomic analysis of quantitative trait loci associated with micronutrient contents, grain quality, and agronomic traits in wheat (*Triticum aestivum* L). *Front. Plant Sci.* 12, 709817. doi:10.3389/fpls.2021.709817
- Soskic, B., Cano-Gamez, E., Smyth, D. J., Rowan, W. C., Nacic, N., Esparza-Gordillo, J., et al. (2019). Chromatin activity at GWAS loci identifies T cell states driving complex immune diseases. *Nat. Genet.* 51 (10), 1486–1493. doi:10.1038/s41588-019-0493-9
- Spitz, D. R., and Oberley, L. W. (1989). An assay for superoxide dismutase activity in mammalian tissue homogenates. *Anal. Biochem.* 179 (1), 8–18. doi:10.1016/0003-2697(89)90192-9
- Sun, D., Cen, H., Weng, H., Wan, L., Abdalla, A., El-Manawy, A. I., et al. (2019). Using hyperspectral analysis as a potential high throughput phenotyping tool in GWAS for protein content of rice quality. *Plant Methods* 15 (1), 54–16. doi:10.1186/s13007-019-0432-x
- Tan, W., Li, W., Li, J., Liu, D., and Xing, W. (2023). Drought resistance evaluation of sugar beet germplasm by response of phenotypic indicators. *Plant Signal. Behav.* 18 (1), 2192570. doi:10.1080/15592324.2023.2192570
- Uhrig, R. G., Labandera, A.-M., Muhammad, J., Samuel, M., and Moorhead, G. B. (2016). Rhizobiales-like phosphatase 2 from Arabidopsis thaliana is a novel phospho-

- tyrosine-specific phospho-protein phosphatase (PPP) family protein phosphatase. *J. Biol. Chem.* 291 (11), 5926–5934. doi:10.1074/jbc.M115.683656
- Wang, Q., Sajja, U., Rosloski, S., Humphrey, T., Kim, M. C., Bomblies, K., et al. (2007). HUA2 caused natural variation in shoot morphology of *A. thaliana*. *Curr. Biol.* 17 (17), 1513–1519. doi:10.1016/j.cub.2007.07.059
- Wang, M., Fang, Z., Yoo, B., Bejerano, G., and Peltz, G. (2021a). The effect of population structure on murine genome-wide association studies. *Front. Genet.* 12, 745361. doi:10.3389/fgene.2021.745361
- Wang, X., Niu, Y., and Zheng, Y. (2021b). Multiple functions of MYB transcription factors in abiotic stress responses. *Int. J. Mol. Sci.* 22 (11), 6125. doi:10.3390/ijms22116125
- Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J. (2013). Glucose-TOR signalling reprograms the transcriptome and activates meristems. *Nature* 496 (7444), 181–186. doi:10.1038/nature12030
- Xu, Y., Yang, T., Zhou, Y., Yin, S., Li, P., Liu, J., et al. (2018). Genome-wide association mapping of starch pasting properties in maize using single-locus and multi-locus models. *Front. Plant Sci.* 9, 1311. doi:10.3389/fpls.2018.01311
- Yang, H., and Wang, K. (2015). Genomic variant annotation and prioritization with ANNOVAR and wANNOVAR. *Nat. Protoc.* 10 (10), 1556–1566. doi:10.1038/nprot.2015.105
- Yang, K., Chen, G., Xian, J., and Chen, W. (2022a). Varying relationship between vascular plant leaf area and leaf biomass along an elevational gradient on the eastern qinghai-tibet plateau. *Front. Plant Sci.* 13, 824461. doi:10.3389/fpls.2022.824461
- Yang, Y., Dong, S., Miao, H., Liu, X., Dai, Z., Li, X., et al. (2022b). Genome-wide association studies reveal candidate genes related to stem diameter in cucumber (*cucumis sativus* L.). *Genes* 13 (6), 1095. doi:10.3390/genes13061095
- Zeng, A., Chen, P., Korth, K., Hancock, F., Pereira, A., Brye, K., et al. (2017). Genome-wide association study (GWAS) of salt tolerance in worldwide soybean germplasm lines. *Mol. Breed.* 37 (3), 30–14. doi:10.1007/s11032-017-0634-8
- Zhang, Z., Ersoz, E., Lai, C.-Q., Todhunter, R. J., Tiwari, H. K., Gore, M. A., et al. (2010). Mixed linear model approach adapted for genome-wide association studies. *Nat. Genet.* 42 (4), 355–360. doi:10.1038/ng.546
- Zheng, J., Zhang, Z., Gong, Z., Liang, Y., Sang, Z., Xu, Y., et al. (2021). Genome-wide association analysis of salt-tolerant traits in terrestrial cotton at seedling stage. *Plants* 11 (1), 97. doi:10.3390/plants11010097