



Vitamin D Counteracts *Mycobacterium tuberculosis*-Induced Cathelicidin Downregulation in Dendritic Cells and Allows Th1 Differentiation and IFN γ Secretion

Anna K. O. Rode[†], Martin Kongsbak[†], Marie M. Hansen, Daniel Villalba Lopez, Trine B. Levring, Anders Woetmann, Niels Ødum, Charlotte M. Bonefeld and Carsten Geisler*

Faculty of Health and Medical Sciences, Department of Immunology and Microbiology, University of Copenhagen, Copenhagen, Denmark

OPEN ACCESS

Edited by:

Norbert Reiling,
Research Center Borstel, Germany

Reviewed by:

Mario M. D'Ellos,
University of Florence, Italy
Luciana Balboa,
Consejo Nacional de Investigaciones
Científicas y Técnicas (CONICET),
Argentina

*Correspondence:

Carsten Geisler
cge@sund.ku.dk

[†]These authors have contributed
equally to the work.

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 10 January 2017

Accepted: 18 May 2017

Published: 31 May 2017

Citation:

Rode AKO, Kongsbak M,
Hansen MM, Lopez DV, Levring TB,
Woetmann A, Ødum N, Bonefeld CM
and Geisler C (2017) Vitamin D
Counteracts *Mycobacterium*
tuberculosis-Induced Cathelicidin
Downregulation in Dendritic Cells and
Allows Th1 Differentiation and IFN γ
Secretion.
Front. Immunol. 8:656.
doi: 10.3389/fimmu.2017.00656

Tuberculosis (TB) presents a serious health problem with approximately one-third of the world's population infected with *Mycobacterium tuberculosis* in a latent state. Experience from the pre-antibiotic era and more recent clinical studies have established a beneficial role of sunlight and vitamin D in patients with TB. At the same time, experimental data have shown that Th1 cells through production of IFN γ are crucial for cathelicidin release by macrophages, bacterial killing, and containment of *M. tuberculosis* in granulomas. Paradoxically, vitamin D has repeatedly been ascribed an immune-suppressive function inhibiting Th1 differentiation and production of IFN γ in T cells. The aim of this study was to investigate this apparent paradox. We studied naïve human CD4⁺ T cells activated either with CD3 and CD28 antibodies or with allogeneic dendritic cells (DC) stimulated with heat-killed *M. tuberculosis* (HKMT) or purified toll-like receptor (TLR) ligands. We show that vitamin D does not block differentiation of human CD4⁺ T cells to Th1 cells and that interleukin (IL)-12 partially counteracts vitamin D-mediated inhibition of IFN γ production promoting production of equal amounts of IFN γ in Th1 cells in the presence of vitamin D as in T cells activated in the absence of vitamin D and IL-12. Furthermore, we show that HKMT and TLR2 ligands strongly downregulate cathelicidin expression in DC and that vitamin D counteracts this by upregulating cathelicidin expression. In conclusion, we demonstrate that vitamin D counteracts *M. tuberculosis*-induced cathelicidin downregulation and allows Th1 differentiation and IFN γ secretion.

Keywords: vitamin D, tuberculosis, T cells, dendritic cells, Th1, IFN γ , cathelicidin

INTRODUCTION

Tuberculosis (TB) presents a serious health problem with more than 10 million new cases of active TB responsible for 1.8 million deaths in 2015 (1). The disease is caused by the pathogen *Mycobacterium tuberculosis*. It is presumed that approximately one-third of the world's population is infected with *M. tuberculosis* in an asymptomatic latent state and that 5–10% of infected individuals develops

active TB at some point in their lives either shortly after initial infection or as a progression from a latent infection (2). Children and individuals with an impaired immune system generally have a higher risk of developing active TB (3).

M. tuberculosis is transmitted through aerosol droplets to the lung alveoli, where the pathogen infects alveolar macrophages and subsequently dendritic cells (DC), neutrophils, and macrophages in the lung interstitium (4). Activation of naïve TB-specific T cells depends on the migration of infected DC from the lungs to the mediastinal lymph nodes (5–7). DC present *M. tuberculosis* antigens for the T cells and dependent on the cytokines present during the T cell receptor-mediated activation process, the CD4⁺ T helper (Th) cells differentiate to different types of effector cells in the lymph nodes (8, 9). Macrophages and DC become activated by stimulation through the toll-like receptors (TLR)2, 4, and 9 in response to *M. tuberculosis* and start production of pro-inflammatory cytokines such as tumor necrosis factor α (TNF α), interleukin (IL)-1 β , and IL-12 (10). IL-12 plays a major role in the differentiation of Th cells to interferon γ (IFN γ)-producing Th1 effector cells by signaling through the IL-12 receptor (IL-12R), which results in phosphorylation and dimerization of signal transducer and activator of transcription (STAT)4. STAT4 then translocates to the nucleus and binds to regulatory elements of target genes, including the Th1 master transcription factor *TBX21*, *IL12R β 2*, and *IFNG*. IFN γ induces phosphorylation of STAT1 that further activates transcription of *TBX21* and *IL12R β 2*, which in turn act in a positive-feedback loop to amplify Th1 differentiation (8, 9). The resulting Th1 effector cells migrate out of the lymph nodes and are then recruited to the sites of infection.

In the latent state, *M. tuberculosis* is contained by the immune system in granulomas consisting mainly of infected macrophages surrounded by IFN γ -producing Th1 cells (2, 11). The importance of IFN γ in the immune response to mycobacteria is well established in both humans and mice. Thus, individuals with a mutation in genes related to the production of or response to IFN γ have a high susceptibility to mycobacterial infection (12–16). Likewise, mice that lack IFN γ or the IFN γ receptor are extremely susceptible to TB (17, 18). IFN γ is crucial for activation of macrophages and the formation and containment of *M. tuberculosis* in granulomas (13, 18). A central role of IFN γ is to enhance the ability of macrophages to kill intracellular pathogens such as *M. tuberculosis* (19, 20). The antimicrobial peptide cathelicidin plays an important role in the ability of macrophages to kill bacteria, and it has been reported that IFN γ increases the expression of cathelicidin in human monocytes and macrophages (21–23). Interestingly, these studies found that vitamin D is required for IFN γ -mediated enhancement of cathelicidin. This is in good agreement with several studies showing that vitamin D deficiency is associated with impaired expression of cathelicidin and increased susceptibility to infectious diseases, including TB (24–30), and it could be an important mechanism explaining the beneficial role of vitamin D in TB prevention and treatment (31, 32). In sharp contrast to this, several studies have shown that vitamin D inhibits the production of IFN γ in T cells (33–46). This creates a significant paradox in which vitamin D is required for efficient innate immune responses against *M. tuberculosis* but at

the same time impairs Th1-mediated immune responses against *M. tuberculosis*.

The aim of this study was to look deeper into this apparent paradox and to determine whether vitamin D actually impedes Th1 differentiation and how vitamin D affects IFN γ production in Th1 cells and cathelicidin production in DC.

MATERIALS AND METHODS

Chemicals

25(OH)D₃ (BML-DM-100-0001) and 1,25(OH)₂D₃ (BML-DM200-0050) were from Enzo Life Sciences, Inc., Ann Arbor, MI, USA. Stock solutions of 2.5 mM 25(OH)D₃ and 2.4 mM 1,25(OH)₂D₃ were prepared in anhydrous ($\geq 99.5\%$) ethanol and stored at -20°C . To determine 1,25(OH)₂D₃ in the supernatants, we used the 1,25-dihydroxy vitamin D EIA kit (AC-62F1) from IDS, Tyne and Wear, UK according to the manufacturer's instructions.

Cell Culture

Mononuclear cells from blood were isolated by Lymphoprep (Axis-Shield, Oslo, Norway) density-gradient centrifugation using SepMate™ tubes (86460, Stemcell Technologies, Grenoble, France) from healthy donors after obtaining informed, written consent in accordance with the Declarations of Helsinki principles for research involving human objects. The study was approved by The Committees of Biomedical Research Ethics for the Capital Region in Denmark (H-16033682). Naïve CD4⁺ T cells were isolated and cultured as previously described (47). The purified naïve CD4⁺ T cells were cultured in serum-free X-VIVO 15 medium (BE02-060F, Lonza, Verviers, Belgium) at 37°C, 5% CO₂ at a cell concentration of 1×10^6 cells/ml in flat-bottomed 24-well tissue culture plates (142475) from Nunc and stimulated with Dynabeads human T-activator CD3/CD28 beads (111.31D, Life Technologies, Grand Island, NY, USA) at a cell to bead ratio of 5:2 for up to 3 days. Cells present in the culture after 3 days were defined as activated T cells. In some experiments, 25(OH)D₃, 1,25(OH)₂D₃, or recombinant human IL-12 (219-IL, R&D Systems) was added at the indicated concentrations to the medium during the stimulation period. In Th1 polarization studies, purified naïve CD4⁺ T cells were cultured and stimulated as described above in the presence of recombinant human IL-12 (10 ng/ml) plus human IL-4 antibody (4 $\mu\text{g/ml}$, MAB204, R&D Systems). In some experiments, cells were counted after 72 h of activation using the automated cell counter NucleoCounter® NC-100™ from Chemometec.

Monocytes used for generation of monocyte-derived DC were isolated from blood mononuclear cells using EasySep Human Monocyte Enrichment Kit (19059, Stemcell Technologies) according to the manufacturer's protocol. Following isolation, monocytes were cultured in flat-bottomed six-well tissue culture plates (140675, Nunc) at a cell concentration of 5×10^5 cells/ml for 6 days in medium (RPMI-1640, R5886, Sigma-Aldrich) with penicillin, streptomycin, L-glutamine, and 10% heat-inactivated FBS (10082-147, Gibco) supplemented with GM-CSF and IL-4 (both 50 ng/ml, AF-HDC, PeproTech) with a re-supplementation

of medium and cytokines on day 3. On day 5, the differentiated DCs were supplemented with GM-CSF (50 ng/ml) and treated with either heat-killed *M. tuberculosis* (HKMT) (10 µg/ml, tlr-hkmt, InvivoGen), Pam3CSK4 (300 ng/ml, tlr-pms, InvivoGen) as TLR2 ligand, lipopolysaccharides (LPS) from *E. coli* (50 ng/ml, L 5668, Sigma) as TLR4 ligand, or left untreated for 24 h, washed, and resuspended in X-VIVO 15 medium for use in mono- and co-cultures or in RPMI-1640 medium for use in 1,25(OH)₂D₃ titration experiments. For monocultures, 2.5–5 × 10⁵ cells/ml DC were plated in flat-bottomed 24-well tissue culture plates for 24 h in the presence or absence of 25(OH)D₃ and with or without recombinant human IFN γ (R&D Systems) or with increasing concentrations of 1,25(OH)₂D₃ for titration studies. For co-culture studies, naïve human CD4⁺ T cells were purified from a different donor as described above and co-cultured with 1 × 10⁵ cells/ml DC in a ratio of 1:10 DC to T cells in flat-bottomed 24-well tissue culture plates for another 6 days in X-VIVO 15 medium.

ELISA

IFN γ and IL-13 concentrations in the supernatants were determined by ELISA according to the manufacturer's protocol (Ready-Set-Go; eBioscience).

QPCR

mRNA expression was measured by real-time RT-PCR (QPCR). For this, cells were lysed in TRI reagent (T9424, Sigma-Aldrich) followed by addition of 1-bromo-3-chloropropane (B9673, Sigma Aldrich) to separate the sample into an aqueous and an organic phase. The RNA was precipitated from the aqueous phase using isopropanol supplemented with glycogen (10814-010, Invitrogen), washed with ethanol, and dissolved in RNase free water. Equal amounts of total RNA were used for complementary DNA (cDNA) synthesis using the high-capacity RNA-to-cDNATM Kit from Applied Biosystems (4387406) according to manufacturer's protocol. For QPCR, 12.5 ng cDNA was mixed with TaqMan[®] Universal Master Mix II with UNG (4440038, Applied Biosystems), the target primer, and the eukaryotic 18 S rRNA primer (1509311, Applied Biosystems). The samples were run on a Stratagene Mx3000PTM real-time PCR machine (Agilent Technologies). The thermal profile was set to 2 min at 50°C, a 10 min hot start at 95°C, followed by cycles of 15 s at 95°C and 1 min at 60°C. Signal intensity was measured at the end of the 60°C step, and the threshold cycle values were related to eukaryotic 18S rRNA. The following primers were used; IFNG (Hs00989291_m1), CYP27B1 (Hs01096154_m1), vitamin D receptor (VDR) (Hs01045840_m1), CYP24A1 (Hs00167999_m1), TBX21 (Hs00203436_m1), IL-12RB2 (Hs00155486_m1), CXCL9 (Hs00171065_m1), CXCL10 (Hs00171042_m1), and CAMP (cathelicidin antimicrobial peptide) (Hs00189038_m1) all from Applied Biosystems.

Western Blot Analysis

For Western blot analysis, whole cell lysates were obtained by treatment of the cells with lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM MgCl₂) supplemented with 1% Triton X-100, 1× Protease inhibitor cocktail (5872S, Cell Signaling Technology, Beverly, MA, USA) and 5 mM EDTA. The samples

were run under reducing conditions on polyacrylamide gels for 2 h at 120 volt in 1× NuPAGE MOPS SDS Running buffer (K855, Amresco, Solon, OH, USA) in an XCell SureLock[®] Mini-Cell Module from Life Technologies. The proteins were transferred to nitrocellulose membrane sheets (LC2001, Life Technologies) in 1× NuPAGE Transfer buffer (NP0006-1, Life Technologies) supplemented with 10% ethanol for 60 min at 50 V in an XCell IITM Blot Module from Life Technologies. The membranes were subsequently blocked for 60 min in Tris-buffered saline supplemented with 5% milk powder (70166, Sigma-Aldrich) and 0.1% Tween 20 (P1379, Sigma-Aldrich) and incubated at 4°C for 24 h with primary antibodies diluted in Tris-buffered saline supplemented with 5% bovine serum albumin (A4503, Sigma-Aldrich) and 0.1% Tween 20. The membranes were washed and the proteins visualized following 60 min incubation at room temperature with secondary HRP-rabbit anti-mouse Ig and HRP-swine anti-rabbit Ig using ECL (RPN2232, Sigma Aldrich) technology. The anti-phospho-STAT1 (9171) and phospho-STAT4 (5267) antibodies were from Cell Signaling Technologies, the anti-STAT1 (SC-346), STAT4 (SC-486), and VDR (SC-13133) antibodies were from Santa Cruz Biotechnology, Santa Cruz, CA, USA and the anti-GAPDH (Ab9485) was from Abcam, Cambridge, UK. For band density quantification, ECL-exposed sheets were analyzed in a ChemiDoc MP Imaging System from Bio-Rad.


Statistical Analysis

Data are shown as mean ± SEM. Statistical analyses were performed using Student's *t*-test with a 5% significance level, paired observations, and equal variance. * indicates *p* ≤ 0.05.

RESULTS

IL-12 Partially Counteracts Vitamin D-Mediated Inhibition of IFN γ Production in CD4⁺ T Cells

Most previous studies on the effect of vitamin D on IFN γ production in human T cells have been performed using heterogeneous cell populations or purified CD4⁺ T cells in cell culture medium supplemented with serum and non-physiological, high concentrations (1–100 nM) of the active form of vitamin D (1,25(OH)₂D) (41–45). Various sources of serum contain different concentrations of the inactive form of vitamin D (25(OH)D), 1,25(OH)₂D, and the vitamin D-binding protein, all of great importance when investigating the effect of vitamin D on T cells (24, 46). To study the direct effect of 25(OH)D₃ and 1,25(OH)₂D₃ on IFN γ production in human Th cells under strictly defined conditions, we isolated naïve CD4⁺ T cells and stimulated them with human T-activator CD3/CD28 beads in serum-free medium in the absence or presence of 25(OH)D₃ or 1,25(OH)₂D₃. We found that 1,25(OH)₂D₃ strongly inhibited IFN γ production (~15-fold) at physiological concentrations (60–110 pM) (**Figure 1A**). Furthermore, we found that 25(OH)D₃ at physiological concentrations (50–125 nM) inhibited IFN γ production to the same degree as 1,25(OH)₂D₃ (**Figure 1A**), which confirms that T cells have the ability to convert the inactive 25(OH)D₃ to the active 1,25(OH)₂D₃ (46, 47). It is well known




that anti-CD3/CD28 bead stimulation promotes expansion of CD4⁺ T cells (48). To investigate whether the lower levels of IFN γ in vitamin D-treated cells could be attributed to fewer cells, we stimulated CD4⁺ T cells for 72 h in the presence or absence of 100 nM 25(OH) D_3 and subsequently counted the cells using an automated cell counter. At this time point, we found that there were approximately 1.5-fold more CD4⁺ T cells in the cultures compared to the initial cell number independently of the absence or presence of vitamin D (**Figure 1B**). To exclude a general toxic effect of vitamin D on the T cells, we subsequently measured secretion of IL-13 in parallel with IFN γ . We found that vitamin D increased IL-13 secretion in parallel with its inhibition of IFN γ

secretion (**Figure 1C**). Thus, the decreased levels of IFN γ were not caused by a reduced cell number or a toxic effect of vitamin D but most likely by a specific effect of vitamin D on IFN γ secretion. Previous studies have reported that 1,25(OH) $_2D_3$ inhibits production of IFN γ and augment the production of IL-4, and based on these observations it has been concluded that vitamin D restrains Th1 differentiation and promotes Th2 differentiation (41–45). However, to our knowledge, the effect of physiological concentrations of vitamin D on IFN γ production in T cells activated under Th1-inducing conditions has not been reported. To study this, we activated naïve CD4⁺ T cells in the absence or presence of IL-12 and 25(OH) D_3 . In the absence of 25(OH) D_3 , IL-12 increased IFN γ production (**Figure 1D**). Interestingly, although 25(OH) D_3 at physiological concentrations did reduce IFN γ production in IL-12 treated cells, these cells still produced equal amounts of IFN γ as T cells activated in the absence of 25(OH) D_3 and IL-12 (**Figure 1D**). From these experiments, we could conclude that 1,25(OH) $_2D_3$ inhibits the production of IFN γ in CD4⁺ effector T cells but that IL-12 partially rescues IFN γ production in the presence of vitamin D allowing production of similar amounts of IFN γ as in T cells activated in the absence of vitamin D and IL-12.

IL-12 Does Not Inhibit 1,25(OH) $_2D_3$ Production or VDR Expression and Function in CD4⁺ T Cells

One way for IL-12 to counteract vitamin D-mediated inhibition of IFN γ production in CD4⁺ T cells could be by limiting the ability of the cells to convert 25(OH) D_3 to 1,25(OH) $_2D_3$. To investigate this, we studied whether IL-12 affected the expression of CYP27B1, the enzyme responsible for converting 25(OH) D_3 to 1,25(OH) $_2D_3$. We activated naïve CD4⁺ T cells in the presence of 100 nM 25(OH) D_3 in the absence or presence of IL-12 and measured the expression levels of CYP27B1 mRNA in the cells. We found that IL-12 did not influence the expression of CYP27B1 (**Figure 2A**). Furthermore, we directly measured the production of 1,25(OH) $_2D_3$ in the supernatants and found that IL-12 did not affect the production of 1,25(OH) $_2D_3$ (**Figure 2B**). Another way IL-12 might counteract vitamin D-mediated inhibition of IFN γ production in CD4⁺ T cells could be by inhibiting the expression or function of the VDR. However, we found that IL-12 slightly increased VDR expression on the mRNA level (**Figure 2C**). To further investigate whether IL-12 affected VDR expression, we determined the effect of IL-12 on VDR protein levels. We did not observe any significant effect of IL-12 on VDR expression at the protein level (**Figure 2D**). CYP24A1 is the enzyme that initiates degradation of 1,25(OH) $_2D_3$, and CYP24A1 expression is dependent on functional 1,25(OH) $_2D_3$ /VDR complexes. Consequently, we measured the CYP24A1 levels in cells treated with or without IL-12. We found that the cells produced very high amounts of CYP24A1 in the presence of 100 nM 25(OH) D_3 (approximately 100.000-fold upregulated compared to unstimulated T cells) although IL-12 slightly reduced CYP24A1 expression (**Figure 2E**). Taken together, these experiments indicated that IL-12 does not counteract vitamin D-mediated inhibition of IFN γ production in CD4⁺ T cells by reducing their ability to



produce 1,25(OH)₂D₃ or by reducing their VDR expression or function.

Vitamin D Does Not Prevent Th1 Differentiation


Due to the inhibition of IFN γ production, it has been suggested that vitamin D impedes differentiation of Th1 cells. However, whether vitamin D actually inhibits Th1 differentiation is still not known. To directly study whether vitamin D affects Th1 differentiation, we stimulated naïve CD4⁺ T cells under classical Th1 conditions with IL-12 and anti-IL-4 in the absence or presence of 25(OH)D₃ and measured the expression levels of TBX21 and IL-12R β 2. Compared to naïve T cells, cells activated for 24 h in the absence of 25(OH)D₃ upregulated TBX21 approximately 120-fold. Although not significantly, 25(OH)D₃ slightly reduced TBX21 upregulation at 24 h but had no inhibitory effect on TBX21 expression after 48 and 72 h (Figure 3A). Likewise, IL-12R β 2 expression was slightly

reduced after 24 h but unaffected by 25(OH)D₃ after 48 and 72 h of stimulation (Figure 3B). While vitamin D did not affect TBX21 and IL-12R β 2 expression in cells stimulated for 48 and 72 h, it clearly reduced IFN γ expression in the same cells (Figure 3C). Thus, these experiments indicated that vitamin D does not block differentiation of Th1 cells but directly affects the transcription of the *IFNG* gene. Although reduced, IFN γ expression in Th1 cells treated with vitamin D was still highly upregulated (1000- to 2000-fold, Figure 3C) and they produced high levels of IFN γ (Figure 1D). In line with the strong production of IFN γ , we found that vitamin D did not affect STAT1 phosphorylation significantly during T cell activation (Figure 3D). Likewise, STAT4 phosphorylation was not significantly affected of vitamin D (Figure 3E).

IFN γ itself plays an important role during Th1 differentiation as STAT1 activated by IFN γ further activates transcription of *TBX21* and *IL12R β 2* (8, 9). We have previously demonstrated that naïve human CD4⁺ T cells do not express the VDR but that they start to express it 24–48 h after TCR/CD28 stimulation (46, 49). This suggests that vitamin D cannot affect IFN γ production in the early stages of T cell activation due to the lack of VDR. To study the effect of 25(OH)D₃ and Th1-inducing conditions on the early kinetics of VDR expression, we stimulated naïve CD4⁺ T cells for 12, 24, and 48 h in the presence or absence of 25(OH)D₃ and Th1-inducing conditions. We could confirm that only very low levels of VDR are present after 12 and 24 h of stimulation, and that substantial VDR expression levels are found after 48 h of stimulation (Figure 4A). In agreement, we found that at 12 h of stimulation, vitamin D did not influence IFN γ production neither in cells activated in the absence or presence of Th1 conditions (Figure 4B). In contrast, Th1 conditions acted rapidly with increased production of IFN γ already 12 h after stimulation (Figure 4B). After 24 h of stimulation, a minor inhibitory effect of vitamin D on IFN γ production was seen (Figure 4C), and after 48 h vitamin D substantially inhibited IFN γ production (Figure 4D). These experiments indicated that IL-12-mediated signaling is initiated rapidly after initial T cell stimulation with increased IFN γ production as a consequence, and that this takes place before the VDR is expressed at sufficiently high levels to inhibit IFN γ production. These observations support that CD4⁺ T cells activated under Th1-inducing conditions start differentiation toward Th1 cells even in the presence of vitamin D.

Vitamin D Is Required for Concomitant Production of IFN γ and Cathelicidin in DC–T Cell Co-Cultures


Activation of naïve TB-specific T cells depends on the migration of infected DC from the lungs to the mediastinal lymph nodes where the DC present TB-derived antigens for the T cells (5–7). It has been suggested that antigen-presenting cells mediate the vitamin D-induced inhibition of IFN γ production in T cells (36, 38, 40), and the absence of antigen-presenting cells in our experimental set up could maybe explain why we observed IFN γ production in purified T cells stimulated in the presence of vitamin D and IL-12. To investigate this possibility, we cultured



naïve CD4⁺ T cells with allogeneic DC for 6 days with or without 25(OH)D₃ in the absence or presence of Th1-inducing conditions and subsequently measured IFNγ expression and production. Vitamin D strongly inhibited IFNγ mRNA expression (~15-fold) (Figure 5A) and production (~20-fold) (Figure 5B) in the absence of Th1-inducing conditions. Th1-inducing conditions increased IFNγ expression and production and counteracted the inhibitory effect of vitamin D. Thus, vitamin D only inhibited IFNγ mRNA expression ~2-fold and IFNγ production ~3-fold in the presence of Th1-inducing conditions (Figures 5A,B). Interestingly, IFNγ expression and production in cells cultured with vitamin D under Th1-inducing conditions were at least as high as in cells cultured without vitamin D and Th1-inducing conditions (Figures 5A,B). This indicated that DC do not shut down IFNγ production in T cells in the presence of vitamin D. Taken together, we could conclude that vitamin D strongly inhibits IFNγ expression and production in T cells in the absence of Th1-inducing conditions, and that Th1-inducing conditions partially rescue IFNγ production in the presence of vitamin D independently on the presence or absence of DC.

Cellular responses mediated by IFNγ are primarily caused by gene expression modulation. Primary IFNγ-responsive genes are induced early by binding of STAT1 dimers to gamma-activating sequences in the promoter of target genes such as IRF1, CXCL9, and CXCL10 (50). To study whether the levels of IFNγ produced in Th1 cells in the presence of vitamin D were sufficient to affect IFNγ-responsive genes, we measured the expression of CXCL9 and CXCL10 in the same co-cultures as described above. We found that vitamin D reduced CXCL9 and CXCL10 expression ~110 and ~25-fold in Th0 cells, respectively (Figures 5C,D). In contrast, vitamin D only reduced CXCL9 and CXCL10 expression ~2- and ~3-fold in Th1 cells compared to untreated Th0 cells. This demonstrated that the cellular responses to the levels of IFNγ produced in Th1 cells in the presence of vitamin D almost equaled the responses to the levels of IFNγ produced in Th0 cells in the absence of vitamin D.

As the importance of IFNγ in the immune response to mycobacteria is well established, it is puzzling why vitamin D, associated with a beneficial role in TB prevention and treatment,




reduces IFN γ production. Cathelicidin plays an important role in the killing of especially intracellular pathogens such as *M. tuberculosis* in macrophages (51). Furthermore, it has been shown that cathelicidin expression is dependent on vitamin D, and that IFN γ increases the synthesis of cathelicidin in macrophages (21–23). To study how cathelicidin is regulated in DC, we cultured naïve CD4⁺ T cells with allogeneic DC for 6 days as described above with or without 25(OH)D₃ in the absence or presence of Th1-inducing conditions and subsequently measured cathelicidin expression. We found that cathelicidin upregulation was absolutely dependent on vitamin D but independent of Th1-inducing conditions (Figure 5E). This indicated that vitamin D without IFN γ is sufficient to induce cathelicidin upregulation, and that IFN γ without vitamin D does not affect cathelicidin expression in DC.

Vitamin D Counteracts the Inhibitory Effect of *M. tuberculosis* on Cathelicidin Expression in DC

Previous studies have demonstrated that IFN γ acts in synergy with TLR2 ligands to upregulate cathelicidin expression in human monocytes and macrophages (21, 22). To determine how

vitamin D, IFN γ , and HKMT/TLR2 ligands affect cathelicidin expression in DC, we incubated DC in the absence or presence of 25(OH)D₃, IFN γ , and HKMT or Pam3CSK4, and measured the expression of cathelicidin. Surprisingly, we found that HKMT and Pam3CSK4 significantly downregulated cathelicidin expression both in the absence and presence of vitamin D (Figure 6A). In contrast, vitamin D strongly upregulated cathelicidin expression both in the absence and presence of HKMT or Pam3CSK4. Although HKMT and Pam3CSK4 downregulated cathelicidin expression ~13- and ~10-fold in the presence of vitamin D, respectively, the expression of cathelicidin was still ~215- and ~120-fold higher in DC stimulated with HKMT/Pam3CSK4 in the presence of vitamin D than in the absence of vitamin D (Figure 6A). IFN γ by itself did not affect cathelicidin expression.

To determine whether the HKMT/Pam3CSK4-induced inhibition of cathelicidin expression was caused by a reduced ability to produce or respond to 1,25(OH)₂D₃, we incubated DC in the absence or presence of 25(OH)D₃, IFN γ , and HKMT/Pam3CSK4 and measured the expression of CYP27B1, VDR, and CYP24A1, and the production of 1,25(OH)₂D₃. We found that HKMT/Pam3CSK4 increased CYP27B1 expression, especially in the presence of 25(OH)D₃ (Figure 6B). In



accordance, although untreated DC produced high amounts of 1,25(OH) $_2D_3$, HKMT/Pam3CSK4 increased the 1,25(OH) $_2D_3$ production four- to fivefold (Figure 6C). In addition, HKMT slightly increased VDR expression in the presence of 25(OH) D_3 (Figure 6D). This suggested that HKMT/Pam3CSK4-treated DC should be at least as efficient as untreated DC to respond to 1,25(OH) $_2D_3$. This was supported by the expression of comparable levels of the 1,25(OH) $_2D_3$ /VDR-dependent CYP24A1 in DC treated or untreated with HKMT/Pam3CSK4 in the presence of 25(OH) D_3 (Figure 6E). These observations indicated that the inhibitory effect of HKMT/Pam3CSK4 on cathelicidin expression was caused by a vitamin D/VDR-independent mechanism. To test this hypothesis, we incubated DC in the absence or presence of HKMT or Pam3CSK4 and increasing concentrations of 1,25(OH) $_2D_3$. In all concentrations of 1,25(OH) $_2D_3$ tested, we found that HKMT/Pam3CSK4 inhibited cathelicidin expression (Figure 6F). Interestingly, stimulation with the more complex HKMT resulted in a ~23-fold reduction in cathelicidin expression, whereas the pure TLR2 ligand Pam3CSK4 caused an ~8-fold reduction in cathelicidin expression at all concentrations of 1,25(OH) $_2D_3$ tested. This suggested that HKMT affected the DC through additional receptors than TLR2. In agreement, we found that the TLR4 ligand LPS also inhibited cathelicidin expression in DC (data not shown). In contrast to the effect on cathelicidin expression, HKMT/Pam3CSK4 did not affect CYP24A1 expression, supporting that HKMT/Pam3CSK4 did not inhibit

the ability of DC to respond to 1,25(OH) $_2D_3$ /VDR complexes (Figure 6G). Taken together, these data indicated that *M. tuberculosis* strongly inhibits cathelicidin expression in DC through TLR signaling but that vitamin D increases cathelicidin expression and thereby more than overcome the inhibitory effect of *M. tuberculosis*.

DISCUSSION

In this study, we show that vitamin D does not block differentiation of naïve human CD4 $^+$ T cells to Th1 cells and that IL-12 partially rescues IFN γ production in T cells exposed to vitamin D. Furthermore, we demonstrate that HKMT/TLR2 ligands strongly downregulate cathelicidin expression in DC and that vitamin D counteracts this by upregulating cathelicidin expression.

Since the first studies on the effect of vitamin D on IFN γ production in human T cells (33, 34) several studies have confirmed that vitamin D inhibits IFN γ production in both human and mouse T cells (35–46). Some studies found that this was an indirect effect mediated via antigen-presenting cells (36, 38, 40), whereas others found a direct effect of vitamin D on T cells (41–43). We found that both the inactive and the active form of vitamin D inhibit IFN γ production in purified CD4 $^+$ T cells. Thus, our study confirms that activated T cells can convert inactive vitamin D to active vitamin D, and that vitamin D has a direct effect on human T cells. This is in good accordance with expression of both CYP27B1 and the VDR in activated




FIGURE 6 | Vitamin D counteracts the inhibitory effect of *M. tuberculosis* on cathelicidin production in DC. Relative cathelicidin (A), CYP27B1 (B), VDR (D), and CYP24A1 (E) expression and 1,25(OH)₂D₃ production (C) in untreated DC (no TLR ligand) or DC treated with HKMT or Pam3CSK4 and subsequently incubated for 24 h in the absence or presence of 100 nM 25(OH)D₃ and 10 ng/ml IFN γ . Data are normalized to the values obtained from untreated DC in the absence of 25(OH)D₃ and IFN γ (mean \pm SEM, $n \geq 3$), * indicates $p \leq 0.05$. Relative cathelicidin (F) and CYP24A1 (G) expression in untreated DC (no TLR ligand) or DC treated with HKMT or Pam3CSK4 incubated for 24 h with the indicated concentrations of 1,25(OH)₂D₃. Data are normalized to untreated DC in the absence of 1,25(OH)₂D₃ (mean \pm SEM, $n \geq 3$), * indicates $p \leq 0.05$ for no TLR ligand untreated versus 1,25(OH)₂D₃ treated, # indicates $p \leq 0.05$ for HKMT versus no TLR ligand, \square indicates $p \leq 0.05$ for Pam3CSK4 versus no TLR ligand.

human T cells (46, 47, 49, 52). Based on the impaired IFN γ production and increased IL-4 production in T cells exposed to vitamin D, it has been concluded that vitamin D restrains Th1 differentiation and promotes Th2 differentiation. However, only few studies have investigated whether vitamin D actually affects the master transcription factors TBX21 and GATA3 regulating Th1 and Th2 differentiation. One study found that although 1,25(OH)₂D₃ inhibited IFN γ production, it had no effect on the levels of TBX21 and GATA3 in mouse T cells activated

under Th0, Th1, or Th2 conditions (37). Another study found a very discrete increase in GATA3 expression in mouse T cells activated in the presence of 1,25(OH)₂D₃ under Th0 conditions, but they did not study TBX21 expression nor the effect of Th1 conditions (53). We found that vitamin D did not significantly affect the expression of TBX21 in human T cells activated under Th1 conditions. Likewise, although IL-12R β 2 upregulation was slightly inhibited at 24 h, vitamin D did not affect IL-12R β 2 expression at later time points. This indicated that vitamin D

does not restrain Th1 differentiation. STAT4 and STAT1 signaling play critical roles in the differentiation of Th1 cells (8). We are not aware of previous studies that have investigated the effect of vitamin D on STAT1 and STAT4 activation in human T cells. In the present study, we found that vitamin D did not significantly affect phosphorylation of STAT1 and STAT4 in T cells activated under Th1 conditions, which further supports that vitamin D does not block differentiation of Th1 cells. Thus, vitamin D most likely directly affects transcription of the *IFNG* gene and not the differentiation of naïve CD4⁺ T cells to Th1 cells. This is in accordance with previous studies, which found that 1,25(OH)₂D₃/VDR complexes directly bind to the *IFNG* promoter and inhibit *IFNG* transcription (54) and that the fraction of IFN γ -producing T cells is not reduced in T cells activated in the presence of vitamin D under Th1 conditions (55). The kinetic studies illustrated in **Figures 3** and **4** in this study also strongly support that vitamin D does not inhibit Th1 differentiation. The VDR is not or only weakly expressed in T cells for the first 24 h of stimulation, and consequently vitamin D cannot affect IFN γ expression or production at these early time points. In contrast, IL-12 signaling is efficient already shortly after T cell activation and augments IFN γ production and initiates Th1 differentiation.

It has been demonstrated that the promoter of the human cathelicidin gene contains a consensus vitamin D response element, and that cathelicidin is upregulated by vitamin D (56, 57). Shortly after these studies, it was found that signaling through TLR2 induced upregulation of CYP27B1 and the VDR in human monocytes and macrophages but not in DC (24). Furthermore, upregulation of cathelicidin and growth inhibition of *M. tuberculosis* were seen if the cell culture media was supplemented with 25(OH)D₃ (24). A subsequent study confirmed that the vitamin D-mediated antimicrobial activity against *M. tuberculosis* in the human monocytic cell line THP-1 was dependent on cathelicidin (51). In accordance with Liu et al. (24), we found that HKMT/TLR2 triggering did not significantly affect VDR expression in DC. In contrast to VDR, we found that the TLR2 ligand Pam3CSK4 and especially HKMT upregulated CYP27B1 expression and 1,25(OH)₂D₃ production in the presence of 25(OH)D₃. However, although HKMT/Pam3CSK4 upregulated CYP27B1, untreated DC still expressed high levels of CYP27B1. Accordingly, we found that even untreated DCs have a high capacity to convert 25(OH)D₃ to 1,25(OH)₂D₃, and to respond to 1,25(OH)₂D₃ with strong upregulation of cathelicidin and CYP24A1 expression. In sharp contrast to studies finding that TLR2 ligands upregulated cathelicidin in human monocytes and macrophages in a vitamin D-dependent manner (21, 22, 24), we found that HKMT profoundly downregulated cathelicidin in human DC. Although HKMT is described as a primary TLR2 ligand, it has been reported that *M. tuberculosis* stimulates a range of other receptors, including TLR4 and 9 (10). However, we found that the TLR2 ligand Pam3CSK4 mimicked HKMT in regard to cathelicidin downregulation, although not quite as efficiently as HKMT. TLR4 triggering also resulted in cathelicidin downregulation and we could conclude that *M. tuberculosis* can downregulate cathelicidin expression in DC through signaling via TLR2 and

most likely additional TLR. As for cathelicidin expression, different responses between monocytes/macrophages and DC to TLR2 signaling have been reported concerning MHC class II expression. Thus, whereas TLR2 signaling inhibits MHC class II expression in macrophages, it enhances MHC class II expression in DC (58). Interestingly, although the paradigm is that TLR2 signaling increases cathelicidin expression in human monocytes and macrophages (21, 22, 24), two papers have actually found that TLR2 signaling reduces cathelicidin expression in human monocytes in accordance with what we find in human DC (59, 60).

M. tuberculosis uses various strategies to evade antimicrobial mechanisms of macrophages, including inhibition of phagosome-lysosome fusion, antigen presentation, MHC class II expression, autophagy, and cytokine production (61). Interestingly, many of these immune evasion mechanisms are driven by TLR2 signaling (62–65). In the present study, we show that HKMT can use TLR2 signaling to strongly inhibit cathelicidin expression in DC, and as for autophagy (22, 24), we find that this attempt to evade the immune system by *M. tuberculosis* can be counteracted by vitamin D. Thus, although vitamin D reduces IFN γ production, we could conclude that vitamin D does not block Th1 development, and that Th1 cells produce significant amounts of IFN γ in the presence of vitamin D. At the same time, vitamin D is required to upregulate and counteract downregulation of cathelicidin by *M. tuberculosis*. We believe that these findings add to the understanding of the beneficial role of vitamin D in TB prevention and treatment.

ETHICS STATEMENT

The study was approved by The Committees of Biomedical Research Ethics for the Capital Region in Denmark (H-16033682). Blood samples were obtained from healthy donors after obtaining informed, written consent in accordance with the Declarations of Helsinki principles for research involving human objects.

AUTHOR CONTRIBUTIONS

AR, MK, MH, DL, and TL performed the laboratory experiments. AR, MK, AW, NØ, CB, and CG conceived and designed the experiments. AR, MK, and CG analyzed the data and wrote the paper. All authors revised and approved the final manuscript.

FUNDING

This work was supported by The Danish Medical Research Council and Laege Sofus Carl Emil Friis og hustru Olga Doris Friis' Legat.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://journal.frontiersin.org/article/10.3389/fimmu.2017.00656/full#supplementary-material>.

REFERENCES

- WHO. *Global Tuberculosis Report 2016*. (2016). Available from: www.who.int/tb/publications/global_report/en/
- Nunes-Alves C, Booty MG, Carpenter SM, Jayaraman P, Rothchild AC, Behar SM. In search of a new paradigm for protective immunity to TB. *Nat Rev Microbiol* (2014) 12:289–99. doi:10.1038/nrmicro3230
- Getahun H, Matteelli A, Chaisson RE, Raviglione M. Latent *Mycobacterium tuberculosis* infection. *N Engl J Med* (2015) 372:2127–35. doi:10.1056/NEJMra1405427
- Srivastava S, Ernst JD, Desvignes L. Beyond macrophages: the diversity of mononuclear cells in tuberculosis. *Immunol Rev* (2014) 262:179–92. doi:10.1111/immr.12217
- Humphreys IR, Stewart GR, Turner DJ, Patel J, Karamanou D, Snelgrove RJ, et al. A role for dendritic cells in the dissemination of mycobacterial infection. *Microbes Infect* (2006) 8:1339–46. doi:10.1016/j.micinf.2005.12.023
- Wolf AJ, Linas B, Trevejo-Nunez GJ, Kincaid E, Tamura T, Takatsu K, et al. *Mycobacterium tuberculosis* infects dendritic cells with high frequency and impairs their function in vivo. *J Immunol* (2007) 179:2509–19. doi:10.4049/jimmunol.179.4.2509
- Reiley WW, Calayag MD, Wittmer ST, Huntington JL, Pearl JE, Fountain JJ, et al. ESAT-6-specific CD4 T cell responses to aerosol *Mycobacterium tuberculosis* infection are initiated in the mediastinal lymph nodes. *Proc Natl Acad Sci U S A* (2008) 105:10961–6. doi:10.1073/pnas.0801496105
- Zhu J, Yamane H, Paul WE. Differentiation of effector CD4 T cell populations. *Annu Rev Immunol* (2010) 28:445–89. doi:10.1146/annurev-immunol-030409-101212
- Tripathi SK, Lahesmaa R. Transcriptional and epigenetic regulation of T-helper lineage specification. *Immunol Rev* (2014) 261:62–83. doi:10.1111/immr.12204
- Kleinnijenhuis J, Oosting M, Joosten LA, Netea MG, Van CR. Innate immune recognition of *Mycobacterium tuberculosis*. *Clin Dev Immunol* (2011) 2011:405310. doi:10.1155/2011/405310
- Tsai MC, Chakravarty S, Zhu G, Xu J, Tanaka K, Koch C, et al. Characterization of the tuberculous granuloma in murine and human lungs: cellular composition and relative tissue oxygen tension. *Cell Microbiol* (2006) 8:218–32. doi:10.1111/j.1462-5822.2005.00612.x
- Newport MJ, Huxley CM, Huston S, Hawrylowicz CM, Oostra BA, Williamson R, et al. A mutation in the interferon-gamma-receptor gene and susceptibility to mycobacterial infection. *N Engl J Med* (1996) 335:1941–9. doi:10.1056/NEJM199612263352602
- Jouanguy E, Altare F, Lamhamedi S, Revy P, Emile JF, Newport M, et al. Interferon-gamma-receptor deficiency in an infant with fatal bacille Calmette-Guerin infection. *N Engl J Med* (1996) 335:1956–61. doi:10.1056/NEJM199612263352604
- Casanova JL, Abel L. Genetic dissection of immunity to mycobacteria: the human model. *Annu Rev Immunol* (2002) 20:581–620. doi:10.1146/annurev.immunol.20.081501.125851
- Rosenzweig SD, Holland SM. Defects in the interferon-gamma and interleukin-12 pathways. *Immunol Rev* (2005) 203:38–47. doi:10.1111/j.0105-2896.2005.00227.x
- Bustamante J, Boisson-Dupuis S, Abel L, Casanova JL. Mendelian susceptibility to mycobacterial disease: genetic, immunological, and clinical features of inborn errors of IFN-gamma immunity. *Semin Immunol* (2014) 26:454–70. doi:10.1016/j.smim.2014.09.008
- Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR. An essential role for interferon gamma in resistance to *Mycobacterium tuberculosis* infection. *J Exp Med* (1993) 178:2249–54. doi:10.1084/jem.178.6.2249
- Kamijo R, Le J, Shapiro D, Havell EA, Huang S, Aguet M, et al. Mice that lack the interferon-gamma receptor have profoundly altered responses to infection with *Bacillus Calmette-Guerin* and subsequent challenge with lipopolysaccharide. *J Exp Med* (1993) 178:1435–40. doi:10.1084/jem.178.4.1435
- Nathan CF, Murray HW, Wiebe ME, Rubin BY. Identification of interferon-gamma as the lymphokine that activates human macrophage oxidative metabolism and antimicrobial activity. *J Exp Med* (1983) 158:670–89. doi:10.1084/jem.158.3.670
- Schroder K, Hertzog PJ, Ravasi T, Hume DA. Interferon- γ : an overview of signals, mechanisms and functions. *J Leukoc Biol* (2004) 75:163–89. doi:10.1189/jlb.0603252
- Edfeldt K, Liu PT, Chun R, Fabri M, Schenk M, Wheelwright M, et al. T-cell cytokines differentially control human monocyte antimicrobial responses by regulating vitamin D metabolism. *Proc Natl Acad Sci U S A* (2010) 107:22593–8. doi:10.1073/pnas.1011624108
- Fabri M, Stenger S, Shin DM, Yuk JM, Liu PT, Realegeno S, et al. Vitamin D is required for IFN-gamma-mediated antimicrobial activity of human macrophages. *Sci Transl Med* (2011) 3:104ra102. doi:10.1126/scitranslmed.3003045
- Klug-Micu GM, Stenger S, Sommer A, Liu PT, Krutzik SR, Modlin RL, et al. CD40 ligand and interferon-gamma induce an antimicrobial response against *Mycobacterium tuberculosis* in human monocytes. *Immunology* (2013) 139:121–8. doi:10.1111/imm.12062
- Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* (2006) 311:1770–3. doi:10.1126/science.1123933
- Gibney KB, MacGregor L, Leder K, Torresi J, Marshall C, Ebeling PR, et al. Vitamin D deficiency is associated with tuberculosis and latent tuberculosis infection in immigrants from sub-Saharan Africa. *Clin Infect Dis* (2008) 46:443–6. doi:10.1086/525268
- Nnoaham KE, Clarke A. Low serum vitamin D levels and tuberculosis: a systematic review and meta-analysis. *Int J Epidemiol* (2008) 37:113–9. doi:10.1093/ije/dym247
- Talat N, Pery S, Parsonnet J, Dawood G, Hussain R. Vitamin d deficiency and tuberculosis progression. *Emerg Infect Dis* (2010) 16:853–5. doi:10.3201/eid1605.091693
- Bhan I, Camargo CA Jr, Wenger J, Ricciardi C, Ye J, Borregaard N, et al. Circulating levels of 25-hydroxyvitamin D and human cathelicidin in healthy adults. *J Allergy Clin Immunol* (2011) 127:1302–4. doi:10.1016/j.jaci.2010.12.1097
- Dixon BM, Barker T, McKinnon T, Cuomo J, Frei B, Borregaard N, et al. Positive correlation between circulating cathelicidin antimicrobial peptide (hCAP18/LL-37) and 25-hydroxyvitamin D levels in healthy adults. *BMC Res Notes* (2012) 5:575. doi:10.1186/1756-0500-5-575
- Rahman S, Rehn A, Rahman J, Andersson J, Svensson M, Brighenti S. Pulmonary tuberculosis patients with a vitamin D deficiency demonstrate low local expression of the antimicrobial peptide LL-37 but enhanced FoxP3+ regulatory T cells and IgG-secreting cells. *Clin Immunol* (2015) 156:85–97. doi:10.1016/j.clim.2014.12.003
- Salahuddin N, Ali F, Hasan Z, Rao N, Aqeel M, Mahmood F. Vitamin D accelerates clinical recovery from tuberculosis: results of the SUCCINCT study [Supplementary cholecalciferol in recovery from tuberculosis]. A randomized, placebo-controlled, clinical trial of vitamin D supplementation in patients with pulmonary tuberculosis. *BMC Infect Dis* (2013) 13:22. doi:10.1186/1471-2334-13-22
- Selvaraj P, Harishankar M, Afsal K. Vitamin D: immuno-modulation and tuberculosis treatment. *Can J Physiol Pharmacol* (2015) 93:377–84. doi:10.1139/cjpp-2014-0386
- Reichel H, Koeffler HP, Tobler A, Norman AW. 1 α ,25-dihydroxyvitamin D₃ inhibits γ -interferon synthesis by normal human peripheral blood lymphocytes. *Proc Natl Acad Sci U S A* (1987) 84:3385–9. doi:10.1073/pnas.84.10.3385
- Rigby WF, Denome S, Fanger MW. Regulation of lymphokine production and human T lymphocyte activation by 1,25-dihydroxyvitamin D₃. Specific inhibition at the level of messenger RNA. *J Clin Invest* (1987) 79:1659–64. doi:10.1172/JCI113004
- Muscettola M, Grasso G. Effect of 1,25-dihydroxyvitamin D₃ on interferon gamma production in vitro. *Immunol Lett* (1988) 17:121–4. doi:10.1016/0165-2478(88)90079-X
- Penna G, Adorini L. 1 alpha,25-dihydroxyvitamin D₃ inhibits differentiation, maturation, activation, and survival of dendritic cells leading to impaired alloreactive T cell activation. *J Immunol* (2000) 164:2405–11. doi:10.4049/jimmunol.164.5.2405
- Staeva-Vieira TP, Freedman LP. 1,25-dihydroxyvitamin D₃ inhibits IFN-gamma and IL-4 levels during in vitro polarization of primary murine CD4+ T cells. *J Immunol* (2002) 168:1181–9. doi:10.4049/jimmunol.168.3.1181
- van Halteren AG, van EE, De Jong EC, Bouillon R, Roep BO, Mathieu C. Redirection of human autoreactive T-cells upon interaction with dendritic cells modulated by TX527, an analog of 1,25 dihydroxyvitamin D(3). *Diabetes* (2002) 51:2119–25. doi:10.2337/diabetes.51.7.2119
- Thien R, Baier K, Pietschmann P, Peterlik M, Willheim M. Interactions of 1 alpha,25-dihydroxyvitamin D₃ with IL-12 and IL-4 on cytokine expression of

- human T lymphocytes. *J Allergy Clin Immunol* (2005) 116:683–9. doi:10.1016/j.jaci.2005.05.013
40. Penna G, Amuchastegui S, Giarratana N, Daniel KC, Vulcano M, Sozzani S, et al. 1,25-Dihydroxyvitamin D3 selectively modulates tolerogenic properties in myeloid but not plasmacytoid dendritic cells. *J Immunol* (2007) 178:145–53. doi:10.4049/jimmunol.178.1.145
 41. Bartels LE, Jorgensen SP, Agnholt J, Kelsen J, Hvas CL, Dahlerup JF. 1,25-Dihydroxyvitamin D3 and dexamethasone increase interleukin-10 production in CD4+ T cells from patients with Crohn's disease. *Int Immunopharmacol* (2007) 7:1755–64. doi:10.1016/j.intimp.2007.09.016
 42. Jeffery LE, Burke F, Mura M, Zheng Y, Qureshi OS, Hewison M, et al. 1,25-Dihydroxyvitamin D3 and IL-2 combine to inhibit T cell production of inflammatory cytokines and promote development of regulatory T cells expressing CTLA-4 and FoxP3. *J Immunol* (2009) 183:5458–67. doi:10.4049/jimmunol.0803217
 43. Correale J, Ysrraelit MC, Gaitan MI. Immunomodulatory effects of vitamin D in multiple sclerosis. *Brain* (2009) 132:1146–60. doi:10.1093/brain/awp033
 44. Baek F, Takiishi T, Korf H, Gysemans C, Mathieu C. Vitamin D: modulator of the immune system. *Curr Opin Pharmacol* (2010) 10(4):482–96. doi:10.1016/j.coph.2010.04.001
 45. Peelen E, Knippenberg S, Muris AH, Thewissen M, Smolders J, Tervaert JW, et al. Effects of vitamin D on the peripheral adaptive immune system: a review. *Autoimmun Rev* (2011) 10:733–43. doi:10.1016/j.autrev.2011.05.002
 46. Kongsbak M, von Essen MR, Levring TB, Schjerling P, Woetmann A, Odum N, et al. Vitamin D-binding protein controls T cell responses to vitamin D. *BMC Immunol* (2014) 15:35. doi:10.1186/s12865-014-0035-2
 47. Kongsbak M, von Essen MR, Boding L, Levring TB, Schjerling P, Lauritsen JP, et al. Vitamin D up-regulates the vitamin D receptor by protecting it from proteasomal degradation in human CD4+ T cells. *PLoS One* (2014) 9:e96695. doi:10.1371/journal.pone.0096695
 48. Li Y, Kurlander RJ. Comparison of anti-CD3 and anti-CD28-coated beads with soluble anti-CD3 for expanding human T cells: differing impact on CD8 T cell phenotype and responsiveness to restimulation. *J Transl Med* (2010) 8:104. doi:10.1186/1479-5876-8-104
 49. von Essen MR, Kongsbak M, Schjerling P, Olgaard K, Odum N, Geisler C. Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nat Immunol* (2010) 11:344–9. doi:10.1038/ni.1851
 50. Saha B, Jyothi PS, Chandrasekar B, Nandi D. Gene modulation and immunoregulatory roles of interferon gamma. *Cytokine* (2010) 50:1–14. doi:10.1016/j.cyt.2009.11.021
 51. Liu PT, Stenger S, Tang DH, Modlin RL. Cutting edge: vitamin D-mediated human antimicrobial activity against *Mycobacterium tuberculosis* is dependent on the induction of cathelicidin. *J Immunol* (2007) 179:2060–3. doi:10.4049/jimmunol.179.4.2060
 52. Kongsbak M, Levring TB, Geisler C, von Essen MR. The vitamin D receptor and T cell function. *Front Immunol* (2013) 4:148. doi:10.3389/fimmu.2013.00148
 53. Boonstra A, Barrat FJ, Crain C, Heath VL, Savelkoul HF, O'Garra A. 1alpha,25-dihydroxyvitamin d3 has a direct effect on naive CD4(+) T cells to enhance the development of Th2 cells. *J Immunol* (2001) 167:4974–80. doi:10.4049/jimmunol.167.9.4974
 54. Cippitelli M, Santoni A. Vitamin D3: a transcriptional modulator of the interferon-gamma gene. *Eur J Immunol* (1998) 28:3017–30. doi:10.1002/(SICI)1521-4141(199810)28:10<3017::AID-IMMU3017>3.3.CO;2-Y
 55. Ikeda U, Wakita D, Ohkuri T, Chamoto K, Kitamura H, Iwakura Y, et al. 1alpha,25-dihydroxyvitamin D3 and all-trans retinoic acid synergistically inhibit the differentiation and expansion of Th17 cells. *Immunol Lett* (2010) 134:7–16. doi:10.1016/j.imlet.2010.07.002
 56. Wang TT, Nestel FP, Bourdeau V, Nagai Y, Wang Q, Liao J, et al. Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression. *J Immunol* (2004) 173:2909–12. doi:10.4049/jimmunol.173.5.2909
 57. Gombart AF, Borregaard N, Koeffler HP. Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D3. *FASEB J* (2005) 19:1067–77. doi:10.1096/fj.04-3284com
 58. Harding CV, Boom WH. Regulation of antigen presentation by *Mycobacterium tuberculosis*: a role for toll-like receptors. *Nat Rev Microbiol* (2010) 8:296–307. doi:10.1038/nrmicro2321
 59. Martineau AR, Wilkinson KA, Newton SM, Floto RA, Norman AW, Skolimowska K, et al. IFN-gamma- and TNF-independent vitamin D-inducible human suppression of mycobacteria: the role of cathelicidin LL-37. *J Immunol* (2007) 178:7190–8. doi:10.4049/jimmunol.178.11.7190
 60. Adams JS, Ren S, Liu PT, Chun RF, Lagishetty V, Gombart AF, et al. Vitamin D-directed rheostatic regulation of monocyte antibacterial responses. *J Immunol* (2009) 182:4289–95. doi:10.4049/jimmunol.0803736
 61. Hmam Z, Pena-Diaz S, Joseph S, Av-Gay Y. Immuno-evasion and immunosuppression of the macrophage by *Mycobacterium tuberculosis*. *Immunol Rev* (2015) 264:220–32. doi:10.1111/immr.12268
 62. Gopalakrishnan A, Salgame P. Toll-like receptor 2 in host defense against *Mycobacterium tuberculosis*: to be or not to be—that is the question. *Curr Opin Immunol* (2016) 42:76–82. doi:10.1016/j.coi.2016.06.003
 63. Richardson ET, Shukla S, Sweet DR, Wearsch PA, Tschlis PN, Boom WH, et al. Toll-like receptor 2-dependent extracellular signal-regulated kinase signaling in *Mycobacterium tuberculosis*-infected macrophages drives anti-inflammatory responses and inhibits Th1 polarization of responding T cells. *Infect Immun* (2015) 83:2242–54. doi:10.1128/IAI.00135-15
 64. Bradfute SB, Castillo EF, Arko-Mensah J, Chauhan S, Jiang S, Mandell M, et al. Autophagy as an immune effector against tuberculosis. *Curr Opin Microbiol* (2013) 16:355–65. doi:10.1016/j.mib.2013.05.003
 65. Kim JK, Yuk JM, Kim SY, Kim TS, Jin HS, Yang CS, et al. MicroRNA-125a inhibits autophagy activation and antimicrobial responses during mycobacterial infection. *J Immunol* (2015) 194:5355–65. doi:10.4049/jimmunol.1402557

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Rode, Kongsbak, Hansen, Lopez, Levring, Woetmann, Odum, Bonefeld and Geisler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.