



Corrigendum: Natural Killer Cell-Derived Exosomal miR-3607-3p Inhibits Pancreatic Cancer Progression by Targeting IL-26

Hongwei Sun^{1†}, Keqing Shi^{2†}, Kai Qi^{3†}, Hongru Kong¹, Jie Zhang¹, Shengjie Dai¹, Wen Ye¹, Tuo Deng¹, Qiye He^{4,5} and Mengtao Zhou^{6*}

OPEN ACCESS

Edited and reviewed by:

Katy Rezvani,
University of Texas MD Anderson
Cancer Center, United States

*Correspondence:

Mengtao Zhou
qianazi@yeah.net

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cancer Immunity and Immunotherapy,
a section of the journal
Frontiers in Immunology

Received: 01 January 2020

Accepted: 04 February 2020

Published: 21 February 2020

Citation:

Sun H, Shi K, Qi K, Kong H, Zhang J,
Dai S, Ye W, Deng T, He Q and
Zhou M (2020) Corrigendum: Natural
Killer Cell-Derived Exosomal
miR-3607-3p Inhibits Pancreatic
Cancer Progression by Targeting
IL-26. *Front. Immunol.* 11:277.
doi: 10.3389/fimmu.2020.00277

¹ Department of Hepatobiliary Surgery, Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China, ² Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, Center of Precision Medicine, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China, ³ Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, Shanghai, China, ⁴ Singlera Genomics Inc., San Diego, CA, United States, ⁵ Singlera Genomics (Shanghai) Ltd., Shanghai, China, ⁶ Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, Precision Medical Center Laboratory, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China

Keywords: natural killer, pancreatic cancer, extracellular vesicles, miR-3607-3p, IL-26

A Corrigendum on

Natural Killer Cell-Derived Exosomal miR-3607-3p Inhibits Pancreatic Cancer Progression by Targeting IL-26

by Sun, H., Shi, K., Qi, K., Kong, H., Zhang, J., Dai, S., et al. (2019). *Front. Immunol.* 10:2819. doi: 10.3389/fimmu.2019.02819

In the original article, there was a mistake in **Figure 1** and **Figure 5** as published. Figures 1E and Figure 5F were wrongly placed. The corrected **Figure 1** and **Figure 5** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Sun, Shi, Qi, Kong, Zhang, Dai, Ye, Deng, He and Zhou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 | NK cells co-culture inhibited tumor progression of pancreatic cancer both *in vitro* and *in vivo*. **(A,B)** CCK-8 assay showed cell viability of Mia PaCa-2 and PANC-1 cells co-cultured with (NK cells+) or without (NK cells-) natural killer cells. **(C)** Colony formation assay showed cell proliferation of Mia PaCa-2 and PANC-1 cells co-cultured with (NK cells+) or without (NK cells-) natural killer cells. **(D,E)** Transwell assays showed cell migratory and invasive ability of Mia PaCa-2 and PANC-1 cells co-cultured with (NK cells+) or without (NK cells-) natural killer cells. **(F,G)** PANC-1 cells were implanted into the flank of mice ($n = 4$ each group), without (NK cell-) or with co-injection of natural killer cells (NK cell+), respectively, followed by growth curve evaluation on the indicated day after injection. **(H-J)** Representative *in vivo* images showed tumor colonization in the lungs of mice ($n = 5$ each group) following tail vein injection of PANC-1 cells, without (NK cell-) or with co-injection of natural killer cells (NK cell+), respectively, H&E staining of lung sections of mice (metastatic nodules were indicated by yellow arrow, 200 \times) and incidence of lung metastasis in mice following tail vein injection of the respective PANC-1 cells. The data represent the mean \pm SD from three independent experiments. $**P < 0.01$; $***P < 0.001$, two-way ANOVA for **(A,B,G)**, χ^2 test for **J**, Student's *t*-test for others.

