



ramR mutations affecting fluoroquinolone susceptibility in epidemic multidrug-resistant *Salmonella enterica* serovar Kentucky ST198

Sylvie Baucheron^{1,2}, Simon Le Hello³, Benoît Doublet^{1,2}, Etienne Giraud^{1,2}, François-Xavier Weill³ and Axel Cloeckaert^{1,2*}

¹ INRA, UMR1282 Infectiologie et Santé Publique, Nouzilly, France

² Université François Rabelais de Tours, UMR1282 Infectiologie et Santé Publique, Tours, France

³ Institut Pasteur, Unité des Bactéries Pathogènes Entériques, Centre National de Référence des *Escherichia coli*, *Shigella* et *Salmonella*, Paris, France

Edited by:

Michel S. Zygmunt, Institut National de la Recherche Agronomique, France

Reviewed by:

Kunihiko Nishino, Osaka University, Japan

Seamus Fanning, University College Dublin, Ireland

*Correspondence:

Axel Cloeckaert, Unité Infectiologie et Santé Publique site 213, Institut National de la Recherche Agronomique, 37380 Nouzilly, France
e-mail: axel.cloeckaert@tours.inra.fr

A screening for non-target mutations affecting fluoroquinolone susceptibility was conducted in epidemic multidrug-resistant *Salmonella enterica* serovar Kentucky ST198. Among a panel of representative isolates ($n = 27$), covering the epidemic, only three showed distinct mutations in *ramR* resulting in enhanced expression of genes encoding the AcrAB-TolC efflux system and low increase in ciprofloxacin MIC. No mutations were detected in other regulatory regions of this efflux system. Ciprofloxacin resistance in serovar Kentucky ST198 is thus currently mainly due to multiple target gene mutations.

Keywords: *Salmonella*, ciprofloxacin resistance, efflux pump, regulation, ram

INTRODUCTION

Fluoroquinolones, together with extended-spectrum cephalosporins, are the treatment of choice for nontyphoid salmonellosis, as stable resistance to the most common members of different families of antimicrobial agents (ampicillin, chloramphenicol, streptomycin, sulfonamides, and tetracycline) has developed during the 1990s with the epidemic *Salmonella enterica* serovar Typhimurium phage type DT104 (Cloeckaert and Schwarz, 2001). Emerging resistance to fluoroquinolones in *Salmonella* spp. has been reported for both human and animal cases and is thus threatening to become a serious public health problem (Cloeckaert and Chaslus-Dancla, 2001; Piddock, 2002; Velge et al., 2005; Giraud et al., 2006). Of particular concern is the international spread of ciprofloxacin-resistant serovar Kentucky ST198 (Le Hello et al., 2011). This clone is not only highly resistant to ciprofloxacin but also multidrug-resistant (MDR) due to the presence of the *Salmonella* genomic island 1 (SGI1) carrying a multiple antibiotic resistance gene cluster, mostly variant SGI1-K carrying another resistance gene cluster (Doublet et al., 2008; Le Hello et al., 2011). SGI1 was initially identified in MDR serovar Typhimurium DT104 (Boyd et al., 2001), but nor the MDR serovar Typhimurium DT104 clone neither other MDR *S. enterica* serovars carrying SGI1 or variants of it, have to our knowledge been reported to display this high-level ciprofloxacin resistance.

In *Salmonella* spp., quinolone/fluoroquinolone resistance is mostly attributed to point mutations in the quinolone resistance-determining regions (QRDRs) of the target genes *gyrA*, *gyrB*, *parC*, and *parE*. For the *gyrA* gene, coding for the A subunit of

DNA gyrase, mutations resulting in amino acid changes at Ser83 (to Phe, Tyr, or Ala) or at Asp87 (to Gly, Asn, or Tyr) are the most frequently observed in nalidixic acid-resistant strains (Cloeckaert and Chaslus-Dancla, 2001; Piddock, 2002; Velge et al., 2005; Giraud et al., 2006). High-level fluoroquinolone resistance has been reported in several *S. enterica* serovars (Choleraesuis, Schwarzengrund, Typhimurium) and is essentially due to the combination of several target gene mutations of which the most frequent are double mutations resulting in modifications of both residues 83 and 87 of GyrA together with one mutation leading to the amino acid change Ser80Ile in the ParC subunit of topoisomerase IV (Baucheron et al., 2002, 2004; Chu et al., 2005). In addition two main other mechanisms have been reported consisting of active efflux mediated by the chromosomally-encoded AcrAB-TolC efflux system and target protection by Qnr proteins which are mostly encoded by plasmids acquired by horizontal transfer (Giraud et al., 2006). However, according to the literature over 15 years, these mechanisms appear less frequently and thus from an epidemic point of view seem of lesser importance than multiple target gene mutations to reach high-level ciprofloxacin resistance and compromise treatment.

In the case of ciprofloxacin resistance in serovar Kentucky ST198, three combinations of multiple target modifications, acquired in a possible sequential way, have been reported consisting of a first GyrA Ser83Phe modification, followed by three different situations of a second GyrA modification at position 87, i.e., Asp87Asn, Asp87Gly, or Asp87Tyr, and finally the ParC modification Ser80Ile (Le Hello et al., 2011). Qnr proteins have not been reported yet as additional mechanism for this epidemic

Table 1 | *Salmonella enterica* serovar Kentucky ST198 strains analyzed in this study.

Strain	Country	Year of isolation	Antimicrobial resistance profile	SGI1	PFGE type	CIP MIC ($\mu\text{g/ml}$)	Substitution(s) in the QRDR of:		Mutation(s) in efflux pump regulatory regions	AcrA production ratio
							GyrA	ParC		
00 1059	Egypt	2000	AMX NAL	+	(SGI1-P1) XKEN-1a	0.125	S83F	None	–	3
01 2100	Egypt	2001	AMX STR SPT GEN SUL TET NAL	+	(SGI1-K1) XKEN-1a	0.125	S83F	None	–	2
02 2818	Egypt	2002	AMX STR SPT GEN SUL TET NAL	+	XKEN-1i	0.5	S83F	None	+	(ramR) 5
02 2691	Egypt	2002	AMX STR SPT GEN SUL TET NAL	+	(SGI1-K3) XKEN-1a	0.125	S83F	None	–	1
02 8051	Egypt	2002	AMX STR SPT GEN SUL TET NAL	+	XKEN-1a	0.25	S83F	None	–	1
02 8141	Egypt	2002	AMX STR SPT GEN SUL TET NAL	+	(SGI1-K1) XKEN-1m	0.5	S83F	None	+	(ramR) 5
02 9866	Egypt	2002	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1a	8	S83F, D87N	S80I	–	2
03 9270	India	2003	NAL	–	XKEN-2d	0.125	S83F	None	–	1
04 2049	Egypt	2004	NAL CIP	+	XKEN-1b	8	S83F, D87G	S80I	–	2
04 4567	Egypt	2004	AMX STR SPT GEN SUL TET NAL CIP	+	(SGI1-K1) XKEN-1g	4	S83F, D87G	S80I	–	2
04 6248	Egypt	2004	STR SPT GEN SUL TET NAL CIP	+	XKEN-1a	8	S83F, D87G	S80I	–	1
04 7734	Egypt	2004	AMX STR SPT GEN SUL TET NAL	+	(SGI1-K1) XKEN-1h	0.5	S83F	None	–	1
04 8262	Egypt	2004	STR SPT GEN SUL NAL CIP	+	(SGI1-K5) XKEN-1a	8	S83F, D87N	S80I	–	1
04 9384	Egypt	2004	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1g	4	S83F, D87G	S80I	–	1
05 0490	Egypt	2005	STR SPT GEN SUL TET NAL CIP	+	XKEN-1a	4	S83F, D87G	S80I	–	2
05 0520	Egypt	2005	AMX NAL CIP	+	(SGI1-P2) XKEN-1a	4	S83F, D87Y	S80I	–	1
05 1016	Kenya	2005	NAL CIP	+	(SGI1-Q2) XKEN-1a	4	S83F, D87Y	S80I	–	1
05 1199	Egypt	2005	STR SPT GEN SUL NAL CIP	+	(SGI1-Q3) XKEN-1a	4	S83F, D87G	S80I	–	1
05 2131	Egypt	2005	AMX NAL CIP	+	(SGI1-Q1) XKEN-1a	4	S83F, D87N	S80I	–	3
05 2354	Kenya/ Tanzania	2005	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1c	8	S83F, D87Y	S80I	–	3
05 3290	Egypt	2005	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1c	4	S83F, D87G	S80I	–	1
05 3883	Kenya/ Tanzania	2005	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1d	4	S83F, D87Y	S80I	–	2
05 4680	Sudan	2005	STR SPT GEN SUL TET NAL CIP	+	(SGI1-K4) XKEN-1i	4	S83F, D87G	S80I	–	2
05 7714	Unknown	2005	AMX NAL CIP	+	XKEN-1b	4	S83F, D87N	S80I	–	2
05 8560	Tunisia	2005	AMX STR SPT GEN SUL TET NAL CIP	+	XKEN-1d	16	S83F, D87N	S80I	+	(ramR) 6
05 236	Egypt	2005	AMX NAL CIP	+	XKEN-1c	4	S83F, D87N	S80I	–	1
05 5111	Libya	2005	AMX SUL TET NAL CIP	+	(SGI1-K2) XKEN-1a	4	S83F, D87N	S80I	–	1

clone, and active efflux has been suspected in a previous study due to a moderate increase of production in some isolates of the AcrA protein belonging to the AcrAB-TolC efflux system (Weill et al., 2006).

In the present study we assessed the frequency of enhanced efflux by AcrAB-TolC in a subset of serovar Kentucky ST198 strains of the 2000–2005 period of the epidemic. In case of significant increased production of AcrAB-TolC we investigated more deeply the regulatory mechanisms behind this overproduction, in particular the involvement of the *ram*, *sox*, and *mar* regulatory loci (Abouzeed et al., 2008; Kehrenberg et al., 2009). Among these loci, the *ramRA* locus appears to be the most important in regulating AcrAB-TolC expression in *Salmonella* spp. (Abouzeed et al., 2008; Kehrenberg et al., 2009). *ramR* encodes a repressor protein (RamR) belonging to the TetR family of repressor proteins, and has been shown to be the local repressor protein of *ramA* transcription (Abouzeed et al., 2008; Baucheron et al., 2012); while *ramA* encodes a transcriptional activator protein (Rama) belonging to the AraC/XylS family of regulatory proteins (Nikaido et al., 2008). The latter is involved in upregulating expression

of the AcrAB-TolC system (Nikaido et al., 2008). Several mutations in *ramR* or its binding site upstream of *ramA*, affecting expression of this efflux system, have been detected in clinical isolates of serovar Typhimurium and of minor serovars Hadar, Infantis, Livingstone, or Schwarzengrund (Abouzeed et al., 2008; Kehrenberg et al., 2009; Hentschke et al., 2010; Akiyama and Khan, 2012).

MATERIALS AND METHODS

The 27 serovar Kentucky ST198 strains selected for this study are shown in **Table 1**. Bacterial isolates were selected for this study, based on their evolutionary history following the emergence of target gene mutations initially in *gyrA* at the commencement of the epidemic in 2000–2002, followed by isolates with additional mutations (in *gyrA* and *parC*) toward the end in 2002–2005 and which demonstrated a higher MIC toward ciprofloxacin. An additional criterion for selection consisted of the differences observed in ciprofloxacin MICs suggestive for another resistance mechanism than target gene mutation. MICs were determined as described previously (Baucheron et al., 2002, 2004). SGI1

Table 2 | Primers used for PCRs.

Primer used and target region	Primer	Nucleotide position relative to the LT2 strain genome*	Oligonucleotide sequences(s) (5' to 3')	Size (bp)	Annealing temp (°C)	References
DETECTION OF MUTATIONS						
<i>ramR-ramA</i>	ram5	638085	TCGGTAAAAGGCAGTTCCAG	958	60	This study
	ramA6	639042	GTCGATAACCTGAGCGGAAA			
<i>acrR-acrA</i>	acrR1	533463	CAGTGGTTCCGTTTTAGTG	992	58	Olliver et al., 2005
	acrR2	534454	ACAGAATAGCGACACAGAAA			
<i>marC-marO-marR-marA</i>	marR1	1597459	CAGTGTTCGCTCTGGACATC	787	60	This study
	marR2	1598245	GCTAACGGGAGCAGTACGAC			
<i>soxS-soxR</i>	sox1	4503970	CTACAGGCGGTGACGGTAAT	915	60	This study
	sox2	4504884	CGGCGCTTTAGTTTTAGGTG			
<i>acrS-acrE</i>	acrS1	3560054	TTGGCATTAAATGCCTCACA	1094	62	This study
	acrS2	3561128	ATGATGAATGAGGGCAGGAG			
qRT-PCR						
<i>gmk</i>	gmk-f	3933294	TTGGCAGGGAGGCGTTT	62	60	Baucheron et al., 2012
	gmk-r	3933355	GCGCGAAGTGCCGTAGTAAT			
<i>gyrB</i>	gyrB-f	4040275	TCTCCTCACAGACCAAGATAAGCT	81	60	Baucheron et al., 2012
	gyrB-r	4040195	CGCTCAGCAGTTCGTTTCATC			
<i>rrs</i>	rrs-f	NA**	CCAGCAGCCGCGGTAAT	57	60	Baucheron et al., 2012
	rrs-r	NA**	TTTACGCCAGTAATCCGATT			
<i>ramA</i>	ramA-f	639180	GCGTGAACGGAAGCTAAAAC	167	60	Baucheron et al., 2012
	ramA-r	639346	GGCCATGCTTTTCTTACGA			
<i>acrA</i>	acrA-f	533120	GAAACCGCACGTATCAACCT	220	60	Baucheron et al., 2012
	acrA-r	532901	CCTGTTTCAGCGAACCATTT			
<i>tolC</i>	tolC-f	3349107	GCCCGTGCGCAATATGAT	67	60	Baucheron et al., 2012
	tolC-r	3349173	CCGCGTTATCCAGGTTGTTG			

* GenBank NC_003197.1.

**NA: Not Applicable due to the number of copies of this gene in *Salmonella*.

detection and characterization were performed as described previously (Boyd et al., 2001; Doublet et al., 2008). Efflux pump production was assessed by Dot blot using an anti-AcrA polyclonal antibody as described previously (Abouzeed et al., 2008). Occurrence of mutations affecting *acrAB* and *tolC* expression was determined by PCR and sequencing the regulatory regions *ramR-ramA*, *acrR-acrA*, *marC-marO-marR-marA*, *soxS-soxR*, and *acrS-acrE* using primers listed in **Table 2**. Transcription levels of *ramA*, *acrA*, and *tolC* were determined by qRT-PCR as described previously (Giraud et al., 2013).

RESULTS AND DISCUSSION

As shown in the **Table 1** most of the strains selected carried SGI1 or variants of it and were thus MDR. They were all from human cases in France who acquired their infection during travel to Africa or India. As assessed by Dot blot, most of the strains ($n = 24$) did not show significant increased production of AcrA relative to susceptible serovar Kentucky reference strain 98K (AcrA production ratios from 1 to 2; **Table 1**). Relative to strain 98K, three strains showed a 3-fold increased AcrA production, and more suggestive for increased active efflux three strains a 5- to 6-fold increased production of AcrA (**Table 1**). Among these regulatory regions, mutations were detected only in the *ramR* open reading frame and in only three strains of this

study (**Table 3**). The mutations were distinct frame shift mutations and consisted of a GATC duplication for strain 02-2818, a G insertion for strain 05-8560, and a 91 bp deletion for strain 02-8141 (**Figure 1**). The role of these mutations in upregulating *acrAB* and *tolC* expression, and consecutive enhanced efflux-mediated resistance, was further assessed by: (i) complementing with the wild-type *ramR* gene (using plasmid pRamR Abouzeed et al., 2008); (ii) determining the MICs of ciprofloxacin and unrelated antibiotic florfenicol shown to be substrate of AcrAB-TolC (Baucheron et al., 2002); and (iii) measuring expression of *ramA*, *acrA*, and *tolC* by qRT-PCR (Giraud et al., 2013). The results shown in **Table 3** are in agreement with data published previously for other *S. enterica* serovars (Abouzeed et al., 2008; Kehrenberg et al., 2009), i.e. *ramR* mutations observed account for a 2- to 4-fold increased resistance level by active efflux through enhanced expression of AcrAB-TolC. As also observed in previous studies, the effect of such mutations on *ramA* transcription level was significantly higher than on *acrA* or *tolC* transcription levels. It is somehow expected considering the direct local repressor activity of RamR on *ramA* transcription and the distant RamA transcriptional activator activity on *acrAB* and *tolC* (Abouzeed et al., 2008; Baucheron et al., 2012; Giraud et al., 2013).

Non-target mutations as assessed in this study confirm they are infrequent in *Salmonella* spp. but seem nevertheless


Table 3 | Characteristics of the *Salmonella enterica* serovar Kentucky ST198 strains carrying ramR mutations.

Strain	Source	Geographic origin	Antimicrobial resistance profile ^a			PFGE type	SGI1 (variant) ^b	MIC of indicated antibiotic (μ g/ml)			Substitution(s) in the QRDR of:		Mutation in ramR	Transcription levels of:									
			AMX	STR	SPT			GEN	SUL	TET	NAL	CIP		FFC ^c	GyrA	ParC	ramA	acrAtoIC					
MDR STRAINS																							
05-8560	Human	Tunisia	AMX	STR	SPT	GEN	SUL	TET	NAL	CIP	XKEN-1d	+	(Ks)	>1024	16	16	S83F	D87Y	S80I	1 bp insertion (position 506)	24.6	7.2	2.6
05-8560(pRamR)														>1024	4	4					2.4	1.7	1.7
02-8141	Human	Egypt	AMX	STR	SPT	GEN	SUL	TET	NAL		XKEN-1m	+	(K1)	512	0.50	16	S83F			91 bp insertion (position 42)	106.1	10.4	7.8
02-8141(pRamR)														512	0.125	8					1.6	1.1	1.2
02-2818	Human	Egypt	AMX	STR	SPT	GEN	SUL	TET	NAL		XKEN-1i	+	(Ks)	512	0.50	16	S83F			4 bp duplication (position 508)	29.1	5.3	4.7
02-2818(pRamR)														256	0.25	4					1.9	0.9	1.6
02-9866	Human	Egypt	AMX	STR	SPT	GEN	SUL	TET	NAL	CIP	XKEN-1a	+	(Ks)	>1024	8	4	S83F	D87N	S80I		2.9	1.2	1.6
02-9866(pRamR)														>1024	4	4					1.8	1.6	2.4
REFERENCE STRAIN																							
98K	Chicken	USA									XKEN-4			1	0.004	2					1.0	1.0	1.0
98K(pRamR)														1	0.004	2					2.1	1.3	1.5

^a AMX, amoxicillin; STR, streptomycin; SPT, spectinomycin; GEN, gentamicin; SUL, sulfonamides; TET, tetracycline; NAL, nalidixic acid; CIP, ciprofloxacin.

^b Ks: subgroup of SGI1-K.

^c FFC, florfenicol.



mostly restricted to the *ram* regulatory region. Most mutations in the *ramR-ramA* region reported to date, as also shown in this study, are distinct and found in single isolates. To our knowledge only independent isolates of the epidemic ciprofloxacin-resistant serovar Typhimurium DT204 clone from the 1990s have been shown to carry the same mutation in *ramR* consisting of an insertion by an *IS1* element (Abouzeed et al., 2008). We may nevertheless expect the further global spread of ciprofloxacin-resistant serovar Kentucky ST198 and its resistance evolution will possibly, like in the

case of serovar Typhimurium DT204, result in successful *ramR*-mutation-carrying subclones.

ACKNOWLEDGMENTS

We are grateful to Isabelle Monchaux and Laetitia Fabre for excellent technical assistance. We would like to thank all corresponding laboratories of the French National Reference Center for *E. coli*, *Shigella*, and *Salmonella*. The French National Reference Center for *E. coli*, *Shigella*, and *Salmonella* is funded by the Institut Pasteur and the Institut de Veille Sanitaire.

REFERENCES

- Abouzeed, Y. M., Baucheron, S., and Cloeckaert, A. (2008). *ramR* mutations involved in efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 52, 2428–2434. doi: 10.1128/AAC.00084-08
- Akiyama, T., and Khan, A. A. (2012). Molecular characterization of strains of fluoroquinolone-resistant *Salmonella enterica* serovar Schwarzengrund carrying multidrug resistance isolated from imported foods. *J. Antimicrob. Chemother.* 67, 101–110. doi: 10.1093/jac/dkr414
- Baucheron, S., Chaslus-Dancla, E., and Cloeckaert, A. (2004). Role of TolC and *parC* mutation in high-level fluoroquinolone resistance in *Salmonella enterica* serotype Typhimurium DT204. *J. Antimicrob. Chemother.* 53, 657–659. doi: 10.1093/jac/dkh122
- Baucheron, S., Coste, F., Canepa, S., Maurel, M. C., Giraud, E., Culard, F., et al. (2012). Binding of the RamR repressor to wild-type and mutated promoters of the *ramA* gene involved in efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 56, 942–948. doi: 10.1128/AAC.05444-11
- Baucheron, S., Imberechts, H., Chaslus-Dancla, E., and Cloeckaert, A. (2002). The AcrB multidrug transporter plays a major role in high-level fluoroquinolone resistance in *Salmonella enterica* serovar Typhimurium phage type DT204. *Microb. Drug Resist.* 8, 281–289. doi: 10.1089/10766290260469543
- Boyd, D. A., Peters, G. A., Cloeckaert, A., Boumedine, K. S., Chaslus-Dancla, E., Imberechts, H., et al. (2001). Complete nucleotide sequence of a 43-kilobase genomic island associated with the multidrug resistance region of *Salmonella enterica* serovar Typhimurium DT104 and its identification in phage type DT120 and serovar Agona. *J. Bacteriol.* 183, 5725–5732. doi: 10.1128/JB.183.19.5725-5732.2001
- Chu, C., Su, L. H., Chu, C. H., Baucheron, S., Cloeckaert, A., and Chiu, C. H. (2005). Resistance to fluoroquinolones linked to *gyrA* and *parC* mutations and overexpression of *acrAB* efflux pump in *Salmonella enterica* serotype Choleraesuis. *Microb. Drug Resist.* 11, 248–253. doi: 10.1089/mdr.2005.11.248
- Cloeckaert, A., and Chaslus-Dancla, E. (2001). Mechanisms of quinolone resistance in *Salmonella*. *Vet. Res.* 32, 291–300. doi: 10.1051/vetres:2001105
- Cloeckaert, A., and Schwarz, S. (2001). Molecular characterization, spread and evolution of multidrug resistance in *Salmonella enterica* typhimurium DT104. *Vet. Res.* 32, 301–310. doi: 10.1051/vetres:2001126
- Doublet, B., Praud, K., Bertrand, S., Collard, J. M., Weill, F. X., and Cloeckaert, A. (2008). Novel insertion sequence- and transposon-mediated genetic rearrangements in genomic island SG11 of *Salmonella enterica* serovar Kentucky. *Antimicrob. Agents Chemother.* 52, 3745–3754. doi: 10.1128/AAC.00525-08
- Giraud, E., Baucheron, S., and Cloeckaert, A. (2006). Resistance to fluoroquinolones in *Salmonella*: emerging mechanisms and resistance prevention strategies. *Microbes Infect.* 8, 1937–1944. doi: 10.1016/j.micinf.2005.12.025
- Giraud, E., Baucheron, S., Virlogeux-Payant, I., Nishino, K., and Cloeckaert, A. (2013). Effects of natural mutations in the *ramRA* locus on invasiveness of epidemic fluoroquinolone-resistant *Salmonella enterica* serovar Typhimurium isolates. *J. Infect. Dis.* 207, 794–802. doi: 10.1093/infdis/jis755
- Hentschke, M., Christner, M., Sobottka, I., Aepfelbacher, M., and Rohde, H. (2010). Combined *ramR* mutation and presence of a Tn1721-associated *tet(A)* variant in a clinical isolate of *Salmonella enterica* serovar Hadar resistant to tigecycline. *Antimicrob. Agents Chemother.* 54, 1319–1322. doi: 10.1128/AAC.00993-09
- Kehrenberg, C., Cloeckaert, A., Klein, G., and Schwarz, S. (2009). Decreased fluoroquinolone susceptibility in mutants of *Salmonella* serovars other than Typhimurium: detection of novel mutations involved in modulated expression of *ramA* and *soxS*. *J. Antimicrob. Chemother.* 64, 1175–1180. doi: 10.1093/jac/dkp347
- Le Hello, S., Hendriksen, R. S., Doublet, B., Fisher, I., Nielsen, E. M., Whichard, J. M., et al. (2011). International spread of an epidemic population of *Salmonella enterica* serotype Kentucky ST198 resistant to ciprofloxacin. *J. Infect. Dis.* 204, 675–684. doi: 10.1093/infdis/jir409
- Nikaido, E., Yamaguchi, A., and Nishino, K. (2008). AcrAB multidrug efflux pump regulation in *Salmonella enterica* serovar Typhimurium by RamA in response to environmental signals. *J. Biol. Chem.* 283, 24245–24253. doi: 10.1074/jbc.M804544200

- Olliver, A., Vallé, M., Chaslus-Dancla, E., and Cloeckaert, A. (2005). Overexpression of the multidrug efflux operon acrEF by insertional activation with IS1 or IS10 elements in *Salmonella enterica* serovar typhimurium DT204 acrB mutants selected with fluoroquinolones. *Antimicrob. Agents Chemother.* 49, 289–301. doi: 10.1128/AAC.49.1.289-301.2005
- Piddock, L. J. V. (2002). Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiol. Rev.* 26, 3–16.
- Velge, P., Cloeckaert, A., and Barrow, P. (2005). Emergence of *Salmonella* epidemics: the problems related to *Salmonella enterica* serotype Enteritidis and multiple antibiotic resistance in other major serotypes. *Vet. Res.* 36, 267–288. doi: 10.1051/vetres:2005005
- Weill, F. X., Bertrand, S., Guesnier, F., Baucheron, S., Cloeckaert, A., and Grimont, P. A. (2006). Ciprofloxacin-resistant *Salmonella* Kentucky in travelers. *Emerg. Infect. Dis.* 12, 1611–1612. doi: 10.3201/eid1210.060589
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 21 June 2013; accepted: 09 July 2013; published online: 31 July 2013.
Citation: Baucheron S, Le Hello S, Doublet B, Giraud E, Weill F-X and Cloeckaert A (2013) ramR mutations affecting fluoroquinolone susceptibility in epidemic multidrug-resistant *Salmonella enterica* serovar Kentucky ST198. *Front. Microbiol.* 4:213. doi: 10.3389/fmicb.2013.00213
- This article was submitted to *Frontiers in Antimicrobials, Resistance and Chemotherapy*, a specialty of *Frontiers in Microbiology*. Copyright © 2013 Baucheron, Le Hello, Doublet, Giraud, Weill and Cloeckaert. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.