



A Review of SHV Extended-Spectrum β -Lactamases: Neglected Yet Ubiquitous

Apostolos Liakopoulos¹, Dik Mevius^{1,2} and Daniela Ceccarelli^{1*}

¹ Department of Bacteriology and Epidemiology, Central Veterinary Institute of Wageningen UR, Lelystad, Netherlands,

² Faculty of Veterinary Medicine, Utrecht University, Utrecht, Netherlands

OPEN ACCESS

Edited by:

Axel Cloeckaert,
French National Institute for
Agricultural Research (INRA), France

Reviewed by:

Sebastian Guenther,
Free University of Berlin, Germany
David Wareham,
Queen Mary University of London, UK

*Correspondence:

Daniela Ceccarelli
daniela.ceccarelli@wur.nl

Specialty section:

This article was submitted to
Antimicrobials, Resistance and
Chemotherapy,
a section of the journal
Frontiers in Microbiology

Received: 20 July 2016

Accepted: 19 August 2016

Published: 05 September 2016

Citation:

Liakopoulos A, Mevius D and
Ceccarelli D (2016) A Review of SHV
Extended-Spectrum β -Lactamases:
Neglected Yet Ubiquitous.
Front. Microbiol. 7:1374.
doi: 10.3389/fmicb.2016.01374

β -lactamases are the primary cause of resistance to β -lactams among members of the family Enterobacteriaceae. SHV enzymes have emerged in Enterobacteriaceae causing infections in health care in the last decades of the Twentieth century, and they are now observed in isolates in different epidemiological settings both in human, animal and the environment. Likely originated from a chromosomal penicillinase of *Klebsiella pneumoniae*, SHV β -lactamases currently encompass a large number of allelic variants including extended-spectrum β -lactamases (ESBL), non-ESBL and several not classified variants. SHV enzymes have evolved from a narrow- to an extended-spectrum of hydrolyzing activity, including monobactams and carbapenems, as a result of amino acid changes that altered the configuration around the active site of the β -lactamases. SHV-ESBLs are usually encoded by self-transmissible plasmids that frequently carry resistance genes to other drug classes and have become widespread throughout the world in several Enterobacteriaceae, emphasizing their clinical significance.

Keywords: β -lactamase, ESBL, *bla*_{SHV}, SHV-2, SHV-5, SHV-12, plasmid, Enterobacteriaceae

INTRODUCTION

Thanks to their ability to inhibit cell wall biosynthesis, β -lactams remained the first-line defense against bacterial infections for over 20 years, before resistant bacteria appeared in clinical practice.

Resistance to this class of drugs can be the result of antibiotic target site alteration, prevention of antibiotic access by altered permeability or forced efflux, or antibiotic degradation (Wilke et al., 2005). The latter, represents the primary resistance mechanism in Gram-negative bacteria producing β -lactamase enzymes able to covalently bind the carbonyl moiety of the β -lactam ring and hydrolyze its amide bond (Fisher et al., 2005). Naturally occurring chromosomally located β -lactamases are quite common in Gram-negative bacteria; likely evolved from penicillin-binding proteins, when produced in small quantity they do not significantly contribute to antibiotic resistance. It was the appearance of the first plasmid-mediated β -lactamase TEM-1 (Datta and Kontomichalou, 1965) to designate the beginning of an unstoppable phenomenon in the 1960s. Ever since, the introduction of new natural or synthetic drugs to replace old ones in an attempt to limit the insurgence of antibiotic resistant bacteria triggered a chain reaction providing bacteria with a constant selective pressure driving the expansion of different resistance mechanisms (Medeiros, 1997).

In recent years β -lactamases have extensively diversified in response to the clinical use of new generations of β -lactams (penicillin, cephalosporins, carbapenems, and monobactams) leading

to the need of classification schemes. Based on primary structure (Ambler, 1980), enzymatic properties and biochemical attributes (Bush et al., 1995), and the increasingly available amino acid sequences (Bush and Jacoby, 2010) four major classes (A, B, C, D) can be acknowledged. Serine β -lactamases belonging to class A are the most abundant (Philippon et al., 2016), with more than 500 enzymes, including the most clinically significant extended spectrum β -lactamases (ESBL) variants, i.e., CTX-M-, TEM-, and SHV-type enzymes (Bush and Fisher, 2011).

Although, SHV enzymes did not undergo the explosive dissemination observed for CTX-M-type variants (Canton et al., 2012), in recent years they have been found in several Enterobacteriaceae outside of the typical clinical hosts *Klebsiella pneumoniae* and *Escherichia coli*, with a rising allele variability (<http://www.lahey.org/studies>), and in different environmental niches. Many admirable works describing the biochemistry, the genetics and the evolution of SHV β -lactamases have appeared over the last years. The aim of this review is to provide the readers with an updated overview on SHV β -lactamases, their amino acid variants and spectrum of activity, and to describe the occurrence of plasmid-associated SHV enzymes in Enterobacteriaceae and their epidemiological significance.

ORIGIN AND DIVERSITY OF THE SHV FAMILY

The first *bla*_{SHV-1} gene was identified in the 1970s in *E. coli* (Pitton, 1972). The encoded enzyme SHV-1 (sulfhydryl reagent variable) proved its activity against penicillins and first generation cephalosporins (Matthew et al., 1979) and was confirmed part of the conjugative plasmid p453 (Barthélémy et al., 1988; **Table 1**). The most likely ancestor of the plasmid-mediated SHV-1 is a chromosomal species-specific penicillinase detected in fecal *K. pneumoniae* isolates from neonates (Haeggman et al., 1997). The enzyme showed a typical antibiogram with penicillin rather than cephalosporin resistance and a marked inhibition by clavulanic acid. How *bla*_{SHV-1} moved from the chromosome to the plasmid does not have a conclusive explanation since the proposed association with a transposable element (Nugent and Hedges, 1979) has not been confirmed.

As of today, 189¹ SHV allelic variants have been described, having developed resistance to 3rd generation cephalosporin (Tzouveleki and Bonomo, 1999), monobactam and carbapenems (Poirel et al., 2003). Only a small proportion is biochemically and/or genetically characterized (<http://www.lahey.org/studies>). SHV β -lactamases can be divided into three subgroups on the basis of molecular characteristics or functional properties: (i) subgroup 2b ($n = 37$), able to hydrolyze penicillins and early cephalosporins (cephaloridine and cephalothin) and strongly inhibited by clavulanic acid and tazobactam; (ii) subgroup 2br ($n = 7$), broad-spectrum β -lactamases that acquired resistance to clavulanic acid; and (iii) subgroup 2be ($n = 46$), comprises ESBLs that can also hydrolyze one or more oxymino β -lactams (cefotaxime, ceftazidime, and aztreonam).

¹Of the 194 variants available on line (<http://www.lahey.org/studies>), 5 were withdrawn or invalidated as only partial sequence.

More than half of these variants ($n = 99$) has not been classified yet due to absence of biochemical characterization.

Figure 1 illustrates a phylogenetic analysis of 149 out of the 189 SHV β -lactamase variants whose amino acid sequences were available online (<http://www.lahey.org/studies>), as of July 2016. Unlike other β -lactamase families (D'andrea et al., 2013; Evans and Amyes, 2014), there is no clear clustering of the different subgroups, as also mirrored by gene based analysis (Supplementary Figure S1). Among the majority of unclassified variants, subgroup 2b and the few 2br variants are scattered all over the tree. Subgroup 2be showed clustering of most of the ESBL variants (including SHV-2a, SHV-5, and SHV-12), together with few non-classified enzymes (SHV-29, SHV-152, SHV-153, SHV-160, and SHV-165). It has been proposed that SHV β -lactamases descended from an unidentified ancestor holding an extended spectrum phenotype (2be) and that subgroup 2b derived from it (Hall and Barlow, 2004). Our analysis showed that several of SHV ESBL variants were scattered along the tree with short branch lengths with neighboring 2b or unknown variants within the SHV phylogeny (i.e., SHV-40, SHV-11, and SHV-35; **Figure 1**), supporting the hypothesis that they evolved from multiple variants, probably within the antibiotic era. Among the non-ESBL variants, *bla*_{SHV-11} represents one of the most successful and, together with *bla*_{SHV-1}, the likely source of evolution for the existing SHV ESBL variants. *bla*_{SHV-11} was first identified as plasmid-encoded in clinical *K. pneumoniae* from Switzerland (Nüesch-Inderbinen et al., 1997) and ever since has been isolated worldwide.

Although, nearly displaced, together with TEM, by CTX-M enzymes over the years (Canton et al., 2012), 46 ESBL *bla*_{SHV} genes have been described so far (**Table 1**). The first report of SHV-mediated resistance to third-generation cephalosporins was in 1983 with the isolation and characterization of *bla*_{SHV-2}, encoded by plasmid pBP60 in a German clinical isolate of *Klebsiella ozaenae* and showing only a few nucleotide mismatches with *bla*_{SHV-1} (Kliebe et al., 1985). In a few years four other ESBL variants were identified as plasmid-encoded in clinical *K. pneumoniae*, showing variable gene homologies with the *bla*_{SHV-1} and *bla*_{SHV-2} sequences (50–90%): *bla*_{SHV-2a} encoded by conjugative plasmid pZMPI (Podbielski et al., 1991); *bla*_{SHV-3} on pUD18 (Nicolas et al., 1989); *bla*_{SHV-4}, widely disseminated from France as a result of a single *K. pneumoniae* clone diffusion (Arlet et al., 1990, 1994); and *bla*_{SHV-5} able to hydrolyze broad-spectrum cephalosporins and monobactams (Gutmann et al., 1989). Of these first variants, the most epidemiologically successful were *bla*_{SHV-2a} and *bla*_{SHV-5}, which will be further discussed, together with *bla*_{SHV-2} and *bla*_{SHV-12}, in a dedicated paragraph (Section Expansion toward New Ecological Niches). Interestingly, *bla*_{SHV-3} and *bla*_{SHV-4} have been only sporadically detected since their first description. *bla*_{SHV-3} seems to be geographically restricted to the USA where it was detected in *E. coli* of animal origin, associated with other antibiotic resistance genes such as *bla*_{CTX-M-15}, *bla*_{CTX-M-24}, *bla*_{CMY-2}, and/or *bla*_{TEM-1} (Shaheen et al., 2011). *bla*_{SHV-4} was identified also in *Enterobacter aerogenes* and *Citrobacter diversus* in different countries (Arpin et al., 1996; El Harrif-Heraud et al., 1997; Pitout et al., 1998).

TABLE 1 | SHV-type extended-spectrum β -lactamases.

Gene ^s	Accession Number	pI	Isolation		Bacterial Species	Genetic background				References
			Location	Year ^a		Genetic Location [‡]	Conjugative plasmid	Plasmid (Kb)	Other Ab genes	
<i>blaSHV-1</i> **	AF148850	7.6	NA	1972	<i>E. coli</i>	p453	Yes	ND	ND	Pitton, 1972; Matthew et al., 1979
<i>blaSHV-2</i>	AF148851	7.6	Germany	1983	<i>K. ozaenae</i>	pBP60	Yes	45	ND	Kliebe et al., 1985
<i>blaSHV-2a</i>	X98102	7.6	Germany	1987–1988	<i>K. pneumoniae</i>	pZMP1	Yes	66	ND	Podbielski et al., 1991
<i>blaSHV-3</i>	KX092356	7.0	France	1986	<i>K. pneumoniae</i>	pUD18	Yes	180	ND	Nicolas et al., 1989
<i>blaSHV-4</i>	NA	7.8	France	1987	<i>K. pneumoniae</i>	P	Yes	180	ND	Péduzzi et al., 1989; Arlet et al., 1990
<i>blaSHV-5</i>	X55640	8.2	Chile	1987	<i>K. pneumoniae</i>	pAFF1	No	150	ND	Gutmann et al., 1989
<i>blaSHV-6</i>	Y11069.1	7.6	France	1991	<i>K. pneumoniae</i>	pSLH06	Yes	180	ND	Arlet et al., 1991
<i>blaSHV-7</i>	U20270	7.6	USA	1993	<i>K. pneumoniae</i>	P	Yes	10	ND	Bradford et al., 1995
<i>blaSHV-8</i>	U92041	7.6	USA	1990	<i>E. coli</i>	C	–	–	–	Rasheed et al., 1997
<i>blaSHV-9</i>	S82452.1	8.2	Greece	1995	<i>E. coli</i> ; <i>K. pneumoniae</i> ; <i>S. marcescens</i>	pK318-1; pE77-1; pS24-1	Yes	ND	ND	Prinarakis et al., 1996
<i>blaSHV-11</i> **	X98101	8.2	Switzerland	1993–1995	<i>K. pneumoniae</i>	P	Yes	80	ND	Nüesch-Inderbinen et al., 1997
<i>blaSHV-12</i>	JX268741	8.2	Switzerland	1993–1995	<i>E. coli</i> ; <i>K. pneumoniae</i>	P	Yes	80	ND	Nüesch-Inderbinen et al., 1997
<i>blaSHV-13</i>	AF164577	7.6	Netherlands	1994	<i>K. pneumoniae</i>	P	Yes	170	ND	Yuan et al., 2000
<i>blaSHV-15</i>	AJ011428.2	ND	India	1998	<i>E. coli</i>	ND	ND	ND	ND	http://www.liahey.org/studies/
<i>blaSHV-16</i>	AF072684.2	7.6	France	1996	<i>K. pneumoniae</i>	P	Yes	>100	–	Arpin et al., 2001
<i>blaSHV-18</i>	AF132290	7.8	USA	1994	<i>K. pneumoniae</i>	P	Yes	80	ND	Rasheed et al., 2000
<i>blaSHV-23</i>	AF117747	ND	South Africa	1990	<i>K. pneumoniae</i>	ND	ND	ND	ND	Essack et al., 2004
<i>blaSHV-24</i>	AB023477	7.5	Japan	1996	<i>E. coli</i>	pCAZFR001	Yes	150	ND	Kurokawa et al., 2000
<i>blaSHV-27</i>	AF293345.1	8.2	Brazil	1999	<i>K. pneumoniae</i>	C	–	–	ND	Corkill et al., 2001
<i>blaSHV-30</i>	AY661885	6.7	USA	2003	<i>E. cloacae</i>	P	ND	9.4	AmpC, <i>blaTEM-1</i> and <i>blaSHV-7</i>	Szabó et al., 2005
<i>blaSHV-31</i>	AY277255	7.8	Netherlands	2001	<i>K. pneumoniae</i>	C	–	–	–	Mazzariol et al., 2007
<i>blaSHV-34</i>	AY036620	ND	USA	1998–2000	<i>C. koseri</i> ; <i>E. coli</i> ; <i>K. pneumoniae</i>	pOZ185	Yes	>100	ND	Heritage et al., 2003
<i>blaSHV-38</i>	AY079099	7.6	France	2001	<i>K. pneumoniae</i>	C	–	–	–	Poirel et al., 2003
<i>blaSHV-40</i>	AF535128	7.6	Canada	1999–2000	<i>K. pneumoniae</i>	ND	ND	ND	ND	Mulvey et al., 2004
<i>blaSHV-41</i>	AF535129	7.6	Canada	1999–2000	<i>K. pneumoniae</i>	ND	ND	ND	ND	Mulvey et al., 2004
<i>blaSHV-42</i>	AF535130	7.6	Canada	1999–2000	<i>K. pneumoniae</i>	ND	ND	ND	ND	Mulvey et al., 2004
<i>blaSHV-45</i>	AF547625	8.2	Brazil	NA	<i>K. pneumoniae</i>	IncA/C	ND	97–145	<i>blaCTX-M-2</i> and <i>blaSHV-27</i>	Dropa et al., 2015

(Continued)

TABLE 1 | Continued

Gene [§]	Accession Number	pI	Isolation		Bacterial Species	Genetic background			References	
			Location	Year*		Genetic Location [‡]	Conjugative plasmid	Plasmid (Kb)		Other Ab genes
<i>blaSHV-46</i>	AY210887	8.2	New York	1998	<i>K. oxytoca</i>	P	Yes	70	<i>blaTEM-1</i> ; <i>blaOXY-2</i> ; <i>blaKPC-2</i> ; <i>blaOXA (?)</i>	Yigit et al., 2003
<i>blaSHV-55</i>	DQ054528	ND	Portugal	NA	<i>K. pneumoniae</i>	ND	No	-	TEM1	Mendonça et al., 2006
<i>blaSHV-57</i>	AY223863	8.3	Taiwan	1998	<i>E. coli</i>	pMTY512	Yes	40-60	ND	Ma et al., 2005
<i>blaSHV-64</i>	DQ174304	ND	China	2000-2002	<i>K. pneumoniae</i>	ND	ND	ND	ND	Zuo et al., 2006
<i>blaSHV-66</i>	DQ174306	ND	China	2000-2002	<i>K. pneumoniae</i>	ND	ND	ND	ND	Zuo et al., 2006
<i>blaSHV-70</i>	DQ013287	7.6	China	2003-2004	<i>E. cloacae</i>	pEC04	Yes	ND	ND	Ling et al., 2006
<i>blaSHV-86</i>	DQ328802	8.2	Colombia	2003	<i>K. pneumoniae</i>	P	Yes	ND	ND	Espinal et al., 2010
<i>blaSHV-90</i>	NA	8.2	Portugal	2003	<i>K. pneumoniae</i>	ND	ND	ND	ND	Machado et al., 2007
<i>blaSHV-91</i>	NA	7.6	Portugal	2003	<i>K. pneumoniae</i>	ND	ND	ND	ND	Machado et al., 2007
<i>blaSHV-98</i>	AM941844	7.6	Algeria	2005	<i>K. pneumoniae</i>	ND	ND	ND	ND	Ramdani-Bougouessa et al., 2011
<i>blaSHV-99</i>	AM941845	7.8	Algeria	2005	<i>K. pneumoniae</i>	ND	ND	ND	ND	Ramdani-Bougouessa et al., 2011
<i>blaSHV-100</i>	AM941846	7.2	Algeria	2005	<i>K. pneumoniae</i>	ND	ND	ND	ND	Ramdani-Bougouessa et al., 2011
<i>blaSHV-102</i>	EU024485	ND	Spain	2003-2004	<i>K. pneumoniae</i>	ND	ND	ND	ND	Vinué et al., 2008
<i>blaSHV-104</i>	EU274581	7.3/8.6	Tunisia	2004	<i>E. coli</i>	pML2011	Yes	50	ND	Ben Achour et al., 2014
<i>blaSHV-105</i>	FJ194944	ND	USA	NA	<i>K. pneumoniae</i>	ND	ND	ND	<i>blaSHV-1</i> ; <i>blaSHV-5</i>	Jones et al., 2009
<i>blaSHV-106</i>	AM941847	7.6	Portugal	1999	<i>K. pneumoniae</i>	ND	ND	ND	<i>blaTEM-1</i> ; <i>blaCTX-M-32</i>	Mendonça et al., 2009
<i>blaSHV-128</i>	GU932590	8.6	Tunisia	2009	<i>E. cloacae</i>	IncFII (IS26)	Yes	100	ND	Bourouis et al., 2015
<i>blaSHV-129</i>	GU827715	ND	Italy	2008	<i>E. coli</i>	pEc6-66	ND	ND	ND	Lascols et al., 2012
<i>blaSHV-134</i>	HM559945	ND	Spain	2009	<i>K. pneumoniae</i>	IncFIIA (IS26)	Yes	75	<i>blaIM-1</i> ; <i>aac(6)-Ib</i> ; <i>dhfrII</i> ; <i>aadA1</i> ; <i>catB2</i> ;	Sánchez-Romero et al., 2012
<i>blaSHV-183</i>	HG934764	ND	NA	NA	<i>E. cloacae</i>	ND	ND	ND	<i>blaTEM-1</i> ; <i>aac(3)-IIa</i>	http://www.lahey.org/studies/

[§]Gene *blaSHV-115* was not included in the table because no information is available (<http://www.lahey.org/studies/>).
^{*}Isolation or first description.

^{**}Non ESBL genes *blaSHV-1* and *blaSHV-11* are provided as reference.

[‡]P, plasmid; C, Chromosome; when known plasmid name or Inc/rep group, and Insertion Sequences are indicated. NA, not available; ND, not determined.




FIGURE 1 | Maximum likelihood amino acid tree of 149 SHV-type β -lactamases. Variants whose sequence has not been released in GenBank as of July 2016, that show partial sequence or are identical to others (<http://www.lahey.org/studies/>) were not included in the analysis. SHV-180 and SHV-181 share the same sequence as well as SHV-121 and SHV-136. The tree was implemented in Mega version 6.06 (Tamura et al., 2013). Solid circles represent: red, extended-spectrum β -lactamases (2be; $n = 46$); green, broad-spectrum β -lactamases (2br, $n = 5$); and blue, penicillinases (2b, $n = 30$). Unclassified alleles are reported in black ($n = 68$).

The last two decades witnessed the appearance of several new variants (*bla*_{SHV-7}, *bla*_{SHV-8}, *bla*_{SHV-9}, *bla*_{SHV-31}, *bla*_{SHV-38}, *bla*_{SHV-40}, *bla*_{SHV-41}, and *bla*_{SHV-42}) whose dissemination was restricted to limited cases (Supplementary Table S1). A few

variants seem to be geographically constrained: (i) *bla*_{SHV-106}, only described in Portuguese isolates of *K. pneumoniae* together with *bla*_{TEM-1}, and/or *bla*_{CTX-M-32} (Mendonça et al., 2009); (ii) *bla*_{SHV-55}, in Portugal (Mendonça et al., 2006; Machado

et al., 2007) and recently in Brazil (Dropa et al., 2015); and (iii) *bla*_{SHV-57}, in *E. coli* isolates from Taiwan and China (Ma et al., 2005; Tian et al., 2012). A variant worth to mention is *bla*_{SHV-27} (Corkill et al., 2001), that has been detected on different plasmids in *E. coli*, *K. pneumoniae* and *Enterobacter cloacae*, associated with a vast array of antibiotic resistance genes (*bla*_{DHA-1}, *bla*_{TEM-1}, *bla*_{TEM-1b}, *bla*_{CMY-2}, *bla*_{IMP}, *bla*_{CTX-M-14}, *bla*_{CTX-M-15}, *bla*_{SHV-12}, *bla*_{SHV-45}, *bla*_{OXA-1}, *dfrA5*, *ereA2*; Muratani et al., 2006; Abbassi et al., 2008; Kiratisin et al., 2008; Duval et al., 2009; Hammami et al., 2011).

Most of SHV ESBLs (25 out of 46) are unique cases, with only one report so far. Seventeen variants are exclusively found in clinical *K. pneumoniae*: *bla*_{SHV-6}, *bla*_{SHV-13}, *bla*_{SHV-16}, *bla*_{SHV-18}, *bla*_{SHV-23}, *bla*_{SHV-45}, *bla*_{SHV-64}, *bla*_{SHV-66}, *bla*_{SHV-86}, *bla*_{SHV-90}, *bla*_{SHV-91}, *bla*_{SHV-98}, *bla*_{SHV-99}, *bla*_{SHV-100}, *bla*_{SHV-104}, *bla*_{SHV-105}, and *bla*_{SHV-134}. These variants have been described worldwide (Brazil, Portugal, Algeria, USA, Tunisia, Netherlands, France, South Africa, Colombia, and China) and are mostly associated to plasmids (Table 1). Some of these variants are sporadically accompanied by other antibiotic resistance genes like in the case of: (i) *bla*_{SHV-45} encoded by an IncA/C plasmid together with *bla*_{CTX-M-2} and *bla*_{SHV-27} (Dropa et al., 2015); (ii) *bla*_{SHV-134} encoded by an IncFII plasmid accompanied by a second plasmid carrying *bla*_{VIM-1} (Sánchez-Romero et al., 2012); (iii) and *bla*_{SHV-105}, conferring reduced susceptibility to ceftazidime, ceftriaxone, and aztreonam together with *bla*_{SHV-1}, and *bla*_{SHV-5} (Jones et al., 2009). One of the oldest variants, *bla*_{SHV-6}, was only described in France in 1991 in a *K. pneumoniae* clinical case (Arlet et al., 1991). It might be speculated that the 180 kb plasmid encoding *bla*_{SHV-6} and conferring decreased susceptibility to ceftazidime and aztreonam was not stable or it reduced bacterial strain fitness preventing a successful dissemination.

Four variants have been described only in clinical *E. coli*: (i) *bla*_{SHV-15}, described together with *bla*_{CMY-2} in a strain imported from India into the United Kingdom (<http://www.lahey.org/studies/>); (ii) *bla*_{SHV-24}, identified in Japan on a transferable 150 Kb plasmid conferring high-level resistance to ceftazidime but not cefotaxime and ceftazidime (Kurokawa et al., 2000); emergence of SHV-24 might have been driven by the extensive use of ceftazidime in Japan, enabling bacterial survival in high concentrations of this drug; (iii) *bla*_{SHV-102}, recovered in a Spanish hospital and hydrolyzing cefotaxime and ceftazidime (Vinué et al., 2008); (iv) and *bla*_{SHV-129}, detected in an abscess specimen from a patient hospitalized in Italy in 2008 (Lascols et al., 2012).

*bla*_{SHV-46} was only described on a 70 Kb conjugative plasmid also carrying *bla*_{TEM-1} and *bla*_{KPC-2} in a carbapenem-resistant strain of *Klebsiella oxytoca* from the urine of a hospitalized patient in New York (USA) in 1998 (Yigit et al., 2003). Finally, *bla*_{SHV-34} is an interesting example of extended-spectrum β -lactamase encoded by an epidemic plasmid circulating among *Citrobacter koseri*, *E. coli*, and *K. pneumoniae* in the same US hospital between 1998 and 2000 (Heritage et al., 2003).

Majority of SHV ESBLs have been detected in *K. pneumoniae* or *E. coli* (Table 1). *bla*_{SHV-30} was the first variant to be detected

in an *E. cloacae* isolate from a blood culture from a solid-organ transplant recipient in the USA in 2003 (Szabó et al., 2005). The gene, previously described in *K. pneumoniae* and *Salmonella* (Mulvey et al., 2004; Whichard et al., 2007), was located on a 9.4 Kb plasmid and contributed together with chromosomal *ampC*, *bla*_{SHV-7}, and *bla*_{TEM-1} to the antibiotic resistance profile of the *E. cloacae* isolate, the first of its kind producing two different SHV enzymes. Three other novel ESBL variants have been solely identified as plasmid-encoded in clinical *E. cloacae*: (i) *bla*_{SHV-70}, from a Chinese patient with history of ceftazidime treatment (Ling et al., 2006) and observed in other clinical Chinese settings (Liu et al., 2008); (ii) *bla*_{SHV-128}, isolated in Tunisia in 2009, located on an IncFII conjugative plasmid, and conferring resistance to all β -lactams except imipenem (Bourouis et al., 2015); (iii) and *bla*_{SHV-183}, for which additional description is not available (<http://www.lahey.org/studies/>).

SHV EXTENDED-SPECTRUM β -LACTAMASES: CATALYTIC PROPERTIES AND RESISTANCE PHENOTYPE

Extended-spectrum SHV β -lactamases belong to functional group 2be, while very recently they were assigned to subclass A1 of serine β -lactamases, clustering with TEM and CTX-M enzymes among other clinically relevant β -lactamases (Bush, 2013; Philippon et al., 2016). SHV ESBLs consist of two subdomains: an α/β that includes an antiparallel five-stranded β -sheet flanked by α -helices, and an all- α -helical subdomain (Matagne et al., 1998). Similar to TEM β -lactamases (Jelsch et al., 1993), the active site is located within the cleft created by the subdomains and it contains the Ser⁷⁰ residue that mediates the nucleophilic attack on the carbonyl group of the β -lactam ring. In the vicinity of this serine residue, several conserved structural and functional amino acid motifs have been identified. These include the Ser⁷⁰XXLYs ("SXXK" motif, with X representing variable amino acids), the Ser¹³⁰AspAsn ("SDN" motif), the Glu¹⁶⁶XXLYsAsn ("EXXLN" motif), and the Lys²³⁴Thr/SerGly ("KTG" motif) (Bush, 2013).

Each SHV ESBL has one (SHV-2, SHV-6, SHV-8, SHV-24, SHV-27, SHV-38, SHV-41, SHV-57, SHV-98, SHV-99, SHV-102, and SHV-104) to six (SHV-128) amino acid substitutions when compared to SHV-1 (Table 2), indicating that even a single amino acid substitution is enough to convey an extended-spectrum phenotype. Therefore, we can speculate that other SHV ESBLs may still evolve from a parental SHV β -lactamase due to single spectrum-extending substitutions, although the majority of them have possibly emerged through a stepwise acquisition of several mutations (substitutions, deletions and/or insertions) from pre-existing extended-spectrum SHV variants.

Among SHV ESBLs, amino acid substitutions are predominantly located at positions Leu³⁵, Gly²³⁸, and Glu²⁴⁰, while other less frequent but critical substitutions for the extended-spectrum phenotype occur on several amino acids including Ile⁸, Arg⁴³, Glu⁶⁴, Gly¹⁵⁶, Asp¹⁷⁹, and Arg²⁰⁵ (Table 2). Although, most of these residues are not involved directly in β -lactams hydrolysis, they result in the enhancement or relaxation

TABLE 2 | Amino acid polymorphisms in SHV-type extended-spectrum β -lactamases.

	7	8	10	14	18	20	25	35	43	54	61	64	75	80	89	96	97	104	122	123	129	140	142	146	148	154	156	163	169	179	186	187	188	192	193	195	202	205	238	240	243	271	274	275	276	282	286				
	Amino acid position																																																		
SHV-1	Y	I	L	S	T	P	A	L	R	G	R	E	V	V	E	H	Y	D	L	C	M	A	V	A	L	Q	G	D	L	D	M	A	A	K	L	T	R	R	S	E	A	S	E	R	N	I	L				
SHV-2		
SHV-2A	Q	
SHV-3
SHV-4
SHV-5
SHV-6
SHV-7	.	F	S	
SHV-8
SHV-9	Del.	R	
SHV-11**	Q
SHV-12	Q
SHV-13	Q
SHV-15	Q	.	M	K
SHV-16	T	H
SHV-18	.	F	S	
SHV-23*	F
SHV-24
SHV-27
SHV-30	.	F	S
SHV-31	Q
SHV-34	.	F	S	.	G
SHV-38
SHV-40	Q
SHV-41	F	
SHV-42	S	V	
SHV-45
SHV-46	
SHV-55	F	
SHV-57
SHV-64	Q	L	
SHV-66	Q	Q	
SHV-70	Q	
SHV-86	F	Q		
SHV-90	

(Continued)

TABLE 3 | Kinetic parameters of available SHV-type extended-spectrum β -lactamases.

Enzyme	Parameter	PEN	AMP	AMX	TIC	PIP	CER	CEF	CAZ	CTX	FEP	ATM	CLA	SUL	TZB
SHV-1**	K_{cat}	455		900	60	570	170	10	NH	NH	>100	NH			
	K_m	20		90	22	60	110	26	ND	ND	>3000	ND			
	K_{cat}/K_m	23,000		10,000	2700	10,000	1500	400	ND	ND	>35	ND			
	V_{max}/K_m	100					4	1					0.19	1.70	0.057
	K_i												0.057	7.50	0.150
SHV-2	K_i												0.16	0.36	0.04
	IC ₅₀												0.020	0.57	0.049
	V_{max}	100							6.5	70		1			
	K_m	3.5	12			ND		ND	24	18	NA	10			
	K_{cat}		206			ND		ND		11	NA				
SHV-2a	K_{cat}/K_m		17			ND		ND		0.6	0.008				
	K_i		13					5	72	4		3	0.08	0.47	0.027
SHV-4	IC ₅₀		100					53	1	10			0.018	0.68	0.038
	V_{max}	100							52	115		5			
SHV-5	K_m	3.5							60	25		0.5			
	K_i												0.10	0.18	0.036
SHV-5	IC ₅₀												0.005	0.40	0.022
	K_m	15	11					3	23	7		0.02			
	V_{max}/K_m	100	100					51	4	7					
	V_{max}								24	11		13			
SHV-7	K_m							2.7	13	30		3.3			
	V_{max}							35	13						
SHV-9	V_{max}	100	215					58	10	24					
	K_m	18	12					5	18	9					
	IC ₅₀												0.14	0.43	
SHV-13	K_m	10	28			18			91	11		77			
	V_{max}	100	178			136			0.38	12		0.66			
	V_{max}/K_m	100	64			76			0.42	11		0.86			
	V_{max}														
SHV-18	V_{max}	100					200		13.5	26.9		<1			
	V_{max}/K_m	100					53		1.5	24		ND			

(Continued)

TABLE 3 | Continued

Enzyme	Parameter	PEN	AMP	AMX	TIC	PIP	CER	CEF	CAZ	CTX	FEP	ATM	CLA	SUL	TZB
SHV-24	V_{max}		2				2.37		0.043			0.735			
	K_m		32				210		30			500			
	V_{max}/K_m		0.0625				0.0113		0.000143			0.00147			
	K_i		57				ND		37			ND			
SHV-38 [§]	K_{cat}	100		1800	10	100	40	5	110	1	3	3			
	K_m	13		35	14	80	150	100	3800	800	1600	5500			
	K_{cat}/K_m	7700		51,000	700	1300	270	50	30	1	2	0.5			
	K_m	5 \pm 0.51		10 \pm 0.14	6 \pm 0.02	8 \pm 0.37	9 \pm 0.68	9 \pm 0.68	58 \pm 7.40	21 \pm 0.13	149 \pm 2.61	5 \pm 0.62			
SHV-55*	K_{cat}	23 \pm 0.76		23 \pm 0.17	8 \pm 0.00	27 \pm 1.53	38 \pm 3.94	38 \pm 3.94	9 \pm 0.21	24 \pm 0.34	30 \pm 3.10	<0.1			
	K_{cat}/K_m	5.3 \pm 0.42		2.5 \pm 0.002	1.5 \pm 0.00	3.7 \pm 0.03	4.4 \pm 0.78	4.4 \pm 0.78	0.2 \pm 0.02	1.1 \pm 0.01	0.2 \pm 0.02	ND			
	IC_{50}												0.02		
	K_m	67							30.9						
SHV-57	K_{cat}	3.8 $\times 10^{-3}$							8.6 $\times 10^{-4}$						
	K_{cat}/K_m	5.67 $\times 10^{-5}$							2.78 $\times 10^{-5}$						
	K_i												27 $\times 10^3$		1.16 $\times 10^3$
	K_m	12 \pm 0.11		11 \pm 0.26	5 \pm 0.93	13 \pm 1.43	102 \pm 11.38	102 \pm 11.38	136 \pm 4.09	183 \pm 0.72		196 \pm 0.60			
SHV-99*	K_{cat}	778 \pm 616		563 \pm 8	58 \pm 2	563 \pm 13	37 \pm 2	37 \pm 2	<0.1	<0.1		0.5 \pm 0.001			
	K_{cat}/K_m	62.3 \pm 4.4		49.6 \pm 1.8	13 \pm 2.4	43.5 \pm 6.5	0.37 \pm 0.04	0.37 \pm 0.04	<0.001	<0.001		0.003			
	IC_{50}												0.02		0.03
	K_{cat}	55		80			30	30							
SHV-104	K_m	94		10			68	68							
	K_{cat}/K_m	0.6		8			0.44	0.44							
	K_{cat}	22.8 \pm 11		1688 \pm 4			26 \pm 1	26 \pm 1	3.1 \pm 1.5	4.8 \pm 3.4	4.5 \pm 0.5				
	K_m	46.8 \pm 24		25 \pm 9			12.1 \pm 3.7	12.1 \pm 3.7	24 \pm 3	26.7 \pm 5.5	52 \pm 3.5				
SHV-129 [#] *	K_{cat}/K_m	0.5 \pm 0.7		7 \pm 0.4			2.2 \pm 0.3	2.2 \pm 0.3	0.13 \pm 0.5	0.2 \pm 0.5	0.09 \pm 0.01				
	K_i												0.4	0.4	0.04

Antibiotic: Penicillin (PEN); Ampicillin (AMP); Amoxicillin (AMX); Ticarcillin (TIC); Piperacillin (PIP); Cephaloridine (CER); Cephalothin (CEF); Cefotaxime (CTX); Cefepime (FEP); Aztreonam (ATM); Clavulanic Acid (CLA); Sulbactam (SUL); Tazobactam (TZB).
 Parameters are expressed as μM (K_i , IC_{50} , K_m), s^{-1} (K_{cat}), and $\mu M/min$ (K_{cat}/K_m , V_{max}). Only antibiotics for which 3 or more SHV enzyme values were available are reported.
 NA, not able to determine the rate of hydrolysis and affinity; NH, not hydrolyzed; ND, not determined.
 **Non ESBL SHV-1 is provided as reference.
 # K_{cat}/K_m values are expressed as mM/s.
 # K_{cat}/K_m values are expressed as $\mu M/s$.
 *Values (Except IC_{50}) represent mean \pm standard deviation.
 References: SHV-1 (Gutmann et al., 1989; Poinel et al., 2003); SHV-2 (Gutmann et al., 1989; Bradford et al., 1995; Winkler and Bonomo, 2016); SHV-2a (Podbielski et al., 1991); SHV-4 (Péduzzi et al., 1989); SHV-5 (Gutmann et al., 1989); SHV-7 (Bradford et al., 1995); SHV-9 (Pinarakis et al., 1995); SHV-13 (Yuan et al., 2000); SHV-18 (Rasheed et al., 2000); SHV-24 (Kurokawa et al., 2000); SHV-38 (Poinel et al., 2000); SHV-55 (Mendonça et al., 2006); SHV-57 (Ila et al., 2005); SHV-99 (Ramdani-Bouguessa et al., 2011); SHV-104 (Ben Achour et al., 2014); SHV-129 (Winkler and Bonomo, 2016).

other amino acid substitutions (Table 2) have been described as either responsible for or possibly contributing to the ESBL phenotype, the detailed description of which exceeds the scope of this review.

Apart from point mutations leading to amino acid substitutions, frame shift mutations have been observed with very low occurrence among SHV ESBLs resulting in amino acid insertions (Arpin et al., 2001; Ramdani-Bouguessa et al., 2011) or deletions (Prinarakis et al., 1996). However, their role in the rising of the extended-spectrum phenotype remains unclear. SHV ESBL variants falling in this category are: (i) SHV-9, with the deletion of Gly⁵⁴ (Prinarakis et al., 1996); (ii) SHV-16, with a 5-amino acid sequence duplication (Asp^{163a}ArgGluTrpGluThr-Asp^{163b}ArgGluTrpGluThr) of the amino acids between 163 and 167, including Glu¹⁶⁶ in the Ω loop (Arpin et al., 2001); (iii) SHV-100, with a 13-amino acid insertion (SerGluSerGlnLeuSerGlyArgValGlyMetIleGlu) between amino acids 35 and 36 (Ramdani-Bouguessa et al., 2011); and (iv) SHV-183, with an Ala insertion between amino acids 186 and 187 (<http://www.lahey.org/studies>). Of note, the duplication observed in SHV-16 was shown to increase the conformational flexibility of the catalytic region facilitating the access of bulkier cephalosporins, such as ceftazidime, but resulted in enzymatic instability (Arpin et al., 2001). This finding could explain the low incidence of frame shift mutations among extended-spectrum SHV β -lactamases, due to a deleterious effect on the enzymes.

Overall, the available SHV ESBL kinetic parameters show that most of the substitutions lead to more efficient hydrolysis of oxyimino- β -lactams than penicillins, as depicted by the low K_{cat} values for penicillins (Table 3). While they retain their ability to hydrolyze penicillins, they are not catalytically so efficient compared to SHV-1 (Bush and Singer, 1989) and this is due to the decreased strength of the crucial hydrogen-bonding network needed for penicillin catalysis (turnover). As a consequence, since β -lactam inhibitors (clavulanic acid, sulbactam, and tazobactam) are structurally very similar to penicillin substrates, SHV ESBLs also exhibit increased susceptibility to β -lactam inhibitors compared to SHV-1 (Table 3) leading to less inhibitor required for inactivation (lower K_i and IC_{50} ; Tzouveleakis and Bonomo, 1999).

DETECTION

There are at least 46 known SHV-ESBL genes together with more than 150 non-ESBL or unclassified alleles to date (<http://www.lahey.org/studies/>). Accurate identification of these variants is essential for surveillance and for epidemiological studies of transmission mode, particularly in clinical setting, where appropriate antimicrobial therapy is critical.

A panel of different phenotypic confirmatory tests is available to determine the presence of extended-spectrum β -lactamases, including SHV-variants: minimum inhibitory concentration (MIC) determination of β -lactam with and without clavulanic acid, double disk synergy test (DDST), inhibitor potentiated

disk diffusion test (IPDDT), three-dimensional test (TDT) and commercially available methods (Etest for ESBLs, Vitek ESBL cards, MicroScan panels, and BD Phoenix Automated Microbiology System (Bradford, 2001; Paterson and Bonomo, 2005)). Standard microbiological procedures can take up to several days for culture, isolation and characterization and many comparative studies have shown that PCR-based methods have higher sensitivity (Bedenic et al., 2001, 2007; Singh et al., 2012), mostly due to variable levels of gene expression. Therefore, PCR and nucleotide sequence analysis (Stürenburg et al., 2003), together with various PCR-based methods, remain the gold standard for extended-spectrum β -lactamase SHV-variants identification.

Chanawong and colleagues developed a PCR-restriction fragment length polymorphism (PCR-RFLP) method to allow the identification of new SHV β -lactamases variants through detection of known mutations that alter recognition sites of restriction endonucleases (Chanawong et al., 2000). PCR-RFLP complements pre-existing PCR-single strand conformational polymorphism (PCR-SSCP) limited by partial gene amplification, thus missing potential mutation sites (M'Zali et al., 1998). PCR-RFLP can also be used in combination with restriction site insertion-PCR (RSI-PCR), a method based on primers mismatches, allowing the unambiguous identification of up to 27 SHV variants by point mutation (Chanawong et al., 2001a). Fluorescently labeled hybridization probes followed by melting curve analysis can also be used to discriminate between ESBL and non-ESBL *bla*_{SHV} genes (Randegger and Hächler, 2001). This method, termed the SHV melting curve mutation detection method, is also able to categorize SHV ESBL producers into phenotypically relevant subgroups: (i) weak ceftazidime resistance (SHV-6 and SHV-8); (ii) significant resistance to cefotaxime and ceftriaxone and moderate resistance to ceftazidime (SHV-2, SHV-2a, and SHV-3); and (iii), most effective against all expanded-spectrum cephalosporins (SHV-4, SHV-5, SHV-9, and SHV-12). Combined systems can also be developed *ad hoc* to rapidly screen local epidemiological settings (Chia et al., 2005). A modified SHV melting-curve mutation detection method able to distinguish between prevalent Taiwanese *bla*_{SHV} genes (SHV-1, SHV-2, SHV-2a, SHV-5, SHV-11, and SHV-12) was combined with a multiplex PCR to identify different β -lactamases genes (*bla*_{SHV}, *bla*_{CTX-M-3}-like, and *bla*_{CTX-M-14}). The design of this method can be easily adapted to other geographic areas where different ESBLs are prevalent. Multiplex real-time PCR assays for the fast detection of extended-spectrum β -lactamase and carbapenemase genes were developed with differential melting curves able to recognize up to 120 different SHV allelic variants (Singh et al., 2016).

New techniques for ESBL detection are employed alongside PCR-based methods these days. Loop-mediated isothermal amplification (LAMP) was applied to detect SHV- and other ESBL-producing bacteria in meat and proved to be more specific and sensitive than MacConkey agar or cefpodoxime disc methods (Anjum et al., 2013). Commercial DNA microarrays are also proving themselves to be accurate, with sensitivity and specificity values for ESBL detection being high. Up to

53 SHV-variants can be covered on a same array (Leinberger et al., 2010), but on the other hand some alleles may fail to be detected (i.e., SHV-12), as previously reported (Stuart et al., 2012). Because arrays have major limitations to detect novel genes or variants, PCR and sequencing remains essential. Matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry (MS) is routinely used for bacteria identification and has been recently applied to detect ESBL-producing Enterobacteriaceae from positive blood cultures in clinical practice (Jung et al., 2014; Oviaño et al., 2014). Although, this methodology has yet to be fully validated, preliminary results show 99% sensitivity and 100% specificity, and denote a novel approach to categorize bacteria as ESBL producers.

Pyrosequencing combines standard PCR and sequencing by synthesis to rapidly determine the sequence of a target DNA region; it has been extensively used for the detection of bacterial resistance genes and bacterial community composition (Tang et al., 2016; Tian et al., 2016). This technique has been used to perform mutation analysis of *bla*_{SHV} to resolve heterogeneous sequences in clinical isolates of *K. pneumoniae* containing more than one SHV variant (Haanperä et al., 2008). An alternative protocol for pyrosequencing is the single-nucleotide polymorphism (SNP), ideal for the sequencing of mixed templates and determination of SNPs at the position of interest. This protocol has been applied to discriminate between eight *bla*_{SHV} variants from clinical isolates of *E. coli* and *K. pneumoniae*, reporting great reproducibility and ability to discriminate between sequences (Jones et al., 2009). A multiplex pyrosequencing assay coupled with qPCR amplification has also been recently developed to enable rapid and accurate detection of *bla*_{SHV} and *bla*_{TEM}-producing Enterobacteriaceae (Deccache et al., 2015). Overall, pyrosequencing can be a useful epidemiological tool for the exact identification of *bla*_{SHV} as a prerequisite for analyzing the spread of certain SHV variants.

Finally, the advent of whole genome sequencing (WGS) has taken differentiation of bacterial strains and identification of the associated antibiotic resistance gene cargo to another level. Aside from the phylogenetic analysis that WGS provides, the complete resistome of a strain can be unraveled as well as its mobilome, i.e., the mobile genetic elements that are associated with antibiotic resistance diffusion. Only this information can provide us with full understanding of complex genomic structures as observed, for example, in clinical *K. pneumoniae* genomes carrying (i) nineteen antibiotic resistance genes including *bla*_{OXA-1} and *bla*_{SHV-28} in the chromosome, *bla*_{NDM-1} in a plasmid, and *bla*_{OXA-232} in a second plasmid (Kwon et al., 2016); (ii) β -lactamase genes *bla*_{KPC-2}, *bla*_{SHV-11}, *bla*_{TEM-169}, and *bla*_{OXA-9}, together with *aac*(6')-Ib, *aadA2*, and *aph*(3')-Ia as aminoglycoside resistance encoding genes, *mph*(A) for macrolides, *oqx*A and *oqx*B for quinolone, *cat*A1 for phenicol, *sul*1 for sulfonamide, and *dfr*A12 for trimethoprim (Lee et al., 2014); or (iii) six different plasmids, adding up to 0.43 Mbp, coding for six β -lactamases (*bla*_{SHV-12}, *bla*_{OXA-9}, *bla*_{TEM-1}, *bla*_{CTX-M-2}, and *bla*_{KPC-2}), together with *bla*_{SHV-110} and adhesin-related gene clusters on the chromosome (Perreira Ramos et al., 2014).

EXPANSION TOWARD NEW ECOLOGICAL NICHES

Over the last years the presence of antibiotics as well as antibiotic resistant bacteria has been shown outside the clinical environment, including water, soil and, most notably, food producing animals. When looking at SHV-variants distribution it is evident that in recent years, as for most extended-spectrum β -lactamases (Canton et al., 2012), their presence has been confirmed in virtually all ecological niches (Supplementary Table S1), making it more challenging to restrain antibiotic resistance diffusion. The most representative cases and variants will be discussed.

Aquatic Environment

In an effort to control the release of antibiotics and antibiotic resistant bacteria in the environment, aquatic environments are being investigated worldwide, whether they be natural, drinking or wastewaters. The latter are particularly worrisome given the high prevalence of *bla*_{SHV} alleles, as observed in untreated hospital wastewater in Australia (Gündogdu et al., 2013), their possible association with determinants of quinolone and other β -lactamase resistance (Calhau et al., 2015; Osinska et al., 2016), and their relatively easy transmission to surface water through waste water treatment plant discharges (Marti et al., 2013). Studies showed that SHV types, together with CTX-M and OXA genes can be significantly decreased by biological treatments such as activated sludge processing and anaerobic digestion, although not all can be effectively eliminated (Yi et al., 2015).

Urban waters are also exposed to relatively high population densities and therefore are often unprotected from biological contaminants, with people playing a crucial role in antibiotic resistance dissemination in the environment. Unusual finding of SHV-producing *Stenotrophomonas maltophilia* in a swimming recreational Serbian lake and its transient presence during summer months can be considered as a proof of its anthropogenic origin, given its nature of emerging nosocomial pathogen (Novovic et al., 2015). Similar conclusions can be drawn for SHV-producing *K. pneumoniae* and *E. cloacae* isolated from a Bangladeshi lake, which receives waste water from surrounding residents, commercial buildings and clinics in Dhaka city (Haque et al., 2014), as well as for artificial water reservoirs in Poland (Wolny-Koladka and Lenart-Boron, 2016), or urban surface waters in Malaysia (Tissera and Lee, 2013). In recent surveillance studies of different rivers and lakes in Switzerland, *bla*_{SHV-12}-producing Enterobacteriaceae were isolated only in 4% of the cases (Zurfluh et al., 2013), although this variant is predominant in clinical Swiss isolates (Nüesch-Inderbinen et al., 1997). *bla*_{SHV-12} was also detected in Enterobacteriaceae from seawater, together with *tet*(A) and *sul*2 in Portugal (Alves et al., 2014), and plasmid-encoded together with *bla*_{TEM-1} and/or *bla*_{CTX-M-1} in Croatia (Maravic et al., 2015). Finally, data on ESBL-producing Enterobacteriaceae isolated from drinking water is also increasing, reporting SHV alleles in rural water reservoirs in China (Zhang et al., 2015), or drinking water sources for First Nation communities in Canada (Fernando et al., 2016).

Food Producing Animals

Food producing animals have become subject of increasing interest after several studies demonstrated that resistant strains of animal origin can be associated to human infections, possibly through the food chain (Hasman et al., 2005). Majority of SHV variants in this reservoir belong to *bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, and *bla*_{SHV-12} (Supplementary Table S1) owing to their successful association with conjugative plasmids (see Section Plasmid epidemiology of *bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, and *bla*_{SHV-12}).

Surveillance activities in healthy animals worldwide are generating a tremendous amount of data on ESBL distribution. Most SHV β -lactamase producers are *E. coli* from swine and broiler fecal samples as observed in China (Tian et al., 2012); in Spain, with *bla*_{SHV-2} associated with *bla*_{CTX-M-9} and *bla*_{SHV-12} with *bla*_{CTX-M-1}, in pigs and broilers respectively (Blanc et al., 2006); in layers, cattle, and broilers but not in swine in Japan (Hiki et al., 2013; Kameyama et al., 2013); and in the Netherlands, where healthy broilers carried *bla*_{SHV-2} in combination with *bla*_{TEM-1} or *bla*_{TEM-135} (Dierikx et al., 2010). Other Enterobacteriaceae like *K. pneumoniae* and *Citrobacter freundii* were positive for *bla*_{SHV-2} or *bla*_{SHV-12} from poultry and swine, respectively (Machado et al., 2008).

Finding ESBL producers in food producing animals is also mirrored by positive food samples worldwide, mostly retail chicken meat, as reported in Tunisia, with *E. coli* carrying *bla*_{SHV-5} isolated from different butcheries, supermarkets, and local markets (Jouini et al., 2013), or *Salmonella enterica* carrying *bla*_{SHV-12} in Japan (Noda et al., 2015). The cross-contamination between food producing animals and retail meat has been internationally demonstrated due to the detection of plasmid-borne SHV variants, such as *bla*_{SHV-2} and *bla*_{SHV-2a} from Canadian chicken meat and abattoir chicken cecum (Pouget et al., 2013) or *bla*_{SHV-2} and *bla*_{TEM-1} in Japan (Hiroi et al., 2011), presenting the potential for horizontal transfer between Enterobacteriaceae as a high public health concern.

SHV β -lactamase producing Enterobacteriaceae have been detected also in diseased animals, as reported for septicemic broilers due to avian pathogenic *E. coli* encoding a remarkable array of antibiotic resistance genes (*dfxA17-aadA5*, *bla*_{TEM-1}, *bla*_{CTX-M-15}, *bla*_{OXA-1}, *bla*_{SHV-2}, *tet(A)*, *tet(E)*, *qnrB2*, *aac(6)-Ib-cr*) (Ahmed et al., 2013); for *K. pneumoniae* isolated from bovine mastitis in the United Kingdom (Timofte et al., 2014) and Egypt (Ahmed and Shimamoto, 2011); and for multidrug resistant *S. enterica* serotypes Enteritidis and Typhimurium isolated from diarrheic calves (Ahmed et al., 2009).

Finally, *bla*_{SHV-27} is the only other SHV variant frequently reported as chromosomally located in *K. pneumoniae* from swine, in association with *bla*_{SHV-11} and *bla*_{CTX-M-1} in China (Zou et al., 2011); in *E. coli* isolated from farmed fish together with non ESBLs *bla*_{SHV-1}, *bla*_{SHV-11}, *bla*_{SHV-25}, and *bla*_{SHV-26} (Jiang et al., 2012); and in opportunistic pathogens asymptotically colonizing healthy milk cows (Hammad and Shimamoto, 2011).

Wildlife, Companion Animals, and Vegetables

ESBL diffusion has been studied extensively in Enterobacteriaceae from humans and livestock, whereas information on antibiotic resistance in the environment is still limited. Yet, the dissemination success of *bla*_{SHV-12} is confirmed by its introduction into the wildlife, notably in birds, as reported in Spain (Alcalá et al., 2015), the Netherlands (Veldman et al., 2013), Poland (Literak et al., 2010), and the Czech Republic (Dolejská et al., 2009). This success is likely associated to predominant avian clones and to efficient plasmids (Table 4, Figure 3) of the IncN incompatibility group, described to be more frequent in pathogenic than in commensal avian and human *E. coli* strains (Johnson et al., 2007). *bla*_{SHV-5} was also detected in *E. coli* from several birds of prey in Portugal, alone or in associations with *bla*_{TEM-1b} (Pinto et al., 2010).

Emergence of Enterobacteriaceae producing β -lactamases in companion animals have been gradually reported, with CTX-M enzymes being prevalent as observed in the human scenario (Rubin and Pitout, 2014). Few studies, on both healthy and diagnostic clinical canine and feline samples, report finding other ESBL variants including *bla*_{SHV-3} in the USA (Shaheen et al., 2011), *bla*_{SHV-2} in Mexico (Rocha-Gracia et al., 2015), *bla*_{SHV-12} in Italy and Poland (Carattoli et al., 2005b; Rzewuska et al., 2015) and *bla*_{SHV-12} in association with *bla*_{OXA-48}, *bla*_{CMY-2}, *bla*_{TEM-1}, *aac(6)-Ib-cr*, and *qnrB2* in Germany (Stolle et al., 2013).

Lastly, SHV variants have been detected in imported vegetables in Switzerland together with *bla*_{SHV-12} for the first time in the opportunistic foodborne pathogen *Cronobacter sakazakii* whose potential to cause bacteremia and meningitis is an actual concern (Zurfluh et al., 2015). Similar results were observed in vegetables collected in South Korea (Kim et al., 2015), salads in the Netherlands (Reuland et al., 2014), and Spain (Egea et al., 2011), displaying a new route of introduction for ESBLs and pathogenic Enterobacteriaceae.

PLASMID EPIDEMIOLOGY OF *bla*_{SHV-2}, *bla*_{SHV-2A}, *bla*_{SHV-5}, AND *bla*_{SHV-12}

The role of plasmids in the successful spread of β -lactamase genes has been extensively described (Carattoli, 2009, 2013) and, among the SHV family, it finds its best examples in *bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, and *bla*_{SHV-12}. Combination of these alleles with different dissemination machineries has brought the enzymes to reach diverse niches worldwide (Figure 2). Plasmids belonging to seven replicon types (A/C, F, HI2, I1, L/M, N, X3) have been shown to drive the epidemiology of these four predominant SHV ESBLs, although their distribution varies on the plasmid families (Table 4). Other rep families that have been only incidentally associated with extended-spectrum SHV β -lactamases include the ColE, K, P, and R (Table 4).

IncA/C

*bla*_{SHV-12} has been identified on mostly conjugative broad-host range IncA/C plasmids in a variety of bacterial species, including

TABLE 4 | Plasmid epidemiology of SHV-type extended-spectrum β -lactamases.

Inc Group	Plasmid Size (Kb)*	<i>bla</i> _{SHV} allele [§]	Other Antibiotic Resistance Genes	Bacterial Species [#]	Country	References
IncA/C	ND (C)	SHV-12 or SHV-2a	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
	150 (C)	SHV-12	<i>bla</i> _{VIM-1} , <i>aac</i> (6')-Ib', <i>aadA1b</i> , <i>catB2</i> , <i>sul1</i> , <i>dfrA14</i>	<i>A. caviae</i> (H)	Italy	Antonelli et al., 2016
	150 (NC)	SHV-12 (IS26)	<i>bla</i> _{CTX-M-14} , <i>bla</i> _{DHA-1}	<i>P. mirabilis</i> (H)	Korea	Song et al., 2011
	ND	SHV-2, SHV-5 or SHV-12	ND	<i>E. coli</i> (H)	France	Marcadé et al., 2009
	130 (C)	SHV-5 (IS26)	<i>bla</i> _{VEB-1} , <i>bla</i> _{VIM-1} , <i>aacA7</i> , <i>dfrA1</i> , <i>aadA1</i> , <i>bla</i> _{OXA-1} , <i>bla</i> _{TEM-1} , <i>aadB</i> , <i>arr2</i> , <i>cmIA5</i>	<i>P. stuartii</i> (H)	Greece	Giakkoupi et al., 2015
	97–145 63.5–209	SHV-45 SHV-55	<i>bla</i> _{CTX-M-2} ; <i>bla</i> _{SHV-27} <i>bla</i> _{CTX-M-2} ; <i>bla</i> _{SHV-28}	<i>K. pneumoniae</i> (H) <i>K. pneumoniae</i> (H)	Brazil Brazil	Dropa et al., 2015 Dropa et al., 2015
IncA/C-IncR	220 (C)	SHV-5	<i>bla</i> _{VEB-1} , <i>bla</i> _{VIM-1} , <i>rmtB</i> , <i>aacA7</i> , <i>dfrA1</i> , <i>aadA1</i>	<i>P. stuartii</i> (H)	Greece	Oikonomou et al., 2016
IncF	125	SHV-5 (IS26)	ND	<i>E. coli</i> (H)	Poland	Zienkiewicz et al., 2013
IncFIA-FIB	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
IncFIB	ND	SHV-12	<i>sul3</i>	<i>E. coli</i> (A)	Italy	Bortolaia et al., 2010
	95–200 (C)	SHV-2	<i>aadA1</i>	<i>E. coli</i> (A)	Canada	Pouget et al., 2013
	>23	SHV-2	ND	<i>K. pneumoniae</i> (H)	China	Wang et al., 2012
	ND	SHV-2	ND	<i>E. coli</i> (H)	France	Marcadé et al., 2009
	ND (C)	SHV-5	-	<i>E. coli</i> (H)	Uruguay	
IncFIB10	ND	SHV-12 (IS26)	<i>bla</i> _{TEM-1}	<i>E. coli</i> (H)	UK	Doumith et al., 2012
IncFIC	ND	SHV-5	<i>aac</i> (6')-Ib', <i>aadA1</i>	<i>S. marcescens</i> (H)	Uruguay	García-Fulgueiras et al., 2011
IncF-N	ND	SHV-2	<i>aac</i> (6')-Ib'	<i>K. pneumoniae</i> (H)	Uruguay	García-Fulgueiras et al., 2011
IncFII	ND	SHV-2 or SHV-12	ND	<i>E. coli</i> (H)	France	Marcadé et al., 2009
	70–80 (C)	SHV-2a (IS26)	ND	<i>K. pneumoniae</i> (H)	Tunisia	Elhani et al., 2010
	100 (C)	SHV-128 (IS26)	ND	<i>E. cloacae</i> (H)	Tunisia	Bourouis et al., 2015
IncFII-FIA	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
IncFII-FIA-FIB	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
IncFII-FIB	ND	SHV-2	ND	<i>E. coli</i> (H)	France	Marcadé et al., 2009
	ND (C)	SHV-2a	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
IncFIIk1	200–220	SHV-2, SHV-55 or SHV-106	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
IncFIIk5	220	SHV-55	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
IncHI2	ND (C)	SHV-2a or SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
	95 (C)	SHV-12 (IS26)	ND	<i>K. pneumoniae</i> (H)	Tunisia	Elhani et al., 2010
	310 (C)	SHV-12	<i>qnrB2</i> , <i>bla</i> _{TEM-1} , <i>sul1</i> , <i>dfrA19</i> , <i>tet</i> (D), <i>strA</i> , <i>strB</i> , <i>aac</i> (60)-Ib	<i>S. Senftenberg</i> (H)	Netherlands	Veldman et al., 2010
	200 (NC)	SHV-12	<i>tet</i> (D)	<i>S. Concord</i> (H)	Netherlands	Veldman et al., 2010

(Continued)

TABLE 4 | Continued

Inc Group	Plasmid Size (Kb)*	<i>bla</i> _{SHV} allele [§]	Other Antibiotic Resistance Genes	Bacterial Species [#]	Country	References
	290 (C)	SHV-12	<i>qnrB2</i> , <i>bla</i> _{TEM-1} , <i>sul1</i> , <i>sul2</i> , <i>dfrA19</i> , <i>tet(D)</i> , <i>strA</i> , <i>strB</i> ,	<i>S. Concord</i> (H)	Netherlands	Veldman et al., 2010
	180, 350, 380	SHV-12	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
	400	SHV-12	ND	<i>E. cloacae</i> (H)	Portugal	Rodrigues et al., 2014
	320 (C)	SHV-12	<i>qnrB2</i> , <i>strA/B</i> , <i>tet(D)</i> , <i>clmA</i> , <i>sul1</i>	<i>S. Bredeney</i> (H)	Spain	Herrera-Leon et al., 2011
	ND (C)	SHV-12 (IS26)	<i>bla</i> _{CTX-M-14}	<i>E. cloacae</i> (H)	Taiwan	Chen C. M. et al., 2015
	ND (C)	SHV-12 (IS26)	<i>bla</i> _{CTX-M-3}	<i>E. cloacae</i> (H)	Taiwan	Chen C. M. et al., 2015
InclH2 (ST1)	300 (C)	SHV-2	ND	<i>S. Agona</i> or <i>Keurmassar</i> (H)	Senegal	Harrois et al., 2014
Incl1	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Bulgaria	Markovska et al., 2014
	ND	SHV-12	ND	<i>E. coli</i> (H)	France	Marcadé et al., 2009
	ND	SHV-12	<i>sul3</i>	<i>E. coli</i> (A)	Italy	Bortolaia et al., 2010
	19 (C)	SHV-12	–	<i>E. coli</i> (A)	Italy	Bortolaia et al., 2011
	340 (C)	SHV-12	ND	<i>S. Concord</i> (H)	Norway (Ethiopia)	Fabre et al., 2009
	95 (C)	SHV-12	–	<i>E. coli</i> (A)	Poland	Literak et al., 2010
	10 (NC)	SHV-12	–	<i>S. enteritidis</i> (H)	Spain	de Toro et al., 2013
	60 (C)	SHV-12	<i>bla</i> _{VIM-1} - <i>aacA4</i> - <i>dfrI</i> - <i>aadA1</i> - <i>catB2</i>	<i>K. pneumoniae</i> , <i>E. coli</i> (H)	Spain	Tato et al., 2007
	ND (C)	SHV-12 (IS26)	<i>bla</i> _{CTX-M-3}	<i>E. cloacae</i> (H)	Taiwan	Chen C. M. et al., 2015
	95–200 (C)	SHV-2	<i>aadA1</i>	<i>E. coli</i> , <i>S. Heidelberg</i> (A)	Canada	Pouget et al., 2013
	95–200 (C)	SHV-2	–	<i>E. coli</i> (A)	Canada	Pouget et al., 2013
	95–200 (C)	SHV-2a	<i>aadA1</i> , <i>dfrA1</i>	<i>E. coli</i> , <i>S. Kiambu</i> (A)	Canada	Pouget et al., 2013
	95–200 (C)	SHV-2a	<i>aadA1</i>	<i>E. coli</i> (A)	Canada	Pouget et al., 2013
Incl1 (ST26)	ND	SHV-12	ND	<i>E. coli</i> (H)	Italy	Accogli et al., 2013
	ND	SHV-12 (IS26)	ND	<i>E. coli</i> (A)	Portugal	Jones-Dias et al., 2016
Incl1 (ST27, CC26)	115 (C)	SHV-2	<i>aadA2</i>	<i>S. Livingstone</i> (H)	Spain	de Toro et al., 2013
Incl1 (ST29/CC26)	ND (C)	SHV-12 (IS26)	ND	<i>E. coli</i> (E)	Portugal	Jones-Dias et al., 2016
Incl1 (ST3)	ND	SHV-12	ND	<i>E. coli</i> (A)	Italy	Accogli et al., 2013
	104 (C)	SHV-12	-	<i>E. coli</i> (A)	Italy	Bortolaia et al., 2011
InclK	ND (NC)	SHV-12	ND	<i>K. pneumoniae</i> (A)	England	Timofte et al., 2014
	155	SHV-2	–	<i>E. coli</i> (A)	Netherlands	Dierikx et al., 2010
InclL/M	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
	65	SHV-12	<i>bla</i> _{KPC-2} , <i>rmtB</i>	<i>K. pneumoniae</i> (H)	China	Liu et al., 2015
	65	SHV-2	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
	ND (C)	SHV-2a	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
	60–70 (C)	SHV-2a (IS26)	ND	<i>K. pneumoniae</i> (H)	Tunisia	Elhani et al., 2010
	ND (C)	SHV-5 (IS26)	<i>aacA4</i> , <i>aacC1</i> , <i>aadA1</i> , <i>sul1</i>	<i>S. Typhimurium</i> (H)	Italy	Villa et al., 2000
	90 (C)	SHV-5	<i>tet(A)</i> , <i>aadA1</i> , <i>aacC1</i> , <i>aacA4</i> , <i>dfrA1</i>	<i>K. oxytoca</i> (H)	USA	Preston et al., 2014

(Continued)


TABLE 4 | Continued


Inc Group	Plasmid Size (Kb)*	<i>bla</i> _{SHV} allele [§]	Other Antibiotic Resistance Genes	Bacterial Species [#]	Country	References
IncN	ND (C)	SHV-12	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
	ND (C)	SHV-12	ND	<i>K. pneumoniae</i> (H)	Bulgaria	Markovska et al., 2014
	50	SHV-12	<i>bla</i> _{VIM-1} , <i>qnrS</i>	<i>K. pneumoniae</i> , <i>E. coli</i> (H)	Norway	Naseer et al., 2012
	50 (C)	SHV-12	<i>bla</i> _{VIM-1} , <i>qnrS</i>	<i>K. pneumoniae</i> (H)	Norway	Samuelsen et al., 2011
	>23	SHV-2 (IS26)	ND	<i>K. pneumoniae</i> (H)	China	Wang et al., 2012
	ND (C)	SHV-2a	ND	<i>E. coli</i> (H)	Tunisia	Mnif et al., 2013
IncN (ST1)	ND	SHV-12	<i>aadA2</i>	<i>E. coli</i> (H)	Netherlands	Dierikx et al., 2013
IncN (ST16)	50 (C)	SHV-2	ND	<i>S. Miami</i> (U)	Senegal	Harrois et al., 2014
IncP	ND (C)	SHV-12 (IS26)	-	<i>E. cloacae</i> (H)	Taiwan	Chen C. M. et al., 2015
	95–200 (C)	SHV-2a	<i>aadA1</i> , <i>dfrA1</i>	<i>E. coli</i> (A)	Canada	Pouget et al., 2013
IncX3	50 (C)	SHV-12	<i>bla</i> _{KPC-2}	<i>K. pneumoniae</i> (H)	Australia	Partridge et al., 2015
	54 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>K. pneumoniae</i> (H)	China	Wang et al., 2014
	54 (C)	SHV-12 (IS26)	<i>bla</i> _{NDM-1}	<i>K. pneumoniae</i> , <i>C. freundii</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. coli</i> (H)	China	Ho et al., 2012
	60 (C)	SHV-12	<i>bla</i> _{NDM-1} , <i>bla</i> _{TEM-1}	<i>E. coli</i> (H)	China	Huang et al., 2016
	60 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>E. coli</i> (H)	China	Huang et al., 2016
	54 (C)	SHV-12 (IS26)	<i>bla</i> _{NDM-1}	<i>E. coli</i> (H)	China	Feng et al., 2015
	54 (C)	SHV-12 (IS26)	<i>bla</i> _{NDM-1}	<i>C. freundii</i> (H)	China	Du et al., 2013
	50	SHV-12	<i>qnrB7</i>	<i>E. coli</i> (A)	Czech Republic	Dobiasova and Dolejska, 2016
	40	SHV-12	<i>qnrS1</i>	<i>E. coli</i> (E)	Czech Republic	Dobiasova and Dolejska, 2016
	53	SHV-12 (IS26)	<i>bla</i> _{KPC-2}	<i>K. pneumoniae</i> (H)	France	Kassib-Chikhani et al., 2013
	50 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>E. cloacae</i> (H)	UAE	Sonnevend et al., 2013
	50 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>E. coli</i> (H)	UAE	Sonnevend et al., 2013
	50 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>C. freundii</i> (H)	UAE	Sonnevend et al., 2013
	43 (C)	SHV-12 (IS26)	-	<i>E. cloacae</i> (H)	USA	Hargreaves et al., 2015
IncX3-N	80	SHV-12	<i>bla</i> _{TEM-1} , <i>qnrS1</i>	<i>E. coli</i> (A)	Germany	Dobiasova and Dolejska, 2016
ColETp	10 (NC)	SHV-12	<i>qnrS1</i>	<i>S. Typhimurium</i> (H)	Spain	Herrera-Leon et al., 2011
R	70	SHV-12	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
R+IncFIIk1	300	SHV-2	ND	<i>K. pneumoniae</i> (H)	Portugal	Rodrigues et al., 2014
Untypable	90–140 (C)	SHV-12 (IS26)	ND	<i>K. pneumoniae</i> (H)	Tunisia	Elhani et al., 2010
	ND	SHV-12 (IS26)	-	<i>E. coli</i> (H)	UK	Doumith et al., 2012
	ND	SHV-12 (IS26)	<i>bla</i> _{TEM-1}	<i>E. coli</i> (H)	UK	Doumith et al., 2012
	ND	SHV-12 (IS26)	<i>bla</i> _{TEM-1} , <i>bla</i> _{OXA-1} , <i>qnrS1</i>	<i>E. coli</i> (H)	UK	Doumith et al., 2012
	50 (C)	SHV-12	<i>bla</i> _{NDM-1}	<i>K. pneumoniae</i> (H)	UAE	Sonnevend et al., 2013

*C, conjugative; NC, non-conjugative; when blank is because not determined.

[§] When present, IS26 is indicated in parenthesis.

[#]H, human; A, animal (mostly poultry, turkey and broilers; check reference for full description); E, environment. ND, not determined.





E. coli, *Proteus mirabilis* and *Aeromonas caviae*, isolated from clinical samples in Tunisia, France, Korea and Italy (Marcadé et al., 2009; Song et al., 2011; Mnif et al., 2013; Antonelli et al., 2016). *E. coli* isolates recovered from clinical specimens encoding *bla*_{SHV-2}, *bla*_{SHV-2a}, and *bla*_{SHV-5} have been also identified in Tunisia and France (Marcadé et al., 2009; Mnif et al., 2013), whereas *Providencia stuartii* isolates encoding *bla*_{SHV-5} on either IncA/C or multireplicon IncA/C-R plasmids have been reported from different outbreaks in Greece (Giakkoupi et al., 2015; Oikonomou et al., 2016). Interestingly, these IncA/C plasmids (130–220 Kb) often carried multiple resistance genes, conferring multidrug resistant phenotypes (Giakkoupi et al., 2015; Antonelli et al., 2016; Oikonomou et al., 2016), resulting in the proliferation of the SHV ESBLs by co-selection.

IncF

Plasmids belonging to the narrow-host range IncF group, including plasmids with fused replicons, have been reported to accommodate *bla*_{SHV-12} among clinical *E. coli* isolates from France (IncFII), Tunisia (IncFIA-FIB, IncFII-FIA, IncFII-FIA-FIB) and United Kingdom (IncFIB), but also among food-producing animals from Italy (IncFIB) (Marcadé et al., 2009; Bortolaia et al., 2010; Doumith et al., 2012; Mnif et al., 2013). IncF plasmids account for the dissemination of *bla*_{SHV-2} gene among *E. coli* from both clinical specimens in France (IncFIB, IncFII, IncFII-FIB) and food-producing animals (avian and porcine sources) in Canada (IncFIB), as well as in clinical *K. pneumoniae* isolates belonging to ST654 and ST15 from China (IncFIB) and Portugal (IncFII), respectively (Marcadé et al., 2009; Wang et al., 2012; Pouget et al., 2013; Rodrigues et al., 2014). Finally, clinical *E. coli* and *K. pneumoniae* from Tunisia were found to encode *bla*_{SHV-2a} (Elhani et al., 2010; Mnif et al., 2013), clinical *E. coli* from Poland encoded *bla*_{SHV-5} on IncF plasmids, as well as clinical *K. pneumoniae* and *Serratia marcescens* from Uruguay

(García-Fulgueiras et al., 2011; Zienkiewicz et al., 2013), whereas the same plasmids have been associated with less prevalent SHV ESBLs (*bla*_{SHV-55} and *bla*_{SHV-106}) in clinical *K. pneumoniae* isolates from Portugal (Rodrigues et al., 2014).

IncHI2

In contrast with the IncA/C and IncF plasmids, the broad-host range IncHI2 group is responsible mainly for the dissemination of *bla*_{SHV-12}, although this group has been found incidentally to also accommodate *bla*_{SHV-2a} (Mnif et al., 2013). Plasmids of this group varying in sizes (95–400 Kb) have been reported to encode *bla*_{SHV-12} in various bacterial species, such as *E. coli*, *K. pneumoniae*, *E. cloacae*, and at least three *S. enterica* serotypes (Bredeney, Concord, and Senftenberg) from human specimens with diverse geographical origin (Netherlands, Portugal, Spain, Taiwan, Tunisia; Elhani et al., 2010; Veldman et al., 2010; Herrera-Leon et al., 2011; Mnif et al., 2013; Rodrigues et al., 2014; Chen C. M. et al., 2015). Apart from *bla*_{SHV-12}, some of these conjugative plasmids have been reported to co-encode for other resistance genes, including additional SHV ESBLs (*bla*_{CTX-M-3}, *bla*_{CTX-M-14}; Veldman et al., 2010; Chen C. M. et al., 2015).

IncI1

The IncI1 group, consisting of narrow-host range mostly conjugative plasmids, ranks amongst the top facilitators of *bla*_{SHV-2}, *bla*_{SHV-2a}, and *bla*_{SHV-12} genes. The range of bacterial species they have encountered is limited to *E. coli*, *K. pneumoniae*, *E. cloacae*, and the *S. enterica* serotypes Concord, Enteritidis, Heidelberg and Kiambu. Nevertheless, IncI1 plasmids (19–340 Kb) occur in very diverse settings: *bla*_{SHV-2}- and *bla*_{SHV-12}-encoding isolates from human infections (Bulgaria, France, Italy, Spain, Taiwan; Tato et al., 2007; Marcadé et al., 2009; Accogli et al., 2013; de Toro et al., 2013; Markovska et al., 2014; Chen C. M. et al., 2015) and colonization (Ethiopia) (Fabre et al., 2009); *bla*_{SHV-2}-, *bla*_{SHV-2a}-, and *bla*_{SHV-12}-encoding isolates

from poultry (Canada, Italy, Portugal; Bortolaia et al., 2010, 2011; Accogli et al., 2013; Pouget et al., 2013; Jones-Dias et al., 2015), *bla*_{SHV-2}- and *bla*_{SHV-2a}-encoding isolates from pigs (Canada) (Pouget et al., 2013); *bla*_{SHV-12}-encoding isolates from aquatic birds (Poland) (Literak et al., 2010); and *bla*_{SHV-12}-encoding isolates from farming soil (Portugal) (Jones-Dias et al., 2016). Remarkably, *bla*_{SHV-12} on IncI1 plasmids belonging to pST26 have been identified among *E. coli* isolates of human and animal origin (Accogli et al., 2013; Jones-Dias et al., 2015), indicating the potential transmission of these *bla*_{SHV-12}-encoding vehicles from human to animals and/or vice versa.

IncL/M and IncN

The broad-host range IncL/M and IncN plasmids contribute to a lesser extent to the epidemiology of *bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, and *bla*_{SHV-12} than the above-mentioned families. IncL/M plasmids (60–90 Kb) carrying SHV ESBL genes have been reported only among *E. coli*, *K. pneumoniae*, *K. oxytoca*, and *S. enterica* serotype Typhimurium of human origin in Portugal (*bla*_{SHV-2}), Tunisia (*bla*_{SHV-2a}, *bla*_{SHV-12}), Italy (*bla*_{SHV-5}), USA (*bla*_{SHV-5}), and recently in China (*bla*_{SHV-12}) (Villa et al., 2000; Elhani et al., 2010; Mnif et al., 2013; Preston et al., 2014; Rodrigues et al., 2014; Liu et al., 2015). The same bacterial species mostly from human sources carry IncN plasmids (~50 Kb) encoding *bla*_{SHV-2} (China, Senegal), *bla*_{SHV-2a} (Tunisia) or *bla*_{SHV-12} (Bulgaria, Netherlands, Norway, Tunisia; Samuelsen et al., 2011; Naseer et al., 2012; Wang et al., 2012; Dierikx et al., 2013; Mnif et al., 2013; Harrois et al., 2014; Markovska et al., 2014). Interestingly, the presence of IncN (pST1) plasmids encoding *bla*_{SHV-12} has been reported among *E. coli* from human and animal sources (Dierikx et al., 2013), mirroring the situation for IncI1 plasmids and underscoring the contribution of this plasmid family in the transmission of *bla*_{SHV-12} within or between these niches.

IncX3

The IncX3 plasmid subgroup consists of narrow-host range plasmids and plays an important role in the exclusive dissemination of *bla*_{SHV-12}. Conjugative plasmids (40–60 Kb) of this subgroup have been identified in diverse bacterial species (*E. coli*, *K. pneumoniae*, *C. freundii*, *E. aerogenes*, *E. cloacae*), sources (human, animal, environment) and geographical areas (Australia, China, Czech Republic, France, United Arab Emirates, US; Ho et al., 2012; Du et al., 2013; Kassis-Chikhani et al., 2013; Sonnevend et al., 2013; Wang et al., 2014; Feng et al., 2015; Hargreaves et al., 2015; Partridge et al., 2015; Dobiasova and Dolejska, 2016; Huang et al., 2016). Interestingly, the majority of these plasmids appear to co-harbor carbapenemase genes (*bla*_{KPC-2}, *bla*_{NDM-1}), whereas the co-localization of SHV ESBL and carbapenemase genes was reported only on IncA/C or IncA/C-R (*bla*_{VIM-1}), IncL/M (*bla*_{KPC-2}), and IncN (*bla*_{VIM-1}) plasmids (Samuelsen et al., 2011; Naseer et al., 2012; Giakkoupi et al., 2015; Oikonomou et al., 2016), enhancing the plasmid potential maintenance among bacterial populations and the subsequent preservation and dissemination of the SHV ESBL genes.

Miscellaneous Plasmids

*bla*_{SHV-12} has been incidentally found on: (i) a ColE plasmid from *S. enterica* serotype Typhimurium DT104b in Spain (Herrera-Leon et al., 2011); (ii) an IncK plasmid from *K. pneumoniae* in the United Kingdom (Timofte et al., 2014); (iii) an IncP plasmid from *E. cloacae* in Taiwan (Chen C. M. et al., 2015); and (iv) a plasmid assigned to the R replicon type from *K. pneumoniae* in Portugal (Rodrigues et al., 2014). *E. coli* and *K. pneumoniae* encoding *bla*_{SHV-2} on IncK plasmids were recovered from animal and human sources in the Netherlands and in Uruguay, respectively (Dierikx et al., 2010; García-Fulgueiras et al., 2011). IncP plasmids encoding *bla*_{SHV-2a} from animals in Canada and *bla*_{SHV-5} from human in Uruguay have also been reported (García-Fulgueiras et al., 2011; Pouget et al., 2013). Finally, a number of reports highlight the presence of *bla*_{SHV-12} on mostly conjugative non-typeable plasmids, according to the PCR-based replicon-typing scheme (Carattoli et al., 2005a). These plasmids of human origin, varying between 50 and 140 Kb in size, were mostly detected among *E. coli* from the United Kingdom (Doumith et al., 2012) and *K. pneumoniae* from Tunisia (Elhani et al., 2010) and United Arab Emirates (Sonnevend et al., 2013), underscoring that their dissemination is wider than we know.

IS26 Role in *bla*_{SHV} Mobilization

Analysis of the sequences bracketing several SHV ESBL genes (*bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, *bla*_{SHV-12}, *bla*_{SHV-106}, and *bla*_{SHV-134}) among Gram-negative bacteria, including Enterobacteriaceae and non-fermenters, revealed that these β -lactamase genes are mostly associated with the IS26 element (Table 4, Supplementary Table S1). Beside SHV ESBLs, this member of the IS6 insertion sequence family (Mahillon and Chandler, 1998), has been associated with a plethora of resistance genes (Allard et al., 1993; Miriagou et al., 2005; Post and Hall, 2009; Cain et al., 2010; Hordijk et al., 2011) and has been found to contribute to their expression by supplying a promoter -35 box that can be coupled with a -10 box in the adjacent DNA (Lee et al., 1990; Cain and Hall, 2011). In contrast to other insertions sequences, it has been suggested that IS26 transposes preferentially within plasmids rather than into the chromosome (He et al., 2015), possibly explaining the linkage of IS26 and the four predominant SHV ESBL genes with IncA/C (*bla*_{SHV-5}, *bla*_{SHV-12}), IncF (*bla*_{SHV-2a}, *bla*_{SHV-5}, *bla*_{SHV-12}), IncHI2 (*bla*_{SHV-12}), IncI1 (*bla*_{SHV-12}), IncL/M (*bla*_{SHV-2a}, *bla*_{SHV-5}), IncN (*bla*_{SHV-2}), IncP (*bla*_{SHV-12}), IncX3 (*bla*_{SHV-12}), and non-typeable (*bla*_{SHV-12}) plasmids (Table 4). Similarly to most antibiotic resistance genes, IS26-mediated mobilization of SHV ESBL genes on conjugative plasmids facilitated their subsequent intra- and inter-species dissemination (Table 4). Available sequences of transposons flanked by copies of intact and/or truncated IS26 elements (Figure 3) and coding for SHV ESBL genes show the presence of other co-linear genes originating from the chromosome of *K. pneumoniae* (i.e., *fucA*, *ygbI*, *ygbK*, *ygbJ*, *ygbM*, *deoR*; Ho et al., 2012; Wang et al., 2012; Du et al., 2013; Kassis-Chikhani et al., 2013; Preston et al., 2014; Chen C. M. et al., 2015; Feng et al., 2015; Giakkoupi et al., 2015), likely underscoring the involvement of IS26 in the mobilization

of *bla*_{SHV} from the chromosome of *K. pneumoniae*, as previously suggested (Haeggman et al., 1997).

OUTSIDE OF THE ENTEROBACTERIACEAE AND A FEW PECULIAR SHV ESBLs

SHV β -lactamases have virtually invaded all human, environmental and animal sceneries, mostly associated to Enterobacteriaceae. In recent years, the first reports of alternative bacterial hosts have been described, notably in Aeromonads, ubiquitous in aquatic habitats and occasionally able to cause human infections. *bla*_{SHV-12} was detected, in association with *bla*_{FOX-2} and *bla*_{CTX-M-15}, on the chromosome of the foodborne pathogens *A. caviae* and *Aeromonas hydrophila* from wild-growing mussels from Croatia (Maravić et al., 2013). The first identification of plasmid-encoded SHV-12, together with VIM-1, occurred in clinical *A. caviae* accountable for a newborn bloodstream infection (Antonelli et al., 2016). The coproduction of these enzymes highlights the potential risks for public health and the role of Aeromonads as reservoirs and dissemination tools of resistance determinants in both environmental and clinical settings.

Occasionally, SHV ESBL-producing *Pseudomonas aeruginosa* can be detected in nosocomial settings and can pose a serious threat as healthcare-associated infection in many regions of the world. *bla*_{SHV-2a} was first identified on the chromosome of a 1995 clinical *P. aeruginosa* strain, with high sequence homology to plasmid pMPA2a from *K. pneumoniae* indicating a likely enterobacterial gene origin (Naas et al., 1999). Subsequent studies demonstrated the insertion of *bla*_{SHV} alleles into *P. aeruginosa* chromosome: *bla*_{SHV} in China (Chen Z. et al., 2015) and Iran (Shahcheraghi et al., 2009); *bla*_{SHV-5} (Poirel et al., 2004) and *bla*_{SHV-12} (Neonakis et al., 2003) in Greece; and *bla*_{SHV-2a} in Tunisia (Mansour et al., 2009) and France (Hocquet et al., 2010; Jeannot et al., 2013). The role of IS26 in the mobilization of *bla*_{SHV-12} was demonstrated by the chromosomal insertion of an IS26 composite transposon (>24 kb) thanks to the co-mobilization of antibiotic resistance *aac*(6′)-Ib, which confers amikacin resistance, likely occurred during the clinical course of a burn infection, immediately after amikacin administration (Uemura et al., 2010). *bla*_{SHV-5}, *bla*_{SHV-11}, *bla*_{SHV-12} were also detected in different combinations, together with *bla*_{TEM-1b}, on various plasmids in Thailand (Chanawong et al., 2001b).

Finally, one of the most effective associations outside of the Enterobacteriaceae is with *Acinetobacter baumannii*, contributing to the worrisome spread of ESBL-producing strains especially in clinical outbreaks (Blackwell et al., 2016). Plasmid transfer from nosocomial SHV-encoding Enterobacteriaceae seems to be responsible for this phenomenon, as observed for SHV-12 in the Netherlands (Naiemi et al., 2005), or SHV-5 in the USA, a country where this variant is the most prevalent ESBL gene in Enterobacteriaceae (Naas et al., 2007).

Among all ESBL SHV β -lactamases, few enzymes deserve special consideration because of their unique enzymatic features.

SHV-38 is a unique allelic variant of the SHV family to have an expanded-spectrum to carbapenems. It was first described in *K. pneumoniae* from France (Poirel et al., 2003) and it holds a point mutation (Ala¹⁴⁶Val) compared to the chromosome-encoded SHV-1. Among all 46 available SHV ESBL variants, only SHV-38 possess the Ala¹⁴⁶Val substitution (Table 2), likely inducing subtle structural conformational changes favoring imipenem but not meropenem hydrolysis (Walther-Rasmussen and Høiby, 2007).

SHV-129 is a novel clinically acquired variant identified in 2012 from an Italian *E. coli* isolate (Table 1; Lascols et al., 2012) and it represents an interesting example of enzyme evolution due to antibiotic pressure. Alongside two well-known amino acid substitutions (Gly²³⁸Ser, Glu²⁴⁰Lys), SHV-129 contains new substitutions, Arg²⁷⁵Leu and Asn¹⁴⁶Asp). The latter was recently demonstrated to be the first global suppressor substitution identified in the SHV β -lactamase family (Winkler and Bonomo, 2016), likely helping in protein stabilization and functionality, as well as in the ability of the enzyme to acquire additional substitutions. It is also proposed that due to the increasing clinical use of cefepime, SHV-129 might have evolved from SHV-2 or SHV-5 in an alternative conformation to expand its spectrum to hydrolyze cefepime, as mirrored by the kinetic parameters of the three enzymes (Table 3).

Finally, SHV-2 can be located on both chromosome and self-transmissible plasmids (Table 4; Supplementary Table S1). Association of *bla*_{SHV-2} with RCS47, a P1-like bacteriophage that infects and lysogenizes *E. coli* and several other enteric bacteria, was recently reported (Billard-Pomares et al., 2014). *bla*_{SHV-2} is flanked by two IS26 elements that likely drove the insertion in the phage backbone. The P1-like prophages were found with high prevalence in natural *E. coli* of both animal and human origin, including ESBL-producing isolates. This kind of association was already reported for other β -lactamases (*bla*_{TEM}, *bla*_{CTX-M}, and *mecA*) from river and urban sewage water (von Wintersdorff et al., 2016), suggesting that bacteriophages might play a wider role in favoring horizontal transfer of antibiotic resistance determinants than initially thought (Muniesa et al., 2013).

CONCLUDING REMARKS

Tzouveleki and Bonomo suggested that “it will not be surprising if (SHV) enzymes will continue to expand their substrate spectrum as long as the current antibiotics, or novel ones derived from the basic β -lactam structure, are used” (Tzouveleki and Bonomo, 1999). In the last two decades we observed the appearance of multiple SHV-type variants, with few ones significantly expanding their substrate. One exception is represented by SHV-38, the only known SHV allelic variant able to hydrolyze carbapenems (Poirel et al., 2003), a feature that has not been associated with any TEM or CTX-M enzyme. In this image resides the fate of SHV extended β -lactamases, unable to undergo the dominant propagation observed, for instance, for the CTX-M family but yet contributing to β -lactam resistance in a not negligible way.

The persistence of SHV enzymes in the bacterial community might also be secured by co-selection with emerging resistance genes. Association of *bla*_{SHV-12} with IncX3 plasmids carrying carbapenemase genes *bla*_{KPC-2} and *bla*_{NDM} has been observed in recent years (Table 4) and it seems to be a phenomenon occurring in clinical carbapenem-resistant Enterobacteriaceae worldwide (Kassis-Chikhani et al., 2013; Sonnevend et al., 2013; Partridge et al., 2015; Huang et al., 2016). As highlighted in this review, the association of successful variants *bla*_{SHV-2}, *bla*_{SHV-2a}, *bla*_{SHV-5}, and *bla*_{SHV-12} with different families of conjugative plasmids (IncA/C, IncF, IncHI2) might also underlie the colonization of virtually all ecological niches encompassing food producing animals, aquatic environment, wildlife, companion animals, and vegetables. Plasmid mediated transfer from nosocomial Enterobacteriaceae enabled SHV dispersion toward alternative bacterial hosts such as the emerging nosocomial pathogens of aquatic origin *S. maltophilia* and *A. caviae*, or contributed to the worrisome spread of ESBL-producing strains of *A. baumannii* and *P. aeruginosa*. Most interestingly, the ubiquitous presence of SHV ESBL genes and plasmids is suggestive for transmission in human, animals, and the environment, most likely through the food chain, highlighting the potential risks for public health and endorsing a one health research approach.

Overall, SHV ESBL enzymes have kept a stable role in antibiotic resistance over the years. Allele diversification is still

occurring, the latest variant being identified in *E. cloacae* in 2014 (*bla*_{SHV-183}), and effective associations with new genetic platforms are taking place helping expansion toward novel bacterial hosts and reservoirs.

AUTHOR CONTRIBUTIONS

The paper was written by AL and DC, and reviewed by DM. All authors discussed, read, contributed to and approved the final manuscript.

ACKNOWLEDGMENTS

The authors would like to thank Mr. Bo Derks for valuable technical help in Figure 2 production. AL was partially supported by the ESBLAT project (BO-22.04-008-001). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.01374>

REFERENCES

- Abbassi, M. S., Torres, C., Achour, W., Vinué, L., Sáenz, Y., Costa, D., et al. (2008). Genetic characterisation of CTX-M-15-producing *Klebsiella pneumoniae* and *Escherichia coli* strains isolated from stem cell transplant patients in Tunisia. *Int. J. Antimicrob. Agents* 32, 308–314. doi: 10.1016/j.ijantimicag.2008.04.009
- Accogli, M., Fortini, D., Giufrè, M., Graziani, C., Dolejska, M., Carattoli, A., et al. (2013). IncI1 plasmids associated with the spread of CMY-2, CTX-M-1 and SHV-12 in *Escherichia coli* of animal and human origin. *Clin. Microbiol. Infect.* 19, E238–E240. doi: 10.1111/1469-0691.12128
- Ahmed, A. M., and Shimamoto, T. (2011). Molecular characterization of antimicrobial resistance in Gram-negative bacteria isolated from bovine mastitis in Egypt. *Microbiol. Immunol.* 55, 318–327. doi: 10.1111/j.1348-0421.2011.00323.x
- Ahmed, A. M., Shimamoto, T., and Shimamoto, T. (2013). Molecular characterization of multidrug-resistant avian pathogenic *Escherichia coli* isolated from septicemic broilers. *Int. J. Med. Microbiol.* 303, 475–483. doi: 10.1016/j.ijmm.2013.06.009
- Ahmed, A. M., Younis, E. E., Ishida, Y., and Shimamoto, T. (2009). Genetic basis of multidrug resistance in *Salmonella enterica* serovars Enteritidis and Typhimurium isolated from diarrheic calves in Egypt. *Acta Trop.* 111, 144–149. doi: 10.1016/j.actatropica.2009.04.004
- Alcalá, L., Alonso, C. A., Simón, C., González-Esteban, C., Orós, J., Rezusta, A., et al. (2015). Wild birds, frequent carriers of Extended-Spectrum β -Lactamase (ESBL) producing *Escherichia coli* of CTX-M and SHV-12 types. *Microb. Ecol.* doi: 10.1007/s00248-015-0718-0. [Epub ahead of print].
- Allard, J. D., Gibson, M. L., Vu, L. H., Nguyen, T. T., and Bertrand, K. P. (1993). Nucleotide sequence of class D tetracycline resistance genes from *Salmonella* ordenez. *Mol. Gen. Genet.* 237, 301–305. doi: 10.1007/bf00282811
- Alves, M. S., Pereira, A., Araújo, S. M., Castro, B. B., Correia, A. C., and Henriques, I. (2014). Seawater is a reservoir of multi-resistant *Escherichia coli*, including strains hosting plasmid-mediated quinolones resistance and extended-spectrum β -lactamases genes. *Front. Microbiol.* 5:426. doi: 10.3389/fmicb.2014.00426
- Ambler, R. P. (1980). The structure of β -lactamases. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 289, 321–331. doi: 10.1098/rstb.1980.0049
- Anjum, M. F., Lemma, F., Cork, D. J., Meunier, D., Murphy, N., North, S. E., et al. (2013). Isolation and detection of extended spectrum β -lactamase (ESBL)-producing Enterobacteriaceae from meat using chromogenic agars and isothermal loop-mediated amplification (LAMP) assays. *J. Food Sci.* 78, M1892–M1898. doi: 10.1111/1750-3841.12297
- Antonelli, A., D'andrea, M. M., Montagnani, C., Bartolesi, A. M., Di Pilato, V., Fiorini, P., et al. (2016). Newborn bacteraemia caused by an *Aeromonas caviae* producing the VIM-1 and SHV-12 β -lactamases, encoded by a transferable plasmid. *J. Antimicrob. Chemother.* 71, 272–274. doi: 10.1093/jac/dkv304
- Arlet, G., Rouveau, M., Bengoufa, D., Nicolas, M. H., and Philippon, A. (1991). Novel transferable extended-spectrum β -lactamase (SHV-6) from *Klebsiella pneumoniae* conferring selective resistance to ceftazidime. *FEMS Microbiol. Lett.* 81, 57–62. doi: 10.1016/0378-1097(91)90471-1
- Arlet, G., Rouveau, M., Casin, I., Bouvet, P. J., Lagrange, P. H., and Philippon, A. (1994). Molecular epidemiology of *Klebsiella pneumoniae* strains that produce SHV-4 β -lactamase and which were isolated in 14 French hospitals. *J. Clin. Microbiol.* 32, 2553–2558.
- Arlet, G., Sanson-le Pors, M. J., Rouveau, M., Fournier, G., Marie, O., Schlemmer, B., et al. (1990). Outbreak of nosocomial infections due to *Klebsiella pneumoniae* producing SHV-4 β -lactamase. *Eur. J. Clin. Microbiol. Infect. Dis.* 9, 797–803. doi: 10.1007/BF01967377
- Arpin, C., Coze, C., Rogues, A. M., Gachie, J. P., Bebear, C., and Quentin, C. (1996). Epidemiological study of an outbreak due to multidrug-resistant *Enterobacter aerogenes* in a medical intensive care unit. *J. Clin. Microbiol.* 34, 2163–2169.
- Arpin, C., Labia, R., Andre, C., Frigo, C., El Harrif, Z., and Quentin, C. (2001). SHV-16, a β -lactamase with a pentapeptide duplication in the omega loop. *Antimicrob. Agents Chemother.* 45, 2480–2485. doi: 10.1128/AAC.45.9.2480-2485.2001
- Barthélémy, M., Peduzzi, J., and Labia, R. (1988). Complete amino acid sequence of p453-plasmid-mediated PIT-2 β -lactamase (SHV-1). *Biochem. J.* 251, 73–79. doi: 10.1042/bj2510073

- Bedenic, B., Randegger, C., Boras, A., and Haechler, H. (2001). Comparison of five different methods for detection of SHV extended-spectrum β -lactamases. *J. Chemother.* 13, 24–33. doi: 10.1179/joc.2001.13.1.24
- Bedenic, B., Vranes, J., Mihaljevic, L. J., Tonkic, M., Sviben, M., Plecko, V., et al. (2007). Sensitivity and specificity of various β -lactam antibiotics and phenotypical methods for detection of TEM, SHV and CTX-M extended-spectrum β -lactamases. *J. Chemother.* 19, 127–139. doi: 10.1179/joc.2007.19.2.127
- Ben Achour, N., Belhadj, O., Galleni, M., Ben Moussa, M., and Mercuri, P. S. (2014). Study of a natural mutant SHV-type β -lactamase, SHV-104, from *Klebsiella pneumoniae*. *Int. J. Microbiol.* 2014, 6. doi: 10.1155/2014/548656
- Billard-Pomares, T., Fouteau, S., Jacquet, M. E., Roche, D., Barbe, V., Castellanos, M., et al. (2014). Characterization of a P1-like bacteriophage carrying an SHV-2 extended-spectrum β -lactamase from an *Escherichia coli* strain. *Antimicrob. Agents Chemother.* 58, 6550–6557. doi: 10.1128/AAC.03183-14
- Blackwell, G. A., Hamidian, M., and Hall, R. M. (2016). IncM plasmid R1215 is the source of chromosomally located regions containing multiple antibiotic resistance genes in the globally disseminated *Acinetobacter baumannii* GC1 and GC2 clones. *mSphere* 1:e00117-16. doi: 10.1128/mSphere.00117-16
- Blanc, V., Mesa, R., Saco, M., Lavilla, S., Prats, G., Miró, E., et al. (2006). ESBL- and plasmidic class C β -lactamase-producing *E. coli* strains isolated from poultry, pig and rabbit farms. *Vet. Microbiol.* 118, 299–304. doi: 10.1016/j.vetmic.2006.08.002
- Bonnet, R. (2004). Growing group of extended-spectrum β -lactamases: the CTX-M enzymes. *Antimicrob. Agents Chemother.* 48, 1–14. doi: 10.1128/AAC.48.1.1-14.2004
- Bortolaia, V., Guardabassi, L., Trevisani, M., Bisgaard, M., Venturi, L., and Bojesen, A. M. (2010). High diversity of Extended-Spectrum β -Lactamases in *Escherichia coli* isolates from Italian broiler flocks. *Antimicrob. Agents Chemother.* 54, 1623–1626. doi: 10.1128/AAC.01361-09
- Bortolaia, V., Larsen, J., Damborg, P., and Guardabassi, L. (2011). Potential pathogenicity and host range of Extended-Spectrum β -Lactamase-producing *Escherichia coli* isolates from healthy poultry. *Appl. Environ. Microbiol.* 77, 5830–5833. doi: 10.1128/AEM.02890-10
- Bourouis, A., Ben Moussa, M., and Belhadj, O. (2015). Multidrug-resistant phenotype and isolation of a novel SHV- β -lactamase variant in a clinical isolate of *Enterobacter cloacae*. *J. Biomed. Sci.* 22, 1–7. doi: 10.1186/s12929-015-0131-5
- Bradford, P. A. (2001). Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin. Microbiol. Rev.* 14, 933–951. doi: 10.1128/CMR.14.4.933-951.2001
- Bradford, P. A., Urban, C., Jaiswal, A., Mariano, N., Rasmussen, B. A., Projan, S. J., et al. (1995). SHV-7, a novel cefotaxime-hydrolyzing β -lactamase, identified in *Escherichia coli* isolates from hospitalized nursing home patients. *Antimicrob. Agents Chemother.* 39, 899–905. doi: 10.1128/AAC.39.4.899
- Bush, K. (2013). The ABCD's of β -lactamase nomenclature. *J. Infect. Chemother.* 19, 549–559. doi: 10.1007/s10156-013-0640-7
- Bush, K., and Fisher, J. F. (2011). Epidemiological expansion, structural studies, and clinical challenges of new β -lactamases from Gram-negative bacteria. *Annu. Rev. Microbiol.* 65, 455–478. doi: 10.1146/annurev-micro-090110-102911
- Bush, K., and Jacoby, G. A. (2010). Updated functional classification of β -lactamases. *Antimicrob. Agents Chemother.* 54, 969–976. doi: 10.1128/AAC.01009-09
- Bush, K., Jacoby, G. A., and Medeiros, A. A. (1995). A functional classification scheme for β -lactamases and its correlation with molecular structure. *Antimicrob. Agents Chemother.* 39, 1211–1233. doi: 10.1128/AAC.39.6.1211
- Bush, K., and Singer, S. B. (1989). New plasmid borne β -lactamases - biochemical characteristics of extended broad-spectrum β -lactamases. *Infection* 17, 429–433. doi: 10.1007/BF01645566
- Cain, A. K., and Hall, R. M. (2011). Transposon Tn5393e Carrying the aphA1-Containing Transposon Tn6023 upstream of strAB does not confer resistance to Streptomycin. *Microb. Drug Resist.* 17, 389–394. doi: 10.1089/mdr.2011.0037
- Cain, A. K., Liu, X., Djordjevic, S. P., and Hall, R. M. (2010). Transposons related to Tn1696 in IncHI2 Plasmids from Multiple Antibiotic Resistant *Salmonella enterica* Serovar Typhimurium from Australian Animals. *Microb. Drug Resist.* 16, 197–202. doi: 10.1089/mdr.2010.0042
- Calhau, V., Mendes, C., Pena, A., Mendonça, N., and Da Silva, G. J. (2015). Virulence and plasmidic resistance determinants of *Escherichia coli* isolated from municipal and hospital wastewater treatment plants. *J. Water Health* 13, 311–318. doi: 10.2166/wh.2014.327
- Cantón, R., González-Alba, J. M., and Galán, J. C. (2012). CTX-M enzymes: origin and diffusion. *Front. Microbiol.* 3:110. doi: 10.3389/fmicb.2012.00110
- Carattoli, A. (2009). Resistance plasmid families in Enterobacteriaceae. *Antimicrob. Agents Chemother.* 53, 2227–2238. doi: 10.1128/AAC.01707-08
- Carattoli, A. (2013). Plasmids and the spread of resistance. *Int. J. Med. Microbiol.* 303, 298–304. doi: 10.1016/j.ijmm.2013.02.001
- Carattoli, A., Bertini, A., Villa, L., Falbo, V., Hopkins, K. L., and Threlfall, E. J. (2005a). Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63, 219–228. doi: 10.1016/j.mimet.2005.03.018
- Carattoli, A., Lovari, S., Franco, A., Cordaro, G., Di Matteo, P., and Battisti, A. (2005b). Extended-spectrum β -lactamases in *Escherichia coli* isolated from dogs and cats in Rome, Italy, from 2001 to 2003. *Antimicrob. Agents Chemother.* 49, 833–835. doi: 10.1128/AAC.49.2.833-835.2005
- Chanawong, A., M'Zali, F. H., Heritage, J., Lulitanond, A., and Hawkey, P. M. (2000). Characterisation of extended-spectrum β -lactamases of the SHV family using a combination of PCR-single strand conformational polymorphism (PCR-SSCP) and PCR-restriction fragment length polymorphism (PCR-RFLP). *FEMS Microbiol. Lett.* 184, 85–89. doi: 10.1111/j.1574-6968.2000.tb08995.x
- Chanawong, A., M'Zali, F. H., Heritage, J., Lulitanond, A., and Hawkey, P. M. (2001a). Discrimination of SHV β -lactamase genes by restriction site insertion-PCR. *Antimicrob. Agents Chemother.* 45, 2110–2114. doi: 10.1128/AAC.45.7.2110-2114.2001
- Chanawong, A., M'Zali, F. H., Heritage, J., Lulitanond, A., and Hawkey, P. M. (2001b). SHV-12, SHV-5, SHV-2a and VEB-1 extended-spectrum β -lactamases in Gram-negative bacteria isolated in a university hospital in Thailand. *J. Antimicrob. Chemother.* 48, 839–852. doi: 10.1093/jac/48.6.839
- Chen, C. M., Yu, W. L., Huang, M., Liu, J. J., Chen, I. C., Chen, H. F., et al. (2015). Characterization of IS26-composite transposons and multidrug resistance in conjugative plasmids from *Enterobacter cloacae*. *Microbiol. Immunol.* 59, 516–525. doi: 10.1111/1348-0421.12289
- Chen, Z., Niu, H., Chen, G., Li, M., Li, M., and Zhou, Y. (2015). Prevalence of ESBLs-producing *Pseudomonas aeruginosa* isolates from different wards in a Chinese teaching hospital. *Int. J. Clin. Exp. Med.* 8, 19400–19405.
- Chia, J. H., Chu, C., Su, L. H., Chiu, C. H., Kuo, A. J., Sun, C. F., et al. (2005). Development of a multiplex PCR and SHV melting-curve mutation detection system for detection of some SHV and CTX-M β -lactamases of *Escherichia coli*, *Klebsiella pneumoniae*, and *Enterobacter cloacae* in Taiwan. *J. Clin. Microbiol.* 43, 4486–4491. doi: 10.1128/JCM.43.9.4486-4491.2005
- Corkill, J. E., Cuevas, L. E., Gurgel, R. Q., Greensill, J., and Hart, C. A. (2001). SHV-27, a novel cefotaxime-hydrolyzing β -lactamase, identified in *Klebsiella pneumoniae* isolates from a Brazilian hospital. *J. Antimicrob. Chemother.* 47, 463–465. doi: 10.1093/jac/47.4.463
- D'Andrea, M. M., Arena, F., Pallecchi, L., and Rossolini, G. M. (2013). CTX-M-type β -lactamases: a successful story of antibiotic resistance. *Int. J. Med. Microbiol.* 303, 305–317. doi: 10.1016/j.ijmm.2013.02.008
- Datta, N., and Kontomichalou, P. (1965). Penicillinase synthesis controlled by infectious R factors in Enterobacteriaceae. *Nature* 208, 239–241. doi: 10.1038/208239a0
- Decache, Y., Ireng, L. M., Ambroise, J., Savov, E., Marinescu, D., Chirimwami, R. B., et al. (2015). A qPCR and multiplex pyrosequencing assay combined with automated data processing for rapid and unambiguous detection of ESBL-producers Enterobacteriaceae. *AMB Express* 5:136. doi: 10.1186/s13568-015-0136-1
- de Toro, M., García, P., Rodríguez, I., Rojo-Bezares, B., Helmuth, R., Sáenz, Y., et al. (2013). Characterisation of plasmids implicated in the mobilisation of extended-spectrum and AmpC β -lactamase genes in clinical *Salmonella enterica* isolates and temporal stability of the resistance genotype. *Int. J. Antimicrob. Agents* 42, 167–172. doi: 10.1016/j.ijantimicag.2013.04.016
- Dierikx, C., Van Der Goot, J., Fabri, T., Van Essen-Zandbergen, A., Smith, H., and Mevius, D. (2013). Extended-spectrum- β -lactamase- and AmpC- β -lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *J. Antimicrob. Chemother.* 68, 60–67. doi: 10.1093/jac/dks349
- Dierikx, C., van Essen-Zandbergen, A., Veldman, K., Smith, H., and Mevius, D. (2010). Increased detection of extended spectrum β -lactamase producing *Salmonella enterica* and *Escherichia coli* isolates from poultry. *Vet. Microbiol.* 145, 273–278. doi: 10.1016/j.vetmic.2010.03.019

- Dobiasova, H., and Dolejska, M. (2016). Prevalence and diversity of IncX plasmids carrying fluoroquinolone and β -lactam resistance genes in *Escherichia coli* originating from diverse sources and geographical areas. *J. Antimicrob. Chemother.* 71, 2118–2124. doi: 10.1093/jac/dkw144
- Dolejská, M., Bierošová, B., Kohoutová, L., Literák, I., and Čížek, A. (2009). Antibiotic-resistant *Salmonella* and *Escherichia coli* isolates with integrons and extended-spectrum β -lactamases in surface water and sympatric black-headed gulls. *J. Appl. Microbiol.* 106, 1941–1950. doi: 10.1111/j.1365-2672.2009.04155.x
- Doumith, M., Dhanji, H., Ellington, M. J., Hawkey, P., and Woodford, N. (2012). Characterization of plasmids encoding extended-spectrum β -lactamases and their addiction systems circulating among *Escherichia coli* clinical isolates in the UK. *J. Antimicrob. Chemother.* 67, 878–885. doi: 10.1093/jac/dkr553
- Dropa, M., Balsalobre, L. C., Lincopan, N., Matté, G. R., and Matté, M. H. (2015). Complex class 1 integrons harboring CTX-M-2-encoding genes in clinical Enterobacteriaceae from a hospital in Brazil. *J. Infect. Dev. Ctries* 9, 890–897. doi: 10.3855/jidc.6241
- Du, X. X., Wang, J. F., Fu, Y., Zhao, F., Chen, Y., Wang, H. P., et al. (2013). Genetic characteristics of blaNDM-1-positive plasmid in *Citrobacter freundii* isolate separated from a clinical infectious patient. *J. Med. Microbiol.* 62, 1332–1337. doi: 10.1099/jmm.0.057091-0
- Duval, V., Maiga, I., Maiga, A., Guillard, T., Brasme, L., Forte, D., et al. (2009). High prevalence of CTX-M-type β -lactamases among clinical isolates of Enterobacteriaceae in Bamako, Mali. *Antimicrob. Agents Chemother.* 53, 4957–4958. doi: 10.1128/AAC.00675-09
- Egea, P., López-Cerero, L., Navarro, M. D., Rodríguez-Baño, J., and Pascual, A. (2011). Assessment of the presence of extended-spectrum β -lactamase-producing *Escherichia coli* in eggshells and ready-to-eat products. *Eur. J. Clin. Microbiol. Infect. Dis.* 30, 1045–1047. doi: 10.1007/s10096-011-1168-3
- Elhani, D., Bakir, L., Aouni, M., Passet, V., Arlet, G., Brisse, S., et al. (2010). Molecular epidemiology of extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* strains in a university hospital in Tunis, Tunisia, 1999–2005. *Clin. Microbiol. Infect.* 16, 157–164. doi: 10.1111/j.1469-0691.2009.03057.x
- El Harrif-Heraud, Z., Arpin, C., Benliman, S., and Quentin, C. (1997). Molecular epidemiology of a nosocomial outbreak due to SHV-4-producing strains of *Citrobacter diversus*. *J. Clin. Microbiol.* 35, 2561–2567.
- Espinal, P., Garza-Ramos, U., Reyna, F., Rojas-Moreno, T., Sanchez-Perez, A., Carrillo, B., et al. (2010). Identification of SHV-type and CTX-M-12 extended-spectrum β -lactamases (ESBLs) in multiresistant Enterobacteriaceae from Colombian Caribbean hospitals. *J. Chemother.* 22, 160–164. doi: 10.1179/joc.2010.22.3.160
- Essack, S. Y., Hall, L. M. C., and Livermore, D. M. (2004). *Klebsiella pneumoniae* isolate from South Africa with multiple TEM, SHV and AmpC β -lactamases. *Int. J. Antimicrob. Agents* 23, 398–400. doi: 10.1016/j.ijantimicag.2003.08.010
- Evans, B. A., and Amyes, S. G. (2014). OXA β -lactamases. *Clin. Microbiol. Rev.* 27, 241–263. doi: 10.1128/CMR.00117-13
- Fabre, L., Delauné, A., Espié, E., Nygard, K., Pardos, M., Polomack, L., et al. (2009). Chromosomal integration of the extended-spectrum β -lactamase gene blaCTX-M-15 in *Salmonella enterica* serotype Concord isolates from internationally adopted children. *Antimicrob. Agents Chemother.* 53, 1808–1816. doi: 10.1128/AAC.00451-08
- Feng, Y., Yang, P., Xie, Y., Wang, X., McNally, A., and Zong, Z. (2015). *Escherichia coli* of sequence type 3835 carrying blaNDM-1, blaCTX-M-15, blaCMY-42 and blaSHV-12. *Sci. Rep.* 5:12275. doi: 10.1038/srep12275
- Fernando, D. M., Tun, H. M., Poole, J., Patidar, R., Li, R., Mi, R., et al. (2016). Detection of antibiotic resistance genes in source and drinking water samples from a first nation community in Canada. *Appl. Environ. Microbiol.* 82, 4767–4775. doi: 10.1128/aem.00798-16
- Fisher, J. F., Meroueh, S. O., and Mobashery, S. (2005). Bacterial resistance to β -lactam antibiotics: compelling opportunism, compelling opportunity. *Chem. Rev.* 105, 395–424. doi: 10.1021/cr030102i
- García-Fulgueiras, V., Bado, I., Mota, M. I., Robino, L., Cordeiro, N. F., Varela, A., et al. (2011). Extended-spectrum β -lactamases and plasmid-mediated quinolone resistance in enterobacterial clinical isolates in the paediatric hospital of Uruguay. *J. Antimicrob. Chemother.* 66, 1725–1729. doi: 10.1093/jac/dkr222
- Giakkoupi, P., Tryfinopoulou, K., Polemis, M., Pappa, O., Miriagou, V., and Vatopoulos, A. (2015). Circulation of a multiresistant, conjugative, IncA/C plasmid within the nosocomial *Providencia stuartii* population in the Athens area. *Diagn. Microbiol. Infect. Dis.* 82, 62–64. doi: 10.1016/j.diagmicrobio.2015.02.009
- Gündođdu, A., Jennison, A. V., Smith, H. V., Stratton, H., and Katouli, M. (2013). Extended-spectrum β -lactamase producing *Escherichia coli* in hospital wastewaters and sewage treatment plants in Queensland, Australia. *Can. J. Microbiol.* 59, 737–745. doi: 10.1139/cjm-2013-0515
- Gutmann, L., Ferré, B., Goldstein, F. W., Rizk, N., Pinto-Schuster, E., Acar, J. F., et al. (1989). SHV-5, a novel SHV-type β -lactamase that hydrolyzes broad-spectrum cephalosporins and monobactams. *Antimicrob. Agents Chemother.* 33, 951–956. doi: 10.1128/AAC.33.6.951
- Haanperä, M., Forssten, S. D., Huovinen, P., and Jalava, J. (2008). Typing of SHV extended-spectrum β -lactamases by pyrosequencing in *Klebsiella pneumoniae* strains with chromosomal SHV β -lactamase. *Antimicrob. Agents Chemother.* 52, 2632–2635. doi: 10.1128/AAC.01259-07
- Haeggman, S., Löfdahl, S., and Burman, L. G. (1997). An allelic variant of the chromosomal gene for class A β -lactamase K2, specific for *Klebsiella pneumoniae*, is the ancestor of SHV-1. *Antimicrob. Agents Chemother.* 41, 2705–2709.
- Hall, B. G., and Barlow, M. (2004). Evolution of the serine β -lactamases: past, present and future. *Drug Resist. Updat.* 7, 111–123. doi: 10.1016/j.drug.2004.02.003
- Hammad, A. M., and Shimamoto, T. (2011). Asymptomatic intramammary infection with multidrug-resistant Gram-negative bacteria in a research dairy farm: incidence and genetic basis of resistance. *J. Vet. Med. Sci.* 73, 1089–1092. doi: 10.1292/jvms.10-0361
- Hammami, S., Boubaker, I. B.-B., Saidani, M., Lakhali, E., Hassen, A. B., Kamoun, A., et al. (2011). Characterization and molecular epidemiology of extended spectrum β -lactamase producing *Enterobacter cloacae* isolated from a Tunisian hospital. *Microb. Drug Resist.* 18, 59–65. doi: 10.1089/mdr.2011.0074
- Haque, A., Yoshizumi, A., Saga, T., Ishii, Y., and Tateda, K. (2014). ESBL-producing Enterobacteriaceae in environmental water in Dhaka, Bangladesh. *J. Infect. Chemother.* 20, 735–737. doi: 10.1016/j.jiac.2014.07.003
- Hargreaves, M. L., Shaw, K. M., Dobbins, G., Snippet Vagnone, P. M., Harper, J. E., Boxrud, D., et al. (2015). Clonal dissemination of *Enterobacter cloacae* harboring blaKPC-3 in the upper midwestern United States. *Antimicrob. Agents Chemother.* 59, 7723–7734. doi: 10.1128/AAC.01291-15
- Harrois, D., Breurec, S., Seck, A., Delauné, A., Hello, S. L., Gándara, M. P. D. L., et al. (2014). Prevalence and characterization of extended-spectrum β -lactamase-producing clinical *Salmonella enterica* isolates in Dakar, Senegal, from 1999 to 2009. *Clin. Microbiol. Infect.* 20, O109–O116. doi: 10.1111/1469-0691.12339
- Hasman, H., Mevius, D., Veldman, K., Olesen, I., and Aarestrup, F. M. (2005). β -Lactamases among extended-spectrum β -lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in The Netherlands. *J. Antimicrob. Chemother.* 56, 115–121. doi: 10.1093/jac/dki190
- He, S., Hickman, A. B., Varani, A. M., Siguier, P., Chandler, M., Dekker, J. P., et al. (2015). Insertion sequence IS26 reorganizes Plasmids in clinically isolated multidrug-resistant Bacteria by replicative transposition. *mBio* 6:e00762-15. doi: 10.1128/mBio.00762-15
- Heritage, J., Chambers, P. A., Tyndall, C., and Buescher, E. S. (2003). SHV-34: an extended-spectrum β -lactamase encoded by an epidemic plasmid. *J. Antimicrob. Chemother.* 52, 1015–1017. doi: 10.1093/jac/dkh017
- Herrera-Leon, S., Gonzalez-Sanz, R., Herrera-Leon, L., and Echeita, M. A. (2011). Characterization of multidrug-resistant Enterobacteriaceae carrying plasmid-mediated quinolone resistance mechanisms in Spain. *J. Antimicrob. Chemother.* 66, 287–290. doi: 10.1093/jac/dkq423
- Hiki, M., Usui, M., Kojima, A., Ozawa, M., Ishii, Y., and Asai, T. (2013). Diversity of plasmid replicons encoding the blaCMY-2 gene in broad-spectrum cephalosporin-resistant *Escherichia coli* from livestock animals in Japan. *Foodborne Pathog. Dis.* 10, 243–249. doi: 10.1089/fpd.2012.1306
- Hiroi, M., Harada, T., Kawamori, F., Takahashi, N., Kanda, T., Sugiyama, K., et al. (2011). A survey of β -lactamase-producing *Escherichia coli* in farm animals and raw retail meat in Shizuoka Prefecture, Japan. *Jpn. J. Infect. Dis.* 64, 153–155.
- Ho, P. L., Li, Z., Lo, W. U., Cheung, Y. Y., Lin, C. H., Sham, P. C., et al. (2012). Identification and characterization of a novel incompatibility group X3 plasmid carrying blaNDM-1 in Enterobacteriaceae isolates with epidemiological links

- to multiple geographical areas in China. *Emerg. Microbes Infect.* 1, e39. doi: 10.1038/emi.2012.37
- Hocquet, D., Plesiat, P., Dehecq, B., Mariotte, P., Talon, D., and Bertrand, X. (2010). Nationwide investigation of extended-spectrum β -lactamases, metallo- β -lactamases, and extended-spectrum oxacillinases produced by ceftazidime-resistant *Pseudomonas aeruginosa* strains in France. *Antimicrob. Agents Chemother.* 54, 3512–3515. doi: 10.1128/AAC.01646-09
- Hordijk, J., Bosman, A. B., van Essen-Zandbergen, A., Veldman, K., Dierikx, C., Wagenaar, J. A., et al. (2011). qnrB19 Gene Bracketed by IS26 on a 40-Kilobase IncR Plasmid from an *Escherichia coli* Isolate from a Veal Calf. *Antimicrob. Agents Chemother.* 55, 453–454. doi: 10.1128/AAC.00866-10
- Huang, Y., Yu, X., Xie, M., Wang, X., Liao, K., Xue, W., et al. (2016). Widespread dissemination of carbapenem-resistant *Escherichia coli* sequence tpe 167 strains harboring blaNDM-5 in clinical settings in China. *Antimicrob. Agents Chemother.* 60, 4364–4368. doi: 10.1128/AAC.00859-16
- Huletsky, A., Knox, J. R., and Levesque, R. C. (1993). Role of Ser-238 and Lys-240 in the hydrolysis of third-generation cephalosporins by SHV-type β -lactamases probed by site-directed mutagenesis and three-dimensional modeling. *J. Biol. Chem.* 268, 3690–3697.
- Jeannot, K., Fournier, D., Müller, E., Cholley, P., and Plesiat, P. (2013). Clonal dissemination of *Pseudomonas aeruginosa* isolates producing extended-spectrum β -lactamase SHV-2a. *J. Clin. Microbiol.* 51, 673–675. doi: 10.1128/JCM.02313-12
- Jelsch, C., Mourey, L., Masson, J. M., and Samama, J. P. (1993). Crystal structure of *Escherichia coli* TEM1 β -lactamase at 1.8Å resolution. *Proteins* 16, 364–383. doi: 10.1002/prot.340160406
- Jiang, H.-X., Tang, D., Liu, Y.-H., Zhang, X.-H., Zeng, Z.-L., Xu, L., et al. (2012). Prevalence and characteristics of β -lactamase and plasmid-mediated quinolone resistance genes in *Escherichia coli* isolated from farmed fish in China. *J. Antimicrob. Chemother.* 67, 2350–2353. doi: 10.1093/jac/dks250
- Johnson, T. J., Wannemuehler, Y. M., Johnson, S. J., Logue, C. M., White, D. G., Doetkott, C., et al. (2007). Plasmid replicon typing of commensal and pathogenic *Escherichia coli* isolates. *Appl. Environ. Microbiol.* 73, 1976–1983. doi: 10.1128/AEM.02171-06
- Jones, C. H., Ruzin, A., Tuckman, M., Visalli, M. A., Petersen, P. J., and Bradford, P. A. (2009). Pyrosequencing using the single-nucleotide polymorphism protocol for rapid determination of TEM- and SHV-type extended-spectrum β -lactamases in clinical isolates and identification of the novel β -lactamase genes blaSHV-48, blaSHV-105, and blaTEM-155. *Antimicrob. Agents Chemother.* 53, 977–986. doi: 10.1128/AAC.01155-08
- Jones-Dias, D., Manageiro, V., and Caniça, M. (2016). Influence of agricultural practice on mobile bla genes: IncII-bearing CTX-M, SHV, CMY and TEM in *Escherichia coli* from intensive farming soils. *Environ. Microbiol.* 18, 260–272. doi: 10.1111/1462-2920.13021
- Jones-Dias, D., Manageiro, V., Martins, A. P., Ferreira, E., and Caniça, M. (2015). New class 2 integron Int2-4 among IncII-positive *Escherichia coli* isolates carrying ESBL and PMA β genes from food animals in Portugal. *Foodborne Pathog. Dis.* 13, 36–39. doi: 10.1089/fpd.2015.1972
- Jouini, A., Slama, K. B., Klibi, N., Sallem, R. B., Estepa, V., Vinué, L., et al. (2013). Lineages and virulence gene content among extended-spectrum β -Lactamase producing *Escherichia coli* strains of food origin in Tunisia. *J. Food Prot.* 76, 323–327. doi: 10.4315/0362-028X.JFP-12-251
- Jung, J. S., Popp, C., Sparbier, K., Lange, C., Kostrzewa, M., and Schubert, S. (2014). Evaluation of matrix-assisted laser desorption ionization–time of flight mass spectrometry for rapid detection of β -lactam resistance in Enterobacteriaceae derived from blood cultures. *J. Clin. Microbiol.* 52, 924–930. doi: 10.1128/JCM.02691-13
- Kameyama, M., Chuma, T., Yabata, J., Tominaga, K., Iwata, H., and Okamoto, K. (2013). Prevalence and epidemiological relationship of CMY-2 AmpC β -lactamase and CTX-M extended-spectrum β -lactamase-producing *Escherichia coli* isolates from broiler farms in Japan. *J. Vet. Med. Sci.* 75, 1009–1015. doi: 10.1292/jvms.12-0453
- Kassis-Chikhani, N., Frangeul, L., Drieux, L., Sengelin, C., Jarlier, V., Brisse, S., et al. (2013). Complete nucleotide sequence of the first KPC-2- and SHV-12-encoding IncX plasmid, pKpS90, from *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 57, 618–620. doi: 10.1128/AAC.01712-12
- Kim, H.-S., Chon, J.-W., Kim, Y.-J., Kim, D.-H., Kim, M.-S., and Seo, K.-H. (2015). Prevalence and characterization of extended-spectrum- β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in ready-to-eat vegetables. *Int. J. Food Microbiol.* 207, 83–86. doi: 10.1016/j.ijfoodmicro.2015.04.049
- Kiratisin, P., Apisarnthanarak, A., Laesripa, C., and Saifon, P. (2008). Molecular characterization and epidemiology of extended-spectrum- β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* isolates causing health care-associated infection in Thailand, where the CTX-M family is endemic. *Antimicrob. Agents Chemother.* 52, 2818–2824. doi: 10.1128/AAC.00171-08
- Kliebe, C., Nies, B. A., Meyer, J. F., Tolxdorff-Neutzling, R. M., and Wiedemann, B. (1985). Evolution of plasmid-coded resistance to broad-spectrum cephalosporins. *Antimicrob. Agents Chemother.* 28, 302–307. doi: 10.1128/AAC.28.2.302
- Knox, J. R. (1995). Extended-spectrum and inhibitor-resistant TEM-type β -lactamases: mutations, specificity, and three-dimensional structure. *Antimicrob. Agents Chemother.* 39, 2593–2601. doi: 10.1128/AAC.39.12.2593
- Kurokawa, H., Yagi, T., Shibata, N., Shibayama, K., Kamachi, K., and Arakawa, Y. (2000). A new SHV-derived extended-spectrum β -Lactamase (SHV-24) that hydrolyzes ceftazidime through a single-amino-acid substitution (D179G) in the Ω -Loop. *Antimicrob. Agents Chemother.* 44, 1725–1727. doi: 10.1128/AAC.44.6.1725-1727.2000
- Kwon, T., Jung, Y. H., Lee, S., Yun, M. R., Kim, W., and Kim, D. W. (2016). Comparative genomic analysis of *Klebsiella pneumoniae* subsp. pneumoniae KP617 and PittNDM01, NUHL24835, and ATCC BAA-2146 reveals unique evolutionary history of this strain. *Gut Pathog.* 8, 34. doi: 10.1186/s13099-016-0117-1
- Lascols, C., Hackel, M., Hujer, A. M., Marshall, S. H., Bouchillon, S. K., Hoban, D. J., et al. (2012). Using nucleic acid microarrays to perform molecular epidemiology and detect novel β -lactamases: a snapshot of extended-spectrum β -lactamases throughout the world. *J. Clin. Microbiol.* 50, 1632–1639. doi: 10.1128/JCM.06115-11
- Lee, K. Y., Hopkins, J. D., and Syvanen, M. (1990). Direct involvement of Is26 in an Antibiotic-Resistance Operon. *J. Bacteriol.* 172, 3229–3236.
- Lee, Y., Kim, B. S., Chun, J., Yong, J. H., Lee, Y. S., Yoo, J. S., et al. (2014). Clonality and resistome analysis of KPC-producing *Klebsiella pneumoniae* strain isolated in Korea using whole genome sequencing. *Biomed. Res. Int.* 2014:352862. doi: 10.1155/2014/352862
- Leinberger, D. M., Grimm, V., Rubtsova, M., Weile, J., Schroppe, K., Wichelhaus, T. A., et al. (2010). Integrated detection of extended-spectrum- β -lactam resistance by DNA microarray-based genotyping of TEM, SHV, and CTX-M genes. *J. Clin. Microbiol.* 48, 460–471. doi: 10.1128/JCM.00765-09
- Ling, B.-D., Liu, G., Xie, Y.-E., Zhou, Q.-X., Zhao, T.-K., Li, C.-Q., et al. (2006). Characterisation of a novel extended-spectrum β -lactamase, SHV-70, from a clinical isolate of *Enterobacter cloacae* in China. *Int. J. Antimicrob. Agents* 27, 355–356. doi: 10.1016/j.ijantimicag.2006.02.003
- Literak, I., Dolejska, M., Janoszowska, D., Hrusakova, J., Meissner, W., Rzycka, H., et al. (2010). Antibiotic resistant *Escherichia coli* bacteria, including strains with genes encoding the extended-spectrum β -lactamase and qnrS, in waterbirds on the Baltic Sea coast of Poland. *Appl. Environ. Microbiol.* 76, 8126–8134. doi: 10.1128/AEM.01446-10
- Liu, G., Ling, B. D., Zeng, Y., Lin, L., Xie, Y. E., and Lei, J. (2008). Molecular characterization of extended-spectrum β -lactamases produced by clinical isolates of *Enterobacter cloacae* from a teaching hospital in China. *Jpn. J. Infect. Dis.* 61, 286–289.
- Liu, Y., Wan, L. G., Deng, Q., Cao, X. W., Yu, Y., and Xu, Q. F. (2015). First description of NDM-1-, KPC-2-, VIM-2- and IMP-4-producing *Klebsiella pneumoniae* strains in a single Chinese teaching hospital. *Epidemiol. Infect.* 143, 376–384. doi: 10.1017/S0950268814000995
- Ma, L., Alba, J., Chang, F.-Y., Ishiguro, M., Yamaguchi, K., Siu, L. K., et al. (2005). Novel SHV-derived extended-spectrum β -lactamase, SHV-57, that confers resistance to ceftazidime but not ceftazolin. *Antimicrob. Agents Chemother.* 49, 600–605. doi: 10.1128/AAC.49.2.600-605.2005
- Machado, E., Coque, T. M., Cantón, R., Novais, A., Sousa, J. C., Baquero, F., et al. (2007). High diversity of extended-spectrum β -lactamases among clinical isolates of Enterobacteriaceae from Portugal. *J. Antimicrob. Chemother.* 60, 1370–1374. doi: 10.1093/jac/dkm381
- Machado, E., Coque, T. M., Cantón, R., Sousa, J. C., and Peixe, L. (2008). Antibiotic resistance integrons and extended-spectrum β -lactamases among Enterobacteriaceae isolates recovered from chickens and swine in Portugal. *J. Antimicrob. Chemother.* 62, 296–302. doi: 10.1093/jac/dkn179

- Mahillon, J., and Chandler, M. (1998). Insertion sequences. *Microbiol. Mol. Biol. Rev.* 62, 725–774.
- Mansour, W., Dahmen, S., Poirer, L., Charfi, K., Bettaieb, D., Boujaafar, N., et al. (2009). Emergence of SHV-2a extended-spectrum β -lactamases in clinical isolates of *Pseudomonas aeruginosa* in a university hospital in Tunisia. *Microb. Drug. Resist.* 15, 295–301. doi: 10.1089/mdr.2009.0012
- Maravic, A., Skocibusic, M., Cvjetan, S., Samanic, I., Fredotovic, Z., and Puizina, J. (2015). Prevalence and diversity of extended-spectrum- β -lactamase-producing Enterobacteriaceae from marine beach waters. *Mar. Pollut. Bull.* 90, 60–67. doi: 10.1016/j.marpolbul.2014.11.021
- Maravić, A., Skočibušić, M., Samanić, I., Fredotović, Z., Cvjetan, S., Jutronic, M., et al. (2013). *Aeromonas* spp. simultaneously harbouring blaCTX-M-15, blaSHV-12, blaPER-1 and blaFOX-2, in wild-growing Mediterranean mussel (*Mytilus galloprovincialis*) from Adriatic Sea, Croatia. *Int. J. Food Microbiol.* 166, 301–308. doi: 10.1016/j.ijfoodmicro.2013.07.010
- Marcadé, G., Deschamps, C., Boyd, A., Gautier, V., Picard, B., Branger, C., et al. (2009). Replicon typing of plasmids in *Escherichia coli* producing extended-spectrum β -lactamases. *J. Antimicrob. Chemother.* 63, 67–71. doi: 10.1093/jac/dkn428
- Markovska, R., Schneider, I., Ivanova, D., Mitov, I., and Bauernfeind, A. (2014). Predominance of IncL/M and IncF plasmid types among CTX-M-ESBL-producing *Escherichia coli* and *Klebsiella pneumoniae* in Bulgarian hospitals. *APMIS* 122, 608–615. doi: 10.1111/apm.12204
- Marti, E., Jofre, J., and Balcazar, J. L. (2013). Prevalence of antibiotic resistance genes and bacterial community composition in a river influenced by a wastewater treatment plant. *PLoS ONE* 8:e78906. doi: 10.1371/journal.pone.0078906
- Matagne, A., Lamotte-Brasseur, J., and Frère, J. M. (1998). Catalytic properties of class A β -lactamases: efficiency and diversity. *Biochem. J.* 330(Pt 2), 581–598.
- Matthew, M., Hedges, R. W., and Smith, J. T. (1979). Types of β -lactamase determined by plasmids in gram-negative bacteria. *J. Bacteriol.* 138, 657–662.
- Mazzariol, A., Roelofsens, E., Koncan, R., Voss, A., and Cornaglia, G. (2007). Detection of a new SHV-type Extended-spectrum β -Lactamase, SHV-31, in a *Klebsiella pneumoniae* strain causing a large nosocomial outbreak in The Netherlands. *Antimicrob. Agents Chemother.* 51, 1082–1084. doi: 10.1128/AAC.00909-06
- Medeiros, A. A. (1997). Evolution and dissemination of β -lactamases accelerated by generations of β -lactam antibiotics. *Clin. Infect. Dis.* 24 (Suppl. 1), S19–S45.
- Mendonça, N., Ferreira, E., and Caniça, M. (2006). Occurrence of a novel SHV-type enzyme (SHV-55) among isolates of *Klebsiella pneumoniae* from Portuguese origin in a comparison study for extended-spectrum β -lactamase-producing evaluation. *Diagn. Microbiol. Infect. Dis.* 56, 415–420. doi: 10.1016/j.diagmicrobio.2006.06.023
- Mendonça, N., Ferreira, E., Louro, D., and Caniça, M. (2009). Molecular epidemiology and antimicrobial susceptibility of extended- and broad-spectrum β -lactamase-producing *Klebsiella pneumoniae* isolated in Portugal. *Int. J. Antimicrob. Agents* 34, 29–37. doi: 10.1016/j.ijantimicag.2008.11.014
- Miriagou, V., Carattoli, A., Tzelepi, E., Villa, L., and Tzouveleki, L. S. (2005). IS26-Associated In4-type integrons forming multiresistance loci in enterobacterial plasmids. *Antimicrob. Agents Chemother.* 49, 3541–3543. doi: 10.1128/AAC.49.8.3541-3543.2005
- Mnif, B., Harhour, H., Jdidi, J., Mahjoubi, F., Genel, N., Arlet, G., et al. (2013). Molecular epidemiology of extended-spectrum β -lactamase-producing *Escherichia coli* in Tunisia and characterization of their virulence factors and plasmid addition systems. *BMC Microbiol.* 13:147. doi: 10.1186/1471-2180-13-147
- Mulvey, M. R., Bryce, E., Boyd, D., Ofner-Agostini, M., Christianson, S., Simor, A. E., et al. (2004). Ambler class A extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella* spp. in Canadian hospitals. *Antimicrob. Agents Chemother.* 48, 1204–1214. doi: 10.1128/AAC.48.4.1204-1214.2004
- Muniesa, M., Colomer-Lluch, M., and Jofre, J. (2013). Potential impact of environmental bacteriophages in spreading antibiotic resistance genes. *Future Microbiol.* 8, 739–751. doi: 10.2217/fmb.13.32
- Muratani, T., Kobayashi, T., and Matsumoto, T. (2006). Emergence and prevalence of β -lactamase-producing *Klebsiella pneumoniae* resistant to cepheims in Japan. *Int. J. Antimicrob. Agents* 27, 491–499. doi: 10.1016/j.ijantimicag.2006.03.007
- M'Zali, F. H., Heritage, J., Gascoyne-Binzi, D. M., Snelling, A. M., and Hawkey, P. M. (1998). PCR single strand conformational polymorphism can be used to detect the gene encoding SHV-7 extended-spectrum β -lactamase and to identify different SHV genes within the same strain. *J. Antimicrob. Chemother.* 41, 123–125. doi: 10.1093/jac/41.1.123
- Naas, T., Namdari, F., Réglier-Poupet, H., Poyart, C., and Nordmann, P. (2007). Panresistant extended-spectrum β -lactamase SHV-5-producing *Acinetobacter baumannii* from New York City. *J. Antimicrob. Chemother.* 60, 1174–1176. doi: 10.1093/jac/dkm366
- Naas, T., Philippon, L., Poirer, L., Ronco, E., and Nordmann, P. (1999). An SHV-derived extended-spectrum β -lactamase in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 43, 1281–1284.
- Naiemi, N. A., Duim, B., Savelkoul, P. H. M., Spanjaard, L., de Jonge, E., Bart, A., et al. (2005). Widespread transfer of resistance genes between bacterial species in an intensive care unit: implications for hospital epidemiology. *J. Clin. Microbiol.* 43, 4862–4864. doi: 10.1128/JCM.43.9.4862-4864.2005
- Naseer, U., Eriksen, B. O., Sundsfjord, A., and Samuelsen, Ø. (2012). Fecal colonization of VIM-1-producing *Klebsiella pneumoniae* and *in vivo* transfer of multidrug-resistant IncN plasmid in a renal transplant patient. *Diagn. Microbiol. Infect. Dis.* 72, 363–366. doi: 10.1016/j.diagmicrobio.2011.12.010
- Neonakis, I. K., Scoulica, E. V., Dimitriou, S. K., Gikas, A. I., and Tselentis, Y. J. (2003). Molecular epidemiology of extended-spectrum β -lactamases produced by clinical isolates in a university hospital in Greece: detection of SHV-5 in *Pseudomonas aeruginosa* and prevalence of SHV-12. *Microb. Drug Resist.* 9, 161–165. doi: 10.1089/107662903765826750
- Nicolas, M. H., Jarlier, V., Honore, N., Philippon, A., and Cole, S. T. (1989). Molecular characterization of the gene encoding SHV-3 β -lactamase responsible for transferable cefotaxime resistance in clinical isolates of *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 33, 2096–2100. doi: 10.1128/AAC.33.12.2096
- Noda, T., Murakami, K., Etoh, Y., Okamoto, F., Yatsuyanagi, J., Sera, N., et al. (2015). Increase in resistance to extended-spectrum cephalosporins in *Salmonella* isolated from retail chicken products in Japan. *PLoS ONE* 10:e0116927. doi: 10.1371/journal.pone.0116927
- Novovic, K., Filipic, B., Veljovic, K., Begovic, J., Mirkovic, N., and Jovic, B. (2015). Environmental waters and blaNDM-1 in Belgrade, Serbia: endemicity questioned. *Sci. Total. Environ.* 511, 393–398. doi: 10.1016/j.scitotenv.2014.12.072
- Nüesch-Inderbinen, M. T., Kayser, F. H., and Hächler, H. (1997). Survey and molecular genetics of SHV β -lactamases in Enterobacteriaceae in Switzerland: two novel enzymes, SHV-11 and SHV-12. *Antimicrob. Agents Chemother.* 41, 943–949.
- Nugent, M. E., and Hedges, R. W. (1979). The nature of genetic determinants for the SHV-1 β -lactamase. *Mol. Gen. Genet.* 175, 239–243. doi: 10.1007/BF00397222
- Nukaga, M., Mayama, K., Hujer, A. M., Bonomo, R. A., and Knox, J. R. (2003). Ultrahigh resolution structure of a class A β -lactamase: on the mechanism and specificity of the extended-spectrum SHV-2 enzyme. *J. Mol. Biol.* 328, 289–301. doi: 10.1016/S0022-2836(03)00210-9
- Oikonomou, O., Liakopoulos, A., Phee, L. M., Betts, J., Mevius, D., and Wareham, D. W. (2016). Providencia stuartii isolates from Greece: co-carriage of cephalosporin (blaSHV-5, blaVEB-1), carbapenem (blaVIM-1), and aminoglycoside (rmtB) resistance determinants by a multidrug-resistant outbreak clone. *Microb. Drug Resist.* 22, 379–386. doi: 10.1089/mdr.2015.0215
- Osinska, A., Harnisz, M., and Korzeniewska, E. (2016). Prevalence of plasmid-mediated multidrug resistance determinants in fluoroquinolone-resistant bacteria isolated from sewage and surface water. *Environ. Sci. Pollut. Res. Int.* 23, 10818–10831. doi: 10.1007/s11356-016-6221-4
- Oviaño, M., Fernández, B., Fernández, A., Barba, M. J., Mouriño, C., and Bou, G. (2014). Rapid detection of Enterobacteriaceae producing extended spectrum β -lactamases directly from positive blood cultures by matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry. *Clin. Microbiol. Infect.* 20, 1146–1157. doi: 10.1111/1469-0691.12729
- Partridge, S. R., Ginn, A. N., Wiklendt, A. M., Ellem, J., Wong, J. S., Ingram, P., et al. (2015). Emergence of blaKPC carbapenemase genes in Australia. *Int. J. Antimicrob. Agents* 45, 130–136. doi: 10.1016/j.ijantimicag.2014.10.006
- Paterson, D. L., and Bonomo, R. A. (2005). Extended-spectrum β -lactamases: a clinical update. *Clin. Microbiol. Rev.* 18, 657–686. doi: 10.1128/CMR.18.4.657-686.2005

- Péduzzi, J., Barthélémy, M., Tiwari, K., Mattioni, D., and Labia, R. (1989). Structural features related to hydrolytic activity against ceftazidime of plasmid-mediated SHV-type CAZ-5 β -lactamase. *Antimicrob. Agents Chemother.* 33, 2160–2163. doi: 10.1128/AAC.33.12.2160
- Perreira Ramos, P. L., Picão, R. C., Almeida, L. G., Lima, N. C., Girardello, R., Vivian, A. C., et al. (2014). Comparative analysis of the complete genome of KPC-2-producing *Klebsiella pneumoniae* Kp13 reveals remarkable genome plasticity and a wide repertoire of virulence and resistance mechanisms. *BMC Genomics* 15:54. doi: 10.1186/1471-2164-15-54
- Philippon, A., Slama, P., Dény, P., and Labia, R. (2016). A structure-based classification of class A β -lactamases, a broadly diverse family of enzymes. *Clin. Microbiol. Rev.* 29, 29–57. doi: 10.1128/CMR.00019-15
- Pinto, L., Radhouani, H., Coelho, C., Martins Da Costa, P., Simões, R., Brandão, R. M. L., et al. (2010). Genetic detection of extended-spectrum β -lactamase-containing *Escherichia coli* isolates from birds of prey from Serra da Estrela Natural Reserve in Portugal. *Appl. Environ. Microbiol.* 76, 4118–4120. doi: 10.1128/AEM.02761-09
- Pitout, J. D. D., Thomson, K. S., Hanson, N. D., Ehrhardt, A. F., Coudron, P., and Sanders, C. C. (1998). Plasmid-mediated resistance to expanded-spectrum cephalosporins among *Enterobacter aerogenes* strains. *Antimicrob. Agents Chemother.* 42, 596–600.
- Pitton, J. (1972). "Mechanisms of bacterial resistance to antibiotics," in *Review of Physiology*, ed R. Adirna (Berlin: Springer-Verlag), 15–93.
- Podbielski, A., Schönling, J., Melzer, B., Warnatz, K., and Leusch, H.-G. (1991). Molecular characterization of a new plasmid-encoded SHV-type β -lactamase (SHV-2 variant) conferring high-level cefotaxime resistance upon *Klebsiella pneumoniae*. *J. Gen. Microbiol.* 137, 569–578. doi: 10.1099/00221287-137-3-569
- Poirel, L., Héritier, C., Podglajen, I., Sougakoff, W., Gutmann, L., and Nordmann, P. (2003). Emergence in *Klebsiella pneumoniae* of a chromosome-encoded SHV β -Lactamase that compromises the efficacy of imipenem. *Antimicrob. Agents Chemother.* 47, 755–758. doi: 10.1128/AAC.47.2.755-758.2003
- Poirel, L., Lebossi, E., Castro, M., Fèvre, C., Foustoukou, M., and Nordmann, P. (2004). Nosocomial outbreak of extended-spectrum β -lactamase SHV-5-producing isolates of *Pseudomonas aeruginosa* in Athens, Greece. *Antimicrob. Agents Chemother.* 48, 2277–2279. doi: 10.1128/AAC.48.6.2277-2279.2004
- Post, V., and Hall, R. M. (2009). AbaR5, a large multiple-antibiotic resistance region found in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* 53, 2667–2671. doi: 10.1128/AAC.01407-08
- Pouget, J. G., Coutinho, F. J., Reid-Smith, R. J., and Boerlin, P. (2013). Characterization of blaSHV genes on plasmids from *Escherichia coli* and *Salmonella enterica* isolates from Canadian food animals (2006–2007). *Appl. Environ. Microbiol.* 79, 3864–3866. doi: 10.1128/AEM.00355-13
- Preston, K. E., Hitchcock, S. A., Aziz, A. Y., and Tine, J. A. (2014). The complete nucleotide sequence of the multi-drug resistance-encoding IncL/M plasmid pACM1. *Plasmid* 76, 54–65. doi: 10.1016/j.plasmid.2014.08.005
- Prinarakis, E. E., Tzelepi, E., Gazouli, M., Mentis, A. F., and Tzouveleki, L. S. (1996). Characterization of a novel SHV α -lactamase variant that resembles the SHV-5 enzyme. *FEMS Microbiol. Lett.* 139, 229–234.
- Ramdani-Bougues, N., Manageiro, V., Jones-Dias, D., Ferreira, E., Tazir, M., and Caniça, M. (2011). Role of SHV β -lactamase variants in resistance of clinical *Klebsiella pneumoniae* strains to β -lactams in an Algerian hospital. *J. Med. Microbiol.* 60, 983–987. doi: 10.1099/jmm.0.030577-0
- Randegger, C. C., and Hächler, H. (2001). Real-time PCR and melting curve analysis for reliable and rapid detection of SHV extended-spectrum β -lactamases. *Antimicrob. Agents Chemother.* 45, 1730–1736. doi: 10.1128/AAC.45.6.1730-1736.2001
- Randegger, C. C., Keller, A., Irla, M., Wada, A., and Hächler, H. (2000). Contribution of natural amino acid substitutions in SHV extended-spectrum β -lactamases to resistance against various β -lactams. *Antimicrob. Agents Chemother.* 44, 2759–2763. doi: 10.1128/AAC.44.10.2759-2763.2000
- Rasheed, J. K., Anderson, G. J., Yigit, H., Queenan, A. M., Doménech-Sánchez, A., Swenson, J. M., et al. (2000). Characterization of the extended-spectrum β -lactamase reference strain, *Klebsiella pneumoniae* K6 (ATCC 700603), which produces the novel enzyme SHV-18. *Antimicrob. Agents Chemother.* 44, 2382–2388. doi: 10.1128/AAC.44.9.2382-2388.2000
- Rasheed, J. K., Jay, C., Metchock, B., Berkowitz, F., Weigel, L., Crellin, J., et al. (1997). Evolution of extended-spectrum β -lactam resistance (SHV-8) in a strain of *Escherichia coli* during multiple episodes of bacteremia. *Antimicrob. Agents Chemother.* 41, 647–653.
- Reuland, E. A., Al Naiemi, N., Raadsen, S. A., Savelkoul, P. H., Kluytmans, J. A., and Vandenbroucke-Grauls, C. M. (2014). Prevalence of ESBL-producing Enterobacteriaceae in raw vegetables. *Eur. J. Clin. Microbiol. Infect. Dis.* 33, 1843–1846. doi: 10.1007/s10096-014-2142-7
- Rocha-Gracia, R. C., Cortés-Cortés, G., Lozano-Zarain, P., Bello, F., Martínez-Laguna, Y., and Torres, C. (2015). Faecal *Escherichia coli* isolates from healthy dogs harbour CTX-M-15 and CMY-2 β -lactamases. *Vet. J.* 203, 315–319. doi: 10.1016/j.tvjl.2014.12.026
- Rodrigues, C., Machado, E., Ramos, H., Peixe, L., and Novais, Â. (2014). Expansion of ESBL-producing *Klebsiella pneumoniae* in hospitalized patients: a successful story of international clones (ST15, ST147, ST336) and epidemic plasmids (IncR, IncFIK). *Int. J. Antimicrob. Agents* 304, 1100–1108. doi: 10.1016/j.ijmm.2014.08.003
- Rubin, J. E., and Pitout, J. D. D. (2014). Extended-spectrum β -lactamase, carbapenemase and AmpC producing Enterobacteriaceae in companion animals. *Vet. Microbiol.* 170, 10–18. doi: 10.1016/j.vetmic.2014.01.017
- Rzewuska, M., Stefanska, I., Kizerwetter-Swida, M., Chrobak-Cmiel, D., Szczygielska, P., Lesniak, M., et al. (2015). Characterization of extended-spectrum β -lactamases produced by *Escherichia coli* strains isolated from dogs in Poland. *Pol. J. Microbiol.* 64, 285–288.
- Samuelsen, Ø., Toleman, M. A., Hasseltvedt, V., Fuursted, K., Leegaard, T. M., Walsh, T. R., et al. (2011). Molecular characterization of VIM-producing *Klebsiella pneumoniae* from Scandinavia reveals genetic relatedness with international clonal complexes encoding transferable multidrug resistance. *Clin. Microbiol. Infect.* 17, 1811–1816. doi: 10.1111/j.1469-0691.2011.03532.x
- Sánchez-Romero, I., Asensio, A., Oteo, J., Muñoz-Algarra, M., Isidoro, B., Vindel, A., et al. (2012). Nosocomial outbreak of VIM-1-producing *Klebsiella pneumoniae* isolates of multilocus sequence type 15: molecular basis, clinical risk factors, and outcome. *Antimicrob. Agents Chemother.* 56, 420–427. doi: 10.1128/AAC.05036-11
- Shahcheraghi, F., Nikbin, V. S., and Feizabadi, M. M. (2009). Prevalence of ESBLs genes among multidrug-resistant isolates of *Pseudomonas aeruginosa* isolated from patients in Tehran. *Microb. Drug Resist.* 15, 37–39. doi: 10.1089/mdr.2009.0880
- Shaheen, B. W., Nayak, R., Foley, S. L., Kweon, O., Deck, J., Park, M., et al. (2011). Molecular characterization of resistance to extended-spectrum cephalosporins in clinical *Escherichia coli* isolates from companion animals in the United States. *Antimicrob. Agents Chemother.* 55, 5666–5675. doi: 10.1128/AAC.00656-11
- Singh, K., Mangold, K. A., Wyant, K., Schora, D. M., Voss, B., Kaul, K. L., et al. (2012). Rectal screening for *Klebsiella pneumoniae* carbapenemases: comparison of real-time PCR and culture using two selective screening agar plates. *J. Clin. Microbiol.* 50, 2596–2600. doi: 10.1128/JCM.00654-12
- Singh, P., Pfeifer, Y., and Mustapha, A. (2016). Multiplex real-time PCR assay for the detection of extended-spectrum β -lactamase and carbapenemase genes using melting curve analysis. *J. Microbiol. Methods* 124, 72–78. doi: 10.1016/j.mimet.2016.03.014
- Song, W., Kim, J., Bae, I. K., Jeong, S. H., Seo, Y. H., Shin, J. H., et al. (2011). Chromosome-encoded AmpC and CTX-M extended-spectrum β -lactamases in clinical isolates of *Proteus mirabilis* from Korea. *Antimicrob. Agents Chemother.* 55, 1414–1419. doi: 10.1128/AAC.01835-09
- Sonnevend, A., Al Baloushi, A., Ghazawi, A., Hashmey, R., Girgis, S., Hamadeh, M. B., et al. (2013). Emergence and spread of NDM-1 producer Enterobacteriaceae with contribution of IncX3 plasmids in the United Arab Emirates. *J. Med. Microbiol.* 62, 1044–1050. doi: 10.1099/jmm.0.059014-0
- Sowek, J. A., Singer, S. B., Ohringer, S., Malley, M. F., Dougherty, T. J., Gougoutas, J. Z., et al. (1991). Substitution of lysine at position 104 or 240 of TEM-1pTZ18R β -lactamase enhances the effect of serine-164 substitution on hydrolysis or affinity for cephalosporins and the monobactam aztreonam. *Biochemistry* 30, 3179–3188. doi: 10.1021/bi00227a004
- Stolle, I., Prenger-Berninghoff, E., Stamm, I., Scheufen, S., Hassdenteufel, E., Guenther, S., et al. (2013). Emergence of OXA-48 carbapenemase-producing *Escherichia coli* and *Klebsiella pneumoniae* in dogs. *J. Antimicrob. Chemother.* 68, 2802–2808. doi: 10.1093/jac/dkt259
- Stuart, J. C., Voets, G., Scharringa, J., Fluit, A. C., and Leverstein-Van Hall, M. A. (2012). Detection of carbapenemase-producing Enterobacteriaceae with a commercial DNA microarray. *J. Med. Microbiol.* 61, 809–812. doi: 10.1099/jmm.0.041673-0

- Stürenburg, E., Sobottka, I., Feucht, H.-H., Mack, D., and Laufs, R. (2003). Comparison of BDPhoenix and VITEK2 automated antimicrobial susceptibility test systems for extended-spectrum β -lactamase detection in *Escherichia coli* and *Klebsiella* species clinical isolates. *Diagn. Microbiol. Infect. Dis.* 45, 29–34. doi: 10.1016/S0732-8893(02)00481-9
- Szabó, D., Melan, M. A., Hujer, A. M., Bonomo, R. A., Hujer, K. M., Bethel, C. R., et al. (2005). Molecular analysis of the simultaneous production of two SHV-type extended-spectrum β -lactamases in a clinical isolate of *Enterobacter cloacae* by using single-nucleotide polymorphism genotyping. *Antimicrob. Agents Chemother.* 49, 4716–4720. doi: 10.1128/AAC.49.11.4716-4720.2005
- Tamura, K., Stecher, G., Peterson, D., Filipiński, A., and Kumar, S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729. doi: 10.1093/molbev/mst197
- Tang, J., Bu, Y., Zhang, X. X., Huang, K., He, X., Ye, L., et al. (2016). Metagenomic analysis of bacterial community composition and antibiotic resistance genes in a wastewater treatment plant and its receiving surface water. *Ecotoxicol. Environ. Saf.* 132, 260–269. doi: 10.1016/j.ecoenv.2016.06.016
- Tato, M., Coque, T. M., Ruiz-Garbajosa, P., Pintado, V., Cobo, J., Sader, H. S., et al. (2007). Complex clonal and plasmid epidemiology in the first outbreak of Enterobacteriaceae infection involving VIM-1 metallo- β -Lactamase in Spain: toward endemicity? *Clin. Infect. Dis.* 45, 1171–1178. doi: 10.1086/522288
- Tian, G.-B., Wang, H.-N., Zhang, A.-Y., Zhang, Y., Fan, W.-Q., Xu, C.-W., et al. (2012). Detection of clinically important β -lactamases in commensal *Escherichia coli* of human and swine origin in western China. *J. Med. Microbiol.* 61, 233–238. doi: 10.1099/jmm.0.036806-0
- Tian, Z., Zhang, Y., Yu, B., and Yang, M. (2016). Changes of resistome, mobilome and potential hosts of antibiotic resistance genes during the transformation of anaerobic digestion from mesophilic to thermophilic. *Water Res.* 98, 261–269. doi: 10.1016/j.watres.2016.04.031
- Timofte, D., Maciucă, I. E., Evans, N. J., Williams, H., Wattret, A., Fick, J. C., et al. (2014). Detection and molecular characterization of *Escherichia coli* CTX-M-15 and *Klebsiella pneumoniae* SHV-12 β -lactamases from bovine mastitis isolates in the United Kingdom. *Antimicrob. Agents Chemother.* 58, 789–794. doi: 10.1128/AAC.00752-13
- Tissera, S., and Lee, S. M. (2013). Isolation of Extended Spectrum β -lactamase (ESBL) producing bacteria from urban surface waters in Malaysia. *Malays. J. Med. Sci.* 20, 14–22.
- Tzouveleki, L., and Bonomo, R. (1999). SHV-type β -lactamases. *Curr. Pharm. Des.* 5, 847–864.
- Uemura, S., Yokota, S., Mizuno, H., Sakawaki, E., Sawamoto, K., Maekawa, K., et al. (2010). Acquisition of a transposon encoding extended-spectrum β -lactamase SHV-12 by *Pseudomonas aeruginosa* isolates during the clinical course of a burn patient. *Antimicrob. Agents Chemother.* 54, 3956–3959. doi: 10.1128/AAC.00110-10
- Veldman, K., Dierikx, C., Van Essen-Zandbergen, A., Van Pelt, W., and Mevius, D. (2010). Characterization of multidrug-resistant, *qnrB2*-positive and extended-spectrum- β -lactamase-producing *Salmonella* Concord and *Salmonella* Senftenberg isolates. *J. Antimicrob. Chemother.* 65, 872–875. doi: 10.1093/jac/dkq049
- Veldman, K., van Tulden, P., Kant, A., Testerink, J., and Mevius, D. (2013). Characteristics of cefotaxime-resistant *Escherichia coli* from wild birds in The Netherlands. *Appl. Environ. Microbiol.* 79, 7556–7561. doi: 10.1128/AEM.01880-13
- Villa, L., Pezzella, C., Tosini, F., Visca, P., Petrucca, A., and Carattoli, A. (2000). Multiple-antibiotic resistance mediated by structurally related InC1/M plasmids carrying an extended-spectrum β -lactamase gene and a class 1 integron. *Antimicrob. Agents Chemother.* 44, 2911–2914. doi: 10.1128/AAC.44.10.2911-2914.2000
- Vinué, L., Lantero, M., Sáenz, Y., Somalo, S., de Diego, I., Pérez, F., et al. (2008). Characterization of extended-spectrum β -lactamases and integrons in *Escherichia coli* isolates in a Spanish hospital. *J. Med. Microbiol.* 57, 916–920. doi: 10.1099/jmm.0.047723-0
- von Wintersdorff, C. J. H., Penders, J., van Niekerk, J. M., Mills, N. D., Majumder, S., van Alphen, L. B., et al. (2016). Dissemination of antimicrobial resistance in microbial ecosystems through horizontal gene transfer. *Front. Microbiol.* 7:173. doi: 10.3389/fmicb.2016.00173
- Walther-Rasmussen, J., and Hoiby, N. (2007). Class A carbapenemases. *J. Antimicrob. Chemother.* 60, 470–482. doi: 10.1093/jac/dkm226
- Wang, X. J., Xu, X. L., Li, Z. W., Chen, H. B., Wang, Q., Yang, P. H., et al. (2014). An outbreak of a nosocomial NDM-1-producing *Klebsiella pneumoniae* ST147 at a teaching hospital in mainland China. *Microb. Drug Resist.* 20, 144–149. doi: 10.1089/mdr.2013.0100
- Wang, X. R., Chen, J. C., Kang, Y., Jiang, N., An, S. C., and Gao, Z. C. (2012). Prevalence and characterization of plasmid-mediated *bla*ESBL with their genetic environment in *Escherichia coli* and *Klebsiella pneumoniae* in patients with pneumonia. *Chin. Med. J. (Engl.)* 125, 894–900.
- Whichard, J., Gay, K., Stevenson, J., Joyce, K., Cooper, K., Omondi, M., et al. (2007). Human *Salmonella* and concurrent decreased susceptibility to quinolones and extended-spectrum cephalosporins. *Emerg. Infect. Dis.* 13, 1681. doi: 10.3201/eid1311.061438
- Wilke, M. S., Lovering, A. L., and Strynadka, N. C. (2005). β -lactam antibiotic resistance: a current structural perspective. *Curr. Opin. Microbiol.* 8, 525–533. doi: 10.1016/j.mib.2005.08.016
- Winkler, M. L., and Bonomo, R. A. (2016). SHV-129: a gateway to global suppressors in the SHV β -lactamase family? *Mol. Biol. Evol.* 33, 429–441. doi: 10.1093/molbev/msv235
- Wolny-Koladka, K., and Lenart-Boron, A. (2016). Phenotypic and molecular assessment of drug resistance profile and genetic diversity of waterborne. *Water Air Soil Pollut.* 227, 146. doi: 10.1007/s11270-016-2833-z
- Yi, T., Kim, T. G., and Cho, K. S. (2015). Fate and behavior of extended-spectrum β -lactamase-producing genes in municipal sewage treatment plants. *J. Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.* 50, 1160–1168. doi: 10.1080/10934529.2015.1047673
- Yigit, H., Queenan, A. M., Rasheed, J. K., Biddle, J. W., Domenech-Sanchez, A., Alberti, S., et al. (2003). Carbapenem-resistant strain of *Klebsiella oxytoca* harboring carbapenem-hydrolyzing β -lactamase KPC-2. *Antimicrob. Agents Chemother.* 47, 3881–3889. doi: 10.1128/AAC.47.12.3881-3889.2003
- Yuan, M., Hall, L. M. C., Savelkoul, P. H. M., Vandenbroucke-Grauls, C. M. J. E., and Livermore, D. M. (2000). SHV-13, a novel extended-spectrum β -lactamase, in *Klebsiella pneumoniae* isolates from patients in an intensive care unit in Amsterdam. *Antimicrob. Agents Chemother.* 44, 1081–1084. doi: 10.1128/AAC.44.4.1081-1084.2000
- Zhang, H., Zhou, Y., Guo, S., and Chang, W. (2015). Multidrug resistance found in extended-spectrum β -lactamase-producing Enterobacteriaceae from rural water reservoirs in Guantao, China. *Front. Microbiol.* 6:267. doi: 10.3389/fmicb.2015.00267
- Zienkiewicz, M., Kern-Zdanowicz, I., Carattoli, A., Gniadkowski, M., and Ceglowski, P. (2013). Tandem multiplication of the IS26-flanked amplicon with the *bla*SHV-5 gene within plasmid p1658/97. *FEMS Microbiol. Lett.* 341, 27–36. doi: 10.1111/1574-6968.12084
- Zou, L.-K., Wang, H.-N., Zeng, B., Zhang, A.-Y., Li, J.-N., Li, X.-T., et al. (2011). Phenotypic and genotypic characterization of β -lactam resistance in *Klebsiella pneumoniae* isolated from swine. *Vet. Microbiol.* 149, 139–146. doi: 10.1016/j.vetmic.2010.09.030
- Zuo, B., Liu, Z., Wang, H., Yang, Y., Chen, J., and Ye, H. (2006). Genotype of TEM- and SHV-type β -lactamase producing *Klebsiella pneumoniae* in Guangzhou area [Article in Chinese]. *Zhonghua Yi Xue Za Zhi* 86, 2928–2932.
- Zurfluh, K., Hächler, H., Nüesch-Inderbinen, M., and Stephan, R. (2013). Characteristics of extended-spectrum β -lactamase- and carbapenemase-producing Enterobacteriaceae isolates from rivers and lakes in Switzerland. *Appl. Environ. Microbiol.* 79, 3021–3026. doi: 10.1128/AEM.00054-13
- Zurfluh, K., Nüesch-Inderbinen, M., Morach, M., Zihler Berner, A., Hächler, H., and Stephan, R. (2015). Extended-spectrum- β -lactamase-producing Enterobacteriaceae isolated from vegetables imported from the Dominican Republic, India, Thailand, and Vietnam. *Appl. Environ. Microbiol.* 81, 3115–3120. doi: 10.1128/AEM.00258-15

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Liakopoulos, Mevius and Ceccarelli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.