



Genomic Analysis of *Vulcanisaeta thermophila* Type Strain CBA1501^T Isolated from Solfataric Soil

Joon Yong Kim, Kyung June Yim, Hye Seon Song, Yeon Bee Kim, Dong-Gi Lee, Joseph Kwon, Kyung-Seo Oh and Seong Woon Roh *

Biological Disaster Analysis Group, Korea Basic Science Institute, Daejeon, South Korea

Keywords: *Vulcanisaeta thermophila*, genome sequence, archaea, hyperthermophile, hyperthermophilic enzyme

INTRODUCTION

OPEN ACCESS

Edited by:

Frank T. Robb,
University of Maryland, Blatimore,
USA

Reviewed by:

David L. Bernick,
University of California, Santa Cruz,
USA
Nikolai Ravin,
Research Center for Biotechnology
Russian Academy of Sciences, Russia
Lydia Kreuter,
University of Maryland, Baltimore,
USA

*Correspondence:

Seong Woon Roh
seong18@gmail.com

Specialty section:

This article was submitted to
Evolutionary and Genomic
Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 16 August 2016

Accepted: 30 September 2016

Published: 19 October 2016

Citation:

Kim JY, Yim KJ, Song HS, Kim YB,
Lee D-G, Kwon J, Oh K-S and
Roh SW (2016) Genomic Analysis of
Vulcanisaeta thermophila Type Strain
CBA1501^T Isolated from Solfataric
Soil. *Front. Microbiol.* 7:1639.
doi: 10.3389/fmicb.2016.01639

Hyperthermophilic archaea have been isolated from high-temperature environments such as geothermally heated soils, sulfur-rich hot springs, and submarine volcanic habitats; optimal growth of these organisms occurs above 80°C (Stetter, 1999, 2006, 2013). The genus *Vulcanisaeta* belongs to the family *Thermoproteaceae*, order *Thermoproteales*, phylum *Crenarchaeota*, and was first proposed by Itoh et al. (2002). It currently includes 3 validly named species, that is, *Vulcanisaeta distributa* (Itoh et al., 2002), *V. souniana* (Itoh et al., 2002), and *V. thermophila* (Yim et al., 2015), as per the List of Prokaryotic Names with Standing in Nomenclature database (Parte, 2014). Members of the genus *Vulcanisaeta* are rod-shaped, anaerobic, hyperthermophilic, and acidophilic (Itoh et al., 2002). To date, 15 genomes, including two complete genomes, *V. distributa* and “*Vulcanisaeta moutnovskia*” (Mavromatis et al., 2010; Gumerov et al., 2011), have been reported for the genus *Vulcanisaeta*, as per the NCBI genome database (<http://www.ncbi.nlm.nih.gov/genome/>).

Hyperthermophilic enzymes are stable and active at high temperatures of >70°C (Vieille et al., 1996). These enzymes can be studied using model systems to elucidate enzyme mechanisms and evolution of proteins stable at high temperatures and to determine the higher temperature limit for enzyme stability (Vieille and Zeikus, 2001). In a previous study, *V. thermophila* CBA1501^T (= ATCC BAA-2415^T = JCM 17228^T) was isolated from solfataric soil in the Republic of the Philippines (Yim et al., 2015). It was found to grow at 75–90°C, pH 4.0–6.0, and 0–1.0% (w/v) NaCl, with optimal growth at 85°C, pH 5.0, and 0% (w/v) NaCl. Here, a genome sequence of *V. thermophila* CBA1501^T has been reported and information of hyperthermophilic enzymes of high biotechnological value has been provided.

MATERIALS AND METHODS

Culture Conditions and DNA Extraction

In a previous study, we isolated *V. thermophila* CBA1501^T from the solfataric soil of the Mayon volcano in the Republic of the Philippines (Yim et al., 2015) and cultivated it on modified JCM medium no. 236 (M236) (containing per liter salt base solution: 2.94 g trisodium citrate dihydrate, 0.5 g yeast extract, 10.0 ml trace vitamins, 1.0 mg resazurin, 0.5 g Na₂S·9H₂O, and 20 mM thiosulfate). For DNA extraction, the strain was enriched at 80°C in M236 medium, using a serum bottle. Its genomic DNA was extracted using the G-spin total DNA extraction kit (iNtRON Biotechnology, Korea) and QuickGene DNA tissue kit S (Kurabo, Japan).

Genome Sequencing, Assembly, and Annotation

The genome sequences of *V. thermophila* CBA1501^T were sequenced at a read length of 300 bp using the Illumina MiSeq system, with paired-end library [insert size, 634–1101 bp (average 852 bp),

computed by CLC Genomics Workbench 7.5.1 (CLC bio, Denmark)] constructed using the Nextera DNA Library Prep kit (illumina, USA), according to the manufacturer's instructions (Moon et al., 2015). A total of 6,939,438 reads (with 688-fold coverage) were assembled using CLC Genomics Workbench 7.5.1 with default parameters as follows: masking mode, no masking; mismatch cost, 2; insertion cost, 3; deletion cost, 3; length cost, 3; length fraction, 0.5; similarity fraction, 0.8; global alignment, no; auto-detect paired distances, yes; non-specific match handling map, randomly. To identify ribosomal RNA and transfer RNA, RNAMmer 1.2 (Lagesen et al., 2007) and tRNAscan v. 1.3.1 (Lowe and Eddy, 1997), respectively, were used. Protein coding sequences (CDSs) identification was performed using PRODIGAL v. 2.6.2 (Hyatt et al., 2012), and functional annotation was performed using EggNOG v. 4.1 (Powell et al., 2014), SEED subsystems (Overbeek et al., 2014), Swiss-Prot (UniProt, 2015), and KEGG (Kanehisa et al., 2016) databases with the USEARCH v. 8.0.1517 program (Edgar, 2010).

Phylogenetic Analysis

Similarities based on 16S rRNA gene sequences were calculated using EzBioCloud (<http://www.ezbiocloud.net>). Phylogenetic tree based on 16S rRNA gene sequences was constructed using MEGA5 (Tamura et al., 2011) with the neighbor-joining (Saitou and Nei, 1987), maximum-parsimony (Kluge and Farris, 1969), and maximum-likelihood (Felsenstein, 1981) methods, based on 1000 randomly generated trees.

Comparative Genomic Analysis

For comparative analysis, reference genome sequences of closely related strains of the genus *Vulcanisaeta* were selected using the NCBI genome database (<http://www.ncbi.nlm.nih.gov/genome/>): *V. distributa* JCM 11215 (BBCT00000000), *V. distributa* DSM 14429 (CP002100), *V. distributa* JCM 11217 (BBBJ00000000), *V. souniana* JCM 11219 (BBBK00000000), *V. moutnovskia* 768-28 (CP002529), and *Vulcanisaeta* sp. strains CIS_19 (LOC000000000), JCM 14467 (BBDM000000000), JCM 16159 (BBDN000000000), and JCM 161 (BBDO000000000). To determine the similarity between genome sequences, orthologous average nucleotide identity (OrthoANI) values of CBA1501^T and related strains in the genus *Vulcanisaeta* were calculated using the Orthologous Average Nucleotide Identity Tool (Lee et al., 2015), and a phylogenetic tree based on OrthoANI values was obtained using the EzBioCloud Comparative Genomics Database (EzCgDb; Chunlab; <http://cg.ezbiocloud.net/>). Annotated genomes of CBA1501^T and other related strains were subjected to homology search using the UBLAST program (Ward and Moreno-Hagelsieb, 2014) for pan-genome analysis. Then, pan-genome orthologous groups (POGs) were constructed using EzCgDb.

RESULTS

General Genomic Features of *V. thermophila* CBA1501^T

The draft genome sequence of *V. thermophila* CBA1501^T was 2,022,594 bp in length, with a G+C content of 49.1 mol %

in 10 contigs. The largest contig was 791,731 bp long, and the N50 value was 634,758 bp. The genome was found to contain 2170 CDSs, one 16S-23S-5S rRNA gene operon, and 41 tRNA genes. Genomic features are shown in **Figure 1**. On the basis of information from the EggNOG v. 4.1 database, 1927 genes were categorized into Clusters of Orthologous Groups of proteins (COGs) functional groups. The most abundant COG category was "Function unknown" (S; 729 genes), followed by "Energy production and conversion" (C; 178 genes), "Amino acid transport and metabolism" (E; 168 genes), "Translation, ribosomal structure and biogenesis" (J; 159 genes), "Carbohydrate transport and metabolism" (G; 96 genes), and "Coenzyme transport and metabolism" (H; 87 genes). Among the SEED subsystem categories, "Carbohydrates" (181 genes), "Amino Acids and Derivatives" (171 genes), "Protein Metabolism" (142 genes) and "Cofactors, Vitamins, Prosthetic Groups, Pigments" (116 genes) were the most dominant categories (>10% of a total of 1,094 matched SEED subsystem categories).


Phylogenetic Analysis

V. thermophila CBA1501^T had the greatest 16S rRNA gene sequence similarity with the following (in this order): *V. distributa* DSM 14429^T (98.6%), *Stygiolobus azoricus* DSM 6296^T (98.6%), *V. souniana* IC-059^T (97.5%), *Caldivirga maquilingsensis* IC-167^T (94.6%), *Pyrobaculum ferrireducens* 1860^T (93.8%), *Pyrobaculum islandicum* DSM 4184^T (93.6%), *Thermoproteus uzoniensis* 768-20 (93.6%), *Pyrobaculum organotrophum* JCM 9190^T (93.5%), and *Thermoproteus thermophilus* CBA1502^T (93.2%). The phylogenetic analysis indicated that the strain CBA1501^T clustered with species of the genus *Vulcanisaeta* (Supplementary Figure 1A).

Comparative Genomics Data

V. thermophila CBA1501^T had lesser than 73% orthoANI values with all of the related strains in the genus *Vulcanisaeta* (Supplementary Table 1). In the orthoANI values-based dendrogram, the strain CBA1501^T was located as an outgroup to the other related strains in *Vulcanisaeta* (Supplementary Figure 1B). These results indicate that *V. thermophila* CBA1501^T is evolutionarily distinct from other related strains. The pan-genome analysis showed that 10 genomes in the genus *Vulcanisaeta* have the core genome, comprised of 979 POGs. In contrast, only the genome of strain CBA1501^T had 211 POGs as a singleton. Among these singletons, various enzymes, including arylformamidase, shikimate kinase, formyl-CoA transferase, xanthine dehydrogenase, hydrogensulfite reductase, and amidase, were detected.

In conclusion, the information provided here is useful as the genome of *V. thermophila* CBA1501^T will provide insights into the metabolism of hyperthermophilic archaea and aid in identifying opportunities for biotechnological applications of novel hyperthermophilic enzymes.



DATA ACCESS

The genome sequences of *V. thermophila* CBA1501^T (=ATCC BAA-2415^T =JCM 17228^T) were deposited in the DDBJ under the accession numbers BCL101000001-BCL101000010 (<http://www.ncbi.nlm.nih.gov/Traces/wgs/BCL101>). The annotated data of *V. thermophila* CBA1501^T based on SEED subsystems is accessible on SEED viewer v. 2.0 by logging in with the guest account (Genome ID 6666666.192913, username: guest, password: guest) at the web address: <http://rast.nmpdr.org/seedviewer.cgi?page=Organism&organism=6666666.192913>.

AUTHOR CONTRIBUTIONS

SWR designed and coordinated all the experiments. KJY performed cultivation, DNA extraction and purification. JYK, HSS, YBK, D-GL, JK, and K-SO performed the sequencing, genome assembly, gene prediction, gene annotation and

comparative genomic analysis. JYK, KJY, and SWR wrote manuscript. All authors have read the manuscript and approved.

FUNDING

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science, and Technology (2015R1D1A1A09061039), project fund from the Center for Analytical Research of Disaster Science of the Korea Basic Science Institute (C36703).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.01639>

REFERENCES

- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26, 2460–2461. doi: 10.1093/bioinformatics/btq461
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J. Mol. Evol.* 17, 368–376.
- Gumerov, V. M., Mardanov, A. V., Beletsky, A. V., Prokofeva, M. I., Bonch-Osmolovskaya, E. A., Ravin, N. V., et al. (2011). Complete genome sequence of “*Vulcanisaeta moutnovskia*” strain 768-28, a novel member of the hyperthermophilic crenarchaeal genus *Vulcanisaeta*. *J. Bacteriol.* 193, 2355–2356. doi: 10.1128/JB.00237-11
- Hyatt, D., LoCascio, P. F., Hauser, L. J., and Uberbacher, E. C. (2012). Gene and translation initiation site prediction in metagenomic sequences. *Bioinformatics* 28, 2223–2230. doi: 10.1093/bioinformatics/bts429
- Itoh, T., Suzuki, K., and Nakase, T. (2002). *Vulcanisaeta distributa* gen. nov., sp. nov., and *Vulcanisaeta souiniana* sp. nov., novel hyperthermophilic, rod-shaped crenarchaeotes isolated from hot springs in Japan. *Int. J. Syst. Evol. Microbiol.* 52(Pt 4), 1097–1104. doi: 10.1099/00207713-52-4-1097
- Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2016). KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res.* 44, D457–D462. doi: 10.1093/nar/gkv1070
- Kluge, A. G., and Farris, F. S. (1969). Quantitative phyletics and the evolution of anurans. *Syst. Zool.* 18, 1–32.
- Lagesen, K., Hallin, P., Rødland, E. A., Staerfeldt, H. H., Rognes, T., and Ussery, D. W. (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35, 3100–3108. doi: 10.1093/nar/gkm160
- Lee, I., Kim, Y. O., Park, S. C., and Chun, J. (2015). OrthoANI: An improved algorithm and software for calculating average nucleotide identity. *Int. J. Syst. Evol. Microbiol.* 66, 1100–1103. doi: 10.1099/ijsem.0.000760
- Lowe, T. M., and Eddy, S. R. (1997). tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25, 955–964.
- Mavromatis, K., Sikorski, J., Pabst, E., Teshima, H., Lapidus, A., Lucas, S., et al. (2010). Complete genome sequence of *Vulcanisaeta distributa* type strain (IC-017). *Stand. Genomic Sci.* 3, 117–125. doi: 10.4056/sigs.1113067
- Moon, J. S., Choi, H. S., Shin, S. Y., Noh, S. J., Jeon, C. O., and Han, N. S. (2015). Genome sequence analysis of potential probiotic strain *Leuconostoc lactis* EFEL005 isolated from kimchi. *J. Microbiol.* 53, 337–342. doi: 10.1007/s12275-015-5090-8
- Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T., et al. (2014). The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* 42, D206–D214. doi: 10.1093/nar/gkt1226
- Parte, A. C. (2014). LPSN—list of prokaryotic names with standing in nomenclature. *Nucleic Acids Res.* 42, D613–D616. doi: 10.1093/nar/gkt1111
- Powell, S., Forslund, K., Szklarczyk, D., Trachana, K., Roth, A., Huerta-Cepas, J., et al. (2014). eggNOG v4.0: nested orthology inference across 3686 organisms. *Nucleic Acids Res.* 42, D231–D239. doi: 10.1093/nar/gkt1253
- Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406–425.
- Stetter, K. O. (1999). Extremophiles and their adaptation to hot environments. *FEBS Lett.* 452, 22–25.
- Stetter, K. O. (2006). History of discovery of the first hyperthermophiles. *Extremophiles* 10, 357–362. doi: 10.1007/s00792-006-0012-7
- Stetter, K. O. (2013). A brief history of the discovery of hyperthermophilic life. *Biochem. Soc. Trans.* 41, 416–420. doi: 10.1042/BST20120284
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739. doi: 10.1093/molbev/msr121
- UniProt, C. (2015). UniProt: a hub for protein information. *Nucleic Acids Res.* 43, D204–D212. doi: 10.1093/nar/gku989
- Vieille, C., Burdette, D. S., and Zeikus, J. G. (1996). Thermozyms. *Biotechnol. Annu. Rev.* 2, 1–83.
- Vieille, C. and Zeikus, G. J. (2001). Hyperthermophilic enzymes: sources, uses, and molecular mechanisms for thermostability. *Microbiol. Mol. Biol. Rev.* 65, 1–43. doi: 10.1128/MMBR.65.1.1-43.2001
- Ward, N., and Moreno-Hagelsieb, G. (2014). Quickly finding orthologs as reciprocal best hits with BLAT, LAST, and UBLAST: how much do we miss? *PLoS ONE* 9:e101850. doi: 10.1371/journal.pone.0101850
- Yim, K. J., Cha, I. T., Rhee, J. K., Song, H. S., Hyun, D. W., Lee, H. W., et al. (2015). *Vulcanisaeta thermophila* sp. nov., a hyperthermophilic and acidophilic crenarchaeon isolated from solfataric soil. *Int. J. Syst. Evol. Microbiol.* 65(Pt 1), 201–205. doi: 10.1099/ijms.0.065862-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Kim, Yim, Song, Kim, Lee, Kwon, Oh and Roh. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.