



A Bioengineered Nisin Derivative, M21A, in Combination with Food Grade Additives Eradicates Biofilms of *Listeria monocytogenes*

Muireann K. Smith^{1†}, Lorraine A. Draper^{1†}, Pieter-Jan Hazelhoff², Paul D. Cotter^{1,3}, R. P. Ross¹ and Colin Hill^{1*}

¹ APC Microbiome Institute, School of Microbiology, University College Cork, Cork, Ireland, ² Wageningen University & Research, Wageningen, Netherlands, ³ Teagasc Food Research Centre, Cork, Ireland

OPEN ACCESS

Edited by:

Michael Gänzle,
University of Alberta, Canada

Reviewed by:

Christian U. Riedel,
University of Ulm, Germany
Michael Chikindas,
Rutgers University, USA

*Correspondence:

Colin Hill
c.hill@ucc.ie

[†] These authors share first authorship.

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 13 October 2016

Accepted: 18 November 2016

Published: 30 November 2016

Citation:

Smith MK, Draper LA, Hazelhoff P-J, Cotter PD, Ross RP and Hill C (2016) A Bioengineered Nisin Derivative, M21A, in Combination with Food Grade Additives Eradicates Biofilms of *Listeria monocytogenes*. *Front. Microbiol.* 7:1939. doi: 10.3389/fmicb.2016.01939

The burden of foodborne disease has large economic and social consequences worldwide. Despite strict regulations, a number of pathogens persist within the food environment, which is greatly contributed to by a build-up of resistance mechanisms and also through the formation of biofilms. Biofilms have been shown to be highly resistant to a number of antimicrobials and can be extremely difficult to remove once they are established. In parallel, the growing concern of consumers regarding the use of chemically derived antimicrobials within food has led to a drive toward more natural products. As a consequence, the use of naturally derived antimicrobials has become of particular interest. In this study we investigated the efficacy of nisin A and its bioengineered derivative M21A in combination with food grade additives to treat biofilms of a representative foodborne disease isolate of *Listeria monocytogenes*. Investigations revealed the enhanced antimicrobial effects, in liquid culture, of M21A in combination with citric acid or cinnamaldehyde over its wild type nisin A counterpart. Subsequently, an investigation was conducted into the effects of these combinations on an established biofilm of the same strain. Nisin M21A (0.1 µg/ml) alone or in combination with cinnamaldehyde (35 µg/ml) or citric acid (175 µg/ml) performed significantly better than combinations involving nisin A. All combinations of M21A with either citric acid or cinnamaldehyde eradicated the *L. monocytogenes* biofilm (in relation to a non-biofilm control). We conclude that M21A in combination with available food additives could further enhance the antimicrobial treatment of biofilms within the food industry, simply by substituting nisin A with M21A in current commercial products such as Nisaplin® (Danisco, DuPont).

Keywords: *Listeria monocytogenes*, biofilm, nisin, bioengineered, citric acid, cinnamaldehyde, combinations

INTRODUCTION

The globalization of food distribution and human travel (Tauxe, 2002; Hussain and Dawson, 2013) has meant that foodborne disease can spread further and more rapidly than ever before, bypassing conventional control measures (Koopmans et al., 2003; World Health Organization and Foodborne Disease Burden Epidemiology Reference Group, 2007). The ability of bacterial pathogens to remain viable in food production environments, even with the enforcement of stringent hygiene practices, can be in part due to the accumulation of resistance mechanisms (Holah et al., 2002; Tauxe, 2002)

and through the protection provided via the formation of biofilms (Borucki et al., 2003; Ryu and Beuchat, 2005; Houdt and Michiels, 2010). Surface-attached biofilms are sessile microbial communities that can be irreversibly attached to a substratum and are enclosed in an extracellular polymeric matrix (Donlan, 2002; Olszewska, 2013; Colagiorgi et al., 2016). Once formed, biofilms can be extremely difficult to remove, leading to contamination of processing equipment (Kumar and Anand, 1998; Tan et al., 2014), food spoilage and consequent economic losses (Brooks and Flint, 2008; Houdt and Michiels, 2010). Numerous studies have described the use of novel approaches to control biofilms; these have included the use of biocides (Wood et al., 1996), application of electrical currents (Poortinga et al., 2001), temperature step changes during growth regions (Knight et al., 2004), enzymes (Liu et al., 2014) and for biofilms on or within a food matrix, the use of novel food additives (Furukawa et al., 2010).

The use of bio-preservatives, as an alternative to chemical food preservatives, has become more popular due to customer demand for more natural and minimally processed foods (Lucera et al., 2012). One such group of anti-bacterial preservatives are bacteriocins. Bacteriocins are small, ribosomally synthesized antimicrobials (Cleveland et al., 2001; Cotter et al., 2005; Gálvez et al., 2007; Yang et al., 2014). Those derived from lactic acid bacteria (LAB) have gained particular attention with respect to their use in the food industry. One of the most extensively studied bacteriocins is nisin A, a bactericidal lantibiotic produced by some strains of *Lactococcus lactis* (Rouse et al., 2012; Champion et al., 2013; Healy et al., 2013). It is a 3.5 kDa peptide consisting of 34 amino acids and is a flexible and elongated amphipathic peptide that has the ability to form pores as well as inhibit cell wall biosynthesis (Breukink et al., 1999; Wiedemann et al., 2001; Prudêncio et al., 2015). This dual action of nisin has meant that despite widespread use in the food industry, there have been few reports of resistance development outside of the laboratory (Field et al., 2010b; Molloy et al., 2012). Due to its proven range of antimicrobial effects, nisin has been used as a food preservative for many years (Breukink and de Kruijff, 1999; Elliason and Tatini, 1999; Gálvez et al., 2007). Nisaplin® is an example of a commercially available food additive, containing 2.5% nisin. It is Generally Regarded As Safe (GRAS) for several applications and has been approved for use in the EU as preservative E234 by Directive 95/2/EC (European Food Safety Authority, 2006) by the U.S. Food and Drug Administration (FDA) (Prudêncio et al., 2015) and by the WHO (Sobrino-López and Martín-Belloso, 2008).

Other examples of food additives are organic acids such as lactic acid and citric acid (Young and Foegeding, 1993; Helander and Mattila-Sandholm, 2000; Norhana et al., 2012), salts of organic acids such as trisodium phosphate, potassium sorbate, sodium benzoate and sodium diacetate (Phillips and Duggan, 2001; Norhana et al., 2012) and proteins such as lactoferrin (Branen and Davidson, 2004; Gálvez et al., 2007) and nisin. Enhanced antimicrobial activity has been successfully demonstrated against Gram-positive pathogens when several of these antimicrobials were used in combination

with nisin (Phillips, 1999; McEntire et al., 2003; Branen and Davidson, 2004; Bari et al., 2005; Murdock et al., 2007; Norhana et al., 2012). Plant essential oils are another group of additives that have gained increased attention due to a long history of use for their antimicrobial effects and the increased demand for more “natural” food products (Helander et al., 1998; Burt, 2004; Bakkali et al., 2007; Prudêncio et al., 2015). Essential oils are natural, volatile liquids obtained from plant material (Gálvez et al., 2007) that have aromatic qualities (Burt, 2004). A number of studies have also been carried out that demonstrate the enhanced antimicrobial activity of nisin in combination with essential oils such as cinnamaldehyde (Field et al., 2015a) and carvacrol (Pol and Smid, 1999) against a broad range of pathogens including Gram-positive organisms, such as *Listeria monocytogenes* and *Bacillus cereus* (Pol and Smid, 1999; Field et al., 2015a), and Gram-negative bacteria, including *Salmonella enterica* and *Escherichia coli* (Phillips and Duggan, 2001; Govaris et al., 2010; Norhana et al., 2012; Prudêncio et al., 2015).

The use of nisin in combination with other food preservatives provides a hurdle approach to controlling pathogens while reducing/removing the need for chemically synthesized preservatives (Chalón et al., 2012). These benefits can be further augmented through the use of bioengineered forms of nisin. The gene encoded and ribosomally synthesized nature of nisin has meant that nisin is a perfect candidate for bioengineering, in order to enhance the functionality of the peptide (Field et al., 2008; Molloy et al., 2012, 2013). Indeed, although there are numerous studies that demonstrate the enhanced inhibitory activity of nisin when used in combination with antimicrobial food additives, there is little information pertaining to the use of bioengineered nisin in combination with food additives to inhibit pathogens within the food industry. Here, we demonstrate that the use of a bioengineered derivative of nisin in combination with a series of natural food additives results in enhanced antimicrobial effects against *L. monocytogenes*, including relatively antimicrobial resistant biofilms of this problematic food borne pathogen.

MATERIALS AND METHODS

Growth of Bacterial Strains

Listeria monocytogenes F6854 strain, linked to a single case of human illness traced to contaminated turkey frankfurters in 1988 (Nelson et al., 2004), was grown in Tryptic Soy Broth (TSB) (Merck, Germany) or Tryptic Soy Agar (TSB substituted with 1.5% w/v agar) at 37°C. *L. lactis* strains, *L. lactis* NZ9700, *L. lactis* NZ9800-pCI372 Nisin A, *L. lactis* NZ9800-pCI372 M21A (Field et al., 2008), *L. lactis* NZ9800-pCI372 AAA (Healy et al., 2013) and *L. lactis* NZ9800-pCI372 Nisin V (Rouse et al., 2012), were grown on M17 supplemented with 5% glucose (GM17) broth (Oxoid, England) or GM17 agar (GM17 broth substituted with 1.5% w/v agar) at 30°C. Strains were stocked in 40% glycerol and stored at -20°C.

Deferred Antagonism Agar Diffusion Assay

Agar diffusion assays were performed in triplicate as previously described by Healy et al. (2013). Briefly, *L. lactis* strains producing nisin A, M21A, AAA and Nisin V were spotted onto a GM17 agar plate and allowed to grow overnight. The plate was then subjected to UV radiation for 30 min and then overlaid with GM17 soft agar (0.75% w/v agar) seeded with 2% overnight of *L. monocytogenes* F6854 and allowed to grow overnight at 37°C. Antimicrobial efficiency was determined through measurement of the zone of inhibition using digital calipers.

Nisin Peptide Purification

Purification of nisin peptide was performed as described previously by Campion et al. (2013). Briefly, TY broth (tryptone, yeast extract, $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, MgSO_4) was passed through a column (Length 70 cm, diameter 5 cm) one third packed with Amberlite XAD-16 beads (Sigma). Resulting broth was dispensed in 900 ml volumes and autoclaved. Overnight cultures of nisin producing strains *L. lactis* NZ9800 – pCI372 M21A (for M21A production) or *L. lactis* NZ9700 (for nisin A) were used to inoculate previously filtered TY broth (2 × 900 ml) supplemented with 20% glucose and 20% β -glycerophosphate. After overnight incubation at 30°C, the culture was centrifuged at 8500 g at 4°C for 20 min (Thermo Scientific, Sorvall™ RC 6+ Centrifuge, USA). Culture supernatant was passed dropwise through a column containing 60 g Amberlite XAD-16 beads and discarded. Following a 500 ml wash of 30% ethanol the nisin was eluted from the column using 500 ml of 70% isopropanol (Fisher Scientific, UK) supplemented with 0.1% trifluoroacetic acid (TFA). Concurrently bacterial cell pellets were resuspended in 300 ml 70% isopropanol – 0.1% TFA. Following stirring at room temperature for 3 h the suspension was centrifuged at 8500 g at 4°C for 15 min and supernatant was retained. The column eluant was pooled with the post-centrifugation supernatant and concentrated to a volume of 300 ml using rotary evaporation (BÜCHI Rotavapor R-205, Switzerland). The pH of the sample was adjusted to pH 4.0 and it was subsequently passed through a 60 ml Strata C-18 E column (Phenomenex) that was previously washed with 60 ml methanol (Fisher Scientific, UK) and 60 ml H_2O . After applying 120 ml 30% ethanol, nisin was eluted from the column using 60 ml 70% isopropanol – 0.1% TFA and stored at 4°C. For HPLC purification 10 ml volumes were concentrated to a volume of ~2 ml by rotary evaporation and injected into a HPLC system (Shimadzu) equipped with a Phenomenex C12 Reverse-Phase (RP) column (Jupiter 4 μ proteo 90 Å, 250 mm × 10.0 mm, 4 μ m). Nisin was eluted via a gradient of 30–50% acetonitrile (Fisher) containing 0.1% TFA that was developed over 5–40 min. Nisin containing fractions were pooled, acetonitrile was removed by rotary evaporation and the peptide was lyophilised and stored at –20°C.

Minimum Inhibitory Concentration Determination

Minimum inhibition concentrations (MIC) of peptides and antimicrobials were determined using 96 well microtitre plates

following CLSI (Clinical and Laboratory Standards Institute) guidelines as previously described by Draper et al. (2013) with some modifications. Briefly, each well within the microtitre plate was treated with phosphate buffered saline (PBS) containing 1% (w/v) bovine serum albumin (PBS/BSA) (200 μ l) and incubated for 30 min at 37°C. Following this, the wells were washed with PBS (200 μ l) and allowed to dry. Overnight cultures of *L. monocytogenes* F6854 were then sub-cultured into fresh broth and grown to an OD_{600} of 0.5 [$\sim 10^8$ colony forming units (cfu)/ml] and subsequently diluted to a final concentration of $\sim 10^5$ cfu/ml. Stock concentrations of nisin A (100 μ g/ml), nisin M21A (128 μ g/ml), Nisaplin® (25 mg/ml), and citric acid (150 μ g/ml) were prepared in TSB, while 99%-cinnamaldehyde (10 mg/ml) was prepared in a 50% ethanol solution. Each antimicrobial was subjected to twofold serial dilutions in TSB in the PBS/BSA treated plate and the appropriately diluted F6854 strain was subsequently added (100 μ l). 96 well microtitre plates were incubated overnight at 37°C. Each MIC was performed in triplicate. The MIC was determined as the lowest concentration at which there was no visible growth.

Growth Curves

Growth curves were performed in a checkerboard assay manner in 96 well microtitre plates. A twofold serial dilution of either nisin A or nisin M21A were performed along the abscissa excluding the final row. In a separate 96 well microtitre plate, serial twofold dilutions of the antimicrobials sodium acetate, sodium benzoate (both food grade, Sigma), citric acid, potassium sorbate, carvacrol, lactoferrin and 99% *trans*-cinnamaldehyde (hereafter referred to as cinnamaldehyde) (all research grade, Sigma) and Nisaplin® (Danisco, DuPont Nutrition and Health) were performed separately along the ordinate excluding column 1. These dilutions were directly transferred into the corresponding wells on the original plate. 100 ml of a 2% inoculum from an overnight strain was added to each well on the plate. Cell growth was measured spectrophotometrically over a 24 h period using MWGt Sirius HT plate reader (BioTek Instruments, USA).

Biofilm Growth

Biofilm growth was performed as previously described by Field et al. (2015a), but with modifications as follows. TSB supplemented with 1% D-(+)-glucose (Sigma) (TSBg) was used in order to aid biofilm formation (Choi et al., 2013). 200 μ l of TSBg containing a final concentration of 2% inoculum from an overnight strain was added to test wells of a sterile PBS/BSA treated 96 well microtitre plate (Starstedt, Germany). Additionally, 200 μ l of TSBg was added to control wells as a negative control. Plates were incubated at 37°C for 48 h to allow for biofilm attachment and formation.

Established Biofilm Treatment

Biofilm inhibition assays were carried out as previously described by Field et al. (2015b). Here, nisin A, nisin variant M21A, Nisaplin®, citric acid and 99% cinnamaldehyde were used to treat *L. monocytogenes* F6854 biofilms. By combining a variety of concentrations of the aforementioned antimicrobials,

the minimum amounts needed in combination for effective treatment of an F6854 biofilm was established as follows; 100 μ l of the appropriate concentration for each peptide and/or antimicrobial in TSB were added to the biofilm, both alone and in combination. TSB alone was added to a set of wells as a negative control. A number of biofilm containing wells (labeled untreated) and wells containing TSB alone (labeled control) were left untreated and the plate was incubated at 37°C for 24 h. In order to determine the metabolic activity of the biofilm after antimicrobial treatment an XTT assay was performed as previously described by Field et al. (2015b). The principle of the XTT assay is that viable cells will reduce the tetrazolium salt XTT [2,3-bis(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5 carboxanilide] (Sigma) to an orange-colored water-soluble product measurable via a standard microplate absorbance reader at OD_{492 nm} (Tunney et al., 2004). Briefly, all test and control wells were washed with 200 μ l sterile PBS. 100 μ l of a solution containing 500 mg/l XTT sodium salt and 10 μ M menadione (Sigma) was added to each well. The plates were incubated in darkness at 37°C for 2 h. Absorbance was read at 492 nm.

Statistical Analysis

All aforementioned growth curve experiments were conducted in triplicate. Six replicates of each established biofilm treatment experiment were performed. Statistical analysis was performed using GraphPad Prism 6, GraphPad Software, Inc. Unless otherwise stated a one-way ANOVA was performed followed by Tukey test, where *P*-values are denoted as: **P* > 0.05; ***P* ≤ 0.01; ****P* ≤ 0.001.

RESULTS

Deferred Antagonism Agar Diffusion Assay

In order to evaluate the antimicrobial activity of nisin A and its derivatives against *L. monocytogenes* F6854, a selection of nisin variant producers were assessed via deferred antagonism agar diffusion assay. Of the variant strains tested the observed zone sizes for three derivative producers, AAA (where the three hinge amino acids in positions 20–22 have been replaced with alanines), M21A and M21V were consistently enhanced when compared to the corresponding isogenic nisin A producer (Figure 1). To date a number of studies have been conducted that have highlighted the efficacy of M21V against strains of *L. monocytogenes* (Field et al., 2008, 2015a; Champion et al., 2013). Here, we noted that the antimicrobial efficiency of the M21A producer appeared equivalent to that of M21V, while they both had a greater degree of activity against the indicator strain than that of the producer of the AAA derivative. As a result of these observations the purified nisin M21A peptide was chosen for further investigation via specific activity studies/growth curve experiments.

Minimum Inhibitory Concentrations

The activities of purified nisin A, nisin M21V and M21A were assessed using broth based MIC assays (Table 1). The

MIC recorded for nisin A against *L. monocytogenes* F6854 was 12.5 μ g/ml, a result that is consistent with previously published results for *L. monocytogenes* EGDe::pPL2luxpHELP (Champion et al., 2013). The MIC for both nisin M21V and M21A was 6.25 μ g/ml, i.e., half that observed for nisin A. Indeed, it has previously been established that nisin M21V has twofold greater activity than nisin A against a number of *L. monocytogenes* (Field et al., 2010a; Molloy et al., 2012). In addition to nisin and its variants, the antimicrobial activity of citric acid, sodium acetate, sodium benzoate, potassium sorbate, carvacrol, lactoferrin, cinnamaldehyde and Nisaplin® against F6854 were also assessed (Table 1).

Dual Antimicrobial Inhibition of *L. monocytogenes* F6854

Preliminary checkerboard assays combining nisin A and M21A with each of the other, aforementioned, antimicrobials were performed in order to determine if any of the combinations revealed enhanced antimicrobial activity. While many combinations showed marginally enhanced effects (data not shown), the combination of nisin A, M21A and either citric acid or cinnamaldehyde showed the most dramatic results. Therefore, further triplicate growth curve experiments were conducted using both nisin A and M21A in combination with either citric acid or cinnamaldehyde against *L. monocytogenes* F6854 over a 24 h period (Figures 2 and 3). At 0.78 μ g/ml, neither nisin peptide inhibited the pathogen. Citric acid, at 1170 μ g/ml, extended the lag phase by approximately 1 h and reduced the final OD. Combining these concentrations of nisin A and citric acid extended the lag phase by 14 h and an OD_{600 nm} of 0.3 was only reached after 22 h. When citric acid (1170 μ g/ml) was combined with M21A (0.78 μ g/ml), complete inhibition of growth was observed (Figure 2). An enhanced specific activity of M21A in comparison to nisin A was also noted during combinatorial studies with cinnamaldehyde against *L. monocytogenes* F6854 (Figure 3). 117.2 μ g/ml of cinnamaldehyde alone delayed the lag phase by approximately 2 h but, when combined with M21A (0.78 μ g/ml) the lag phase was extended to 11 h, approximately 4 h longer than that observed for nisin A and cinnamaldehyde.

Biofilm Treatment

These combinations were examined to determine if they also demonstrated enhanced efficacy against established F6854 biofilms (Figures 4 and 5). It was observed that for biofilm treatment the ratio (μ g/ml: μ g/ml) of cinnamaldehyde or citric acid to nisin A or M21A was in some case 150 times the ratio used for planktonic growth inhibition (Table 2). Cinnamaldehyde (35 μ g/ml) and M21A (0.1 μ g/ml) showed significantly enhanced antimicrobial effects against the established biofilm compared to the corresponding combinations of cinnamaldehyde and nisin A. Biofilm treatment with citric acid (175 μ g/ml) and M21A (0.1 μ g/ml) in combination reduced the viable cell numbers significantly compared to the antimicrobial/peptide alone (Figure 5). Most striking was the significant fivefold difference observed between the nisin A-citric acid combination




TABLE 1 | Minimum inhibitory concentrations of nisin peptides and antimicrobials required to treat *L. monocytogenes* F6854.

Antimicrobial	MIC ($\mu\text{g/ml}$)
Nisin A	12.5
Nisin V	6.25
M21A	6.25
Citric Acid	2340
99% cinnamaldehyde	625
Lactoferrin	>2500
Sodium acetate	250,000
Sodium benzoate	25,000
Carvacrol	310
Potassium sorbate	37,500
Nisaplin®	393

and the dual effect of M21A and citric acid. Notably, when used at these concentrations the impact of the nisin derivative M21A in combination with either citric acid or cinnamaldehyde was statistically indistinguishable from the non-biofilm control, indicating complete eradication of the biofilm.

Inhibitory Effect of Nisaplin® in Combination with Antimicrobials on F6854 Biofilm

It was decided to investigate if inhibitory effects were achievable using citric acid and cinnamaldehyde in combination with a readily available nisin A containing food additive Nisaplin®. With the enhanced inhibitory effects of M21A over nisin A evident in the previous combination studies, a greater amount of nisin A and hence Nisaplin in combination would be required to cause eradication of an established biofilm. This would suggest that production of a derivative of Nisaplin®-equivalent containing M21A would enhance the antimicrobial potency of these additives. Since Nisaplin® contains only 2.5% nisin it was of interest to confirm that the remainder of the ingredients within this additive (or a potential M21A containing product) would not inhibit the observed synergy with citric acid and cinnamaldehyde. As seen in **Figure 6** a presumptive synergistic effect was observed between Nisaplin® and cinnamaldehyde against an established

F6854 biofilm (**Figure 6**). Neither Nisaplin® [30 $\mu\text{g/ml}$ (nisin A content: 0.75 $\mu\text{g/ml}$)] nor cinnamaldehyde (15 $\mu\text{g/ml}$) alone had a statistically significant effect on the biofilm in relation to the untreated control. However, the dual effect of both antimicrobials at the same concentrations showed a statistically significant decrease in the biofilm. A similar result was observed when Nisaplin® (30 $\mu\text{g/ml}$ (nisin A content: 0.75 $\mu\text{g/ml}$)) and citric acid (125 $\mu\text{g/ml}$) were combined (**Figure 7**). Neither Nisaplin® nor citric acid alone had a statistically significant effect on the biofilm. However, the combination of both antimicrobials at the same concentrations showed a statistically significant ($P \leq 0.001$) decrease in biofilm viability. When we consider the nisin A content of the Nisaplin® used (30 $\mu\text{g/ml}$) in these combinations it corresponds to 0.75 $\mu\text{g/ml}$ (2.5%). This is >7 times the amount of purified M21A peptide utilized in the corresponding combinations (**Figures 4** and **5**) to achieve biofilm eradication. It must be noted however that the levels of citric acid and cinnamaldehyde utilized were slightly decreased (1.4 and 2.3 times respectively) in comparison to the aforementioned M21A containing biofilm treatments.

DISCUSSION

Listeriosis, the disease associated with the ubiquitous food borne pathogen *L. monocytogenes*, has a low incidence rate (Maertens de Noordhout et al., 2014), but an extremely high mortality rate (20–30%) (Laksanalamai et al., 2012) and is rigorously monitored within the food and beverage industry. Microbial standards within the EU state that <100 cfu/g *L. monocytogenes* is permitted in Ready-To-Eat (RTE) foods that do not have the potential to support listerial growth throughout their shelf life. However, in infant foods, foods for medical purposes and RTE foods that do support listerial growth, the micro-organism must be absent in any 25 g sample at the time of production (The Commission of the European Communities, 2005). The US maintain a zero tolerance policy on *L. monocytogenes* in any RTE product (And and Hoover, 2003). Due to these highly strict regulations, discovering novel and safe means to control of *L. monocytogenes* is of significant importance and benefit to food manufacturers.




FIGURE 2 | Growth curve depicting the effect of both M21A and Nisin A (Nis A) alone and in combination with citric acid (CA) against *L. monocytogenes* F6854. Growth was measured via triplicate readings at OD_{600 nm} over a period of 22 h.





FIGURE 3 | Growth curve depicting the effect of both M21A and Nisin A (Nis A) alone and in combination with cinnamaldehyde (CN) against *L. monocytogenes* F6854. Growth was measured at OD_{600 nm} over a period of 24 h. Results were recorded in triplicate.

The main focus of this study was to determine if an enhanced effect could be obtained by combining a derivative of nisin with other food-grade antimicrobials to inhibit the growth of *L. monocytogenes* F6854 in liquid culture and, in addition, to eradicate established biofilms of this same strain. By including bioengineered versions of nisin in this study we sought to and achieved increased antimicrobial potency. Due to the documented sensitivity of *Listeria* to nisin and the relevance of both to the food industry (Thomas and Wimpenny, 1996; Arqués et al., 2008; Pinto et al., 2010; Field et al., 2015a), strain F6854, which has been linked to contaminated turkey frankfurters, represented an ideal target for nisin and food-grade antimicrobial combinations. The food additives sodium benzoate, sodium acetate, potassium sorbate, lactoferrin, carvacrol, citric acid and cinnamaldehyde were all

tested initially. We noted that the amount of sodium acetate required for inhibition was extremely high as compared to the other antimicrobials in this study. The MIC obtained for both sodium benzoate and potassium sorbate are higher than that previously reported for other *L. monocytogenes* strains (Neetoo et al., 2008). In addition, the concentration of cinnamaldehyde required to inhibit the indicator strain was found to be 625 $\mu\text{g/ml}$. This is approximately 50% greater than the MIC previously published for cinnamaldehyde against a number of different *L. monocytogenes* strains. The MIC for carvacrol was, however, in keeping with results targeting two strains from this same study (Field et al., 2015a). Following initial broth-based growth curves the most notable effects were observed however when citric acid or cinnamaldehyde were used in combination with nisin.



cell function and disruption of the cell membrane (Gill and Holley, 2004; Hyldgaard et al., 2012). Furthermore, a recent study by Field et al. (2015a) also demonstrated that cinnamaldehyde in combination with a nisin variant was more effective than the essential oil or variant alone against a strain of *L. monocytogenes* in planktonic form.

Following the selection of two antimicrobial compounds, cinnamaldehyde and citric acid, we investigated the consequences of their use in combination with nisin A or a bioengineered derivative thereof. From a number of nisin A variants tested, both M21A and M21V demonstrated equally enhanced activity compared to the wild type in both MIC and agar diffusion assays. Nisin A derivatives M21A and M21V both contain a single amino acid changes at position 21. In the case of M21A, a methionine is replaced by an alanine and in M21V, the same methionine is replaced by a valine. M21V has previously demonstrated enhanced activity against strains of *L. monocytogenes* and has been documented in numerous studies (Field et al., 2010b, 2015a; Campion et al., 2013; Healy et al., 2013) and so it was included in MIC studies in order to compare its activity with that of M21A, which until now remained relatively undocumented. This is the first study in which the MIC of the bioengineered derivative M21A has been tested against a strain of *L. monocytogenes*. Having established that the antimicrobial efficacy of M21A against *L. monocytogenes* F6854 was twofold greater than that of nisin A, we went further to demonstrate that M21A possesses enhanced ability when compared to the wild type with respect to eradicating formed *L. monocytogenes* biofilms, a characteristic not attributed to any other bioengineered nisin derivative to date.




Initially inhibition of bacterial growth utilizing the chosen food-grade additives (citric acid and cinnamaldehyde) and lantibiotic peptides nisin A and M21A were investigated using broth based growth curve assays. These studies demonstrated that complete inhibition of this strain of F6854 could be obtained using M21A in combination with citric acid, while nisin A at the same concentrations was unable to achieve such potency. Although a less dramatic result was obtained from the combination of M21A and cinnamaldehyde, M21A exhibited a greater inhibitory effect against *L. monocytogenes* F6854 both alone and in combination with cinnamaldehyde. Hence, the anti-listerial properties of M21A can be further enhanced with the addition of citric acid or cinnamaldehyde. This led to investigation to determine if these enhanced effects could also inhibit established biofilms of the same strain.

Citric acid has a long history of use in food and has GRAS status (Lemmerer and Bernstein, 2010). It is used for its antimicrobial properties, which includes its ability to function as a chelator (Brul and Coote, 1999) and to pass freely through the cell membrane altering the cytoplasmic pH (Cotter and Hill, 2003). It is often used in combination with other preservatives and/or antimicrobials to inhibit pathogens in food. Indeed, the use of citric acid has proven effective in combination with nisin against Gram-negative bacteria (Phillips, 1999) and in combination with nisin against a strain of *Listeria innocua* (Al-Holy et al., 2006). Less is known about the effect of citric acid on *L. monocytogenes*. Cinnamaldehyde has previously been shown to have an antimicrobial effect on strains of *L. monocytogenes* (Gill and Holley, 2004). Although not fully elucidated, cinnamaldehyde is believed to function through inhibition of glucose uptake, inhibition of enzymes involved in

TABLE 2 | Ratio of antimicrobials used in combination to treat *L. monocytogenes* F6854 in both planktonic culture and as established biofilms.

Antimicrobial combinations	Planktonic culture		Established biofilm	
	(µg/ml: µg/ml)	Ratio	(µg/ml: µg/ml)	Ratio
Nisin A: citric acid	0.78:1170	1:150	0.1:175	1:1750
Nisin A: cinnamaldehyde	0.78:117.2	1:150*	0.1:35	1:350
M21A: citric acid	0.78:1170	1:150	0.1:175	1:1750
M21A: cinnamaldehyde	0.78:117.2	1:150*	0.1:35	1:350

These ratios are based on the minimum concentration (µg/ml) of each antimicrobial required in combination to inhibit F6854. *Complete inhibition of *L. monocytogenes* was not achieved in this case.



biofilms and more recently the use of a nisin derivative has been investigated in relation to the inhibition of a *Staphylococcus* biofilm (Field et al., 2015b). However, the current study is the first to our knowledge to show the antimicrobial effects of a bioengineered version of nisin (both alone and in combination) on *L. monocytogenes* biofilms. The demonstrated enhanced antimicrobial effect of M21A in combination with food grade preservatives, citric acid or cinnamaldehyde, resulted in a reduction in viable cells that could not be statistically distinguished from a non-biofilm control.

When we examined the concentrations of antimicrobials required to obtain inhibition we observed the same trend for planktonic and biofilm experiments, with a low level of nisin and a greater amount of citric acid/cinnamaldehyde required. When we examine the ratio of antimicrobial concentrations used for biofilm inhibition the level of cinnamaldehyde and citric acid required to treat the biofilms is up to 150 times that required for planktonic cells. This confirms that determining the ideal proportion of each antimicrobial is vital for optimal synergistic effect and is dependent on the physiological state of bacteria whether it be planktonic or existing as a biofilm. It is also possible that the structural properties of the extracellular polymeric matrix or subtle differences in the behavior of cells within the biofilm occur in comparison to their planktonic counterparts. In fact, it has been shown that lower levels of Nisaplin® are required to inhibit *L. monocytogenes* within a food matrix when compared to broth based assays (Hara et al., 2009).

Currently, nisin A-containing products such as Nisaplin® are commercially available to be used as food preservatives. Here,

Nisin has been used in previous studies with a view to inhibiting bacterial biofilms (Mahdavi et al., 2007; Gou et al., 2011). In addition, nisin in combination with antimicrobials has also been investigated in relation to the inhibition of bacterial

we show that Nisaplin®-citric acid/cinnamaldehyde combinations could significantly reduce and in fact eradicate *Listeria* biofilms. However, >7 times more of the antimicrobial component (i.e., nisin A) was required in comparison to the M21A purified peptide counterpart. Nonetheless due to market availability of Nisaplin® and these food additives, such combinations are readily applicable and may even present a lower cost alternative to the control of *L. monocytogenes* within the food industry. Although, nisin sensitivity to *L. monocytogenes* has been documented, the development of nisin resistant mutants has been reported (Gandhi and Chikindas, 2006). Therefore, the combinations within this study represent an example of hurdle technology whereby bacterial growth struggles to overcome the potency of dual acting antimicrobials. In addition, by combining these antimicrobials it allows for the use of lower concentrations of the individual additives in food items. This may prove beneficial in relation to the use of the essential oil cinnamaldehyde – the use of which has been limited due to the amount needed for an antimicrobial effect affecting the overall organoleptic properties of the food itself. However, sensory evaluation of a chosen food containing this level of the essential oil or citric acid should be conducted in order to distinguish if there are any negative effects to the organoleptic properties of the food.

Ultimately, this study has identified a clear application for M21A in combination with either citric acid or cinnamaldehyde

to inhibit listerial growth, eradicate formed biofilms and hence potentially inhibit biofilm formation. This effect that significantly exceeded the effect of nisin A in the corresponding combination. This study documents the first report of a bioengineered derivative of nisin used with other food additives to control *Listeria* biofilms. We propose that substituting M21A into nisin A-containing products, such as Nisaplin®, would further enhance the antimicrobial potency of such products. Ultimately, such combinations should enhance the microbial safety of foods as well as provide consumers with an effective natural alternative to artificial preservatives.

AUTHOR CONTRIBUTIONS

Conceived and supervised the study: LD, PC, CH, and RR. Designed the experiments: MS, LD, and P-JH. Performed the experiments: MS and P-JH. Analyzed the data: MS, LD, P-JH, PC, RR, and CH. Wrote the paper: MS, LD, P-JH, PC, RR, and CH.

FUNDING

This publication has emanated from research conducted with the financial support of Science Foundation Ireland (SFI) under Grant Number SFI/12/RC/2273.

REFERENCES

- Al-Holy, M., Al-Qadiri, H., Lin, M., and Rasco, B. (2006). Inhibition of *Listeria innocua* in hummus by a combination of nisin and citric acid. *J. Food Prot.* 69, 1322–1327.
- And, C., and Hoover, D. G. (2003). Bacteriocins and their food applications. *Compr. Rev. Food Sci. Food Saf.* 2, 82–100. doi: 10.1111/j.1541-4337.2003.tb00016.x
- Arqués, J. L., Rodríguez, E., Nuñez, M., and Medina, M. (2008). Antimicrobial activity of nisin, reuterin, and the lactoperoxidase system on *Listeria monocytogenes* and *Staphylococcus aureus* in cuajada, a semisolid dairy product manufactured in Spain. *J. Dairy Sci.* 91, 70–75. doi: 10.3168/jds.2007-0133
- Bakkali, F., Averbeck, S., Averbeck, D., and Idaomar, M. (2007). Biological effects of essential oils – A review. *Food Chem. Toxicol.* 46, 446–475. doi: 10.1016/j.fct.2007.09.106
- Bari, M., Ukuku, D., Kawasaki, T., Inatsu, Y., Ishiki, K., and Kawamoto, S. (2005). Combined efficacy of nisin and pediocin with sodium lactate, citric acid, phytic acid, and potassium sorbate and EDTA in reducing the *Listeria monocytogenes* population of inoculated fresh-cut produce. *J. Food Prot.* 68, 1381–1387.
- Borucki, M. K., Peppin, J. D., White, D., Loge, F., and Call, D. R. (2003). Variation in biofilm formation among strains of *Listeria monocytogenes*. *Appl. Environ. Microbiol.* 69, 7336–7342. doi: 10.1128/AEM.69.12.7336-7342.2003
- Branen, J. K., and Davidson, P. M. (2004). Enhancement of nisin, lysozyme, and monolaurin antimicrobial activities by ethylenediaminetetraacetic acid and lactoferrin. *Int. J. Food Microbiol.* 90, 63–74. doi: 10.1016/S0168-1605(03)00172-7
- Breukink, E., and de Kruijff, B. (1999). The lantibiotic nisin, a special case or not? *Biochim. Biophys. Acta* 1462, 223–234. doi: 10.1016/S0005-2736(99)00208-4
- Breukink, E., Wiedemann, I., van Kraaij, C., Kuipers, O. P., Sahl, H. G., and de Kruijff, B. (1999). Use of the cell wall precursor lipid II by a pore-forming peptide antibiotic. *Science* 286, 2361–2364. doi: 10.1126/science.286.5448.2361
- Brooks, J., and Flint, S. (2008). Biofilms in the food industry: problems and potential solutions. *Int. J. Food Sci. Technol.* 43, 2163–2176. doi: 10.1111/j.1365-2621.2008.01839.x
- Brul, S., and Coote, P. (1999). Preservative agents in foods Mode of action and microbial resistance mechanisms. *Int. J. Food Microbiol.* 50, 1–17. doi: 10.1016/S0168-1605(99)00072-0
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods—a review. *Int. J. Food Microbiol.* 94, 223–253. doi: 10.1016/j.ijfoodmicro.2004.03.022
- Campion, A., Casey, P., Field, D., Cotter, P., Hill, C., and Ross, P. (2013). In vivo activity of Nisin A and Nisin V against *Listeria monocytogenes* in mice. *BMC Microbiol.* 13:23. doi: 10.1186/1471-2180-13-23
- Chalón, M., Acuña, L., Morero, R., Minahk, C., and Bellomio, A. (2012). Membrane-active bacteriocins to control *Salmonella* in foods. Are they the definite hurdle? *Food Res. Int.* 45, 735–744. doi: 10.1016/j.foodres.2011.08.024
- Choi, N.-Y., Kim, B.-R., Bae, Y.-M., and Lee, S.-Y. (2013). Biofilm formation, attachment, and cell hydrophobicity of foodborne pathogens under varied environmental conditions. *J. Korean Soc. Appl. Biol. Chem.* 56, 207–220. doi: 10.1007/s13765-012-3253-4
- Cleveland, J., Montville, T., Nes, I., and Chikindas, M. (2001). Bacteriocins: safe, natural antimicrobials for food preservation. *Int. J. Food Microbiol.* 71, 1–20. doi: 10.1016/S0168-1605(01)00560-8
- Colagiorgi, A., Ciccio, P., Zanardi, E., Ghidini, S., and Ianieri, A. (2016). A Look inside the *Listeria monocytogenes* biofilms extracellular matrix. *Microorganisms* 4:22. doi: 10.3390/microorganisms4030022
- Cotter, P., and Hill, C. (2003). Surviving the acid test: responses of gram-positive bacteria to low pH. *Microbiol. Mol. Biol. Rev.* 67, 429–453. doi: 10.1128/MMBR.67.3.429-453.2003
- Cotter, P., Hill, C., and Ross, P. (2005). Bacteriocins: developing innate immunity for food. *Nat. Rev. Microbiol.* 3, 777–788. doi: 10.1038/nrmicro1273
- Donlan, R. (2002). Biofilms: microbial life on surfaces. *Emerg. Infect. Dis.* 8, 881–890. doi: 10.3201/eid0809.020063
- Draper, L., Cotter, P., Hill, C., and Ross, P. (2013). The two peptide lantibiotic lactacin 3147 acts synergistically with polymyxin to inhibit Gram negative bacteria. *BMC Microbiol.* 13:212. doi: 10.1186/1471-2180-13-212
- Elliason, D. J., and Tatini, S. R. (1999). Enhanced inactivation of *Salmonella typhimurium* and verotoxigenic *Escherichia coli* by nisin at 65°C. *Food Microbiol.* 16, 257–267. doi: 10.5803/jfsfm.16.257

- European Food Safety Authority (2006). Opinion of the scientific panel on food additives, flavourings, processing aids and materials in contact with food on a request from the commission related to the use of Nisin (e 234) as a food additive. *EFSA J.* 314, 1–16.
- Field, D., Connor, P., Cotter, P., Hill, C., and Ross, P. (2008). The generation of nisin variants with enhanced activity against specific Gram-positive pathogens. *Mol. Microbiol.* 69, 218–230. doi: 10.1111/j.1365-2958.2008.06279.x
- Field, D., Daly, K., O'Connor, P., Cotter, P., Hill, C., and Ross, P. (2015a). Efficacies of nisin A and nisin V semipurified preparations alone and in combination with plant essential oils for controlling *Listeria monocytogenes*. *Appl. Environ. Microbiol.* 81, 2762–2769. doi: 10.1128/AEM.00070-15
- Field, D., Gaudin, N., Lyons, F., O'Connor, P., Cotter, P., Hill, C., et al. (2015b). A bioengineered nisin derivative to control biofilms of *Staphylococcus pseudintermedius*. *PLoS ONE* 10:e0119684. doi: 10.1371/journal.pone.0119684
- Field, D., Hill, C., Cotter, P., and Ross, P. (2010a). The dawning of a “Golden era” in lantibiotic bioengineering. *Mol. Microbiol.* 78, 1077–1087. doi: 10.1111/j.1365-2958.2010.07406.x
- Field, D., Quigley, L., O'Connor, P., Rea, M., Daly, K., Cotter, P., et al. (2010b). Studies with bioengineered Nisin peptides highlight the broad-spectrum potency of Nisin V. *Microb. Biotechnol.* 3, 473–486. doi: 10.1111/j.1751-7915.2010.00184.x
- Furukawa, S., Akiyoshi, Y., O'Toole, G., Ogihara, H., and Morinaga, Y. (2010). Sugar fatty acid esters inhibit biofilm formation by food-borne pathogenic bacteria. *Int. J. Food Microbiol.* 138, 176–180. doi: 10.1016/j.ijfoodmicro.2009.12.026
- Gálvez, A., Abriouel, H., López, R., and Omar, N. (2007). Bacteriocin-based strategies for food biopreservation. *Int. J. Food Microbiol.* 120, 51–70. doi: 10.1016/j.ijfoodmicro.2007.06.001
- Gandhi, M., and Chikindas, M. (2006). *Listeria*: a foodborne pathogen that knows how to survive. *Int. J. Food Microbiol.* 113, 1–15. doi: 10.1016/j.ijfoodmicro.2006.07.008
- Gill, A., and Holley, R. (2004). Mechanisms of bactericidal action of cinnamaldehyde against *Listeria monocytogenes* and of eugenol against *L. monocytogenes* and *Lactobacillus sakei*. *Appl. Environ. Microbiol.* 70, 5750–5755. doi: 10.1128/AEM.70.10.5750-5755.2004
- Gou, J., Jung, L.-S., Lee, S.-H., and Ahn, J. (2011). Effects of nisin and acid on the inactivation and recovery of *Listeria monocytogenes* biofilms treated by high hydrostatic pressure. *Food Sci. Biotechnol.* 20, 1361–1366. doi: 10.1007/s10068-011-0187-2
- Govaris, A., Solomakos, N., Pexara, A., and Chatzopoulou, P. S. (2010). The antimicrobial effect of oregano essential oil, nisin and their combination against *Salmonella* Enteritidis in minced sheep meat during refrigerated storage. *Int. J. Food Microbiol.* 137, 175–180. doi: 10.1016/j.ijfoodmicro.2009.12.017
- Hara, H., Ohashi, Y., Sakurai, T., Yagi, K., Fujisawa, T., and Igimi, S. (2009). Effect of nisin (Nisaplin) on the growth of *Listeria monocytogenes* in Karashi-mentaiko (Red-pepper Seasoned Cod Roe). *Shokuhin Eiseigaku Zasshi* 50, 173–177. doi: 10.3358/shokueishi.50.173
- Healy, B., Field, D., O'Connor, P., Hill, C., Cotter, P., and Ross, P. (2013). Intensive mutagenesis of the nisin hinge leads to the rational design of enhanced derivatives. *PLoS ONE* 8:e79563. doi: 10.1371/journal.pone.0079563
- Helander, I., Alakomi, H.-L., Latva-Kala, K., Mattila-Sandholm, T., Pol, I., Smid, E., et al. (1998). Characterization of the action of selected essential oil components on gram-negative bacteria. *J. Agric. Food Chem.* 46, 3590–3595. doi: 10.1021/jf980154m
- Helander, I., and Mattila-Sandholm, T. (2000). Permeability barrier of the Gram-negative bacterial outer membrane with special reference to nisin. *Int. J. Food Microbiol.* 60, 153–161. doi: 10.1016/S0168-1605(00)00307-X
- Holah, J. T., Taylor, J. H., Dawson, D. J., and Hall, K. E. (2002). Biocide use in the food industry and the disinfectant resistance of persistent strains of *Listeria monocytogenes* and *Escherichia coli*. *J. Appl. Microbiol.* 92, 111S–120S. doi: 10.1046/j.1365-2672.92.5s1.18.x
- Houdt, V., and Michiels, C. W. (2010). Biofilm formation and the food industry, a focus on the bacterial outer surface. *J. Appl. Microbiol.* 109, 1117–1131. doi: 10.1111/j.1365-2672.2010.04756.x
- Hussain, M., and Dawson, C. (2013). Economic impact of food safety outbreaks on food businesses. *Foods* 2, 585–589. doi: 10.3390/foods2040585
- Hyldgaard, M., Mygind, T., and Meyer, R. (2012). Essential oils in food preservation: mode of action, synergies, and interactions with food matrix components. *Front. Microbiol.* 3:12. doi: 10.3389/fmicb.2012.00012
- Knight, G. C., Nicol, R. S., and McMeekin, T. A. (2004). Temperature step changes: a novel approach to control biofilms of *Streptococcus thermophilus* in a pilot plant-scale cheese-milk pasteurisation plant. *Int. J. Food Microbiol.* 93, 305–318. doi: 10.1016/j.ijfoodmicro.2003.11.013
- Koopmans, M., Vennema, H., Heersma, H., van Strien, E., van Duynhoven, Y., Brown, D., et al. (2003). Early identification of common-source foodborne virus outbreaks in Europe. *Emerg. Infect. Dis.* 9, 1136–1142. doi: 10.3201/eid0909.020766
- Kumar, C. G., and Anand, S. K. (1998). Significance of microbial biofilms in food industry: a review. *Int. J. Food Microbiol.* 42, 9–27. doi: 10.1016/S0168-1605(98)00060-9
- Laksanalamai, P., Joseph, L., Silk, B., Burall, L., Tarr, C., Gerner-Smidt, P., et al. (2012). Genomic characterization of *Listeria monocytogenes* strains involved in a multistate listeriosis outbreak associated with cantaloupe in US. *PLoS ONE* 7:e42448. doi: 10.1371/journal.pone.0042448
- Lemma, A., and Bernstein, J. (2010). The co-crystal of two GRAS substances: (citric acid)(nicotinamide). Formation of four hydrogen bonding heterosynthons in one co-crystal. *CrystEngComm* 12, 2029–2033. doi: 10.1039/b2927422g
- Liu, X., Tang, B., Gu, Q., and Yu, X. (2014). Elimination of the formation of biofilm in industrial pipes using enzyme cleaning technique. *MethodsX* 1, 130–136. doi: 10.1016/j.mex.2014.08.008
- Lucera, A., Costa, C., Conte, A., and Nobile, M. (2012). Food applications of natural antimicrobial compounds. *Front. Microbiol.* 3:287. doi: 10.3389/fmicb.2012.00287
- Maertens de Noordhout, C., Devleeschauwer, B., Angulo, F., Verbeke, G., Haagsma, J., Kirk, M., et al. (2014). The global burden of listeriosis: a systematic review and meta-analysis. *Lancet Infect. Dis.* 14, 1073–1082. doi: 10.1016/S1473-3099(14)70870-9
- Mahdavi, M., Jalali, M., and Kermanshahi, R. K. (2007). The effect of nisin on biofilm forming foodborne bacteria using microtiter plate method. *Res. Pharm. Sci.* 2, 113–118.
- McEntire, J., Montville, T., and Chikindas, M. (2003). Synergy between nisin and select lactates against *Listeria monocytogenes* is due to the metal cations. *J. Food Prot.* 66, 1631–1636.
- Molloy, E., Field, D., Connor, P., Cotter, P., Hill, C., and Ross, P. (2013). Saturation mutagenesis of lysine 12 leads to the identification of derivatives of nisin A with enhanced antimicrobial activity. *PLoS ONE* 8:e58530. doi: 10.1371/journal.pone.0058530
- Molloy, E., Ross, P., and Hill, C. (2012). “Bac” to the future: bioengineering lantibiotics for designer purposes. *Biochem. Soc. Trans.* 40, 1492–1497. doi: 10.1042/BST20120193
- Murdock, C. A., Cleveland, J., Matthews, K. R., and Chikindas, M. L. (2007). The synergistic effect of nisin and lactoferrin on the inhibition of *Listeria monocytogenes* and *Escherichia coli* O157:H7. *Lett. Appl. Microbiol.* 44, 255–261. doi: 10.1111/j.1472-765X.2006.02076.x
- Neetoo, H., Ye, M., and Chen, H. (2008). Potential antimicrobials to control *Listeria monocytogenes* in vacuum-packaged cold-smoked salmon pâté and fillets. *Int. J. Food Microbiol.* 123, 220–227. doi: 10.1016/j.ijfoodmicro.2008.02.001
- Nelson, K., Fouts, D., Mongodin, E., Ravel, J., DeBoy, R., Kolonay, J., et al. (2004). Whole genome comparisons of serotype 4b and 1/2a strains of the food-borne pathogen *Listeria monocytogenes* reveal new insights into the core genome components of this species. *Nucleic Acids Res.* 32, 2386–2395. doi: 10.1093/nar/gkh562
- Norhana, M. N., Poole, S., Deeth, H., and Dykes, G. (2012). Effects of nisin, EDTA and salts of organic acids on *Listeria monocytogenes*, *Salmonella* and native microflora on fresh vacuum packaged shrimps stored at 4°C. *Food Microbiol.* 31, 43–50. doi: 10.1016/j.fm.2012.01.007
- Olshewska, M. (2013). Microscopic findings for the study of biofilms in food environments. *Acta Biochim. Pol.* 60, 531–537.
- Phillips, C. A. (1999). The effect of citric acid, lactic acid, sodium citrate and sodium lactate, alone and in combination with nisin, on the growth of *Arcobacter butzleri*. *Lett. Appl. Microbiol.* 29, 424–428. doi: 10.1046/j.1472-765X.1999.00668.x

- Phillips, C. A., and Duggan, J. (2001). The effect of EDTA and trisodium phosphate, alone and in combination with nisin, on the growth of *Arcobacter butzleri* in culture. *Food Microbiol.* 18, 547–554. doi: 10.1006/fmic.2001.0417
- Pinto, M. S., de Carvalho, A. F., dos Santos Pires, A. C., Souza, A. A. C., da Silva, P. H. F., Sobral, D., et al. (2010). The effects of nisin on *Staphylococcus aureus* count and the physicochemical properties of Traditional Minas Serro cheese. *Int. Dairy J.* 21, 90–96. doi: 10.1016/j.idairyj.2010.08.001
- Pol, L., and Smid, E. (1999). Combined action of nisin and carvacrol on *Bacillus cereus* and *Listeria monocytogenes*. *Lett. Appl. Microbiol.* 29, 166–170. doi: 10.1046/j.1365-2672.1999.00606.x
- Poortinga, A. T., Smit, J., van der Mei, H., and Busscher, H. J. (2001). Electric field induced desorption of bacteria from a conditioning film covered substratum. *Biotechnol. Bioeng.* 76, 395–399. doi: 10.1002/bit.10129
- Prudêncio, C. V., Dos Santos, M. T., and Vanetti, M. C. (2015). Strategies for the use of bacteriocins in Gram-negative bacteria: relevance in food microbiology. *J. Food Sci. Technol.* 52, 5408–5417. doi: 10.1007/s13197-014-1666-2
- Rouse, S., Field, D., Daly, K., O'Connor, P., Cotter, P., Hill, C., et al. (2012). Bioengineered nisin derivatives with enhanced activity in complex matrices. *Microb. Biotechnol.* 5, 501–508. doi: 10.1111/j.1751-7915.2011.00324.x
- Ryu, J. H., and Beuchat, L. R. (2005). Biofilm Formation by *Escherichia coli* O157:H7 on stainless steel: effect of exopolysaccharide and curli production on its resistance to chlorine. *Appl. Environ. Microbiol.* 71, 247–254. doi: 10.1128/AEM.71.1.247-254.2005
- Sobrino-López, A., and Martín-Belloso, O. (2008). Use of nisin and other bacteriocins for preservation of dairy products. *Int. Dairy J.* 18, 329–343. doi: 10.1016/j.idairyj.2007.11.009
- Tan, S., Chew, S., Tan, S., Givskov, M., and Yang, L. (2014). Emerging frontiers in detection and control of bacterial biofilms. *Curr. Opin. Biotechnol.* 26, 1–6. doi: 10.1016/j.copbio.2013.08.002
- Tauxe, R. (2002). Emerging foodborne pathogens. *Int. J. Food Microbiol.* 78, 31–41. doi: 10.1016/S0168-1605(02)00232-5
- The Commission of the European Communities (2005). COMMISSION REGULATION (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs. *Off. J. Eur. Union* L 338/1, 1–26.
- Thomas, L. V., and Wimpenny, J. W. T. (1996). Investigation of the effect of combined variations in temperature, pH, and NaCl concentration on nisin inhibition of *Listeria monocytogenes* and *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 62, 2006–2012.
- Tunney, M., Ramage, G., Field, T., Moriarty, T., and Storey, D. (2004). Rapid colorimetric assay for antimicrobial susceptibility testing of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 48, 1879–1881. doi: 10.1128/AAC.48.5.1879-1881.2004
- Wiedemann, I., Breukink, E., van Kraaij, C., Kuipers, O. P., Bierbaum, G., de Kruijff, B., et al. (2001). Specific binding of nisin to the peptidoglycan precursor lipid II combines pore formation and inhibition of cell wall biosynthesis for potent antibiotic activity. *J. Biol. Chem.* 276, 1772–1779. doi: 10.1074/jbc.M006770200
- Wood, P., Jones, M., Bhakoo, M., and Gilbert, P. (1996). A novel strategy for control of microbial biofilms through generation of biocide at the biofilm-surface interface. *Appl. Environ. Microbiol.* 62, 2598–2602.
- World Health Organization and Foodborne Disease Burden Epidemiology Reference Group (2007). *WHO Initiative to Estimate the Global Burden of Foodborne Diseases. First formal meeting of the Foodborne Disease Burden Epidemiology Reference Group (FERG)*. Geneva: World Health Organization.
- Yang, S.-C., Lin, C.-H., Sung, C., and Fang, J.-Y. (2014). Antibacterial activities of bacteriocins: application in foods and pharmaceuticals. *Front. Microbiol.* 5:241. doi: 10.3389/fmicb.2014.00241
- Young, K. M., and Foegeding, P. M. (1993). Acetic, lactic and citric acids and pH inhibition of *Listeria monocytogenes* Scott A and the effect on intracellular pH. *J. Appl. Bacteriol.* 74, 515–520.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Smith, Draper, Hazelhoff, Cotter, Ross and Hill. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.