



OPEN ACCESS

Edited by:

Haiwei Luo,
The Chinese University of Hong Kong,
Hong Kong

Reviewed by:

You-Hee Cho,
CHA University, South Korea
M. Oves,
King Abdulaziz University, Saudi Arabia
Karl Drlica,
Rutgers University, The State
University of New Jersey,
United States

***Correspondence:**

Marirosa Molina
Molina.marirosa@epa.gov

†Present Address:

Adelumola Oladeinde,
Bacterial Epidemiology and
Antimicrobial Resistance Research
Unit, National Poultry Research
Center, U.S. USDA-ARS, Athens, GA,
United States
Chia-Ying Chen,
Department of Environmental
Engineering, National Chung Hsing
University, Taichung, Taiwan

Specialty section:

This article was submitted to
Evolutionary and Genomic
Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 19 December 2017

Accepted: 17 April 2018

Published: 08 May 2018

Citation:

Oladeinde A, Lipp E, Chen C-Y,
Muirhead R, Glenn T, Cook K and
Molina M (2018) Transcriptome
Changes of *Escherichia coli*,
Enterococcus faecalis, and
Escherichia coli O157:H7 Laboratory
Strains in Response to
Photo-Degraded DOM.
Front. Microbiol. 9:882.
doi: 10.3389/fmicb.2018.00882

Transcriptome Changes of *Escherichia coli*, *Enterococcus faecalis*, and *Escherichia coli* O157:H7 Laboratory Strains in Response to Photo-Degraded DOM

Adelumola Oladeinde^{1,2†}, Erin Lipp², Chia-Ying Chen^{3†}, Richard Muirhead⁴, Travis Glenn², Kimberly Cook⁵ and Marirosa Molina^{6*}

¹ National Exposure Research Laboratory, Student Volunteer, U.S. Environmental Protection Agency, Office of Research and Development, Athens, GA, United States, ² Department of Environmental Health Science, University of Georgia, Athens, GA, United States, ³ National Exposure Research Laboratory, National Research Council Associate, U.S. Environmental Protection Agency, Office of Research and Development, Athens, GA, United States, ⁴ Invermay Research Centre, AgResearch Ltd, Mosgiel, New Zealand, ⁵ Bacterial Epidemiology and Antimicrobial Resistance Research Unit, U.S. National Poultry Research Center, Agricultural Research Service, United States Department of Agriculture, Athens, GA, United States, ⁶ National Exposure Research Laboratory, U.S. Environmental Protection Agency, Office of Research and Development, Athens, GA, United States

In this study, we investigated gene expression changes in three bacterial strains (*Escherichia coli* C3000, *Escherichia coli* O157:H7 B6914, and *Enterococcus faecalis* ATCC 29212), commonly used as indicators of water quality and as control strains in clinical, food, and water microbiology laboratories. Bacterial transcriptome responses from pure cultures were monitored in microcosms containing water amended with manure-derived dissolved organic matter (DOM), previously exposed to simulated sunlight for 12 h. We used RNA sequencing (RNA-seq) and quantitative real-time reverse transcriptase (qRT-PCR) to compare differentially expressed temporal transcripts between bacteria incubated in microcosms containing sunlight irradiated and non-irradiated DOM, for up to 24 h. In addition, we used whole genome sequencing simultaneously with RNA-seq to identify single nucleotide variants (SNV) acquired in bacterial populations during incubation. These results indicate that *E. coli* and *E. faecalis* have different mechanisms for removal of reactive oxygen species (ROS) produced from irradiated DOM. They are also able to produce micromolar concentrations of H₂O₂ from non-irradiated DOM, that should be detrimental to other bacteria present in the environment. Notably, this study provides an assessment of the role of two conjugative plasmids carried by the *E. faecalis* and highlights the differences in the overall survival dynamics of environmentally-relevant bacteria in the presence of naturally-produced ROS.

Keywords: fecal indicator bacteria, gene expression, solar irradiated DOM, oxidative stress genes, RNA-sequencing, reactive oxygen species (ROS), hydrogen peroxide

INTRODUCTION

Survival of bacteria in the environment is dictated by their ability to grow or persist under diverse abiotic and biotic stressors. In aquatic ecosystems, resource availability, sunlight, temperature, pH, and competition have been shown to be important drivers of bacterial population dynamics (Bradford et al., 2013, 2015; Pachepsky et al., 2014). Of these factors, substrate availability, in the form of dissolved organic matter (DOM), is necessary for bacteria to proliferate in surface waters. Further, DOM transformations mediated by sunlight and microbes can have an interacting effect on bacterial survival (Mostofa et al., 2012; Häder et al., 2015).

The colored fraction of DOM (CDOM) is primarily responsible for absorbance of UV light and production of labile nutrients from refractive fractions of DOM, which can be subsequently used for growth (Bushaw and Zepp, 1996; Moran and Zepp, 1997; Häder et al., 2011). These labile nutrients include low molecular weight nitrogen compounds, such as amino acids, that are of great importance to the biogeochemistry of natural waters and are involved in numerous biochemical processes. Another important transient product of absorption of UV/visible light by CDOM is the formation of reactive oxygen species (ROS) from photochemical reactions involving oxygen (Williamson et al., 2017; Wolf et al., 2018). ROS, including hydrogen peroxide (H_2O_2), hydroxyl radical ($\cdot\text{OH}$) singlet oxygen ($^1\text{O}_2$), and super oxide radicals (O_2^-), are photochemically produced reactive intermediates formed during photo-oxidation of CDOM. Further, phototransformation of labile organic compounds, such as thiols present in surface waters can also result in a net increase in ROS steady state (Chu et al., 2016).

Hydrogen peroxide is uncharged and, unlike other ROS, it can easily permeate the bacterial cell surface. It can also persist in natural waters for periods ranging from several hours to days (Kieber et al., 2003; Mostofa and Sakugawa, 2009). Hydrogen peroxide concentration has a diel cycle, with peak concentrations typically noted at noontime during summer months (Clark et al., 2008). At steady state concentrations, H_2O_2 concentrations are reported to range from 6 nM in marine waters up to 3.2 μM in rivers and streams (Mostofa et al., 2013). The concentration of photo-produced H_2O_2 in surface water is dependent on DOM source/type and concentration, sunlight irradiance, and antioxidants present (Mostofa et al., 2013). Antioxidants, such as glutathione and cysteine are important scavengers of peroxides and are the major component of peroxiredoxins used by prokaryotes and eukaryotes for peroxide signaling (Dubbs and Mongkolsuk, 2007; Parsonage et al., 2008). Subsequently, in extracellular environment we expect the steady state of H_2O_2 concentration to be influenced by the concentration of antioxidants or enzymes with antioxidant activity present.

At high extracellular H_2O_2 concentrations, microbes can experience toxicity from significant changes in their cell redox homeostasis. Nevertheless, bacteria have developed elegant systems to relieve themselves of H_2O_2 -induced oxidative stress (Imlay, 2015b). For example, catalase (*kat*) and peroxidase (*ahp*) genes have been demonstrated to be important genes

used by *E. coli* under H_2O_2 exposure (Imlay, 2015b); and mutants of *dps* and iron-sulfur clusters (*suf*) have been shown to suffer Fenton mediated DNA damage (Djaman et al., 2004; Lim et al., 2007). The majority of these studies have used laboratory grade H_2O_2 which may not represent the form of H_2O_2 environmentally-associated microbes are exposed to. In addition, the concentrations used in these studies are higher than reported concentrations in natural waters.

The sources of DOM in aquatic ecosystems are diverse and spatially variable. In watersheds dominated by agriculture, livestock farms or lagoon ponds are a major non-point source of DOM via runoff related processes (Bida et al., 2015; Graeber et al., 2015; Heinz et al., 2015). This form of DOM is of fecal origin and has been broken down by the gut microbiome of warm blooded animals, making them bioavailable for use by extra-intestinal microbes. Microorganisms that can readily utilize this source of nutrients for growth should be at an advantage. Moreover, elevated levels of fecal indicator bacteria, such as *E. coli* and enterococci in streams and rivers have been attributed to these non-point sources of pollution (USEPA, 2017). The survival of indicator bacteria under sunlight exposure have been extensively studied (Rochelle-Newall et al., 2015), however, little is known on the interacting effects of sunlight, DOM, and ROS.

To understand the role of photo-produced ROS on the survival of specific laboratory bacterial strains, we attempted to isolate the effect of H_2O_2 using controlled microcosms of natural water spiked with cattle feces-derived DOM exposed to a solar simulator prior to inoculation with bacteria. Following bacterial inoculation, microcosms were incubated in the dark for up to 24 h. We employed high throughput RNA-seq and qRT-PCR to investigate the expression of transcripts required for H_2O_2 detoxification and oxidative stress. In addition, we used the RNA-seq data to identify Single Nucleotide Variants (SNVs) acquired following inoculation into water amended with DOM. We provide insights into the different survival mechanisms used by two important fecal indicator strains for water quality monitoring, and a zoonotic pathogen of public health interest.

MATERIALS AND METHODS

Cattle Fecal Extract Preparation

Fresh fecal samples were collected from five cows on a commercial farm in northeast Georgia on July 13, 2013. Fecal samples were composited, homogenized, and made into 1:10 fecal slurry in 0.85% KCl and mixed for 1 h in a hand/wrist shaker. The fecal slurry was then centrifuged twice at 4,000 \times g for 10 min and the resulting supernatant was saved and named cattle fecal extract (CFE). CFE was sequentially filtered through 1.2, 0.45, and 0.2 μm pore-sized polycarbonate membrane filters. The dissolved organic carbon (DOC) concentration in CFE was determined using a total organic carbon analyzer (TOC- V_{CPH} , Shimadzu, Kyoto, Japan) equipped with auto samplers.

CFE was spiked into autoclaved phosphate buffered water (PBW) microcosms to concentrations mimicking nutrient inputs from direct fecal deposition into streams (i.e., final DOC concentration of $32.48 \pm 1.52 \text{ mg l}^{-1}$). The absence of bacteria and relevant lytic phages in CFE was confirmed by culturing

100 μ l of CFE in Brain Heart Infusion (BHI) broth and by performing phage double agar overlay assay (Adams, 1959), respectively. For the overlay assay, bacterial strains used in the present study were used as phage hosts. Thereafter, CFE/DOM-spiked water was divided into two volumes, with one exposed to solar radiation [Irradiated DOM-spiked Water (I-DOMW)] and the other serving as a dark control [Non-irradiated DOM-spiked Water (N-DOMW)].

Irradiation of DOM Spiked Water

Solar irradiation was performed in an Atlas SunTest CPS/CPS+ solar simulator (Atlas Materials Testing Technology, Chicago, IL) equipped with a 1 kW xenon arc lamp. Irradiance of the simulator in the UV spectral region was similar to mid-summer, midday natural sunlight at 33.95°N, 83.33°W (Athens, GA, USA). Samples were irradiated for 12 h in 25 ml quartz tubes or in a 1 l jacketed Pyrex beaker (Ace glass, Vineland, NJ) before bacteria inoculation (Figure 1).

Inoculum Preparation

Overnight cultures of *E. coli* C3000 (ATCC 15597)—hereafter referred to as *E. coli*; *Enterococcus faecalis* (ATCC 29212)—hereafter referred to as *E. faecalis*; and *Escherichia coli* O157:H7

B6914—hereafter referred to as *E. coli* O157:H7—were harvested, washed and grown for an additional 1.5 h to exponential phase (OD_{600} of 0.1) in BHI broth. The genotypic characteristics of these strains are reported in Table 1. Each culture was centrifuged at 4,000 \times g for 5 min and washed twice in phosphate-buffered water (PBW). Thereafter, serial dilutions were made to the desired concentrations. Two inocula concentration were tested in separate experiments: ca. 10^6 CFU ml^{-1} for high inoculum and ca. 10^3 CFU ml^{-1} for low inoculum (Supplementary File 1).

Microbiological Analysis

Following irradiation, I-DOMW, N-DOMW, and PBW controls were separately inoculated with mid-logarithmic phase of each bacterium. For low inoculum experiments, 20 ml of each treatment were dispensed into sterile 50 ml centrifuge tubes and incubated in the dark at 25°C in a temperature-controlled incubator shaker (150 rpm) (Innova 4230, New Brunswick Scientific, Edison, NJ). For high inoculum, 50 ml were dispensed in 250 ml Erlenmeyer flasks and covered with sterile aluminum foil before incubation, as mentioned previously. Three replicates derived from one bacterial population were selected at random for analysis, at 0.5, 6, 12, 24, and 48 h. *E. coli*, *E. faecalis* and *E. coli* O157:H7 populations were quantified by culture methods using




FIGURE 1 | Schematic of experimental design. Inset shows spectra of natural sunlight and light emitted by the Xenon lamp in solar simulator used in this study.

TABLE 1 | Characteristics of strains used in this study.

Bacteria (strain)	Plasmid Profile (size) ^a	Prophages present ^b	Unique Single Nucleotide Variants (SNV)	Reference ^c
<i>E. coli</i> (C3000)	IncF ^d	8 defective	89	This study
<i>E. faecalis</i> (ATCC 29212)	repUS11 (66 kb) rep9 (41 kb)	7 defective	58	Minogue et al., 2014
<i>E. coli</i> O157:H7 (B6914)	ND	ND	ND	Uhlich et al., 2017

ND, not determined.

^aDetermined using the Center for Genomic Epidemiology's PlasmidFinder program.

^bDetermined using PHAST phage search tool.

^cPublished complete or draft genome.

^dIncomplete plasmid contig.

modified mTEC agar (EPA method 1603), mEI agar (EPA method 1600), and MUG *E. coli* O157:H7 agar supplemented with 100 µg ml⁻¹ ampicillin (Sigma Aldrich), respectively (Figure 1).

Reactive Oxygen Species (ROS) Measurement

The production and concentration of singlet oxygen (¹O₂) and hydroxyl radicals (.OH) were determined separately, as described by Chen and Jafvert (2010). ¹O₂ was monitored via the loss of furfuryl alcohol (FFA) and the pseudo-state concentration of ¹O₂ was determined. To detect OH, *p*-chlorobenzoic acid (*p*CABA), an OH scavenger was added. *p*CABA was added at a low concentration (2 µM), allowing the pseudo-steady-state concentration of OH to be calculated. Superoxide radicals (O₂⁻) was measured with chemiluminescence reagent 2-methyl-6-(4-methoxyphenyl)-3,7-dihydroimidazo[1,2-a]pyrazin-3(7H)-one (MCLA) in a FeLume chemiluminescence system (Waterville Analytical, LLC, Waterville, Maine) (Rose et al., 2008).

Extracellular H₂O₂ concentrations, before and after bacterial inoculation, were quantified using the copper-DMP (2,9-dimethyl-1, 10-phenanthroline) spectrophotometric method (Kosaka et al., 1998). Samples were filter-sterilized using a 0.22 µm syringe filter to remove bacteria prior to measuring H₂O₂. Absorbance readings in I-DOMW and N-DOMW were normalized against N-DOMW controls (no bacteria inoculation). A calibration curve was constructed by plotting the concentration of known ACS-grade H₂O₂ (Sigma Aldrich) solution vs. the absorbance at 454 nm of the product formed by reaction of the solutions with copper sulfate and DMP. Two separate calibration curves were used throughout the experiment for quality control and to ensure reproducibility of results. A more complete description of the methods used to detect and quantify ROS is provided in the Supplementary Material.

RNA Isolation

Duplicate microcosms from one high inoculum experiment were selected randomly at 0.5, 6, 12, and 24 h, and ~45 ml was

filtered through 0.45 µm pore size isopore membrane (EMD Millipore, Billerica, MA). Filters were folded inwards and saved in a lysing matrix B tube (MP Biomedical, Solon, OH) containing 600 µl RNAlater (Life Technologies, Grand Island, NY). Tubes were kept at -80°C for 2 weeks prior to total RNA extraction. Duplicate filters per time point were removed with sterile forceps and carefully opened to expose the filter surface. Filters were rinsed twice in cold 1X phosphate buffer saline to remove RNAlater, after which 170 µl of 8 mg ml⁻¹ lysozyme were dispensed onto the filter surface. Filters containing lysozyme were incubated at 37°C for 5 min prior to extraction. RNA extraction was performed with the FastRNA spin kit for microbes (MP Biomedical, Solon, OH), according to the manufacturer's instructions, except that the bead-beating step was repeated twice at 6.5 m s⁻¹ for 60 s. Total RNA was eluted twice in 25 µl DEPC-treated water for a final volume of 50 µl. Total RNA was concentrated using a Vacufuge (Eppendorf, NY, USA), with no heat treatment for 2.5 h.

RNA pellets were rehydrated with 20 µl DEPC-treated water and treated with 8U Turbo DNA-free kit (Life Technologies, Grand Island, NY) to remove genomic DNA contamination. Recovered RNA was quantified with a Nanodrop ND 1000 spectrophotometer (Thermo Fisher Scientific, MA, USA), examined for quality on an Agilent Bioanalyzer, and stored at -80°C until used for RNA-seq (between 2 and 3 months).

qRT-PCR

Total RNA (100 ng) from each bacterial population was reverse-transcribed with 100 U of high capacity cDNA reverse transcription kit (Life Technologies, Grand Island, NY), per manufacturer's instructions. Primers were synthesized by Integrated DNA Technologies (Coralville, IA) (Table S1). Quantitative Reverse Transcription PCR (qRT-PCR) was performed on an ABI Prism 7500HT Fast Sequence Detection System (Applied Biosystems, Foster City, CA) with 40 amplification cycles, using Fast SYBR Green PCR Master Mix as a signal reporter. Each reaction was composed of 2 µl of cDNA, 1 µM sense and antisense primers for a total volume of 20 µl. qRT-PCR was run in a Fast 96-well microtiter PCR plate using the following amplification conditions: 1 cycle for 20 s at 95°C; and 40 two-step cycles at 95°C for 3 s and 60°C for 30 s. Specificity of primer pairs was verified by melting curve analysis. qPCR efficiency was tested with serial dilutions of cDNA samples, and all ranged between 1.70 and 2.07.

Expression levels of *idnT*, *cysG*, *hcaT*, *gyrB*, and *gapA* (Michán et al., 1999; Kyle et al., 2010; Zhou et al., 2011a) were evaluated as possible reference controls for the normalization of *E. coli* H₂O₂ scavenging gene expression (Table S2). The constitutive genes were carefully considered to identify the optimal normalization gene (from the set of candidates) by the geNorm finder algorithm available under the Python library-Eleven (Vandesompele et al., 2002). Data were analyzed using the 2^{-ΔΔCT} method described by Livak and Schmittgen (2001). For each gene, the ratio of expression in I-DOMW, compared to that in N-DOMW, was normalized to the expression of the top two ranked constitutive genes in *E. coli* (*gapA* and *gyrB*). For *E. faecalis*, this ratio was normalized to the expression of *rpoB*

and *gyrB* (Riboldi et al., 2014). To assess for reagent and genomic DNA contamination, no-template and no-reverse-transcriptase controls were included.

mRNA Enrichment

Twenty microliters of DNase-treated RNA was pelleted as described earlier and reconstituted in 15 μ l (75 ng to 2.5 μ g) 1X TE buffer. Bacterial 16S and 23S ribosomal RNA removal was completed using MICROBExpress Bacterial mRNA Enrichment Kit (Life Technologies, Grand Island, NY) according to manufacturer's instruction, but with half of the suggested reaction volumes. The recovered mRNA was quantified using Nanodrop ND 1000 spectrophotometer (Thermo Scientific, Waltham, MA).

cDNA Library Preparation

cDNA synthesis was performed on enriched mRNA (~2–160 ng) using the KAPA stranded RNA-seq library preparation kit (Kapa Biosystems, Inc. MA, USA), with a few modifications. Half of the suggested reaction volumes were used throughout. RNA fragmentation was done in a thermocycler at 87.5°C for 6 min. Adapterama I adapters and primers (Glenn et al., 2016) were used for ligation and PCR amplification reactions at a final reaction concentration of 357 and 250 nM, respectively. For PCR amplification, iTru5 forward and iTru7 reverse primers with unique indexes for sample multiplexing were employed (Glenn et al., 2016). The concentrations of amplicons from different samples were quantified using a Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific, MA, USA). The libraries were pooled in equimolar concentrations and sequenced at the Georgia Genomics Facility using an Illumina NextSeq (150 cycles) Mid Output Flow Cell. A 75 bp paired-end sequencing reaction was performed on a NextSeq platform (Illumina, San Diego, CA, USA). cDNA fragments (~339 bp) were obtained for all 24 biosamples. Reads were submitted to NCBI SRA under submission ID SUB1913975 and bioproject PRJNA341849.

RNA-seq Analysis

Unfiltered fastq sequences of each bacterium were aligned to bacterial reference genomes (*E. coli* k-12- NC_000913, *E. coli* O157:H7- NC_002655, and *E. faecalis*- CP008816) using Burrows-Wheeler Aligner (BWA) (Li and Durbin, 2009) with default parameters. Reference genome CP008816 carries two plasmids of sizes 66,548 (CP008815.1) and 41610 bp (CP008814.1). SAM alignment was converted to BAM (samtools view -bs), sorted by coordinates (samtools sort), and PCR duplicates removed using SAMtools (samtools rmdup) (Li et al., 2009). Mapped reads were counted using BEDTools (multiBamCov -bams) (Quinlan and Hall, 2010). Read counts were exported in a tab delimited file for normalization and differential gene expression (DGE) in R (R Developmental Core Team, 2012). DGE analysis was completed with DESeq2 package (Love et al., 2014). Reads with 0 or 1 count were removed before DGE was performed (see Supplementary File 4 for scripts used).

For each RNAseq data set, genes with an absolute fold change ≥ 2 and adjusted *p*-value of <0.01 were used for

further analysis in STRINGDB (Szklarczyk et al., 2014). Proteins corresponding to the obtained gene sets were searched against version 10 of the STRING database to display functional protein-association networks. Interactions with a STRING confidence ≥ 0.4 (medium and high confidence) were considered. A markov cluster algorithm (MCL) (Van Dongen, 2008) of 2 was used for clustering.

Whole Genome Sequencing (WGS)

To identify mutations and plasmids in parental *E. coli* and *E. faecalis* strains used in this study, we performed WGS on DNA extracted from pure colonies. Briefly, archived pure cultures were streaked onto Sheep Blood Agar (Remel Inc, San Diego, CA) and incubated overnight at 37°C for 24 h. Following overnight growth, 5–6 single colonies were randomly chosen and resuspended in 200 μ l molecular grade DEPC-treated water. Genomic DNA was extracted from resuspended cells with FastDNA Spin Kit (Mp Biomedicals, Solon, Ohio) according to manufacturer's instructions and quantified with a Qubit Fluorometer (ThermoFisher Scientific).

WGS libraries were prepared using MiSeq Nextera XT library preparation kit (Illumina, Inc., San Diego, CA). Sequencing was performed on the Illumina MiSeq platform with 250-bp paired end reads using the MiSeq reagent V2 (500 cycles). All isolates were sequenced for an average coverage of 100X. Sequence reads were assembled de novo into contigs using SPAdes assembler (Bankevich et al., 2012). Assembled contigs were submitted to the Center for Genomic Epidemiology's PlasmidFinder (Carattoli et al., 2014) to determine existing plasmid replicon types. Prophages were identified using PHAST (Zhou et al., 2011b).

Single Nucleotide Variants (SNV) Identification

SNVs in parental strains and DOMW-treated populations were identified by mapping unfiltered WGS and RNA-seq reads to closed chromosome and plasmid genome of *E. coli* and *E. faecalis* available in NCBI under accession numbers NC000913 and CP008816, respectively, using BWA. SAM file alignment, sorting and removal of PCR duplicates was done using SAMtools. Genome Analysis Toolkit (McKenna et al., 2010) with a minimum mapping quality of 30 and a minimum base quality of 30 was used for SNV calling.

Statistical Analysis

Bacterial growth rate was derived from the following equation:

$$dN/dt = kN$$

where *N* is the concentration of cells, *t* is the time, and *k* is the growth rate constant. Analyses were performed using R (version 3.4.1). A profile analysis was performed to compare bacterial growth rates (μ_6 vs. $\mu_{0.5}$, μ_{12} vs. μ_6 , μ_{24} vs. μ_{12}) between treatments, after which a multivariate analysis of variance (MANOVA) tested for significant differences in growth rate between bacteria incubated in I-DOMW and N-DOMW microcosms (Table S3). Although small sample size can affect the power and homogeneity of the variance test, profile analysis

still provides more power than univariate tests (Macedo and Waterson, 2016). Graphs were plotted in SigmaPlot (Systat Software, San Jose, CA). Comparisons between treatments were performed using Wilcoxon signed-rank test. A linear regression analysis was used to model the relationship between bacterial concentration (Log CFU ml⁻¹) and extracellular H₂O₂ concentration (μM).

RESULTS

Bacterial Growth Potential Differed in Water Spiked With DOM

To determine the effect of DOM photodegradation on bacterial growth, we inoculated bacteria into DOM-spiked water that was previously irradiated under a sunlight simulator for 12 h. All three strains of bacteria differed in their growth dynamics following inoculation into Irradiated DOM-spiked Water (I-DOMW) and Non-irradiated DOM-spiked Water (N-DOMW). *E. coli* and *E. coli* O157:H7 grew by several orders of magnitude after 12 h of dark incubation in I-DOMW and N-DOMW (Figures 2A,C, Supplementary File 1). At a high inoculum starting concentration (ca. 10⁶ CFU ml⁻¹), *E. coli* population had a lower growth rate in I-DOMW than N-DOMW, however, there was no significant difference in maximum concentration achieved when a high or low starting inoculum was used (Table 2). *E. faecalis* population was lower in I-DOMW than N-DOMW after 24 h of dark incubation when inoculated at both low and high starting inocula concentrations (Figure 2B, Table 2).

Photodegradation of Dissolved Organic and Inorganic Compounds

DOM photodegradation resulted in changes in measured concentration of dissolved organic carbon (DOC), ammonium

(NH₄⁺), nitrate (NO₃⁻), and orthophosphate PO₄³⁻. Following 12 h irradiation of DOMW, the concentration of DOC (35.76 ± 11.6 mg l⁻¹), [NH₄⁺] (0.73 ± 0.2 mg l⁻¹), and [PO₄³⁻] (8.33 ± 1.9 mg l⁻¹) decreased to 31.09 ± 5.3, 0.66 ± 0.1, and 8.03 ± 1.8 mg l⁻¹, respectively (Table S4). For [NO₃⁻] (0.44 ± 0.1 mg l⁻¹), a 1.2-fold increase was observed after DOMW irradiation (Wilcoxon rank-signed Test, *p*.value = 0.001).

ROS Production and Removal in DOM Spiked Water

The observed significant effect of DOM irradiation on *E. coli* and *E. faecalis* populations (Figures 2A,B), made us question if photo-produced ROS played an inhibitory role. To test this hypothesis, we monitored for the production of singlet oxygen (¹O₂), hydroxyl radicals (.OH), superoxide radicals (O₂⁻), and hydrogen peroxide (H₂O₂) following DOM irradiation. Photoproduction of ¹O₂, O₂⁻, and H₂O₂ was observed during 12 h of irradiation of DOMW before bacterial inoculation. No hydroxyl radicals (.OH) were detected. Sunlight irradiation of DOMW produced ¹O₂ and O₂⁻ at steady states of ca. 2.32⁻¹³ and 1.3⁻⁹ M respectively, and H₂O₂ at a concentration of 15.38 ± 0.81 μM.

Superoxide radicals are reactive and cannot cross bacterial lipid bilayer, thus its extracellular role in *E. coli* growth rate reduction in this study is unlikely (Imley, 2008). Regarding ¹O₂, we exposed each bacterium to ¹O₂ generated from sunlight irradiation of Rose Bengal (See supplementary material for a detailed description of methods used). The steady state of ¹O₂ generated (7.38⁻¹³M) was similar to levels reported in surface waters (Burns et al., 2012) and from DOM irradiation in this study. However, there was no significant difference in concentration between exposed and unexposed populations for the three bacterial strains (Figure S1).




FIGURE 2 | Bacteria concentration and extracellular H₂O₂ concentration in the presence of sunlight irradiated cattle fecal extract. A profile analysis followed by MANOVA was performed to test for significant differences in growth rate between bacteria incubated in I-DOMW and N-DOMW microcosms. Dashed lines represent extracellular H₂O₂ concentration during dark incubation in I-DOMW (green) and N-DOMW (yellow) in the presence of (A) *E. coli* (B) *E. faecalis* (C) *E. coli* O157:H7. Horizontal short dash lines represent method detection limit for H₂O₂ (0.5 μM). Error bars represent standard error for H₂O₂ (*n* = 3 replicates per time point) and standard deviation for bacteria concentration (*n* = 2 replicates per time point). *denotes level of significance for the effect of DOM irradiation on bacteria growth rate per time (**p* < 0.05, ***p* < 0.01).

TABLE 2 | Growth rate comparison at low and high inocula concentrations for *E. coli*, *E. faecalis*, and *E. coli* O157:H7.

Bacteria	Inocula concentration	Treatment	Growth rate (hr ⁻¹) ^a	Doubling time (h)	Concentration after 24 h (Log CFU ml ⁻¹)
<i>E. coli</i>	High	I-DOMW	0.14	5.1	7.56
		N-DOMW	0.17	4.1	7.49
	Low	I-DOMW	0.29	2.4	7.49
		N-DOMW	0.24	2.9	7.3
<i>E. faecalis</i>	High	I-DOMW	0.03	22.6	6.25
		N-DOMW	-0.0024	ND	6.56
	Low	I-DOMW	0.09	7.8	5.31
		N-DOMW	0.28	2.5	6.29
<i>E. coli</i> O157:H7	High	I-DOMW	0.27	2.6	7.62
		N-DOMW	0.24	2.9	7.72
	Low	I-DOMW	0.26	2.7	7.27
		N-DOMW	0.26	2.7	7.18

I-DOMW, Irradiated DOM-spiked Water; N-DOMW, Non-irradiated DOM-spiked Water.

Boldness denotes significant difference in growth rate between I-DOMW and N-DOMW (*p*. value < 0.05).

ND, not determined. High and low inocula concentration represents ca. 10⁶ and ca. 10³ CFU ml⁻¹ of starting bacteria concentration, respectively.

^aDetermined from two separate experiments. Experiments were repeated 8, 3, and 3 X with *E. coli*, *E. faecalis*, and *E. coli* O157:H7, respectively (See Supplementary File 1 for data on experiments not reported in this Table).

Further, we determined the concentration of extracellular H₂O₂ during experiments with high starting concentrations of bacteria. The concentration of photo-produced H₂O₂ was negatively correlated with *E. coli* (Adj. *R*² = 0.74; *p*.value = 0.004) and *E. coli* O157:H7 concentrations (Adj. *R*² = 0.93; *p*.value < 0.001) (Figure S2), with extracellular H₂O₂ decreasing and bacterial populations increasing in I-DOMW during the first 6 h of dark incubation (Figure 2). On the other hand, extracellular H₂O₂ increased significantly in the presence of *E. faecalis*. Extracellular H₂O₂ was also detected in μM concentrations with all three types of bacteria during dark incubation in N-DOMW microcosms (Figure 2). Bacterial-produced H₂O₂ showed no significant correlation with bacteria concentration (Figure S2), however. There was no significant decline in H₂O₂ concentration for I-DOMW controls with no bacteria inoculated (Figure S3).

Gene Expression

Oxidative stress in bacteria is ameliorated by multiple regulatory networks including catalase, peroxidase, cytochrome, and hydrogenase systems. The expression of genes encoding catalase or peroxidase enzymes are regulated based on the ROS level present endogenously and exogenously. For *E. coli*, data showed that steady-state intracellular and extracellular H₂O₂ concentration above 200 nM and 2 μM, respectively, are required to induce the OxyR-response (Khademian and Imlay, 2017). To determine highly expressed genes in this study, we performed RNA-seq and qRT-PCR on populations collected from one high inoculum experiment at time 0.5, 6, 12, and 24 h. The number of reads is documented and >90% of reads were mapped to a reference genome (Supplementary File 2). The average number of reads mapped per sample for *E. coli*, *E. faecalis* and *E. coli* O157:H7 were 1,806,326 ± 144,075, 1,529,044 ± 124,201, and 1,962,671 ± 446,418, respectively. Mapped reads corresponded to an average coverage of 117 X with a range

of 33 to 245 X. The number of genes differentially expressed differed as a function of the type of bacteria and the dark incubation time (Table 3, Supplementary File 3). Differentially expressed transcripts between I-DOMW and N-DOMW decreased temporally during dark incubation for all strains tested.

We used STRINGDB to link differentially expressed transcripts to significant genetic interactions (Szklarczyk et al., 2014). The majority of the genes with increased expression for *E. coli* and *E. coli* O157:H7 in I-DOMW were associated with oxidative stress (Figures 3A,C). These genes include *kat* and *ahp* for the detoxification of H₂O₂, *sufABCD*S for the repair of iron-sulfur clusters (Fe-S), and *dps* for the sequestration of irons (Figures 4A,C). In addition, there was increased expression in transcripts involved in iron uptake and siderophore transport in *E. coli* O157:H7. For N-DOMW microcosms, transcripts with increased expression for *E. coli* and *E. coli* O157 were associated with leucine/isoleucine biosynthesis (*ilvBC* and *leuAB*), biofilm formation (*csgAB*) and tryptophan (*trpBCDE*) biosynthesis (Figures 3A,C; Supplementary Figures 4, 6).

In comparison, the majority of transcripts with increased expression in I-DOMW, relative to N-DOMW for *E. faecalis*, were associated with ATP-binding cassette (ABC) transporters, fatty acid biosynthesis, pheromone binding and beta-lactam resistance (Figure 3B). Comparatively, glycerol metabolism (*glp*) and sugar transport transcripts were upregulated in N-DOMW compared to I-DOMW (Figures 3B, 5). Twelve plasmid-encoded transcripts were also differentially expressed in *E. faecalis* (*p*.value < 0.05; Table 5). Ten of these transcripts were carried on the 66 kb plasmid, and encode genes for replication, recombination, bacteriocin metabolism/transport and thioredoxin (Figure 6). Transcripts associated with replication (*rep*) and bacteriocin production (*cylL*) were the only genes with higher expression in I-DOMW compared to N-DOMW.

Finally, we used qRT-PCR to confirm the relative expression of a few OxyR-like genes (Table 4). *katG* and *ahpCF* genes were the most highly expressed [>2 - Fold change (FC)] after 0.5 h for

E. coli and *E. coli* O157:H7. None of the genes tested for *E. faecalis* were expressed above 2- fold.

TABLE 3 | Number of differentially expressed genes (DEG) ($p_{adj} < 0.05$) between bacteria incubated in I-DOMW and N-DOMW.

Bacteri	Time (h)	Upregulated genes in I-DOMW	Upregulated genes in N-DOMW
<i>E. coli</i>	0.5	76	88
	6	10	5
	12	3	4
	24	0	0
<i>E. faecalis</i>	0.5 [#]	22	36
	6	130	143
	12	25	23
	24	2	19
<i>E. coli</i> O157:H7	0.5	47	6
	6	21	7
	12	0	0
	24	1	0

I-DOMW, Irradiated DOM-spiked Water; N-DOMW, Non-irradiated DOM-spiked Water.
[#]*p*. value reported instead of *p*. adj due to low sample size ($n = 1$) at 0.5 h for *E. faecalis*.

Mutation Changes in Bacterial Populations

To leverage the transcriptome expression with mutations, we identified single nucleotide variants (SNV) on the same data set. In addition, we performed whole genome sequencing (WGS) on parental *E. coli* and *E. faecalis* strains used in this study. Our goal was to differentiate SNVs acquired during dark incubation from those present in parental strains used for inoculation. We did not sequence *E. coli* O157:H7 since transcriptomics results showed that the two *E. coli* strains had similar transcripts differentially expressed and because this strain has been extensively engineered (Fratamico et al., 1997).

Sequenced strains had an average coverage of 113X. *E. coli* and *E. faecalis* differed from the reference genomes used in this study by 89 (NC_000913) and 58 (CP008816) SNVs, respectively, on the chromosome. Additionally, we confirmed that *E. faecalis* carried similar 66 and 41 kb plasmids present in CP008816. These plasmids were identified as repUS11 and rep9, and shared DNA homology with the pTEF3 (accession no: AE016832) and sex pheromone plasmid pAD1 (accession no: L01794) of *Enterococcus*, respectively (Table 1). The number of copies of each plasmid present in a cell was determined using the coverage of assembled chromosomal and plasmid contigs.






FIGURE 3 | Genetic networks of differentially expressed transcripts during dark incubation as determined by RNAseq. RNA-seq was performed on populations collected at 0.5, 6, 12, and 24 h. Center: Volcano plot showing fold-change of gene expression in I-DOMW compared to N-DOMW. Transcripts with significant increase in expression between the groups ($p_{adj} < 0.01$; fold- change ≥ 2) are highlighted in red; lower: STRING analysis for significantly altered genes in each case for (A) *E. coli* (B) *E. faecalis* and (C) *E. coli* O157:H7. Solid colored connecting lines depict protein interactions predicted with high confidence (>0.8), while dashed lines indicate protein interactions predicted with moderate confidence (0.4–0.7).



We identified 17 and 20 high quality SNVs in I-DOMW ($n = 5$) and N-DOMW ($n = 8$) *E. coli* populations, respectively, using RNA-seq data (Figure 7, Table S5A). The mutations in I-DOMW populations were present in genes associated with GTP-binding (*obg*, *typ*), hydrogen cycling (*hyaB*), and Qin prophage genes (*ydfC*). In N-DOMW populations, these mutations were acquired in genes involved in peptidoglycan metabolism (*dacC*, *mltA*), oxidoreduction reactions (*Nuo*, *paoC*, *nrfD*), and siderophore production (*entC*). For *E. faecalis*, I-DOMW populations ($n = 6$) acquired 9 mutations, whereas N-DOMW populations ($n = 4$) had 3 mutations. I-DOMW populations gained mutations in genes with transferase (*purF*, *aspB*- locus tag DR75_777 and DR75_1102) and ATP-dependent helicase activity (*rexB*—locus tag DR75_184). *E. faecalis* population in N-DOMW carried a non-synonymous substitution (Asn81Tyr) in *murB* gene involved in peptidoglycan biosynthesis; and a synonymous substitution (Glu269Glu) in a putative LPXTG-motif cell wall anchor domain protein present in pTEF3_66kb plasmid (Table S5B). A significant fraction of the mutations acquired in *E. coli* and *E. faecalis* were absent in populations recovered after 24 h of incubation (Figure S7).

Based on this calculation (coverage of plasmid contig /coverage of chromosome contig), the pTEF3_66kb and pAD1_41kb plasmids were present at 0.96 ± 0.09 and 1.05 ± 0.19 copies per cell.

*E. coli*_{C3000} carried a multi-replicon IncF plasmid of an unknown size and topology. Two contigs were identified (ca. 14 and 16 kb) from the assembled genome by PlasmidFinder (Carattoli et al., 2014) as carrying the initiation of replication (Rep) protein for IncFIC (FII) and IncFIA plasmids, respectively. DNA sequences from these contigs were subjected to BLAST searches against genomes that have been deposited in GenBank. No BLAST hit matched the complete sequence of the 16 kb plasmid contig, though it shared 99% DNA homology with *E. coli* strain C3026 plasmid (accession no: CP014273). Plasmid C3026 is circular and 213924 base pairs long, suggesting that contigs identified by PlasmidFinder in this study represent only a fragment of a longer IncF plasmid. More importantly, IncF plasmids are usually >100 kb in size (Villa et al., 2010). Due to these factors (unknown size and topology), IncF plasmid will not be discussed further in this manuscript.

DISCUSSION

Photodegradation of the DOM of diverse terrigenous sources has been demonstrated to affect the growth and metabolism of bacteria in aquatic ecosystems (Lindell et al., 1995; Scully et al., 2003; Anesio et al., 2005; Lønborg et al., 2013, 2016; Simsek et al., 2013). However, no study to date has provided bacterial transcriptomic or mutational changes occurring after exposure to sunlight irradiated DOM. In this study, we used cattle fecal extract as an agricultural source of DOM to investigate the role of photodegradation on bacterial species used frequently as indicators of fecal contamination; and as control strains in many studies. Based on the maximum concentration and the rapid growth observed in the presence of I-DOMW and N-DOMW, *E. coli* seems to have a greater potential than *E. faecalis* to grow in environments receiving highly bioavailable DOC concentrations (>30 mg l⁻¹ C). *E. faecalis* in this study exhibited a steady state or limited growth, suggesting very different requirements than

E. coli. For instance, Lleó et al. (1998) showed that oligotrophy was the main factor inhibiting divisional capability and activation survival strategies in response to stress caused by low nutrient concentration in *E. faecalis*, *E. faecium*, and *E. hirae*.

Re-growth of *E. coli* in diverse environmental matrices has been reported previously (Sanders et al., 2013; Giannakis et al., 2014, 2015; Harvey et al., 2014; Oladeinde et al., 2014; Oliver et al., 2016). Overnight re-growth has been implicated by high counts of *E. coli* in surface waters in the early morning (Whitman et al., 2004; Desai and Rifai, 2013). Few studies have reported re-growth for *E. faecalis* (Litton et al., 2010; Kim and Wuertz, 2015;

Dubinsky et al., 2016), implying that this bacterium might have different survival strategies in the environment. Importantly, the potential for re-growth may be further limited by the inhibitory role of direct sunlight and production of ROS from DOM photodegradation. Under such adverse environmental conditions, *Enterococcus* spp. may favor activation of the viable, but not culturable (VBNC) state, and the loss of culturability (Lleó et al., 2005).

Here, photodegradation of DOM resulted in a decrease in concentration of DOC, $[NH_4^+]$, $[PO_4^{3-}]$, and an increase in $[NO_3^-]$. Although we did not measure all the photo-oxidation products from this reaction, these data suggest that ROS were produced in the presence of oxygen and may have limited bacterial growth efficiency. Following the irradiation of DOM-spiked water for 12 h, 1O_2 , O_2^- and H_2O_2 were among the extracellular ROS detected. We discarded the inhibitory role of 1O_2 due to the low concentration detected and short half-life (μs). Additionally, following bacteria exposure to 1O_2 generated from Rose Bengal irradiation, there was no difference in concentration between exposed and unexposed controls.

Maraccini et al. (2016) also reported no correlation between bulk-phase steady state concentrations of 1O_2 and exogenous indirect photo-inactivation rate constants. A few studies have reported on the direct role of 1O_2 in bacterial die-off (Dahl et al., 1989; Sassoubre et al., 2012; Glaeser et al., 2014), but these were done at very high steady state concentrations which may not be representative of environmental relevant doses.

On the other hand, the high reactivity and charge of the radicals limit their potential to cross cell barriers and affect vital cell functions (Kieber et al., 2003). Superoxide radicals cannot cross the lipid bilayers at neutral pH, moreover they are immediately scavenged by superoxide dismutase and reductase enzymes produced by many species of bacteria and cyanobacteria (Imlay, 2008). Hydroxyl radicals is the neutral form of hydroxide ion and it is short-lived. They are formed in a series of Fenton reactions involving iron, H_2O_2 and O_2^- as reactants. Hydroxyl

TABLE 4 | Validation of RNA-seq results with qRT-PCR analysis of selected genes at 0.5 h.

Bacteria/Gene	RNA-seq (Log2 Fold change)	qRT-PCR (Log2 Fold change)
<i>E. coli</i>		
<i>katG</i>	4.32	5.63
<i>ahpF</i>	2.84	3.67
<i>oxyS</i>	2.42	4.62
<i>oxyR</i>	0.40	-0.029
<i>E. faecalis</i>		
<i>katA</i>	-1.35	-0.17
<i>ahpC</i>	0.94	-0.23
<i>npr</i>	-0.34	0.16
<i>tpx</i>	-0.066	-0.24
<i>perR</i>	-0.51	0.65
<i>fur</i>	0.76	-1.55
<i>E. coli</i> O157:H7		
<i>katG</i>	3.62	5.27
<i>ahpF</i>	3.61	6.35
<i>oxyR</i>	-0.28	0.20

Boldness denotes genes with significant difference in expression between I-DOMW and N-DOMW ($p_{adj} < 0.001$).

TABLE 5 | Differentially expressed transcripts in plasmids carried by *E. faecalis*.

Locus Tag	Plasmid type	Gene name/Protein domain	0.5 h	6 h	12 h	24 h
			Log2 fold change			
DR75_2888	pTEF3_66kb	LPxTG Gram positive anchor protein	-0.33	-1.15	-0.84	-1.02
DR75_2890	pTEF3_66kb	Hypothetical	-1.08	-1.69	0.86	0.50
DR75_2893	pTEF3_66kb	Bacteriocin ABC transporter (LagD)	-0.19	-1.53	0.35	0.19
DR75_2895	pTEF3_66kb	Thioredoxin	-0.33	-1.52	0.32	0.15
DR75_2896	pTEF3_66kb	Thioredoxin	ND	-2.00	0.30	0.15
DR75_2908	pTEF3_66kb	Replication protein	0.50	1.36	0.11	0.31
DR75_2932	pTEF3_66kb	Fic/Doc family	-0.12	-0.82	0.21	-0.36
DR75_2933	pTEF3_66kb	Single strand DNA binding protein (ssb)	-0.01	-1.11	-0.01	-0.52
DR75_2934	pTEF3_66kb	putative membrane protein	-0.21	-1.37	0.16	0.02
DR75_2953	pTEF3_66kb	cylL	-0.36	0.71	0.75	0.23
DR75_2989	pAD1_41kb	LagD	-2.10	-0.48	0.30	0.09
DR75_2991	pAD1_41kb	Bacteriocin lic	-1.23	-2.44	0.63	0.26

Boldness denotes significant differentially expressed genes in I-DOMW relative to N-DOMW (p value < 0.05). ND, not determined.




FIGURE 6 | Circular map of *E. faecalis* plasmid pTEF3_66kb (CP008815). Genes differentially expressed are highlighted in purple rectangular boxes and genes with mutations are highlighted in red rectangular boxes.


radicals are the most damaging of all ROS, and can cause nucleic acid mutations, lipid peroxidation and amino-acid and protein oxidative damage (Imlay, 2013, 2015a,b). However, the low concentration (< limit of detection) in this study and observed in surface waters (10^{-19} – 10^{-16} M) (Gligorovski et al., 2015) suggest their extracellular presence is not critical to fecal bacteria survival. For example, Maraccini et al. (2016) reported no correlation between measured exogenous OH⁻ and fecal bacteria inactivation rates in a laboratory-controlled experiment.

The level of H₂O₂ produced from DOM photodegradation in this study was in the micromolar range and was stable for more than 24 h (Figure S3). H₂O₂ production from DOM photodegradation and microbial processes play a significant role in controlling bacterial survival dynamics in environmental waters (Mostofa et al., 2013; Cory et al., 2016; Zhang et al., 2016). Anesio et al. (2005) showed that the suppression of bacterial carbon production was highly correlated with the concentration of photochemically formed H₂O₂ and concluded that extracellular H₂O₂ concentrations of about 2 to 3 μM were inhibitory for bacteria. H₂O₂ can cross bacterial cell membranes and can persist in surface waters for several hours (Imlay, 2008; Mostofa et al., 2013). In this study, *E. coli* efficiently scavenged

photo-produced H₂O₂ within 6 h of dark incubation, while there was an accumulation of H₂O₂ in the *E. faecalis* treatment. This observed buildup suggests this bacterium differs in its production and detoxification of H₂O₂ (Marsico et al., 2015). Moreover, *E. faecalis* produces H₂O₂ as a by-product of aerobic glycerol metabolism, which could have contributed to the observed increase in I-DOMW treatment (Bizzini et al., 2010).

H₂O₂ Photoproduction and OxyR-Response Genes in *E. coli* and *E. coli* O157:H7

The OxyR-regulon in *E. coli* is responsible for sensing oxidative stress and positively regulates the induction of several genes including *katG* and *ahpCF* (Zheng et al., 2001; Ravindra Kumar and Imlay, 2013). Mutants lacking either *katG* or *ahp* have been shown to grow poorly or not at all as laboratory cultures (Khademian and Imlay, 2017). Peroxidase genes have the important role of keeping the steady-state concentration of H₂O₂ at only 50 nM. This low steady state is necessary since H₂O₂ reacts rapidly with vulnerable intracellular enzymes. After bacterial inoculation into I-DOMW, we observed an immediate increase (~0.5 h) in expression of *katG* and *ahpCF* genes (>



2-FC), followed by a concomitant decrease in H_2O_2 . After 6 h of dark incubation, there was ca. 95% reduction in measured extracellular H_2O_2 for *E. coli* and *E. coli* O157:H7.

This does not preclude the presence of hydrogenase encoding genes that can tolerate high oxygen levels and ameliorate H_2O_2 induced oxidative stress (Tremblay and Lovley, 2012). For instance, the beta-subunit Ni-Fe hydrogenase protein (HyaB) was identified with a mutation (Val410Ala) in I-DOMW populations after 12 h of dark incubation. This gene was also upregulated (1.85-FC; p .value = 0.008) in I-DOMW compared to N-DOMW at 12 h. More studies will be required to correlate this non-synonymous substitution to increased H_2O_2 tolerance.

Next, we compared transcripts in *E. faecalis* with a mechanism of action proposed for OxyR by others (Verneuil et al., 2004). Contrary to expectations, peroxidase (*npr*, *ahpC*, *tpx*) and thioredoxin (*trx*) genes in *E. faecalis* were not significantly expressed in I-DOMW following inoculation, despite their

relevant role in peroxide metabolism (Figure 4B). La Carbona et al. (2007) demonstrated that mutations in these genes can limit survival of *E. faecalis* exposed to exogenously added H_2O_2 . In contrast, the heme-catalase (*katA*) transcript, which plays only a partial role in protecting *E. faecalis* against the toxic effect of externally-added H_2O_2 (Baureder et al., 2012), was significantly upregulated (1.5-FC; p .adj < 0.01) in I-DOMW at 6 h (Figure 4B). Further, extracellular H_2O_2 concentration increased by 170% after 6 h of dark incubation in I-DOMW, with no significant reduction after 24 h (Figure 2B).

This accumulation of H_2O_2 by *Enterococcus* has been demonstrated in several studies (Moy et al., 2004; La Carbona et al., 2007; Bizzini et al., 2010; Baureder et al., 2012). La Carbona et al. (2007) showed that extracellular H_2O_2 concentration increased with all carbon sources used for growth, reaching $350 \mu M$ in some strains. Further, they reported no difference in growth rate and external H_2O_2 (10–20 μM) concentrations between wild type and peroxidase-deleted mutants grown on glucose. A possible explanation could be a requirement for an intracellular peroxide concentration threshold before these genes are activated or highly expressed.

To our knowledge, there is no reported H_2O_2 threshold for *E. faecalis*. La Carbona et al. (2007) used >5 mM H_2O_2 in their study with *E. faecalis* peroxidase-deletion mutants. Yan et al. (2015) also reported less than a 2-fold change in expression of oxidative stress genes when exposed to 1.5–2 mM H_2O_2 . These concentrations are ~1000-fold higher than the external levels required for OxyR-response in *E. coli*, and ~100-fold higher than the level produced in this study. It is therefore plausible that in this study, H_2O_2 generated inside *E. faecalis* cells was more deleterious than the low concentrations produced exogenously from DOM irradiation. Although a decrease in concentration was observed in I-DOMW treatments at 6 and 24 h (Figure 2B), *E. faecalis* population was the same in I-DOMW and N-DOMW after 48 h (data not shown).

The ability for *E. coli* and *E. coli* O157:H7 to efficiently induce oxidative stress genes in response to ambient levels of ROS may offer them an advantage in the environment. Upon exposure to photo-produced H_2O_2 , these bacteria are able to remove significant concentrations of H_2O_2 and repair DNA lesions within hours, with a potential to grow overnight if necessary substrates are available. Morris et al. (2016), using diel metatranscriptomic data from five published marine studies spanning a variety of open ocean sites, showed that the abundance of transcripts with catalase and peroxidase activity peaked in the late afternoon, coinciding with the highest concentration of H_2O_2 . Dark production of H_2O_2 , presumably through the accidental autoxidation of redox enzymes, was observed among all three bacteria in this study. Studies have shown that *E. coli* generates about 10 to 15 $\mu M s^{-1}$ of endogenous H_2O_2 during growth in air-saturated glucose medium (Seaver and Imlay, 2004). Intracellular H_2O_2 will accumulate in a closed system by flowing out of the cell rather than into the cytoplasm (Ravindra Kumar and Imlay, 2013). In natural oxic ecosystems (e.g., streams, rivers, and ponds), excreted H_2O_2 would not build up, but would be lost to the environment.

This might not be the case for *E. faecalis*; this study shows that while *E. faecalis* can tolerate micromolar levels of exogenous H_2O_2 , it is unable to reach maximum growth potential. Importantly, such levels of H_2O_2 may further limit recovery in the environment; therefore, *E. faecalis* role in H_2O_2 concentration fluctuations in surface waters may be restricted to production.

Glycerol Metabolism Is Associated With Extracellular H_2O_2 Production in *E. faecalis*

The major structural barrier in *E. faecalis* is its peptidoglycan wall that is anchored by teichoic acids (TA) and lipoteichoic acids (LTA). TA and LTA are polymers of poly-glycerol phosphate that are joined by phosphodiester linkages (Brown et al., 2013). They provide several functions, including scavenging of cations, phosphate reservoir, cell envelope adhesiveness, and immunogenicity. Glycerol metabolism is important for synthesis of glycerol phosphate in a reaction involving glycerol facilitator protein (*glpF*) and phosphorylated glycerol kinase (*glpK*). The phosphoenolpyruvate (PEP): carbohydrate phosphotransferase system (PTS) is responsible for the phosphorylation of *glpK* (Bizzini et al., 2010; Ramsey et al., 2014). Many of the proteins in the PEP: PTS are involved in transport of sugars including lactose, sorbose, mannose, sucrose, cellobiose, and fructose across the cell membrane. The PTS in *E. faecalis* is also involved in inducer expulsion, inducer exclusion, and catabolite repression (Saier et al., 1996). Hydrogen peroxide production is controlled by glycerol-3-P oxidase (*glpO*) via oxidation of glycerol phosphate to dihydroxyacetone phosphate, an intermediate of the glycolytic pathway (Ramsey et al., 2014). This enzyme uses molecular oxygen as the electron sink, that leads to formation of H_2O_2 (Figure 8).

Immediately after inoculation, these glycerolipid transcripts (*glpF*, *glpO*, *glpK*, and PEP: PTS enzymes) were significantly upregulated in N-DOMW (compared to I-DOMW) for up to 12 h (Figures 3B, 5), however, extracellular H_2O_2 concentration did not significantly change for the duration of our experiment in N-DOMW (Figure 2). This suggests, that *E. faecalis* can tolerate moderate levels of H_2O_2 and persist or grow under such conditions. In contrast, H_2O_2 increased for 12 h in I-DOMW which indicates the mechanism for extracellular H_2O_2 production/detoxification may differ between I-DOMW and N-DOMW.

This increase in H_2O_2 in I-DOMW may be due to the activity of *nox*. NADH oxidases (*nox*) use molecular oxygen as an electron acceptor to regenerate NAD^+ from NADH. Regeneration of NADH by *nox* allows bacteria to use mixed acid fermentation pathways when grown in the presence of molecular oxygen (Moy et al., 2004). Two categories of NADH oxidases are present in most bacteria—one produces water and the other produces H_2O_2 . In *E. faecium*, mutation in a Δnox gene eliminated nearly all NADH oxidase activity and reduced H_2O_2 production. *Nox* has been shown to be considerably stable in the presence of H_2O_2 (Villegas and Gilliland, 1998; Marty-Teyssset et al., 2000; Jiang and Bommarius, 2004). Jiang and Bommarius (2004) experimentally determined the inhibition rate constant for H_2O_2 was 12 mM for *Lactococcus lactis*, another lactic acid bacteria. In this study, the expression of the *nox* gene was significantly higher (1.8 -FC, $p_{adj} < 0.001$) in I-DOMW than N-DOMW and was only observed at 6 h (Figure 4B). Further, H_2O_2 concentration at 6 h was ca. $42 \mu M$, which is three orders of magnitude lower than the required threshold for *nox* inhibition. Although this may explain the increased H_2O_2 production observed in I-DOMW, we cannot rule out the possibility that additional H_2O_2 -producing genes or metabolic pathways could also be involved (Condon, 1987; Hertzberger et al., 2014).

A negative feedback mechanism in response to the higher external H_2O_2 concentration resulting from the I-DOMW treatment may explain the lack of *E. faecalis* growth in the irradiated treatment. A reduction in the expression of *glpK* and *glpF* will restrict the active transport of carbohydrates into the cell and reduce H_2O_2 formed via glycerol phosphate oxidation. Although this approach will protect the cell from additional H_2O_2 build up, it can limit *E. faecalis*'s ability to survive or grow in I-DOMW, which is consistent with the results observed after 24 h of incubation in this treatment.

The sensitivity of *E. faecalis* to light-produced ROS has been reported (Kadir and Nelson, 2014; Nguyen et al., 2015; Maraccini et al., 2016), but its mechanism of survival has not been clearly described. In this study, *E. faecalis* decreased the expression of several proteins involved in glycerol metabolism and transport of carbohydrates, after inoculation into I-DOMW. A consequence of such response was limited growth and accumulation of H_2O_2 in the growth medium. Although, *E. faecalis* represents only a species of the genus *Enterococcus*, these results support the opinion that enterococci have a low re-growth potential in environmental waters.




FIGURE 8 | Pathway for aerobic glycerol metabolism. Modified from Ramsey et al. (2014).

Conjugative Plasmids of *E. faecalis* and Associated Fitness Cost

Plasmids are extra-chromosomal genetic elements coding for a wide range of traits that allow bacteria to adapt to different environmental stressors and can spread horizontally among bacteria by conjugation. However, carriage of multiple plasmids in a host cell has been demonstrated to impose a fitness cost and their long-term stability remains questionable (MacLean and San Millan, 2015). The *E. faecalis* strain used here, carries two low copy conjugative plasmids—pTEF3_66kb and pAD1_41kb; and both encode accessory proteins for class II bacteriocin production (CylL) and a plasmid maintenance system (Par; Fic/Doc, RelB). In addition, pTEF3_66kb carries two thioredoxin peroxidases (*trx*) associated with H₂O₂ removal.

Based on the discrepancy in coverage between RNA-seq (mean = 8.3 X) and WGS (mean = 80 X) reads mapping to reference plasmid genomes, these data suggests that plasmid-carrying cells maybe low in I-DOMW and N-DOMW populations. The majority of differentially expressed plasmid-encoded transcripts (>80%) belong to pTEF3_66kb, which implies that there may be a selection pressure for this plasmid (Figure 6). Furthermore, the coverage determined for pTEF3_66kb (10 ± 7.7 X) was significantly higher than the coverage for pAD1_41kb (6.6 ± 4.3 X) (Table S6, Wilcoxon rank-signed test; *p*-value = 0.01), supporting the notion that *E. faecalis* selected for it.

Transcripts differentially expressed in pTEF3_66kb were upregulated in N-DOMW compared to I-DOMW which was not expected, since H₂O₂ levels were higher in I-DOMW (Table 5). For example, *trx* genes were among the expressed transcripts in N-DOMW after 0.5 h of incubation. A plausible explanation is that pTEF3_66kb *trx* enzyme is more active toward endogenously generated H₂O₂ rather than exogenously added H₂O₂.

The only mutation identified in *E. faecalis* plasmids was in a putative gene encoding LPxTG cell anchor protein of pTEF3_66kb. This synonymous mutation was observed in N-DOMW populations after 0.5 and 12 h of incubation. Additionally, this gene had significantly higher expression in N-DOMW compared to I-DOMW after 6, 12, and 24 h of incubation (Table 5). This result suggests that codon bias may be in play—a phenomenon in which an organism prefers a different set of codons over others. Codon bias plays an important role in controlling a multitude of cellular processes ranging from fine-tuning gene expression to protein folding (Quax et al., 2015). LPxTG cell anchor proteins have been shown to participate in aggregation substance formation in enterococci and are important for conjugative transfer of virulent plasmids (Hendrickx et al., 2009). Here, this mutation may have contributed to the continuous expression of this gene, which could subsequently increase conjugation rates in N-DOMW populations.

Extracellular ROS Signaling Contributes to Adaptive Response

Environmental cues including non-lethal doses of ROS, antimicrobials, nutrient limitation, and temperature can make bacteria alter their transcriptome rapidly (Kyle et al., 2010;

Fernández and Hancock, 2012; Suzuki et al., 2014). Such responses can prolong their survival and enhance resistance to higher doses of the same stressor or other stressors (Dwyer et al., 2014; Djorić and Kristich, 2015). Such adaptive resistance is believed to be transient and usually reverts upon removal of the inducing agent (Fernández and Hancock, 2012).

In this study, multiple genetic networks associated with virulence, quorum sensing, and antibiotic resistance were up-regulated in I-DOMW for *E. coli* O157:H7. These differentially expressed genes included significant increases in genes for outer membrane receptors that facilitate import of iron-chelating siderophores and iron from host organisms which are important for *E. coli* O157:H7 pathogenicity (Figure 3C, Supplementary File 3) (Hagan, 2009). The tryptophan operon is a repressor operon that is turned on or off, based on levels of tryptophan in the environment. Tryptophan is also the primary source of indole production in *E. coli*, an organic compound that plays a role in quorum sensing, biofilm formation and antibiotic resistance (Hu et al., 2010; Kuczynska-Wiśnik et al., 2010). Kuczynska-Wiśnik et al. (2010) showed that the addition of dimethyl sulfoxide (H₂O₂ scavenger) partly restored *E. coli* biofilm formation in the presence of antibiotics and decreased indole production.

Transcripts for tryptophan biosynthesis (*trpEDCBA*) increased significantly following inoculation (0.5 h) into I-DOMW and were downregulated within 6 h (Figure 3C and Figure S4). In addition, extracellular H₂O₂ concentration in I-DOMW at 0.5 h and 6 h was 15.4 ± 0.81 and 1.34 ± 0.26 (SE) μ M, respectively (Figure 2), which was consistent with the gene regulation observed. These results indicate that indole production may be required for *E. coli* O157: H7 survival under increased extracellular peroxide concentration and, consequently, may affect this pathogen's biofilm formation and antibiotic resistance capability.

Curli is a proteinaceous extracellular matrix associated with attachment and biofilm formation, and plays a major role in bacterial pathogenesis (Normark et al., 1992). Moreover, biofilm formation has been associated with oxidative stress, including H₂O₂ exposure (Boles and Singh, 2008; Geier et al., 2008; Fink et al., 2012). Extracellular H₂O₂ measured at 6 h was higher in N-DOMW than I-DOMW for *E. coli* (4.58 ± 0.71 μ M vs. < LOQ) (Figure 2A). Further, the expression of *csgAB* genes encoding curli/amyloid fibers increased significantly in N-DOMW after 6 h of dark incubation (Figure S6). This provides supporting evidence that *E. coli* may increase biofilm formation in response to endogenously-produced H₂O₂ (Jang et al., 2016).

The expression of genes, coding for porins, and efflux pumps, are important in adaptive resistant development (Fernández and Hancock, 2012; Motta et al., 2015). They are efficiently regulated to respond to specific cues, thereby changing the resistance of a bacterium based on growth conditions. For example, Suzuki et al. (2014) demonstrated that antibiotic resistance development in *E. coli* could be quantitatively predicted by the expression changes of a small number of genes. Oligopeptide (*opp*) genes encoding sex pheromones (EF0063, EF1513) and peptide transport (EF0909) significantly increased in *E. faecalis* after 6 h of dark incubation in I-DOMW (Figure 3B). These genes have important roles to play in beta-lactamase resistance and quorum sensing in *E. faecalis*, and their induction in response

to H₂O₂. In the present study, time point 6 h corresponded with a 170% increase in extracellular H₂O₂ (Figure 2B), which could have produced the response to oxidative stress observed in the oligopeptide genes. The potential for sublethal levels of ROS to increase bacterial minimum inhibitory concentration (MIC) to antibiotics requires more research. Dwyer et al. (2014) reported that pretreatment of *E. coli* cells with 1 or 5 mM H₂O₂ for 15 min did not induce any lethality or growth inhibition, however, it resulted in a transient 1-log protection of cells from antibiotic killing.

Genes associated with quorum sensing (QS) were significantly upregulated in *E. coli* (Figure 3A, Figure S5). QS is a form of regulation of gene expression used by a majority of bacteria in response to fluctuations in cell population density. Quorum sensing bacteria produce and release chemical signal molecules called autoinducers that increase in concentration as a function of cell density. Gram-positive and Gram-negative bacteria use QS communication signals to regulate an array of physiological activities (Rutherford and Bassler, 2012). The level of auto-inducer-2 (AI-2) produced extracellularly varies depending with growth conditions, and its transport is mediated by *lsrACDBFGE* operon. In *E. coli*, expression of *lsrBFG* transcripts significantly increased in I-DOMW after 6 h (Figure S5) which corresponds to the time point with the highest growth rate (Figure 2A). Further, upregulation of QS genes may be influenced by H₂O₂ exposure; for instance, Yu et al. (2013) showed that Δ LuxS *Yersinia pestis* mutants (LuxS regulates *lsr*) were more sensitive to killing by H₂O₂ than their wildtype.

The survival dynamics shown by pure cultures of bacteria under controlled and sterile conditions in this study could differ in the environment owing to other abiotic factors (e.g., temperature) and biotic factors including predation (“top-down control”) and/or competition (“bottom-up control”) (de Brauwere et al., 2014; Rochelle-Newall et al., 2015). Several studies have suggested that antagonistic agents and indigenous microbiota are important to the survival of indicator bacteria (Feng et al., 2010; Wanjugi and Harwood, 2013; Korajkic et al., 2014). Furthermore, it is a possibility that other free radicals, such as peroxy radicals and reactive nitrogen species, could have triggered the oxidative stress gene expression exhibited by the bacteria used in this study. Our results suggests that to better describe, understand or model the dynamics of FIB

concentration in environmental systems, it is necessary to integrate the effect of abiotic factors, such as the ones discussed herein, with the biotic factors described by others.

AUTHOR CONTRIBUTIONS

AO was involved in the planning, experimental design, discussion of results, analysis of data, and manuscript writing. MM was involved in the planning, discussion of results, analysis of data, and manuscript editing. RM was involved in the planning, discussion of results, and manuscript editing. EL was involved in the planning, discussion of results, and manuscript editing. C-YC was involved with solar simulation analysis, reactive oxygen species measurement and manuscript editing. TG was involved in preparation of transcriptomics library and bioinformatics analysis. KC contributed to the whole genome sequencing methodology used in this study and also read, revised, and approved submitted manuscript.

ACKNOWLEDGMENTS

We are grateful to Kyler Herrington and Troy Kieran for their help with sample analysis. We thank Roger Burke for assistance with nutrient analyses. We also appreciate Dr. Richard Zepp for valuable advice during the solar simulation experiments and Mary Katherine Crews and Fran Rauschenberg for manuscript review. We thank the Kennedy family at Covenant Valley Farm and the Nixon family at Cane Creek Farm, Inc., for allowing us access to their cattle. This work was supported by the Environmental Protection Agency (EPA) Star Fellowship [FP-91766701 to AO] and EPA's Office of Research and Development. Any opinions expressed in this paper are those of the authors and do not necessarily reflect the official positions and policies of the U.S. EPA and any mention of products or trade names does not constitute recommendation for use. The authors declare no competing commercial interests in relation to the submitted work.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00882/full#supplementary-material>

REFERENCES

- Adams, M. H. (1959). *Bacteriophages*. London: Interscience.
- Anesio, A. M., Graneli, W., Aiken, G. R., Kieber, D. J., and Mopper, K. (2005). Effect of humic substance photodegradation on bacterial growth and respiration in lake water. *Appl. Environ. Microbiol.* 71, 6267–6275. doi: 10.1128/AEM.71.10.6267-6275.2005
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., et al. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* 19, 455–477. doi: 10.1089/cmb.2012.0021
- Baureder, M., Reimann, R., and Hederstedt, L. (2012). Contribution of catalase to hydrogen peroxide resistance in *Enterococcus faecalis*. *FEMS Microbiol. Lett.* 331, 160–164. doi: 10.1111/j.1574-6968.2012.02567.x
- Bida, M. R., Tyler, A. C., and Pagano, T. (2015). Quantity and composition of stream dissolved organic matter in the watershed of Conesus Lake, New York. *J. Great Lakes Res.* 41, 730–742. doi: 10.1016/j.jglr.2015.06.003
- Bizzini, A., Zhao, C., Budin-Verneuil, A., Sauvageot, N., Giard, J.-C., Auffray, Y., et al. (2010). Glycerol is metabolized in a complex and strain-dependent manner in *Enterococcus faecalis*. *J. Bacteriol.* 192, 779–785. doi: 10.1128/JB.00959-09
- Boles, B. R., and Singh, P. K. (2008). Endogenous oxidative stress produces diversity and adaptability in biofilm communities. *Proc. Natl. Acad. Sci. U.S.A.* 105, 12503–12508. doi: 10.1073/pnas.0801499105
- Bradford, S. A., Morales, V. L., Zhang, W., Harvey, R.W., Packman, A. I., Mohanram, A., et al. (2013). Transport and fate of microbial pathogens in agricultural settings. *Crit. Rev. Environ. Sci. Technol.* 43, 775–893. doi: 10.1080/10643389.2012.710449

- Bradford, S. A., Schijven, J., and Harter, T. (2015). Microbial transport and fate in the subsurface environment: introduction to the special section. *J. Environ. Qual.* 44, 1333–1337. doi: 10.2134/jeq2015.07.0375
- Brown, S., Santa Maria, J. P. Jr., and Walker, S. (2013). Wall teichoic acids of gram-positive bacteria. *Annu. Rev. Microbiol.* 67, 313–336. doi: 10.1146/annurev-micro-092412-155620
- Burns, J. M., Cooper, W. J., Ferry, J. L., King, D. W., DiMento, B. P., McNeill, K., et al. (2012). Methods for reactive oxygen species (ROS) detection in aqueous environments. *Aquat. Sci.* 74, 683–734. doi: 10.1007/s00027-012-0251-x
- Bushaw, K. L., and Zepp, R. G. (1996). Photochemical release of biologically available nitrogen from aquatic dissolved organic matter. *Nature* 381, 404–407. doi: 10.1038/381404a0
- Carattoli, A., Zankari, E., Garcia-Fernandez, A., Larsen, M. V., Lund, O., Villa, L., et al. (2014). *In silico* detection and typing of plasmids using plasmidfinder and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* 58, 3895–3903. doi: 10.1128/AAC.02412-14
- Chen, C.-Y., and Jafvert, C. T. (2010). Photoreactivity of carboxylated single-walled carbon nanotubes in sunlight: reactive oxygen species production in water. *Environ. Sci. Technol.* 44, 6674–6679. doi: 10.1021/es101073p
- Chu, C., Erickson, P. R., Lundeen, R. A., Stamatelatos, D., Alaimo, P. J., Latch, D. E., et al. (2016). Photochemical and nonphotochemical transformations of cysteine with dissolved organic matter. *Environ. Sci. Technol.* 50, 6363–6373. doi: 10.1021/acs.est.6b01291
- Clark, C. D., De Bruyn, W. J., Jakubowski, S. D., and Grant, S. B. (2008). Hydrogen peroxide production in marine bathing waters: implications for fecal indicator bacteria mortality. *Mar. Pollut. Bull.* 56, 397–401. doi: 10.1016/j.marpolbul.2007.10.017
- Condon, S. (1987). Responses of lactic acid bacteria to oxygen. *FEMS Microbiol. Lett.* 46, 269–280. doi: 10.1111/j.1574-6968.1987.tb02465.x
- Cory, R., Davis, T., Dick, G., Johengen, T., Denef, V., Berry, M., et al. (2016). Seasonal dynamics in dissolved organic matter, hydrogen peroxide, and cyanobacterial blooms in lake erie. *Front. Mar. Sci.* 3:54. doi: 10.3389/fmars.2016.00054
- Dahl, T. A., Midden, W., and Hartman, P. E. (1989). Comparison of killing of gram-negative and gram-positive bacteria by pure singlet oxygen. *J. Bacteriol.* 171, 2188–2194. doi: 10.1128/jb.171.4.2188-2194.1989
- de Brauwere, A., Ouattara, N. K., and Servais, P. (2014). Modeling fecal indicator bacteria concentrations in natural surface waters: a review. *Crit. Rev. Environ. Sci. Technol.* 44, 2380–2453. doi: 10.1080/10643389.2013.829978
- Desai, A. M., and Rifai, H. S. (2013). *Escherichia coli* concentrations in urban watersheds exhibit diurnal sag: implications for water-quality monitoring and assessment. *JAWRA J. Am. Water Resour. Assoc.* 49, 766–779. doi: 10.1111/jawr.12039
- Djaman, O., Outten, F. W., and Imlay, J. A. (2004). Repair of oxidized iron-sulfur clusters in *Escherichia coli*. *J. Biol. Chem.* 279, 44590–44599. doi: 10.1074/jbc.M406487200
- Djorić, D., and Kristich, C. J. (2015). Oxidative stress enhances cephalosporin resistance of *Enterococcus faecalis* through activation of a two-component signaling system. *Antimicrob. Agents Chemother.* 59, 159–169. doi: 10.1128/AAC.03984-14
- Dubbs, J. M., and Mongkolsuk, S. (2007). Peroxiredoxins in bacterial antioxidant defense. *Subcell. Biochem.* 44, 143–193. doi: 10.1007/978-1-4020-6051-9_7
- Dubinsky, E. A., Butkus, S. R., and Andersen, G. L. (2016). Microbial source tracking in impaired watersheds using PhyloChip and machine-learning classification. *Water Res.* 105, 56–64. doi: 10.1016/j.watres.2016.08.035
- Dwyer, D. J., Belenky, P. A., Yang, J. H., MacDonald, I. C., Martell, J. D., Takahashi, N., et al. (2014). Antibiotics induce redox-related physiological alterations as part of their lethality. *Proc. Natl. Acad. Sci. U.S.A.* 111, E2100–E2109. doi: 10.1073/pnas.1401876111
- Feng, F., Goto, D., and Yan, T. (2010). Effects of autochthonous microbial community on the die-off of fecal indicators in tropical beach sand. *FEMS Microbiol. Ecol.* 74, 214–225. doi: 10.1111/j.1574-6941.2010.00921.x
- Fernández, L., and Hancock, R. E. (2012). Adaptive and mutational resistance: role of porins and efflux pumps in drug resistance. *Clin. Microbiol. Rev.* 25, 661–681. doi: 10.1128/CMR.00043-12
- Fink, R. C., Black, E. P., Hou, Z., Sugawara, M., Sadowsky, M. J., and Diez-Gonzalez, F. (2012). Transcriptional responses of *Escherichia coli* K-12 and O157: H7 associated with lettuce leaves. *Appl. Environ. Microbiol.* 78, 1752–1764. doi: 10.1128/AEM.07454-11
- Fratamico, P. M., Deng, M. Y., Strobaugh, T. P., and Palumbo, S. A. (1997). Construction and characterization of *Escherichia coli* O157: H7 strains expressing firefly luciferase and green fluorescent protein and their use in survival studies. *J. Food Protect.* 60, 1167–1173. doi: 10.4315/0362-028X-60.10.1167
- Geier, H., Mostowy, S., Cangelosi, G. A., Behr, M. A., and Ford, T. E. (2008). Autoinducer-2 triggers the oxidative stress response in *Mycobacterium avium*, leading to biofilm formation. *Appl. Environ. Microbiol.* 74, 1798–1804. doi: 10.1128/AEM.02066-07
- Giannakis, S., Darakas, E., Escalas-Cañellas, A., and Pulgarin, C. (2014). Elucidating bacterial regrowth: effect of disinfection conditions in dark storage of solar treated secondary effluent. *J. Photochem. Photobiol. A Chem.* 290, 43–53. doi: 10.1016/j.jphotochem.2014.05.016
- Giannakis, S., Darakas, E., Escalas-Cañellas, A., and Pulgarin, C. (2015). Solar disinfection modeling and post-irradiation response of *Escherichia coli* in wastewater. *Chem. Eng. J.* 281, 588–598. doi: 10.1016/j.cej.2015.06.077
- Glaeser, S. P., Berghoff, B. A., Stratmann, V., Grossart, H.-P., and Glaeser, J. (2014). Contrasting effects of singlet oxygen and hydrogen peroxide on bacterial community composition in a humic lake. *PLoS ONE* 9:e92518. doi: 10.1371/journal.pone.0092518
- Glenn, T., Nilsen, R., and Kieran, T. (2016). Adapterama I: Universal stubs and primers for thousands of dual-indexed Illumina libraries (iTru & iNext). *bioRxiv* 049114. doi: 10.1101/049114
- Gligorovski, S., Strekowski, R., Barbati, S., and Vione, D. (2015). Environmental implications of hydroxyl radicals (\bullet OH). *Chem. Rev.* 115, 13051–13092. doi: 10.1021/cr500310b
- Graeber, D., Boëchat, I. G., Encina-Montoya, F., Esse, C., Gelbrecht, J., Goyenola, G., et al. (2015). Global effects of agriculture on fluvial dissolved organic matter. *Sci. Rep.* 5:16328. doi: 10.1038/srep16328
- Häder, D.-P., Helbling, E., Williamson, C., and Worrest, R. (2011). Effects of UV radiation on aquatic ecosystems and interactions with climate change. *Photochem. Photobiol. Sci.* 10, 242–260. doi: 10.1039/c0pp90036b
- Häder, D.-P., Williamson, C. E., Wängberg, S.-Å., Rautio, M., Rose, K. C., Gao, K., et al. (2015). Effects of UV radiation on aquatic ecosystems and interactions with other environmental factors. *Photochem. Photobiol. Sci.* 14, 108–126. doi: 10.1039/C4PP90035A
- Hagan, E. C. (2009). *Iron Acquisition by Uropathogenic Escherichia coli: ChuA and Hma heme Receptors as Virulence Determinants and Vaccine Targets*. University of Michigan.
- Harvey, W. H., Underwood, J. I., Lisle, J., Metge, D. W., and Aiken, G. (2014). “Role of surface water dissolved organic carbon in the survival, growth, and transport of *Escherichia coli* in a deep limestone aquifer in South Florida,” in *U.S. Geological Survey Karst Interest Group Proceedings*, eds E.L. Kuniansky and L.E. Spanler (Carlsbad, New Mexico: U.S. Geological Survey Scientific Investigations Report), 129–132.
- Heinz, M., Graeber, D., Zak, D., Zwirnmann, E., Gelbrecht, J., and Pusch, M. T. (2015). Comparison of organic matter composition in agricultural versus forest affected headwaters with special emphasis on organic nitrogen. *Environ. Sci. Technol.* 49, 2081–2090. doi: 10.1021/es505146h
- Hendrickx, A. P., Willems, R. J., Bonten, M. J., and van Schaik, W. (2009). LPxTG surface proteins of enterococci. *Trends Microbiol.* 17, 423–430. doi: 10.1016/j.tim.2009.06.004
- Hertzberger, R., Arents, J., Dekker, H. L., Pridmore, R. D., Gysler, C., Kleerebezem, M., et al. (2014). H₂O₂ production in species of the *Lactobacillus acidophilus* group: a central role for a novel NADH-dependent flavin reductase. *Appl. Environ. Microbiol.* 80, 2229–2239. doi: 10.1128/AEM.04272-13
- Hu, M., Zhang, C., Mu, Y., Shen, Q., and Feng, Y. (2010). Indole affects biofilm formation in bacteria. *Indian J. Microbiol.* 50, 362–368. doi: 10.1007/s12088-011-0142-1
- Imlay, J. A. (2008). Cellular defenses against superoxide and hydrogen peroxide. *Annu. Rev. Biochem.* 77, 755–776. doi: 10.1146/annurev-biochem.77.061606.161055
- Imlay, J. A. (2013). The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nat. Rev. Microbiol.* 11, 443–454. doi: 10.1038/nrmicro3032

- Imlay, J. A. (2015a). Diagnosing oxidative stress in bacteria: not as easy as you might think. *Curr. Opin. Microbiol.* 24, 124–131. doi: 10.1016/j.mib.2015.01.004
- Imlay, J. A. (2015b). Transcription factors that defend bacteria against reactive oxygen species. *Annu. Rev. Microbiol.* 69, 93–108. doi: 10.1146/annurev-micro-091014-104322
- Jang, I.-A., Kim, J., and Park, W. (2016). Endogenous hydrogen peroxide increases biofilm formation by inducing exopolysaccharide production in *Acinetobacter oleivorans* DR1. *Sci. Rep.* 6:21121. doi: 10.1038/srep21121
- Jiang, R. R., and Bommarius, A. S. (2004). Hydrogen peroxide-producing NADH oxidase (nox-1) from *Lactococcus lactis*. *Tetrahedron Asymmetry* 15, 2939–2944. doi: 10.1016/j.tetasy.2004.07.057
- Kadir, K., and Nelson, K. L. (2014). Sunlight mediated inactivation mechanisms of *Enterococcus faecalis* and *Escherichia coli* in clear water versus waste stabilization pond water. *Water Res.* 50, 307–317. doi: 10.1016/j.watres.2013.10.046
- Khademian, M., and Imlay, J. A. (2017). *Escherichia coli* cytochrome c peroxidase is a respiratory oxidase that enables the use of hydrogen peroxide as a terminal electron acceptor. *Proc. Natl. Acad. Sci. U.S.A.* 114, E6922–E6931. doi: 10.1073/pnas.1701587114
- Kieber, D. J., Peake, B. M., and Scully, N. M. (2003). “Reactive oxygen species in aquatic ecosystems,” in *UV Effects in Aquatic Organisms and Ecosystems*, Vol. 1, eds E. W. Helbling and H. Zagarese (London, UK: Royal Society of Chemistry), 251–288.
- Kim, M., and Wuertz, S. (2015). Survival and persistence of host-associated Bacteroidales cells and DNA in comparison with *Escherichia coli* and *Enterococcus* in freshwater sediments as quantified by PMA-qPCR and qPCR. *Water Res.* 87, 182–192. doi: 10.1016/j.watres.2015.09.014
- Korajkic, A., McMinn, B. R., Shanks, O. C., Sivaganesan, M., Fout, G. S., and Ashbolt, N. J. (2014). Biotic interactions and sunlight affect persistence of fecal indicator bacteria and microbial source tracking genetic markers in the upper Mississippi river. *Appl. Environ. Microbiol.* 80, 3952–3961. doi: 10.1128/AEM.00388-14
- Kosaka, K., Yamada, H., Matsui, S., Echigo, S., and Shishida, K. (1998). Comparison among the methods for hydrogen peroxide measurements to evaluate advanced oxidation processes: application of a spectrophotometric method using copper (II) ion and 2, 9-dimethyl-1, 10-phenanthroline. *Environ. Sci. Technol.* 32, 3821–3824. doi: 10.1021/es9800784
- Kuczynska-Wiśnik, D., Matuszewska, E., Furmanek-Blaszczak, B., Leszczynska, D., Grudowska, A., Szczepaniak, P., et al. (2010). Antibiotics promoting oxidative stress inhibit formation of *Escherichia coli* biofilm via indole signalling. *Res. Microbiol.* 161, 847–853. doi: 10.1016/j.resmic.2010.09.012
- Kyle, J. L., Parker, C. T., Goudeau, D., and Brandl, M. T. (2010). Transcriptome analysis of *Escherichia coli* O157: H7 exposed to lysates of lettuce leaves. *Appl. Environ. Microbiol.* 76, 1375–1387. doi: 10.1128/AEM.02461-09
- La Carbona, S., Sauvageot, N., Giard, J. C., Benachour, A., Posteraro, B., Auffray, Y., et al. (2007). Comparative study of the physiological roles of three peroxidases (NADH peroxidase, Alkyl hydroperoxide reductase and Thiol peroxidase) in oxidative stress response, survival inside macrophages and virulence of *Enterococcus faecalis*. *Mol. Microbiol.* 66, 1148–1163. doi: 10.1111/j.1365-2958.2007.05987.x
- Lleó, M., Bonato, B., Benedetti, D., and Canepari, P. (2005). Survival of enterococcal species in aquatic environments. *FEMS Microbiol. Ecol.* 54, 189–196. doi: 10.1016/j.femsec.2005.03.016
- Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25, 1754–1760. doi: 10.1093/bioinformatics/btp324
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics* 25, 2078–2079. doi: 10.1093/bioinformatics/btp352
- Lim, A. C., Mak, K. C., Ng, N. U., and Ng, T. (2007). Multiple modes of protection against hydrogen peroxide-induced oxidative damage in stationary and exponential phase *Escherichia coli* by DNA-binding protein (Dps). *J. Exp. Microbiol. Immunol.* 11, 86–92.
- Lindell, M. J., Granéli, W., and Tranvik, L. J. (1995). Enhanced bacterial growth in response to photochemical transformation of dissolved organic matter. *Limnol. Oceanogr.* 40, 195–199. doi: 10.4319/lo.1995.40.1.0195
- Litton, R. M., Ahn, J. H., Sercu, B., Holden, P. A., Sedlak, D. L., and Grant, S. B. (2010). Evaluation of chemical, molecular, and traditional markers of fecal contamination in an effluent dominated urban stream. *Environ. Sci. Technol.* 44, 7369–7375. doi: 10.1021/es101092g
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻ΔΔCT method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- Lleó, M. M., Tafi, M. C., and Canepari, P. (1998). Nonculturable *Enterococcus faecalis* cells are metabolically active and capable of resuming active growth. *Syst. Appl. Microbiol.* 21, 333–339. doi: 10.1016/S0723-2020(98)80041-6
- Lonborg, C., Martínez-García, S., Teira, E., and Álvarez-Salgado, X. A. (2013). Effects of the photochemical transformation of dissolved organic matter on bacterial physiology and diversity in a coastal system. *Estuar. Coast. Shelf Sci.* 129, 11–18. doi: 10.1016/j.ecss.2013.05.013
- Lonborg, C., Nieto-Cid, M., Hernando-Morales, V., Hernandez-Ruiz, M., Teira, E., and Alvarez-Salgado, X. A. (2016). Photochemical alteration of dissolved organic matter and the subsequent effects on bacterial carbon cycling and diversity. *FEMS Microbiol. Ecol.* 92:fiw048. doi: 10.1093/femsec/fiw048
- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550. doi: 10.1186/s13059-014-0550-8
- Macedo, M., and Waterson, T. (2016). *Profile Analysis [Online]*. Available Online at: <http://userwww.sfsu.edu/efc/classes/biol710/manova/profileanalysis.htm> [Accessed December, 2016 2016].
- MacLean, R. C., and San Millan, A. (2015). Microbial evolution: towards resolving the plasmid paradox. *Curr. Biol.* 25, R764–R767. doi: 10.1016/j.cub.2015.07.006
- Maraccini, P. A., Wenk, J., and Boehm, A. B. (2016). Photoinactivation of eight health-relevant bacterial species: determining the importance of the exogenous indirect mechanism. *Environ. Sci. Technol.* 50, 5050–5059. doi: 10.1021/acs.est.6b00074
- Marsico, R. M., Schneider, R. J., Voelker, B. M., Zhang, T., Diaz, J. M., Hansel, C. M., et al. (2015). Spatial and temporal variability of widespread dark production and decay of hydrogen peroxide in freshwater. *Aquat. Sci.* 77, 523–533. doi: 10.1007/s00027-015-0399-2
- Marty-Teyssat, C., de la Torre, F., and Garel, J. R. (2000). Increased production of hydrogen peroxide by *Lactobacillus delbrueckii* subsp. *bulgaricus* upon aeration: Involvement of an NADH oxidase in oxidative stress. *Appl. Environ. Microbiol.* 66, 262–267. doi: 10.1128/AEM.66.1.262-267.2000
- McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., et al. (2010). The genome analysis toolkit: a mapreduce framework for analyzing next-generation dna sequencing data. *Genome Res.* 20, 1297–1303. doi: 10.1101/gr.107524.110
- Michán, C., Manchado, M., Dorado, G., and Pueyo, C. (1999). *In vivo* transcription of the *Escherichia coli* oxyR regulon as a function of growth phase and in response to oxidative stress. *J. Bacteriol.* 181, 2759–2764.
- Minogue, T. D., Daligault, H. E., Davenport, K. W., Broomall, S. M., Bruce, D. C., Chain, P. S., et al. (2014). Complete Genome Assembly of *Enterococcus faecalis* 29212, a Laboratory Reference Strain. *Genome Announc.* 2:e00968-14. doi: 10.1128/genomeA.00968-14
- Moran, M. A., and Zepp, R. G. (1997). Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter. *Limnol. Oceanogr.* 42, 1307–1316. doi: 10.4319/lo.1997.42.6.1307
- Morris, J. J., Johnson, Z. I., Wilhelm, S. W., and Zinser, E. R. (2016). Diel regulation of hydrogen peroxide defenses by open ocean microbial communities. *J. Plankton Res.* 38, 1103–1114. doi: 10.1093/plankt/fbw016
- Mostofa, K. G., Liu, C.-Q., Sakugawa, H., Vione, D., Minakata, D., and Wu, F. (2013). “Photoinduced and microbial generation of hydrogen peroxide and organic peroxides in natural waters,” in *Photobiogeochemistry of Organic Matter*, eds K. M. G. Mostofa, T. Yoshioka, A. Mottaleb, and D. Vione (Heidelberg: Berlin: Springer), 139–207.
- Mostofa, K. M., and Sakugawa, H. (2009). Spatial and temporal variations and factors controlling the concentrations of hydrogen peroxide and organic peroxides in rivers. *Environ. Chem.* 6, 524–534. doi: 10.1071/EN09070
- Mostofa, K. M., Yoshioka, T., Mottaleb, A., and Vione, D. (2012). *Photobiogeochemistry of Organic Matter - Principles and Practices in Water Environments*. Environmental Science and Engineering Book Series. Berlin: Springer.

- Motta, S. S., Cluzel, P., and Aldana, M. (2015). Adaptive resistance in bacteria requires epigenetic inheritance, genetic noise, and cost of efflux pumps. *PLoS ONE* 10:e0118464. doi: 10.1371/journal.pone.0118464
- Moy, T. I., Mylonakis, E., Calderwood, S. B., and Ausubel, F. M. (2004). Cytotoxicity of hydrogen peroxide produced by *Enterococcus faecium*. *Infect. Immun.* 72, 4512–4520. doi: 10.1128/IAI.72.8.4512-4520.2004
- Nguyen, M. T., Jasper, J. T., Boehm, A. B., and Nelson, K. L. (2015). Sunlight inactivation of fecal indicator bacteria in open-water unit process treatment wetlands: modeling endogenous and exogenous inactivation rates. *Water Res.* 83, 282–292. doi: 10.1016/j.watres.2015.06.043
- Normark S., Ericson C., Jonsson A., and Olsen A. (1992). "Global regulatory control of curli expression and fibronectin binding in Enterobacteria," in *Microbial Adhesion and Invasion*, eds M. Hook and L. Switalski (New York, NY: Springer), 95–99. doi: 10.1007/978-1-4612-2924-7_9
- Oladeinde, A., Bohrmann, T., Wong, K., Purucker, S., Bradshaw, K., Brown, R., et al. (2014). Decay of fecal indicator bacterial populations and bovine-associated source-tracking markers in freshly deposited cow pats. *Appl. Environ. Microbiol.* 80, 110–118. doi: 10.1128/AEM.02203-13
- Oliver, D. M., Bird, C., Burd, E., and Wyman, M. (2016). Quantitative PCR profiling of *E. coli* in livestock faeces reveals increased population resilience relative to culturable counts under temperature extremes. *Environ. Sci. Technol.* 50, 9497–505. doi: 10.1021/acs.est.6b02657
- Pachepsky, Y. A., Blaustein, R. A., Whelan, G., and Shelton, D. R. (2014). Comparing temperature effects on *Escherichia coli*, *Salmonella*, and *Enterococcus* survival in surface waters. *Lett. Appl. Microbiol.* 59, 278–283. doi: 10.1111/lam.12272
- Parsonage, D., Karplus, P. A., and Poole, L. B. (2008). Substrate specificity and redox potential of AhpC, a bacterial peroxiredoxin. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8209–8214. doi: 10.1073/pnas.0708308105
- Quax, T. E., Claassens, N. J., Söll, D., and van der Oost, J. (2015). Codon bias as a means to fine-tune gene expression. *Mol. Cell* 59, 149–161. doi: 10.1016/j.molcel.2015.05.035
- Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26, 841–842. doi: 10.1093/bioinformatics/btq033
- Ramsey, M., Hartke, A., and Huycke, M. (2014). "The physiology and metabolism of Enterococci," in *Enterococci: From Commensals to Leading Causes of Drug Resistant Infection [Internet]*, eds M. S. Gilmore, D. B. Clewell, Y. Ike, and N. Shankar (Boston, MA: Massachusetts Eye and Ear Infirmary).
- Ravindra Kumar, S., and Imlay, J. A. (2013). How *Escherichia coli* tolerates profuse hydrogen peroxide formation by a catabolic pathway. *J. Bacteriol.* 195, 4569–4579. doi: 10.1128/JB.00737-13
- Riboldi, G. P., Bierhals, C. G., Mattos, E. P., d., Frazzon, A. P. G., and Frazzon, J. (2014). Oxidative stress enhances the expression of sulfur assimilation genes: preliminary insights on the *Enterococcus faecalis* iron-sulfur cluster machinery regulation. *Memórias do Instituto Oswaldo Cruz* 109, 408–413. doi: 10.1590/0074-0276140006
- Rochelle-Newall, E., Nguyen, T. M., Le, T. P., Sengtaheuanghoung, O., and Ribolzi, O. (2015). A short review of fecal indicator bacteria in tropical aquatic ecosystems: knowledge gaps and future directions. *Front. Microbiol.* 6:308. doi: 10.3389/fmicb.2015.00308
- Rose, A. L., Moffett, J. W., and Waite, T. D. (2008). Determination of superoxide in seawater using 2-methyl-6-(4-methoxyphenyl)-3,7-dihydroimidazo[1,2-a]pyrazin-3(7H)-one chemiluminescence. *Anal. Chem.* 80, 1215–1227. doi: 10.1021/ac7018975
- Rutherford, S. T., and Bassler, B. L. (2012). Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb. Perspect. Med.* 2:a012427. doi: 10.1101/cshperspect.a012427
- R Developmental Core Team (2012). "R: A Language and Environment for Statistical Computing. Vienna: R Foundation for Statistical Computing.
- Saier, M. H., Chauvaux, S. Jr., Cook, G. M., Deutscher, J., Paulsen, I. T., Reizer, J., et al. (1996). Catabolite repression and inducer control in Gram-positive bacteria. *Microbiology* 142, 217–230. doi: 10.1099/13500872-142-2-217
- Sanders, E. C., Yuan, Y. P., and Pitchford, A. (2013). Fecal coliform and *E. coli* concentrations in effluent-dominated streams of the upper santa cruz watershed. *Water* 5, 243–261. doi: 10.3390/w5010243
- Sassoubre, L. M., Nelson, K. L., and Boehm, A. B. (2012). Mechanisms for Photoinactivation of *Enterococcus faecalis* in Seawater. *Appl. Environ. Microbiol.* 78, 7776–7785. doi: 10.1128/AEM.02375-12
- Scully, N. M., Cooper, W. J., and Tranvik, L. J. (2003). Photochemical effects on microbial activity in natural waters: the interaction of reactive oxygen species and dissolved organic matter. *FEMS Microbiol. Ecol.* 46, 353–357. doi: 10.1016/S0168-6496(03)00198-3
- Seaver, L. C., and Imlay, J. A. (2004). Are respiratory enzymes the primary sources of intracellular hydrogen peroxide? *J. Biol. Chem.* 279, 48742–48750. doi: 10.1074/jbc.M408754200
- Simsek, H., Wadhawan, T., and Khan, E. (2013). Overlapping photodegradable and biodegradable organic nitrogen in wastewater effluents. *Environ. Sci. Technol.* 47, 7163–7170. doi: 10.1021/es400120m
- Suzuki, S., Horinouchi, T., and Furusawa, C. (2014). Prediction of antibiotic resistance by gene expression profiles. *Nat. Commun.* 5:5792. doi: 10.1038/ncomms6792
- Szkarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., et al. (2014). STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic Acids Res.* 43, D447–D452. doi: 10.1093/nar/gku1003
- Tremblay, P. L., and Lovley, D. R. (2012). Role of the NiFe hydrogenase Hya in oxidative stress defense in *Geobacter sulfurreducens*. *J. Bacteriol.* 194, 2248–2253. doi: 10.1128/JB.00044-12
- Uhlich, G. A., Reichenberger, E. R., Cottrell, B. J., Fratamico, P., and Andreozzi, E. (2017). Whole-Genome Sequence of *Escherichia coli* Serotype O157:H7 Strain B6914-ARS. *Genome Announc.* 5:e01191-17. doi: 10.1128/genomeA.01191-17
- USEPA (2017). *National Water Quality Inventory Report to Congress. EPA 841-R-16-011*. US Environmental Protection Agency. Available online at: <https://www.epa.gov/waterdata/2017-national-water-quality-inventory-report-congress>
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., et al. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 3:RESEARCH0034. doi: 10.1186/gb-2002-3-7-research0034
- Van Dongen, S. (2008). Graph clustering via a discrete uncoupling process. *SIAM J. Matrix Anal. Appl.* 30, 121–141. doi: 10.1137/040608635
- Verneuil, N., Sanguinetti, M., Le Breton, Y., Posteraro, B., Fadda, G., Auffray, Y., et al. (2004). Effects of the *Enterococcus faecalis* hypR gene encoding a new transcriptional regulator on oxidative stress response and intracellular survival within macrophages. *Infect. Immun.* 72, 4424–4431. doi: 10.1128/IAI.72.8.4424-4431.2004
- Villa, L., García-Fernández, A., Fortini, D., and Carattoli, A. (2010). Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J. Antimicrob. Chemother.* 65, 2518–2529. doi: 10.1093/jac/dkq347
- Villegas, E., and Gilliland, S. E. (1998). Hydrogen peroxide production by *Lactobacillus delbrueckii* subsp. *lactis* I at 5 degrees C. *J. Food Sci.* 63, 1070-1074. doi: 10.1111/j.1365-2621.1998.tb15857.x
- Wanjugi, P., and Harwood, V. J. (2013). The influence of predation and competition on the survival of commensal and pathogenic fecal bacteria in aquatic habitats. *Environ. Microbiol.* 15, 517–526. doi: 10.1111/j.1462-2920.2012.02877.x
- Whitman, R. L., Nevers, M. B., Korinek, G. C., and Byappanahalli, M. N. (2004). Solar and temporal effects on *Escherichia coli* concentration at a Lake Michigan swimming beach. *Appl. Environ. Microbiol.* 70, 4276–4285. doi: 10.1128/AEM.70.7.4276-4285.2004
- Williamson, C. E., Madronich, S., Lal, A., Zepp, R. G., Lucas, R. M., Overholt, E. P., et al. (2017). Climate change-induced increases in precipitation are reducing the potential for solar ultraviolet radiation to inactivate pathogens in surface waters. *Sci. Rep.* 7:13033. doi: 10.1038/s41598-017-13392-2
- Wolf, R., Thrane, J. E., Hessen, D. O., and Andersen, T. (2018). Modelling ROS formation in boreal lakes from interactions between dissolved organic matter and absorbed solar photon flux. *Water Res.* 132, 331–339. doi: 10.1016/j.watres.2018.01.025
- Yan, X., Budin-Verneuil, A., Verneuil, N., Gilmore, M. S., Artigaud, S., Auffray, Y., et al. (2015). Transcriptomic response of *Enterococcus faecalis* V583 to low hydrogen peroxide levels. *Curr. Microbiol.* 70, 156–168. doi: 10.1007/s00284-014-0691-8

- Yu, J., Madsen, M. L., Carruthers, M. D., Phillips, G. J., Kavanaugh, J. S., Boyd, J. M., et al. (2013). Analysis of autoinducer-2 quorum sensing in *Yersinia pestis*. *Infect. Immun.* 81, 4053–4062. doi: 10.1128/IAI.00880-13
- Zhang, T., Hansel, C. M., Voelker, B. M., and Lamborg, C. H. (2016). Extensive dark biological production of reactive oxygen species in brackish and freshwater ponds. *Environ. Sci. Technol.* 50, 2983–2993. doi: 10.1021/acs.est.5b03906
- Zheng, M., Wang, X., Templeton, L. J., Smulski, D. R., LaRossa, R. A., and Storz, G. (2001). DNA microarray-mediated transcriptional profiling of the *Escherichia coli* response to hydrogen peroxide. *J. Bacteriol.* 183, 4562–4570. doi: 10.1128/JB.183.15.4562-4570.2001
- Zhou, K., Zhou, L., Lim, Q. E., Zou, R., Stephanopoulos, G., and Too, H.-P. (2011a). Novel reference genes for quantifying transcriptional responses of *Escherichia coli* to protein overexpression by quantitative PCR. *BMC Mol. Biol.* 12:18. doi: 10.1186/1471-2199-12-18
- Zhou, Y., Liang, Y., Lynch, K. H., Dennis, J. J., and Wishart, D. S. (2011b). PHAST: a fast phage search tool. *Nucleic Acids Res.* 39, W347–W352. doi: 10.1093/nar/gkr485

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Oladeinde, Lipp, Chen, Muirhead, Glenn, Cook and Molina. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.