



# Dynamic Regulation of Extracellular Superoxide Production by the Coccolithophore *Emiliana huxleyi* (CCMP 374)

## OPEN ACCESS

Sydney Plummer<sup>1†</sup>, Alexander E. Taylor<sup>2</sup>, Elizabeth L. Harvey<sup>1</sup>, Colleen M. Hansel<sup>3</sup> and Julia M. Diaz<sup>1\*†</sup>

### Edited by:

Ondrej Prasil,  
Institute of Microbiology (ASCR),  
Czechia

### Reviewed by:

Allen Milligan,  
Oregon State University,  
United States  
Gustaaf Marinus Hallegraeff,  
University of Tasmania,  
Australia

### \*Correspondence:

Julia M. Diaz  
j2diaz@ucsd.edu;  
julia.diaz@skio.uga.edu

### †Present address:

Sydney Plummer,  
Scripps Institution of Oceanography,  
University of California San Diego,  
La Jolla, CA, United States  
Julia M. Diaz,  
Scripps Institution of Oceanography,  
University of California San Diego,  
La Jolla, CA, United States

### Specialty section:

This article was submitted to  
Aquatic Microbiology,  
a section of the journal  
Frontiers in Microbiology

Received: 25 February 2019

Accepted: 20 June 2019

Published: 12 July 2019

### Citation:

Plummer S, Taylor AE, Harvey EL,  
Hansel CM and Diaz JM (2019)  
Dynamic Regulation of Extracellular  
Superoxide Production by  
the Coccolithophore  
*Emiliana huxleyi* (CCMP 374).  
*Front. Microbiol.* 10:1546.  
doi: 10.3389/fmicb.2019.01546

<sup>1</sup>Department of Marine Sciences, Skidaway Institute of Oceanography, University of Georgia, Savannah, GA, United States, <sup>2</sup>Department of Chemistry, University of Vermont, Burlington, VT, United States, <sup>3</sup>Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, United States

In marine waters, ubiquitous reactive oxygen species (ROS) drive biogeochemical cycling of metals and carbon. Marine phytoplankton produce the ROS superoxide ( $O_2^-$ ) extracellularly and can be a dominant source of  $O_2^-$  in natural aquatic systems. However, the cellular regulation, biological functioning, and broader ecological impacts of extracellular  $O_2^-$  production by marine phytoplankton remain mysterious. Here, we explored the regulation and potential roles of extracellular  $O_2^-$  production by a noncalcifying strain of the cosmopolitan coccolithophorid *Emiliana huxleyi*, a key species of marine phytoplankton that has not been examined for extracellular  $O_2^-$  production previously. Cell-normalized extracellular  $O_2^-$  production was the highest under presumably low-stress conditions during active proliferation and inversely related to cell density during exponential growth phase. Removal of extracellular  $O_2^-$  through addition of the  $O_2^-$  scavenger superoxide dismutase (SOD), however, increased growth rates, growth yields, cell biovolume, and photosynthetic efficiency ( $F_v/F_m$ ) indicating an overall physiological improvement. Thus, the presence of extracellular  $O_2^-$  does not directly stimulate *E. huxleyi* proliferation, as previously suggested for other phytoplankton, bacteria, fungi, and protists. Extracellular  $O_2^-$  production decreased in the dark, suggesting a connection with photosynthetic processes. Taken together, the tight regulation of this stress independent production of extracellular  $O_2^-$  by *E. huxleyi* suggests that it could be involved in fundamental photophysiological processes.

**Keywords:** reactive oxygen species, superoxide, *Emiliana huxleyi*, photophysiology, oxidative stress, redox homeostasis, biogeochemical cycling

## INTRODUCTION

Ubiquitous reactive oxygen species (ROS) within marine waters help drive global biogeochemical cycling. ROS include intermediates in the reduction of molecular oxygen ( $O_2$ ) to water ( $H_2O$ ), which consist of superoxide ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radical ( $OH\cdot$ ). These ROS shape the transformation of metal nutrients including iron (Rose, 2012) and manganese (Wuttig et al., 2013), as well as carbon (Heller and Croot, 2010b) due to

their ability to act as both oxidants and reductants. Within aquatic environments, ROS are produced through both abiotic (e.g., *via* photodegradation of organic matter) and biotic means (Zinser, 2018). Biotic mechanisms include active extracellular production by marine microorganisms, which can be a dominant source of  $O_2^-$  in natural waters (Rose et al., 2008; Hansard et al., 2010).

While ROS are formed intracellularly within all aerobic organisms as metabolic by-products, a plethora of marine microorganisms actively produce ROS extracellularly as well. Although intracellular  $O_2^-$  can be released into the marine environment upon cell lysis, these rates cannot account for the steady-state concentrations that have been measured in natural waters (Rose, 2012). Furthermore, within cells,  $O_2^-$  exists in equilibrium with its conjugate acid, the hydroperoxyl radical ( $HOO\bullet$ ); however, with a  $pK_a$  of 4.8, the  $O_2^-$  anion is the dominant form at physiological pH (Bielski et al., 1985). Unlike  $H_2O_2$ ,  $O_2^-$  cannot readily diffuse across membranes due to its negative charge, short lifetime ( $\sim 10^{-5}$  s), and limited diffusive distance ( $\sim 10^{-7}$  m) (Lesser, 2006; Brown and Griendling, 2009; Diaz and Plummer, 2018). Thus, the majority of microbially derived  $O_2^-$  within the extracellular environment must be created on or near the cell surface via active extracellular  $O_2^-$  production mechanisms (Diaz and Plummer, 2018).

The ability to produce extracellular  $O_2^-$  has been documented among heterotrophic bacteria (Diaz et al., 2013) and phytoplankton, including cyanobacteria (Rose et al., 2005, 2008; Godrant et al., 2009; Hansel et al., 2016), diatoms (Kustka et al., 2005; Hansel et al., 2016; Schneider et al., 2016), dinoflagellates (Saragosti et al., 2010; Zhang et al., 2016a), nontoxic microalgae (Marshall et al., 2005a), and harmful microalgae (Oda et al., 1997; Marshall et al., 2005a,b; Portune et al., 2010; Diaz and Plummer, 2018). Despite advancements in identifying the extensive presence and environmental relevance of extracellular  $O_2^-$  production by marine microflora, the mechanisms of extracellular  $O_2^-$  production and its biological roles are not well understood. The freshwater chlorophyte *Chlamydomonas reinhardtii* (Anderson et al., 2015) and marine raphidophytes *Chattonella marina* and *Chattonella ovata* (Kim et al., 2000, 2007) are either confirmed or thought to produce extracellular  $O_2^-$  via cell membrane associated enzymes known as NADPH oxidases (Nox). These enzymes transfer electrons from cytosolic NADPH pools across cell membranes to reduce  $O_2$  in the surrounding aqueous environment, thus creating extracellular  $O_2^-$ . The presence of these enzymes has been implicated in diatoms and dinoflagellates as well (Kim et al., 2000; Kustka et al., 2005; Hervé et al., 2006; Saragosti et al., 2010). Extracellular  $O_2^-$  production is light dependent in several phytoplankton taxa (Kim et al., 1999; Marshall et al., 2002; Milne et al., 2009; Saragosti et al., 2010; Hansel et al., 2016; Schneider et al., 2016), which has led to speculation that photosynthesis may serve an indirect role in extracellular  $O_2^-$  production by supplying NADPH to cell surface-associated NADPH-oxidizing enzymes such as Nox (Marshall et al., 2002; Saragosti et al., 2010; Schneider et al., 2016).

Proposed biological roles of phytoplankton-derived extracellular  $O_2^-$  are diverse (Diaz and Plummer, 2018). For instance, extracellular  $O_2^-$  production has been implicated in

harmful algal bloom toxicity (Tanaka et al., 1992; Yang et al., 1995; Kim et al., 1999; Marshall et al., 2003; Kim and Oda, 2010; Dorantes-Aranda et al., 2013, 2015; Mardones et al., 2015), metal nutrient acquisition (Rose et al., 2005; Garg et al., 2007; Liu et al., 2007; Rose, 2012; Roe and Barbeau, 2014), allelopathy (Oda et al., 1992, 1997; Marshall et al., 2005b), and defense against grazing (Martel, 2009; Flores et al., 2012). Model phytoplankton species generate abundant extracellular  $O_2^-$  even under ideal growth conditions in the absence of any obvious stressors (Kustka et al., 2005; Rose et al., 2005; Marshall et al., 2005a,b; Godrant et al., 2009; Portune et al., 2010; Diaz et al., 2013; Hansel et al., 2016; Schneider et al., 2016), suggesting an association with basal functioning. For example, extracellular  $O_2^-$  regulates growth and morphology in the prolific ROS producer, *C. marina* (Oda et al., 1995), as well as growth in bacteria (Saran, 2003; Hansel et al., 2019) and differentiation in microbial eukaryotes (Aguirre et al., 2005). In these microorganisms, extracellular  $O_2^-$  production rates are the highest during active growth and at low cell densities consistent with beneficial cell signaling and autocrine growth regulation, as also seen in plants (Mittler et al., 2011) and animals (Brown and Griendling, 2009; Aguirre and Lambeth, 2010). The accumulation of studies showing that extracellular  $O_2^-$  production by diverse phytoplankton is similarly dependent on cell density (Marshall et al., 2005a; Hansel et al., 2016; Diaz et al., 2018) and growth phase (Oda et al., 1995; Kim et al., 1999; Portune et al., 2010) has spurred speculation that extracellular  $O_2^-$  production may be involved in phytoplankton cell signaling and/or growth regulation in species other than *C. marina*, although this possibility remains largely untested (Hansel et al., 2016; Diaz and Plummer, 2018).

Clarifying the cellular regulation and biological function of active extracellular  $O_2^-$  production by phytoplankton is critical to understand the effects of ROS on ocean redox balance, biogeochemical cycling, and ecological interactions in marine waters. Among phytoplankton, coccolithophores are one of the most prevalent groups in the global ocean. Further, the original report of extracellular  $H_2O_2$  production by the coccolithophorid species *Pleurochrysis carterae* pioneered the recognition of microorganisms as significant sources of ROS in aquatic systems (Palenik et al., 1987). Despite this discovery, however, coccolithophorids are under explored in terms of extracellular ROS production. Therefore, this study was conducted to investigate the dynamics, cellular regulation, and biological function of extracellular  $O_2^-$  production by a noncalcifying strain of *Emiliana huxleyi* (CCMP 374), the most prevalent coccolithophore species in modern oceans (Westbroek et al., 1989; Brown and Yoder, 1994).

## MATERIALS AND METHODS

### Cultivation of *E. huxleyi*, Growth Tracking, and Cell Counts

Axenic cultures of *E. huxleyi* CCMP 374 were obtained from the National Center for Marine Algae and Microbiota (NCMA) at Bigelow Laboratory for Ocean Sciences (East Boothbay, ME).

Cultures of *E. huxleyi* were inoculated into *f/2* growth media prepared without the addition of silicic acid (Guillard and Ryther, 1962) using 0.2  $\mu\text{m}$  filtered natural seawater collected from the South Atlantic Bight. Media were prepared and autoclaved (121°C, 20 min) at least 1 day prior to inoculating cultures. Cultures were begun with exponential phase inocula, unless otherwise stated. Cultures were either grown in borosilicate culture tubes with caps or Erlenmeyer flasks of various sizes with aluminum foil or an acid washed plastic beaker covering the mouth of the flask at 18°C under cool, white light ( $\sim 130 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , 14:10 light dark cycle). Growth was monitored by observing *in vivo* chlorophyll fluorescence using an AquaFluor® handheld fluorometer (Turner Designs, San Jose, CA) or a 10-AU™ fluorometer (Turner Designs, San Jose, CA). *In vivo* fluorescence values were normalized to measurements taken on day 0. Exponential growth phase was defined as the log-linear portion of the *in vivo* fluorescence data versus time ( $R^2 \geq 0.98$  in all cultures). Stationary phase was determined to be the time between the end of the log linear portion of the growth curve and until the end of the growth curve. Specific growth rates during exponential growth phase were found by calculating the slope of the regression of the natural log-normalized *in vivo* fluorescence versus time. Culture pH was monitored using an Accumet AB 15/15+ pH meter (Thermo Fisher Scientific, Waltham, MA) in cultures grown from stationary phase inocula in 25 mm borosilicate tubes (Thermo Fisher Scientific, 14-961-34). *E. huxleyi* cell abundances (cells  $\text{ml}^{-1}$ ) were obtained using a Guava® easyCyte flow cytometer (Millipore Sigma, Merck KGaA, Darmstadt, Germany) and analyzed with Guava InCyte™ 3.1 software. Flow cytometry samples were preserved with a final concentration of 0.5% glutaraldehyde, as well as 1% peptone to prevent cell adsorption to sample tubes. Flow cytometry samples were stored at  $-80^\circ\text{C}$  prior to processing. To process samples, 50–200  $\mu\text{l}$  of each sample was pipetted into 96-well plates, diluted with filtered seawater as needed, and run at a low flow rate (0.24  $\mu\text{l s}^{-1}$ ) for 3 min. For analysis, concentrations of healthy cell populations (cells  $\text{ml}^{-1}$ ) were determined based on gates of red fluorescence and forward scatter signals from previously run samples of exponentially growing cultures.

In two separate experiments, extracellular  $\text{O}_2^-$  was removed from *E. huxleyi* cultures by adding superoxide dismutase (SOD, Millipore Sigma 574,594-50KU), an enzyme that specifically degrades  $\text{O}_2^-$ . To begin this experiment, 7.5 ml of media were inoculated with 300  $\mu\text{l}$  stationary phase culture to give an initial concentration of  $\sim 2.3 \times 10^5$  cells  $\text{ml}^{-1}$  and grown under the conditions above in 13 mm borosilicate glass tubes (Thermo Fisher Scientific, 14-962-26D). Treatments included three different concentrations of SOD added at three different volumes and a deionized water (DI) control. A 10 kU  $\text{ml}^{-1}$  stock of SOD was prepared using DI. Then, treatments were performed on triplicate cultures, where each tube received daily additions of 20.25  $\mu\text{l}$ , 37.5  $\mu\text{l}$ , or 75  $\mu\text{l}$  of the 10 kU  $\text{ml}^{-1}$  SOD stock, or 75  $\mu\text{l}$  DI to give final concentrations of 27 U  $\text{ml}^{-1}$  SOD, 50 U  $\text{ml}^{-1}$  SOD, 100 U  $\text{ml}^{-1}$  SOD, or 0 U  $\text{ml}^{-1}$  SOD, respectively. A subsequent control experiment was performed with diafiltered SOD to ensure that SOD was responsible for potential changes

observed in the cultures. For this control experiment, 7.5 ml of media were inoculated with 300  $\mu\text{l}$  stationary phase culture to give an initial concentration of  $\sim 1.8 \times 10^5$  cells  $\text{ml}^{-1}$  and grown under the conditions above in 13 mm borosilicate glass tubes. To create the dialyzed SOD, a 10 kU  $\text{ml}^{-1}$  SOD stock prepared with DI was diafiltered by passing the SOD solution through an Amicon ultrafiltration device (10 kDa molecular weight cut-off, Millipore) at 3200  $\times g$  for 20 min at 4°C, thereby removing the enzyme from the solution. Treatments were performed on triplicate cultures, where each tube received daily additions of 75  $\mu\text{l}$  dialyzed SOD, 75  $\mu\text{l}$  DI, and 75  $\mu\text{l}$  SOD (100 U  $\text{ml}^{-1}$  SOD final concentration).

## Cell Imaging and Analysis

Individual *E. huxleyi* cells were imaged using a FlowCam® (Fluid Imaging Inc., Scarborough, ME), a continuous flow-through microscope fitted with a color and monochromatic camera. To calibrate the FlowCam® for imaging, dilutions of *E. huxleyi* cultures from 0 to 95% using 0.2  $\mu\text{m}$  filtered seawater were analyzed in varying context settings. Final settings (Supplementary Table S1) were chosen based on their ability to differentiate cell shape and color and provide useable images for morphological and volume assessments. Culture samples of 100  $\mu\text{l}$  were diluted with 500  $\mu\text{l}$  of 0.2  $\mu\text{m}$  filtered seawater prior to analysis on the FlowCam®. Images were analyzed with VisualSpreadsheet® (Fluid Imaging Inc., Scarborough, ME). The spherical biovolume was determined using three separate measurements of cell diameter. The FlowCam® software calculates three different diameters for each cell image [area based diameter (ABD), equivalent spherical diameter (ESD), filled or full diameter (FD)], thus giving three alternative biovolume assessments for every imaged cell. Further analysis and filtering of FlowCam® images as well as details on diameter calculations are described in the Supplementary Material.

## Photophysiology

Photophysiological health of *E. huxleyi* was monitored using the Atlantic fluorescence induction and relaxation (FIRE) fluorometer system (Sea-bird Scientific, Halifax, NS Canada). Prior to analysis, samples were allowed to dark adapt for approximately 30 min and were then diluted using 0.2  $\mu\text{m}$  filtered seawater as necessary to avoid detector saturation. Samples were illuminated with a single turnover flash of blue excitation for a duration of 80  $\mu\text{s}$ . The measured fluorescence response was used to calculate the maximum efficiency of photosystem II (PSII), or  $F_v/F_m$ , using the equation:

$$F_v/F_m = \frac{(F_m - F_o)}{F_m}$$

where  $F_m$  is the maximum fluorescence yield, and  $F_o$  is the minimum fluorescence yield. For each sample, 20 acquisitions were obtained.

## Extracellular $\text{O}_2^-$ Production

Net extracellular  $\text{O}_2^-$  production by *E. huxleyi* was measured using the flow-through FeLume (II) analytical system

(Waterville Analytical, Waterville, ME) by detecting chemiluminescence emitted through the reaction of  $O_2^-$  and the specific probe methyl *Cypridina* luciferin analog (MCLA), as previously described (Diaz et al., 2013; Schneider et al., 2016). This method allows for manipulation of experimental settings (e.g., light levels, increasing cell density) during analysis to detect changes in  $O_2^-$  on immediate timescales (i.e., seconds). It has been used to measure  $O_2^-$  production in pure cultures (Kustka et al., 2005; Diaz et al., 2013; Schneider et al., 2016; Zhang et al., 2016a; Hansel et al., 2019) and natural waters (Diaz et al., 2016; Hansel et al., 2016; Zhang et al., 2016b). Following the procedures of Diaz et al. (2013), MCLA blanks generated in the absence of SOD were used to determine biologically derived  $O_2^-$  concentrations, in order to avoid overestimation of biological  $O_2^-$  production rates due to MCLA auto-oxidation. Briefly, cells were deposited onto an inline filter (0.22  $\mu\text{m}$ ), continuously rinsed (2 ml  $\text{min}^{-1}$ ) with a phosphate buffer (20 mM; pH = 7.6) that matched the salinity of the seawater media base (38 psu), and  $O_2^-$  was quantified in the cell-free effluent upon reaction with the MCLA reagent [4  $\mu\text{M}$  MCLA, 0.1 M MES, 75  $\mu\text{M}$  diethylenetriamine pentaacetic acid (DTPA), pH = 6] in a spiral flow cell adjacent to a photomultiplier tube. Chemiluminescent signals from the phosphate buffer and MCLA baselines as well as biological samples were allowed to stabilize ( $\leq 4\%$  CV) for at least 1 min, allowing calculation of a steady-state  $O_2^-$  concentration. In this way, obtaining a stable chemiluminescent signal from the MCLA reacting with effluent from biological samples demonstrated that the  $O_2^-$  being detected was not a rapid, short-lived release of intracellular  $O_2^-$  due to cell rupture. SOD was added at the end of each analysis (final concentration of  $\sim 800 \text{ U l}^{-1}$ ) to confirm the signal acquired on the FeLume (II) system was due to  $O_2^-$ . In comparison to the aforementioned study by Diaz et al. (2013), the following exceptions were implemented here. First, cells deposited on the filter were exposed to ambient light ( $\sim 5 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) unless they were covered in a dark photography film-changing bag to obtain dark (0  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) measurements, as indicated. Second, only net production rates were determined, and finally, calibration was performed using standard additions of potassium superoxide ( $\text{KO}_2$ ) by the method of Schneider et al. (2016). A preliminary cell concentration was obtained by microscopy using a hemocytometer counting chamber to help ensure that the same number of cells was loaded onto the in-line filter from each biological replicate and on each day within an experiment. The biotic steady-state  $O_2^-$  concentration was calculated by subtracting blank signals generated from the mixture of MCLA and phosphate buffer with a syringe filter inline and in the absence of SOD. Then, net  $O_2^-$  production rates were calculated by multiplying the biotic steady-state  $O_2^-$  concentration (pM) by the flow rate (2 ml  $\text{min}^{-1}$ ), dividing that value by the number of cells loaded onto the inline filter (either found using microscopy or flow cytometry), and converting to final units of  $\text{amol cell}^{-1} \text{ h}^{-1}$ . All chemicals used to measure  $O_2^-$  were obtained

from Millipore Sigma, except for MCLA, which was obtained from Tokyo Chemical Industry Co., Ltd.


## Statistical Analyses

All statistical analyses were performed using JMP Pro 13.0.0 (SAS Institute Inc., Cary, NC). Regression analyses of cell-normalized  $O_2^-$  production rates as a function of time across the growth curve of *E. huxleyi* and cell density were performed using Spearman's rank-order correlation. This regression analysis indicates the presence or absence of monotonic relationships based on the correlation coefficient ( $\rho$ ) and its level of significance ( $p$ ). An independent two sample Student's  $t$ -test was used to determine potential differences between mean  $F_i/F_m$  values measured on various days throughout the growth curve of *E. huxleyi*. To determine the effect of SOD additions on *in vivo* fluorescence, a mixed factor repeated measures ANOVA was used. To determine the effect of dilution on per-cell  $O_2^-$  production and the effect of SOD addition on growth rates, cell abundances, cell biovolume, and  $F_i/F_m$  values, a comparison of means using an independent two sample Student's  $t$ -test was employed for each parameter interrogated. A one-sample Student's  $t$ -test was used to determine the potential difference between *E. huxleyi*  $O_2^-$  production in the presence and absence of light. For all statistical analyses, the significance threshold ( $\alpha$ ) was set to 0.05.

## RESULTS

### Extracellular $O_2^-$ Production as a Function of Growth Phase

To assess per-cell extracellular  $O_2^-$  modulation across different average metabolic states, cell-normalized extracellular  $O_2^-$  production by *E. huxleyi* was measured throughout the growth curve of batch cultures. To rule out potential cell density effects, the number of cells analyzed at each time point was kept constant [average  $\pm$  SE was  $8.32 \times 10^5 \pm 5.98 \times 10^4$  cells ( $n = 30$ )]. The highest per-cell net extracellular  $O_2^-$  production rates were observed during early exponential growth when  $F_i/F_m$  values were the highest (**Supplementary Figure S1**) and significantly declined as *E. huxleyi* grew over time (Spearman's  $\rho = -0.58$ ;  $p < 0.0001$ ) (**Figure 1**). For example, *E. huxleyi* produced maximum amounts of  $O_2^-$  (average  $\pm$  SE) at the first time point measured in early exponential phase ( $4,478 \pm 611 \text{ amol cell}^{-1} \text{ h}^{-1}$ ,  $n = 3$ ; day 2). These maximal rates were nearly 78 times higher than net production rates measured between day 8 and day 25, when average production declined to  $58 \pm 55 \text{ amol cell}^{-1} \text{ h}^{-1}$  ( $n = 30$ ). Flow cytometry analyses revealed this decline was not due to an increase in senescent cells (i.e., cells with low chlorophyll). After day 8, net cell-normalized  $O_2^-$  production rates were occasionally negative, fluctuating between  $-253 \pm 167$  (day 22;  $n = 3$ ) and  $209 \pm 360$  (day 17;  $n = 3$ ). Net per-cell  $O_2^-$  production rates account for the simultaneous production and decay of  $O_2^-$  at the cell surface. Because auto-oxidation of the MCLA probe results in a small amount of  $O_2^-$  production (Fujimori et al., 1993), the negative net per-cell  $O_2^-$  production rates between day 8 and day 25 reflect degradation of  $O_2^-$  originating from the MCLA reagent.



## Extracellular $O_2^-$ Production as a Function of Cell Density

Two approaches were undertaken to determine the potential effect of cell density on extracellular  $O_2^-$  production. First, short-term effects (sec-min) were tested by measuring cell-normalized  $O_2^-$  production rates and total  $O_2^-$  concentrations while increasing the number of cells loaded on the FeLume filter in both exponential and stationary growth phase. The total  $O_2^-$  concentration increased significantly with increasing cell density during both exponential (Spearman's  $\rho = 0.92$ ;  $p < 0.0001$ ) and stationary phase (Spearman's  $\rho = 0.89$ ;  $p < 0.0001$ ). Conversely, net per-cell  $O_2^-$  production rates decreased significantly with increasing cell density during exponential phase (Spearman's  $\rho = -0.74$ ;  $p < 0.001$ ) but not during stationary phase (Spearman's  $\rho = -0.20$ ;  $p = 0.45$ ) (Figure 2). In exponential phase, average net per-cell  $O_2^-$  production rates decreased by more than 200% from the highest ( $9.2 \times 10^6$ ) to lowest ( $4.6 \times 10^5$ ) number of cells analyzed.

To assess longer-term effects (min-h) of cell density on extracellular  $O_2^-$  production, exponentially growing cells were preconditioned to lower cell densities by diluting cultures with 0.22  $\mu\text{m}$  filtered, autoclaved seawater and incubating for 0–6.5 h prior to conducting  $O_2^-$  measurements. In this experiment, the number of cells loaded on the FeLume filter at each dilution level was kept constant [average  $\pm$  SE was  $1.32 \times 10^6 \pm 8.47 \times 10^4$  cells ( $n = 26$ )]. A 10- and 100-fold dilution of *E. huxleyi* resulted in a 51 and 172% increase in cell-normalized net extracellular  $O_2^-$  production rates, respectively.


Although average rates increased at both dilution levels, only the 100-fold dilution resulted with significantly more extracellular  $O_2^-$  per cell than the undiluted control ( $t$ -test;  $p < 0.05$ ) (Figure 3). The 0.22  $\mu\text{m}$  filtered, autoclaved seawater diluent was also measured for  $O_2^-$  to ensure the  $O_2^-$  measured in the diluted samples was not due to the seawater diluent. At maximum,  $O_2^-$  produced in the seawater diluent could only account for 3.0–4.6% of the total steady-state  $O_2^-$  concentrations measured in experiments with *E. huxleyi*, confirming that the contribution from the diluent was negligible.

## SOD Addition Experiments

To assess how extracellular  $O_2^-$  influences growth and physiology, *E. huxleyi* cultures were grown with a range of SOD concentrations and monitored until the end of exponential phase. Since SOD is a large enzyme (>31 kDa) (Cass, 1985), it cannot passively cross cell membranes and therefore selectively targets  $O_2^-$  within the extracellular milieu. Overall, the addition of SOD stimulated growth (Figure 4; Supplementary Figure S2; Supplementary Table S2). For example, *in vivo* fluorescence was significantly different in cultures with various SOD concentrations (mixed factor repeated measures ANOVA;  $p < 0.001$ ) and became more significant over time (mixed factor repeated measures ANOVA;  $p < 0.0001$ ) (Figure 4A; Supplementary Table S2). In addition, specific growth rates were between 15 and 22% higher ( $t$ -test;  $p < 0.05$ ) in the presence of SOD (Figure 4B; Supplementary Table S2). Cell abundances from cultures grown with 100 U  $\text{ml}^{-1}$  SOD were significantly higher ( $t$ -test;  $p < 0.05$ ) than cell abundances from cultures grown without SOD beginning on day 7 (Figure 4C; Supplementary Table S2), when cell concentrations in the highest SOD addition were 41% higher than the unamended treatment.


To confirm whether these results were specifically due to SOD, a control experiment was performed in which SOD was removed via diafiltration and only the low molecular weight fraction (<10 kDa) of the SOD suspension was added to cultures. *In vivo* fluorescence was significantly different between treatments (mixed factor repeated measures ANOVA;  $p < 0.05$ ) with significance increasing over time (mixed factor repeated measures ANOVA;  $p < 0.0001$ ) (Supplementary Figure S2A; Supplementary Table S2). Cultures grown with SOD grew significantly faster ( $t$ -test;  $p < 0.05$ ) than those grown with dialyzed SOD (Figure 2B; Supplementary Table S2). By the end of the control experiment (day 12), cell abundances from cultures grown with dialyzed SOD were significantly lower ( $t$ -test;  $p < 0.05$ ) than those grown with SOD by about 32% (Supplementary Figure S2C; Supplementary Table S2). Thus, the effect of SOD addition on growth rates and growth yields could not be accounted for by the dialyzed SOD control (Supplementary Figure S2; Supplementary Table S2).

In addition to growth yields and growth rates, the cellular biovolume of *E. huxleyi* was monitored in SOD addition experiments. Biovolume was calculated using three different methods, but regardless of the calculation method used, cellular biovolume was larger in cultures grown with SOD ( $t$ -test;  $p < 0.0001$ ) (Figure 5A; Supplementary Figure S3A;



Supplementary Table S2). Adding SOD increased biovolume between 6.3 and 22.1%, depending on the SOD concentration and calculation method used. In the control experiment with dialyzed SOD, biovolume of cells grown with SOD was significantly larger ( $t$ -test;  $p < 0.0001$ ) than those grown with dialyzed SOD by at least 19% (Supplementary Figure S3A; Supplementary Table S2) confirming the response was due to the presence of SOD.


In addition to stimulating growth, SOD improved photophysiological health, as evidenced by increased  $F_v/F_m$  values, which indicated more efficient light assimilation by PSII in the presence of SOD. By day 10, adding SOD at each concentration increased  $F_v/F_m$  between 11 and 18% compared to the 0  $\text{U ml}^{-1}$  SOD control ( $t$ -test;  $p < 0.05$ ) (Figure 5B; Supplementary Table S2). A similar trend was



seen in the control experiment with dialyzed SOD, where  $F_v/F_m$  values from cultures grown with SOD were significantly higher than those from cultures grown with dialyzed SOD starting on day 7 ( $t$ -test;  $p < 0.05$ ) (Supplementary Figure S3B; Supplementary Table S2). Therefore, improvements in photophysiological health are attributed to the effects of SOD.

### Extracellular $O_2^-$ Production in the Presence and Absence of Light


To investigate whether extracellular  $O_2^-$  production by *E. huxleyi* is dependent on light, extracellular  $O_2^-$  production was measured in ambient light and dark conditions. A representative FeLume time series measurement of  $O_2^-$  concentration showed that extracellular  $O_2^-$  production by *E. huxleyi* under ambient light reached and stabilized at  $2,395 \pm 27$  pM (Figure 6). Upon removal of light, there was an immediate decline in production, which stabilized at  $1,085 \pm 29$  pM after  $\sim 160$  s in dark conditions. This result could not be accounted for by abiotic factors, as the removal of light had insignificant effects on  $O_2^-$  production in the absence of *E. huxleyi* cells. The drawdown of signal below the abiotic  $O_2^-$  baseline by SOD confirmed that the biogenic signal was indeed due to  $O_2^-$  production (Figure 6). All biological replicates produced less  $O_2^-$  in dark compared to ambient light conditions ( $t$ -test;  $p < 0.0001$ ), indicating a connection with photophysiological processes (Figure 7). Dark conditions inhibited  $O_2^-$  production by an average of 70%.



## DISCUSSION

Here, we investigated the cellular regulation and potential physiological roles of extracellular  $O_2^-$  production in a noncalcifying strain (CCMP 374) of the cosmopolitan marine coccolithophore *E. huxleyi* to approach a broader understanding of the potential ecological and environmental impacts of phytoplankton-derived extracellular  $O_2^-$ . The ability to produce extracellular  $O_2^-$  is widespread among phytoplankton, yet production rates can vary substantially within and between species (Diaz and Plummer, 2018). For example, compared to *Chattonella* spp., the most prolific microbial ROS producers, maximal extracellular  $O_2^-$  production rates by *E. huxleyi* were about 100–10,000 times lower. Indeed, *E. huxleyi* produced  $O_2^-$  at a rate more similar to nonharmful algae such as *Symbiodinium* spp. and *Thalassiosira* spp. (Diaz and Plummer, 2018).

The extracellular  $O_2^-$  concentrations and production rates by *E. huxleyi* were not due to the release of intracellular  $O_2^-$ . First, the release of intracellular  $O_2^-$  through cell lysis would have been detected as a rapidly decaying pulse of  $O_2^-$ , but this can be ruled out because  $O_2^-$  signals were stable for at least 1–2 min (see section “Materials and Methods”), a significant timeframe compared to the typical half-life of  $O_2^-$  in our analysis (~2 min) (Diaz et al., 2013). Second, the physiochemical nature of the  $O_2^-$  anion prevents it from passively crossing intact cell membranes (Bielski et al., 1985; Lesser, 2006; Brown and Griendling, 2009). Therefore, the production rates measured in this study reflect active production of  $O_2^-$  on or near the surface of *E. huxleyi*. The reported rates of extracellular  $O_2^-$  production reflect the balance of gross production and decay at the cell surface, thus giving a net production rate. Therefore, any change in the net production rate of extracellular  $O_2^-$  could result from a change in gross production, decay, or both. Several abiotic




and biotic factors have the potential to degrade extracellular  $\text{O}_2^-$  at or near the cell surface, such as interactions with trace metals including soluble and mineral-bound iron (Fuji et al., 2006; Heller and Croot, 2010a) and expression of cell surface SODs (Aguirre et al., 2005; Oshikawa et al., 2010; Bauer, 2014).

ROS production is commonly associated with stress; however, *E. huxleyi* produced extracellular  $\text{O}_2^-$  without any added stressors. In fact, cell-normalized  $\text{O}_2^-$  production by *E. huxleyi* was the highest in early exponential phase under presumably the least taxing culture conditions when nutrient concentrations were the highest (Figure 1) and when photosynthetic efficiency was at a maximum (Supplementary Figure S1). These results reflect the rates of extracellular  $\text{O}_2^-$  production by cells that were removed from the prevailing culture conditions and analyzed *ex situ*. The *ex situ* analytical conditions such as pH (7.6) were identical across culture samples of all ages, yet differed from *in situ* levels (Supplementary Figure S4). These results therefore show that as cultures age, there is a shift toward lower  $\text{O}_2^-$  production when cells are analyzed under the same conditions. The potential effect of pH on extracellular  $\text{O}_2^-$  production by *E. huxleyi* is unknown, but an increase in pH stimulates extracellular ROS production by *C. marina* (Liu et al., 2007), which is opposite to the trend reported here (Figure 1; Supplementary Figure S4). The observed decline in extracellular  $\text{O}_2^-$  production with culture age therefore suggests that extracellular  $\text{O}_2^-$  production is physiologically driven and unlikely related to a stress response, as levels of stress would presumably increase with time in batch culture due to the depletion of resources. These findings do not rule out the potential for *E. huxleyi* to upregulate extracellular  $\text{O}_2^-$  under stressful conditions, as seen with extracellular  $\text{H}_2\text{O}_2$  (Evans et al., 2006) and intracellular ROS (Evans et al., 2006; Vardi et al., 2012) during viral infection, but do demonstrate a physiological investment in the production of extracellular  $\text{O}_2^-$  under ideal growth conditions, which suggests some role in basal metabolism. In order to clarify if and how *E. huxleyi* regulates extracellular  $\text{O}_2^-$  in response to biogeochemical variability and stress, future work should consider the influence of factors such as viral infection and pH.


In addition to having growth phase dependence (Figure 1) similar to other phytoplankton (Oda et al., 1995; Kim et al., 1999; Portune et al., 2010), cell-normalized net extracellular  $\text{O}_2^-$  production rates by *E. huxleyi* were also inversely dependent on cell density over a range of timescales during exponential growth phase. For example, this trend occurred when cell density increased on time scales of seconds to minutes (Figure 2A) and when cells were preconditioned to lower cell densities on timescales of minutes to hours (Figure 3). This tight regulation may suggest a dynamic cell density-dependent signaling role for  $\text{O}_2^-$  production in *E. huxleyi*. For instance, extracellular  $\text{O}_2^-$  deriving from one cell may act as a signal between other cells or within the same cell to provide information on surrounding population density (Diaz and Plummer, 2018). Similar cell density dependent trends in extracellular  $\text{O}_2^-$  production have been demonstrated in other phytoplankton (Marshall et al., 2005a; Hansel et al., 2016; Diaz et al., 2018), including *C. marina* (Marshall et al., 2005b).

Although evidence has been accumulating that extracellular  $\text{O}_2^-$  production is directly involved in growth promotion in a range of microbial cell types (Oda et al., 1995; Aguirre et al., 2005), our results show that the presence of extracellular  $\text{O}_2^-$  does not directly stimulate growth of *E. huxleyi*. The potential role of extracellular  $\text{O}_2^-$  in *E. huxleyi* growth was addressed through selective removal of extracellular  $\text{O}_2^-$  from the local environment





**FIGURE 6 |** FeLume time-series of  $O_2^-$  measurements under different light conditions on day 2 of *E. huxleyi* growth (biological replicate B) split into five regions: (1) phosphate buffer solution and MCLA reagent baseline (which is subtracted from the biogenic  $O_2^-$  concentration in regions 3 and 4), (2) shaded region showing loading of *E. huxleyi* cells while the pump was stopped, (3) *E. huxleyi* in ambient light, (4) the second shaded region showing *E. huxleyi* in the dark, and (5) drawdown of the  $O_2^-$  signal below the baseline after addition of SOD (negative  $O_2^-$  concentrations account for SOD driven degradation of  $O_2^-$  originating from auto-oxidation of the MCLA reagent). The average  $\pm$  SE of stable steady-state  $O_2^-$  concentration measurements are indicated by horizontal pink lines.



**FIGURE 7 |**  $O_2^-$  measurements from three biological replicates of *E. huxleyi* in the presence of ambient light ( $\sim 5 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) and in the dark ( $0 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). Asterisks show significant difference ( $p < 0.0001$ ) between the average dark and average ambient light steady-state  $O_2^-$  concentration for each biological replicate. Error bars represent one standard error of the mean ( $n \geq 31$  chemiluminescent counts).

of cells using SOD. Scavenging  $O_2^-$  promoted growth (Figure 4), increased cell biovolume (Figure 5A), and improved photosynthetic efficiency (Figure 5B). Conversely, removal of extracellular  $O_2^-$  from cultures of *Chattonella* spp. attenuates growth (Tanaka et al., 1992; Oda et al., 1995). Specifically, in one prior study, *C. marina* growth was significantly hampered under similar SOD concentrations used in the present study, and the morphological state of cells was altered (Oda et al., 1995). Recently,

Hansel et al. (2019) revealed that the growth of common marine bacteria from the *Roseobacter* clade was inhibited by SOD in a dose-dependent manner (Hansel et al., 2019). Similarly, the removal of extracellular ROS from fungi and the amoebozoan *Dictyostelium discoideum* is also detrimental to development (Aguirre et al., 2005). Taken together, extracellular  $O_2^-$  does not seem to directly stimulate growth in *E. huxleyi*. Rather, these results may highlight a different role for extracellular  $O_2^-$  in *E. huxleyi* that contrasts with the proposed growth-promoting role of extracellular  $O_2^-$  in *C. marina* (Oda et al., 1995), bacteria (Hansel et al., 2019), fungi, and protists (Aguirre et al., 2005). However, the addition of SOD not only removes  $O_2^-$  but produces  $H_2O_2$ , which may also have impacts on *E. huxleyi* physiology. For example, high concentrations of  $H_2O_2$  are harmful to phytoplankton (Dupouy et al., 1985; Morris et al., 2011), but normal growth of *C. marina* is dependent on low levels of extracellular  $H_2O_2$  (Oda et al., 1995). In fact, in a variety of cell types, the dismutation of Nox-derived  $O_2^-$  by cell surface SOD generates extracellular  $H_2O_2$ , which can diffuse into the cell, to elicit gene expression (Shapiguzov et al., 2012), morphogenesis (Rossi et al., 2017), and proliferation (Oshikawa et al., 2010; Bauer, 2014). It remains possible, yet speculative, that the addition of SOD in our experiments accelerated the dismutation of  $O_2^-$  to  $H_2O_2$ , which then may have acted as a growth promoter for *E. huxleyi*. Thus, the role of extracellular  $O_2^-$  in *E. huxleyi* may be contingent on its ability to give rise to  $H_2O_2$ , which should be interrogated in future work.

The fact that there are clear trends in extracellular  $O_2^-$  production as a function of growth phase and cell density, but that *E. huxleyi* growth was not stunted with removal of extracellular  $O_2^-$ , leaves the possibility open that extracellular  $O_2^-$  could be connected to other aspects of *E. huxleyi* physiology and health. To examine whether extracellular  $O_2^-$  production may be involved in photosynthetic physiology, we interrogated  $O_2^-$  production as a function of light and found that  $O_2^-$  production was attenuated within seconds upon transition from light to dark conditions (Figure 6).

This finding adds to a growing body of evidence linking modulation of extracellular  $O_2^-$  production by phytoplankton to light availability and therefore photophysiology. For instance, extracellular  $O_2^-$  production is light dependent in many phytoplankton including *Thalassiosira* spp. (Milne et al., 2009; Schneider et al., 2016), *Trichodesmium* (Hansel et al., 2016), *Symbiodinium* (modulated on the same time scales shown here) (Saragosti et al., 2010), and *Chattonella* spp. (Kim et al., 1999; Marshall et al., 2002; Dorantes-Aranda et al., 2013). Thus,  $O_2^-$  production may somehow be involved with light dependent processes (e.g., photosynthesis, photoacclimation physiology), and this functionality may be conserved across phytoplankton taxa. Interestingly, in a previous study, when *C. marina* was treated with DCMU [3-(3,4-dichlorophenyl)-1,1-dimethylurea], an electron transfer inhibitor between photosystem II and I, extracellular  $O_2^-$  production was quenched to levels observed under dark conditions (Marshall et al., 2002), further illustrating a mechanistic link between extracellular  $O_2^-$  production and photophysiology.

Overall, this study reveals that the stress-independent production of extracellular  $O_2^-$  by *E. huxleyi* is dynamically regulated, and potentially part of a basal process involved with photophysiology. Extracellular  $O_2^-$  production by *E. huxleyi* is conceivably part of healthy cellular functioning for several reasons. First, the fact that cells diverted energy toward making  $O_2^-$  in the absence of a stressor suggests its production can be unassociated with stress and probably related to basal functioning. Indeed, *E. huxleyi* produced maximum amounts of extracellular  $O_2^-$  per cell under ideal growth conditions while cells were most metabolically active (**Figure 1**) and when efficiency of photosynthetic processes was the highest (**Supplementary Figure S1**). Further, the steady-state concentrations of  $O_2^-$  generated by *E. huxleyi* cells are not consistent with concentrations that would be damaging ( $>10^{-6}$  M) but are consistent with concentrations of biological signaling molecules ( $\sim 10^{-12}$  M) (Saran, 2003). Additionally, *E. huxleyi* cells tightly controlled the production of extracellular  $O_2^-$  as a function of light, cell density, and growth phase, and on timescales as short as seconds, consistent with other phytoplankton (Oda et al., 1995; Kim et al., 1999; Marshall et al., 2002, 2005a,b; Milne et al., 2009; Portune et al., 2010; Saragosti et al., 2010; Dorantes-Aranda et al., 2013; Hansel et al., 2016; Schneider et al., 2016; Diaz et al., 2018), suggesting a potential dynamic role in signaling and photophysiology. Indeed, stress-independent extracellular  $O_2^-$  production by *E. huxleyi* is in agreement with several other microorganisms such as phytoplankton and bacteria where its production is prolific under ideal growth conditions (Oda et al., 1995; Kustka et al., 2005; Rose et al., 2005; Marshall et al., 2005a,b; Godrant et al., 2009; Portune et al., 2010; Diaz et al., 2013; Hansel et al., 2016; Schneider et al., 2016). The fact that *E. huxleyi* still produces extracellular  $O_2^-$  in the absence of light suggests there could be additional and/or alternative purposes for this production beyond photophysiology. Indeed, extracellular  $O_2^-$  production can be produced through a variety of subcellular mechanisms and could be produced for a combination of ecophysiological functions (Diaz and Plummer, 2018).

Contrasting with other diverse microorganisms (Tanaka et al., 1992; Oda et al., 1995; Saran, 2003; Aguirre et al., 2005; Hansel et al., 2019), the presence of extracellular  $O_2^-$  does not promote growth in *E. huxleyi*. This finding underscores a potentially unprecedented role for the presence and/or production of extracellular  $O_2^-$ , which should be investigated among other phytoplankton. Finally, non-calcifying *E. huxleyi* cells coexist with calcareous varieties in nature but typically are not dominant (Frada et al., 2012). Several ecological and physiological processes are variable within (Strom et al., 2003; Strom and Bright, 2009; Sunda and Hardison, 2010; Harvey et al., 2015; Poulson-Ellestad et al., 2016) and between (Paasche, 2002; Suggett et al., 2007; Harvey et al., 2015; Poulson-Ellestad et al., 2016) calcifying and noncalcifying strains. Whether the rates, regulation, and roles of extracellular  $O_2^-$  production are different in calcifying versus noncalcifying strains of *E. huxleyi* has yet to be determined but should be considered in future work.

## DATA AVAILABILITY

Any materials and data will be made available to members of the scientific community upon request.

## AUTHOR CONTRIBUTIONS

SP and JD conceived the study. SP conducted the experiments and data analysis, with contributions from AT. All authors contributed to interpretation of results and preparing the manuscript.

## FUNDING

This research was supported by a Junior Faculty Seed Grant from the University of Georgia Research Foundation (JD), a National Science Foundation (NSF) Graduate Research Fellowship (SP), and NSF grant OCE-1355720 (CH). The FlowCam® and FIRE were purchased through a NSF Equipment Improvement Grant (1624593).

## ACKNOWLEDGMENTS

The authors wish to thank Dee King for her assistance with literature searches and reference indexing, Dr. Patricia Yager and Dr. Brian Hopkinson for use of their labs, and Rachel Steffen for assistance in the lab.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2019.01546/full#supplementary-material>

## REFERENCES

- Aguirre, J., and Lambeth, J. D. (2010). Nox enzymes from fungus to fly to fish and what they tell us about Nox function in mammals. *Free Radic. Biol. Med.* 49, 1342–1353. doi: 10.1016/j.freeradbiomed.2010.07.027
- Aguirre, J., Rios-Momberg, M., Hewitt, D., and Hansberg, W. (2005). Reactive oxygen species and development in microbial eukaryotes. *Trends Microbiol.* 13, 111–118. doi: 10.1016/j.tim.2005.01.007
- Anderson, A., Laohavisit, A., Blaby, I. K., Bombelli, P., Howe, C. J., Merchant, S. S., et al. (2015). Exploiting algal NADPH oxidase for biophotovoltaic energy. *Plant Biotechnol. J.* 14, 22–28. doi: 10.1111/pbi.12332
- Bauer, G. (2014). Targeting extracellular ROS signaling of tumor cells. *Anticancer Res.* 34, 1467–1482. Retrieved from: <http://ar.iiarjournals.org>.
- Bielski, B. H. J., Cabelli, D. E., Arudi, R. L., and Ross, A. B. (1985). Reactivity of  $H_2O/O_2^-$  radicals in aqueous-solution. *J. Phys. Chem. Ref. Data* 14, 1041–1100. doi: 10.1063/1.555739
- Brown, D. I., and Griendling, K. K. (2009). Nox proteins in signal transduction. *Free Radic. Biol. Med.* 47, 1239–1253. doi: 10.1016/j.freeradbiomed.2009.07.023
- Brown, C. W., and Yoder, J. A. (1994). Coccolithophorid blooms in the global ocean. *J. Geophys. Res. Oceans* 99, 7467–7482. doi: 10.1029/93JC02156
- Cass, A. E. G. (1985). “Superoxide dismutases” in *Metalloproteins*. ed. P. M. Harrison (London: Palgrave Macmillan UK), 121–156.
- Diaz, J. M., Hansel, C. M., Apprill, A., Brighi, C., Zhang, T., Weber, L., et al. (2016). Species-specific control of external superoxide levels by the coral holobiont during a natural bleaching event. *Nat. Commun.* 7:13801. doi: 10.1038/ncomms13801
- Diaz, J. M., Hansel, C. M., Voelker, B. M., Mendes, C. M., Andeer, P. F., and Zhang, T. (2013). Widespread production of extracellular superoxide by heterotrophic bacteria. *Science* 340, 1223–1226. doi: 10.1126/science.1237331
- Diaz, J. M., and Plummer, S. (2018). Production of extracellular reactive oxygen species by phytoplankton: past and future directions. *J. Plankton Res.* 40, 655–666. doi: 10.1093/plankt/fby039
- Diaz, J. M., Plummer, S., Tomas, C., and Alves-De-Souza, C. (2018). Production of extracellular superoxide and hydrogen peroxide by five marine species of harmful bloom-forming algae. *J. Plankton Res.* 40, 667–677. doi: 10.1093/plankt/fby043
- Dorantes-Aranda, J. J., Nichols, P. D., Waite, T. D., and Hallegraef, G. M. (2013). Strain variability in fatty acid composition of *Chattonella marina* (Raphidophyceae) and its relation to differing ichthyotoxicity toward rainbow trout gill cells. *J. Phycol.* 49, 427–438. doi: 10.1111/jpy.12053
- Dorantes-Aranda, J. J., Seger, A., Mardones, J. I., Nichols, P. D., and Hallegraef, G. M. (2015). Progress in understanding algal bloom-mediated fish kills: the role of superoxide radicals, phycotoxins and fatty acids. *PLoS One* 10:e0133549. doi: 10.1371/journal.pone.0133549
- Dupouy, D., Conter, A., Croute, F., Murat, M., and Planel, H. (1985). Sensitivity of *Synechococcus lividus* to hydrogen peroxide. *Environ. Exp. Bot.* 25, 339–347. doi: 10.1016/0098-8472(85)90031-0
- Evans, C., Malin, G., Mills, G. P., and Wilson, W. H. (2006). Viral infection of *Emiliana huxleyi* (prymnesiophyceae) leads to elevated production of reactive oxygen species. *J. Phycol.* 42, 1040–1047. doi: 10.1111/j.1529-8817.2006.00256.x
- Flores, H. S., Wikfors, G., and Dam, H. (2012). Reactive oxygen species are linked to the toxicity of the dinoflagellate *Alexandrium* spp. to protists. *Aquat. Microb. Ecol.* 66, 199–209. doi: 10.3354/ame01570
- Frada, M. J., Bidle, K. D., Probert, I., and De Vargas, C. (2012). In situ survey of life cycle phases of the coccolithophore *Emiliana huxleyi* (Haptophyta). *Environ. Microbiol.* 14, 1558–1569. doi: 10.1111/j.1462-2920.2012.02745.x
- Fujii, M., Rose, A. L., Waite, T. D., and Omura, T. (2006). Superoxide-mediated dissolution of amorphous ferric oxyhydroxide in seawater. *Environ. Sci. Technol.* 40, 880–887. doi: 10.1021/es051622t
- Fujimori, K., Nakajima, H., Akutsu, K., Mitani, M., Sawada, H., and Nakayama, M. (1993). Chemiluminescence of *Cypridina* luciferin analogues. Part 1. Effect of pH on rates of spontaneous autoxidation of CLA in aqueous buffer solutions. *J. Chem. Soc. Perkin Trans. 2*, 2405–2409.
- Garg, S., Rose, A. L., Godrant, A., and Waite, T. D. (2007). Iron uptake by the ichthyotoxic *Chattonella marina* (Raphidophyceae): impact of superoxide generation. *J. Phycol.* 43, 978–991. doi: 10.1111/j.1529-8817.2007.00394.x
- Godrant, A., Rose, A. L., Sarthou, G., and Waite, T. D. (2009). New method for the determination of extracellular production of superoxide by marine phytoplankton using the chemiluminescence probes MCLA and red-CLA. *Limnol. Oceanogr. Methods* 7, 682–692. doi: 10.4319/lom.2009.7.682
- Guillard, R. R. L., and Ryther, J. H. (1962). Studies of marine planktonic diatoms: I. *Cyclotella nana* Husted, and *detonula confervacea* (Cleve) gran. *Can. J. Microbiol.* 8, 229–239. doi: 10.1139/m62-029
- Hansard, P. S., Vermilyea, A. W., and Voelker, B. M. (2010). Measurements of superoxide radical concentration and decay kinetics in the Gulf of Alaska. *Deep-Sea Res. I Oceanogr. Res. Pap.* 57, 1111–1119. doi: 10.1016/j.dsr.2010.05.007
- Hansel, C. M., Buchwald, C., Diaz, J. M., Ossolinski, J. E., Dyhrman, S. T., Mooy, V., et al. (2016). Dynamics of extracellular superoxide production by *Trichodesmium* colonies from the Sargasso Sea. *Limnol. Oceanogr.* 61, 1188–1200. doi: 10.1002/lno.10266
- Hansel, C. M., Diaz, J. M., and Plummer, S. (2019). Tight regulation of extracellular superoxide points to its vital role in the physiology of the globally relevant *Roseobacter* clade. *MBio*. 10, e02668–e02618. doi: 10.1128/mBio.02668-18
- Harvey, E. L., Bidle, K. D., and Johnson, M. D. (2015). Consequences of strain variability and calcification in *Emiliana huxleyi* on microzooplankton grazing. *J. Plankton Res.* 37, 1137–1148. doi: 10.1093/plankt/fbv081
- Heller, M. I., and Croot, P. L. (2010a). Application of a superoxide ( $O_2^-$ ) thermal source (SOTS-1) for the determination and calibration of  $O_2^-$  fluxes in seawater. *Anal. Chim. Acta* 667, 1–13. doi: 10.1016/j.aca.2010.03.054
- Heller, M. I., and Croot, P. L. (2010b). Kinetics of superoxide reactions with dissolved organic matter in tropical Atlantic surface waters near Cape Verde (TENATSO). *J. Geophys. Res.* 115:C12038. doi: 10.1029/2009JC006021
- Hervé, C., Tonon, T., Collén, J., Corre, E., and Boyen, C. (2006). NADPH oxidases in eukaryotes: red algae provide new hints! *Curr. Genet.* 49, 190–204. doi: 10.1007/s00294-005-0044-z
- Kim, C. S., Lee, S. G., Lee, C. K., Kim, H. G., and Jung, J. (1999). Reactive oxygen species as causative agents in the ichthyotoxicity of the red tide dinoflagellate *Cochlodinium polykrikoides*. *J. Plankton Res.* 21, 2105–2115. doi: 10.1093/plankt/21.11.2105
- Kim, D., Nakamura, A., Okamoto, T., Komatsu, N., Oda, T., Iida, T., et al. (2000). Mechanism of superoxide anion generation in the toxic red tide phytoplankton *Chattonella marina*: possible involvement of NAD(P)H oxidase. *Biochim. Biophys. Acta* 1524, 220–227. doi: 10.1016/s0304-4165(00)00161-6
- Kim, D., Nakashima, T., Matsuyama, Y., Niwano, Y., Yamaguchi, K., and Oda, T. (2007). Presence of the distinct systems responsible for superoxide anion and hydrogen peroxide generation in red tide phytoplankton *Chattonella marina* and *Chattonella ovata*. *J. Plankton Res.* 29, 241–247. doi: 10.1093/plankt/fbm011
- Kim, D., and Oda, T. (2010). “Possible factors responsible for the fish-killing mechanisms of the red tide phytoplankton, *Chattonella marina* and *Cochlodinium polykrikoides*” in *Coastal Environmental and Ecosystem Issues of the East China Sea*, eds. A. Ishimatsu and H.-J. Lie (Tokyo, Japan: TERRAPUB and Nagasaki University), 245–268.
- Kustka, A. B., Shaked, Y., Milligan, A. J., King, D. W., and Morel, F. M. M. (2005). Extracellular production of superoxide by marine diatoms: contrasting effects on iron redox chemistry and bioavailability. *Limnol. Oceanogr.* 50, 1172–1180. doi: 10.4319/lo.2005.50.4.1172
- Lesser, M. P. (2006). Oxidative stress in marine environments: biochemistry and physiological ecology. *Annu. Rev. Physiol.* 68, 253–278. doi: 10.1146/annurev.physiol.68.040104.110001
- Liu, W., Au, D. W. T., Anderson, D. M., Lam, P. K. S., and Wu, R. S. S. (2007). Effects of nutrients, salinity, pH and light:dark cycle on the production of reactive oxygen species in the alga *Chattonella marina*. *J. Exp. Mar. Biol. Ecol.* 346, 76–86. doi: 10.1016/j.jembe.2007.03.007
- Mardones, J. I., Dorantes-Aranda, J. J., Nichols, P. D., and Hallegraef, G. M. (2015). Fish gill damage by the dinoflagellate *Alexandrium catenella* from Chilean fjords: synergistic action of ROS and PUFA. *Harmful Algae* 49, 40–49. doi: 10.1016/j.hal.2015.09.001
- Marshall, J.-A., De Salas, M., Oda, T., and Hallegraef, G. (2005a). Superoxide production by marine microalgae: I. Survey of 37 species from 6 classes. *Mar. Biol.* 147, 553–540. doi: 10.1007/s00227-005-1596-7
- Marshall, J.-A., De Salas, M., Oda, T., and Hallegraef, G. (2005b). Superoxide production by marine microalgae II. Towards understanding ecological

- consequences and possible functions. *Mar. Biol.* 147, 533–540. doi: 10.1007/s00227-005-1597-6
- Marshall, J.-A., Hovenden, M., Oda, T., and Hallegraef, G. M. (2002). Photosynthesis does influence superoxide production in the ichthyotoxic alga *Chattonella marina* (Raphidophyceae). *J. Plankton Res.* 24, 1231–1236. doi: 10.1093/plankt/24.11.1231
- Marshall, J.-A., Nichols, P. D., Hamilton, B., Lewis, R. J., and Hallegraef, G. M. (2003). Ichthyotoxicity of *Chattonella marina* (Raphidophyceae) to damselfish (*Acanthochromis polycanthus*): the synergistic role of reactive oxygen species and free fatty acids. *Harmful Algae* 2, 273–281. doi: 10.1016/S1568-9883(03)00046-5
- Martel, C. M. (2009). Conceptual bases for prey biorecognition and feeding selectivity in the microplanktonic marine phagotroph *Oxyrrhis marina*. *Microb. Ecol.* 57, 589–597. doi: 10.1007/s00248-008-9421-8
- Milne, A., Davey, M. S., Worsfold, P. J., Achterberg, E. P., and Taylor, A. R. (2009). Real-time detection of reactive oxygen species generation by marine phytoplankton using flow injection-chemiluminescence. *Limnol. Oceanogr. Methods* 7, 706–715. doi: 10.4319/lom.2009.7.706
- Mittler, R., Vanderauwera, S., Suzuki, N., Miller, G., Tognetti, V. B., Vandepoel, K., et al. (2011). ROS signaling: the new wave? *Trends Plant Sci.* 16, 300–309. doi: 10.1016/j.tplants.2011.03.007
- Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., and Zinser, E. R. (2011). Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS One* 6:e16805. doi: 10.1371/journal.pone.0016805
- Oda, T., Ishimatsu, A., Shimada, M., Takeshita, S., and Muramatsu, T. (1992). Oxygen-radical-mediated toxic effects of the red tide flagellate *Chattonella marina* on *Vibrio alginolyticus*. *Mar. Biol.* 112, 505–509. doi: 10.1007/BF00356297
- Oda, T., Moritomi, J., Kawano, I., Hamaguchi, S., Ishimatsu, A., and Muramatsu, T. (1995). Catalase- and superoxide dismutase-induced morphological changes and growth inhibition in the red tide phytoplankton *Chattonella marina*. *Biosci. Biotechnol. Biochem.* 59, 2044–2048.
- Oda, T., Nakamura, A., Shikayama, M., Kawano, I., Ishimatsu, A., and Muramatsu, T. (1997). Generation of reactive oxygen species by raphidophycean phytoplankton. *Biosci. Biotechnol. Biochem.* 61, 1658–1662.
- Oshikawa, J., Urao, N., Kim, H. W., Kaplan, N., Razvi, M., McKinney, R., et al. (2010). Extracellular SOD-derived H<sub>2</sub>O<sub>2</sub> promotes VEGF signaling in caveolae/lipid rafts and post-ischemic angiogenesis in mice. *PLoS One* 5:e10189. doi: 10.1371/journal.pone.0010189
- Paasche, E. (2002). A review of the coccolithophorid *Emiliana huxleyi* (Prymnesiophyceae), with particular reference to growth, coccolith formation, and calcification-photosynthesis interactions (Phycological Reviews 20). *Phycologia* 40, 503–529. doi: 10.2216/i0031-8884-40-6-503.1
- Palenik, B., Zafriou, O. C., and Morel, F. M. M. (1987). Hydrogen peroxide production by a marine phytoplankter 1. *Limnol. Oceanogr.* 32, 1365–1369. doi: 10.4319/lo.1987.32.6.1365
- Portune, K. J., Cary, S. C., and Warner, M. E. (2010). Antioxidant enzyme response and reactive oxygen species production in marine raphidophytes. *J. Phycol.* 46, 1161–1171. doi: 10.1111/j.1529-8817.2010.00906.x
- Poulson-Ellestad, K. L., Harvey, E. L., Johnson, M. D., and Mincer, T. J. (2016). Evidence for strain-specific exometabolic responses of the coccolithophore *Emiliana huxleyi* to grazing by the dinoflagellate *Oxyrrhis marina*. *Front. Mar. Sci.* 3:1. doi: 10.3389/fmars.2016.00001
- Roe, K. L., and Barbeau, K. A. (2014). Uptake mechanisms for inorganic iron and ferric citrate in *Trichodesmium erythraeum* IMS101. *Metallomics* 6, 2042–2051. doi: 10.1039/c4mt00026a
- Rose, A. L. (2012). The influence of extracellular superoxide on iron redox chemistry and bioavailability to aquatic microorganisms. *Front. Microbiol.* 3:124. doi: 10.3389/fmicb.2012.00124
- Rose, A. L., Salmon, T. P., Lukondeh, T., Neilan, B. A., and Waite, T. D. (2005). Use of superoxide as an electron shuttle for iron acquisition by the marine cyanobacterium *Lyngbya majuscula*. *Environ. Sci. Technol.* 39, 3708–3715. doi: 10.1021/es048766c
- Rose, A. L., Webb, E. A., Waite, T. D., and Moffett, J. W. (2008). Measurement and implications of nonphotochemically generated superoxide in the equatorial Pacific Ocean. *Environ. Sci. Technol.* 42, 2387–2393. doi: 10.1021/es7024609
- Rossi, D. C. P., Gleason, J. E., Sanchez, H., Schatzman, S. S., Culbertson, E. M., Johnson, C. J., et al. (2017). *Candida albicans* FRE8 encodes a member of the NADPH oxidase family that produces a burst of ROS during fungal morphogenesis. *PLoS Pathog.* 13:e1006763. doi: 10.1371/journal.ppat.1006763
- Saragosti, E., Tchernov, D., Katsir, A., and Shaked, Y. (2010). Extracellular production and degradation of superoxide in the coral *Stylophora pistillata* and cultured *Symbiodinium*. *PLoS One* 5:e12508. doi: 10.1371/journal.pone.0012508
- Saran, M. (2003). To what end does nature produce superoxide? NADPH oxidase as an autocrine modifier of membrane phospholipids generating paracrine lipid messengers. *Free Radic. Res.* 37, 1045–1059. doi: 10.1080/10715760310001594631
- Schneider, R. J., Roe, K. L., Hansel, C. M., and Voelker, B. M. (2016). Species-level variability in extracellular production rates of reactive oxygen species by diatoms. *Front. Chem.* 4:5. doi: 10.3389/fchem.2016.00005
- Shapiguzov, A., Vainonen, J., Wrzaczek, M., and Kangasjärvi, J. (2012). ROS-talk – how the apoplast, the chloroplast, and the nucleus get the message through. *Front. Plant Sci.* 3. doi: 10.3389/fpls.2012.00292
- Strom, S. L., and Bright, K. J. (2009). Inter-strain differences in nitrogen use by the coccolithophore *Emiliana huxleyi*, and consequences for predation by a planktonic ciliate. *Harmful Algae* 8, 811–816. doi: 10.1016/j.hal.2007.10.005
- Strom, S., Wolfe, G., Holmes, J., Stecher, H., Shimeneck, C., Lambert, S., et al. (2003). Chemical defense in the microplankton I: feeding and growth rates of heterotrophic protists on the DMS-producing phytoplankter *Emiliana huxleyi*. *Limnol. Oceanogr.* 48, 217–229. doi: 10.4319/lo.2003.48.1.0217
- Suggett, D. J., Le Flo'ch, E., Harris, G. N., Leonardos, N., and Geider, R. J. (2007). Different strategies of photoacclimation by two strains of *Emiliana huxleyi* (Haptophyta). *J. Phycol.* 43, 1209–1222. doi: 10.1111/j.1529-8817.2007.00406.x
- Sunda, W. G., and Hardison, D. R. (2010). Evolutionary tradeoffs among nutrient acquisition, cell size, and grazing defense in marine phytoplankton promote ecosystem stability. *Mar. Ecol. Prog. Ser.* 401, 63–76. doi: 10.3354/meps08390
- Tanaka, K., Yoshimatsu, S., and Shimada, M. (1992). Generation of superoxide anions by *Chattonella antiqua*: possible causes for fish death caused by 'Red Tide'. *Experientia* 48, 888–890. doi: 10.1007/BF02118427
- Vardi, A., Haramaty, L., Van Mooy, B. A., Fredricks, H. F., Kimman, S. A., Larsen, A., et al. (2012). Host–virus dynamics and subcellular controls of cell fate in a natural coccolithophore population. *Proc. Natl. Acad. Sci.* 109, 19327–19332. doi: 10.1073/pnas.1208895109
- Westbroek, P., Young, J. R., and Linschooten, K. (1989). Coccolith production (Biominalization) in the marine alga *Emiliana huxleyi*. *J. Protozool.* 36, 368–373. doi: 10.1111/j.1550-7408.1989.tb05528.x
- Wuttig, K., Heller, M. I., and Croot, P. L. (2013). Reactivity of inorganic Mn and Mn desferrioxamine B with O<sub>2</sub>, O<sub>2</sub><sup>-</sup>, and H<sub>2</sub>O<sub>2</sub> in seawater. *Environ. Sci. Technol.* 47, 10257–10265. doi: 10.1021/es4016603
- Yang, C. Z., Albright, L. J., and Yousif, A. N. (1995). Oxygen-radical-mediated effects of the toxic phytoplankter heterosigma-carterae on juvenile rainbow-trout *Oncorhynchus mykiss*. *Dis. Aquat. Org.* 23, 101–108. doi: 10.3354/dao023101
- Zhang, T., Diaz, J. M., Brighi, C., Parsons, R. J., McNally, S., Apprill, A., et al. (2016a). Dark production of extracellular superoxide by the coral *Porites astreoides* and representative symbionts. *Front. Mar. Sci.* 3:232. doi: 10.3389/fmars.2016.00232
- Zhang, T., Hansel, C. M., Voelker, B. M., and Lamborg, C. H. (2016b). Extensive dark biological production of reactive oxygen species in brackish and freshwater ponds. *Environ. Sci. Technol.* 50, 2983–2993. doi: 10.1021/acs.est.5b03906
- Zinser, E. R. (2018). The microbial contribution to reactive oxygen species dynamics in marine ecosystems. *Environ. Microbiol. Rep.* 10, 412–427. doi: 10.1111/1758-2229.12626

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Plummer, Taylor, Harvey, Hansel and Diaz. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.