



OPEN ACCESS

EDITED BY

Aurel Popa-Wagner,
University of Medicine and Pharmacy of
Craiova,
Romania

REVIEWED BY

Moises Freitas-Andrade,
Ottawa Hospital Research Institute (OHRI),
Canada
Javier Palazuelos,
Complutense University of Madrid,
Spain

*CORRESPONDENCE

Colin J. Murray
✉ Colinmur@uvic.ca
Marie-Ève Tremblay
✉ evetremblay@uvic.ca

SPECIALTY SECTION

This article was submitted to
Cellular and Molecular Mechanisms of Brain-
aging,
a section of the journal
Frontiers in Aging Neuroscience

RECEIVED 08 December 2022

ACCEPTED 22 February 2023

PUBLISHED 22 March 2023

CITATION

Murray CJ, Vecchiarelli HA and Tremblay M-È
(2023) Enhancing axonal myelination in
seniors: A review exploring the potential impact
cannabis has on myelination in the aged brain.
Front. Aging Neurosci. 15:1119552.
doi: 10.3389/fnagi.2023.1119552

COPYRIGHT

© 2023 Murray, Vecchiarelli and Tremblay. This
is an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Enhancing axonal myelination in seniors: A review exploring the potential impact cannabis has on myelination in the aged brain

Colin J. Murray^{1,2*}, Haley A. Vecchiarelli² and
Marie-Ève Tremblay^{2,3,4,5,6,7,8*}

¹Neuroscience Graduate Program, University of Victoria, Victoria, BC, Canada, ²Division of Medical Sciences, University of Victoria, Victoria, BC, Canada, ³Département de Médecine Moléculaire, Université Laval, Québec City, QC, Canada, ⁴Axe Neurosciences, Center de Recherche du CHU de Québec, Université Laval, Québec City, QC, Canada, ⁵Neurology and Neurosurgery Department, McGill University, Montréal, QC, Canada, ⁶Department of Biochemistry and Molecular Biology, University of British Columbia, Vancouver, BC, Canada, ⁷Centre for Advanced Materials and Related Technology (CAMTEC), University of Victoria, Victoria, BC, Canada, ⁸Institute for Aging and Lifelong Health, University of Victoria, Victoria, BC, Canada

Consumption of cannabis is on the rise as public opinion trends toward acceptance and its consequent legalization. Specifically, the senior population is one of the demographics increasing their use of cannabis the fastest, but research aimed at understanding cannabis' impact on the aged brain is still scarce. Aging is characterized by many brain changes that slowly alter cognitive ability. One process that is greatly impacted during aging is axonal myelination. The slow degradation and loss of myelin (i.e., demyelination) in the brain with age has been shown to associate with cognitive decline and, furthermore, is a common characteristic of numerous neurological diseases experienced in aging. It is currently not known what causes this age-dependent degradation, but it is likely due to numerous confounding factors (i.e., heightened inflammation, reduced blood flow, cellular senescence) that impact the many cells responsible for maintaining overall homeostasis and myelin integrity. Importantly, animal studies using non-human primates and rodents have also revealed demyelination with age, providing a reliable model for researchers to try and understand the cellular mechanisms at play. In rodents, cannabis was recently shown to modulate the myelination process. Furthermore, studies looking at the direct modulatory impact cannabis has on microglia, astrocytes and oligodendrocyte lineage cells hint at potential mechanisms to prevent some of the more damaging activities performed by these cells that contribute to demyelination in aging. However, research focusing on how cannabis impacts myelination in the aged brain is lacking. Therefore, this review will explore the evidence thus far accumulated to show how cannabis impacts myelination and will extrapolate what this knowledge may mean for the aged brain.

KEYWORDS

aged brain, myelination, myelin repair, cannabis, microglia, oligodendrocyte, oligodendrocyte progenitor cell, astrocyte

1. Introduction

Research looking into the potential therapeutic benefits offered by cannabis has drastically increased since its legalization (in Canada, medicinal: 2001, recreational: 2018) in many countries around the world. The proposed therapeutic benefits of cannabis consumption are numerous, ranging from pain management to a potential aid in multiple sclerosis (MS; Whiting et al., 2015; Paes-Colli et al., 2022). The possible benefit offered in MS—an autoimmune disease characterized by demyelination—introduces an interesting association between cannabinoids, the biologically active compounds found in cannabis, and myelination (Longoria et al., 2022).

Myelination is an essential process that involves the efficient and deft wrapping of myelin—a lipid rich sheath—around the axons of neurons by oligodendrocytes in the central nervous system (CNS). This wrapping facilitates rapid propagation of electrical signals and is essential for neuronal synchronization and proper communication between discrete regions of the brain. In adulthood, the total net level of myelin in the brain is relatively constant, but the myelin sheaths themselves turnover in a slow conserved cycle between degradation and regeneration, a process facilitated by oligodendrocytes (Buscham et al., 2019; Aber et al., 2022; Meschkat et al., 2022). However, this homeostatic cycle is lost in the aged brain, leading to an abnormal deposition of myelin and a net decline in myelin content (Peters, 2009; Rivera et al., 2022). This decline is evident in the healthy aged brain and is furthermore a common characteristic of many neurodegenerative diseases associated with aging (Guttmann et al., 1998; Bartzokis, 2004; Cox et al., 2016; Coelho et al., 2021; Furber et al., 2022). Importantly, this loss is tightly linked to cognitive decline (Bartzokis, 2004; Bennett and Madden, 2014; Wang et al., 2020; Coelho et al., 2021).

Due to the prevalence of demyelination in the aged brain and its association with cognitive decline, there is an urgent need to better understand and alleviate the burdens of this process. Certain lifestyle factors such as diet and exercise have recently emerged as a promising way to improve cognition and brain health throughout the lifespan. A controversial lifestyle factor that has relatively unknown cellular effects on the brain during aging is cannabis use.

In recent years, seniors (aged 65+) have increased their use of cannabis faster than any other demographic in North America, possibly as a result of some combination of destigmatization, legalization, and increased accessibility (Salas-Wright et al., 2017; Han and Palamar, 2020; Keethakumar et al., 2021). However, research focusing on cannabis use in seniors is scarce, not to mention research specific to its impact on myelination. Therefore, it is prudent to identify what impact cannabis use has on the integrity of myelin in the aged brain in order to propose harm reduction strategies if needed. Alternatively, the potential for cannabinoids to therapeutically target demyelinating diseases points to an ability for cannabis to regulate the myelination process in a beneficial way. This lack of evidence regarding the beneficial or detrimental outcomes of cannabis on myelination in the aged brain highlights a clear gap in the literature. By excluding seniors, the currently available research is omitting a large portion of the population that would not only benefit from increased research, but also requests more information about the outcomes of cannabis use (Bobitt et al., 2019).

This review will outline how cannabis influences the myelination process in the CNS by examining its impact on different cell types, and


will discuss how cannabis may alter the relationships between neurons and glial cells, particularly in the aged brain. The aim of this review is to highlight the potential for cannabis to modulate myelination in the aged brain and to emphasize the paucity of research in this area in order to stimulate future research.

1.1. The importance of myelin

In 1854, Rudolf Ludwig Virchow coined a term for a ubiquitous substance in the brain—myelin (Boullerne, 2016). It would take another 100 years before the central function performed by this lipid-rich sheath—saltatory conduction—was agreed upon. The rapid propagation of electrical information within the myelinated axon during saltatory conduction is possible due to the insulating properties of myelin and the nodes of Ranvier (Huxley and Stämpeli, 1949). The nodes of Ranvier are unmyelinated sections of the axon that have a high density of voltage-gated sodium (Na_v) channels that respond rapidly to alterations in charge and function to propagate signals necessary for the depolarization of the next node during an action potential (Arancibia-Carcamo and Attwell, 2014). Myelin is therefore deposited around the axon in sections, known as internodes, which creates the boundaries for the nodes of Ranvier (Figure 1). The internodes effectively insulate the axon by increasing the resistance of the axonal membrane and by reducing the capacitance of the axon (Bakiri et al., 2011; Stadelmann et al., 2019).

Mounting evidence indicates that myelination does not become fixed after development, but is experience-driven and remains adaptive well into adulthood (Young et al., 2013; Bechler et al., 2015; Fields, 2015; Ford et al., 2015; Hill et al., 2018; Hughes et al., 2018; Jünemann et al., 2022). Various structural modifications can alter the efficiency of this system. For example, the density of Na_v channels within the node and the length of the node itself can both alter conduction velocity with minimal energy expenditure (Arancibia-Carcamo et al., 2017). The diameter of the axon, thickness of the myelin sheath, and the length of the internode can also adjust conduction velocity; with wider axons, thicker sheaths, and longer internodes increasing conduction velocity up to a certain point (Waxman, 1980; Wu et al., 2012; Chapman and Hill, 2020). A large determinant of this adaptability is experience-driven neuronal activity, which has been shown to promote myelination and contribute to the modification of established myelin sheaths (Wake et al., 2011; Gibson et al., 2014; Bechler et al., 2018; Faria et al., 2019). These modifications then alter conduction speed, translating into variations in synapse strength and, therefore, synaptic plasticity (Fields, 2015).

The structural features of the myelin sheath are also important for maintaining the synchronization of action potentials within and between neurons. This synchronization is fundamental to circuit function and proper cognition. Experience-induced changes in myelination can alter this synchronicity and synaptic plasticity to optimize processing time within the circuit contextually, contributing to complex cognitive processes like social behavior, memory, motor learning and sensory experience throughout the lifespan (Liu et al., 2012; Mount and Monje, 2017; Pan et al., 2020). However, the long-range projection neurons that make up the bulk of these white matter pathways are the most vulnerable neurons during aging, and their deterioration can result in negative effects on the aforementioned cognitive processes (Mattson and Magnus, 2006).



1.1.1. Myelin in the aged brain

It is now apparent that myelination does not peak until mid-life, with peak white matter volume occurring between 30 and 50-years of age in humans (Bartzokis et al., 2001; Sowell et al., 2003; Westlye et al., 2010; Buyanova and Arsalidou, 2021). Non-invasive neuroimaging techniques like diffusion tensor imaging (DTI) have been widely utilized to determine the state of white matter in the human brain throughout the lifespan. These studies have found a significant reduction in white matter volume and alterations in the integrity of myelin that suggest deterioration during aging (Pakkenberg and Gundersen, 1997; Guttmann et al., 1998; Liu et al., 2017; Vinke et al., 2018; Faizy et al., 2020). Electron microscopy performed in aged non-human primates and rodents have further clarified this

deterioration by revealing the structural alterations present in the myelin sheath, such as redundant myelination and reduced myelin thickness (Figure 1; Peters, 2009; Shepherd et al., 2012; Attia et al., 2019; Phillips et al., 2019). These alterations also contribute to the disorganization of ion channels at the nodes of Ranvier and on the axonal membrane at the paranode—sections of the internode directly adjacent to the node—which likely have negative consequences on signal transduction (Hinman et al., 2006).

The macro-scale loss and micro-scale alterations in myelin impact the conduction speed and synchronization of action potentials, ultimately causing latencies and overall disruptions of neuronal communication that likely contribute to driving cognitive deficits in the aging population (Bartzokis, 2004; Bowley et al., 2010; Attia et al.,

2019). Cortical disconnection due to myelin loss has been proposed as a likely candidate for reduced cognition in aged individuals for decades. For example, early *in vivo* evidence in support of this hypothesis using DTI concluded that aged individuals (56–85-years of age) show an age-related decline in white matter, especially in the frontal lobe, which was linked with impairments in executive functioning (O'Sullivan et al., 2001). This observed decline has been confirmed by more recent studies showing impairments in memory, processing speed, attention, and general cognition, which were linked to reduced myelin content (Brickman et al., 2012; Cremers et al., 2016; Coelho et al., 2021). It is important to keep in mind that these alterations occur in the “normal” aging brain, but are exacerbated in neurodegenerative diseases like Alzheimer's disease (Papuć and Rejdak, 2020). Normal, or healthy, aging is an ill-defined term referring to the natural aging process that is devoid of significant physical or cognitive impairments and allows for the maintenance of subjective well-being (Wong, 2018). However, non-debilitating impairments are present in healthy aging, and may contribute to the progression of more serious disabilities.

The mechanisms underlying the natural degradation of myelin and the insufficiency of remyelination in the aged brain have not yet been definitively identified. However, it is probable that there are numerous confounding factors, of which many are expected to be tightly associated with the glial cells responsible for the deposition, maintenance and modification of the myelin sheath, as discussed below (Figure 2A).

1.2. Oligodendrocytes and oligodendrocyte progenitor cells

In the CNS, oligodendrocyte processes wrap around axons and form the myelin sheath (Boullerne, 2016). These glial cells are capable of extending many different processes to wrap multiple axons at a time, with the ability to differentially alter the parameters of each extension due to the activity of each individual neuron contacted (Chong et al., 2012; Yeung et al., 2014). Furthermore, mature oligodendrocytes are capable of facilitating remyelination after damage in cats and non-human primates, while maintaining established myelin sheaths (Duncan et al., 2018). One particularly relevant finding is that oligodendrocytes preferentially myelinate active axons in development and adulthood, as found in primary cell cultures, zebrafish and mice (Gibson et al., 2014; Hines et al., 2015; Wake et al., 2015; Mitew et al., 2018; Faria et al., 2019). Furthermore, through DTI, neuronal circuits activated during a task were found to have increased levels of myelination in seniors, showing the ability for activity-dependent myelination in the aged human brain (Scholz et al., 2009; Jünemann et al., 2022).

The myelin sheath further provides a channel for metabolic support to the axon. Oligodendrocytes transport various energy metabolites like lactate from their soma through distinct cytoplasmic channels to the innermost layer of the myelin sheath where they are deposited into the periaxonal space (Figure 1; Fünfschilling et al., 2012; Meyer et al., 2018). The subsequent uptake of metabolites supports axon function and is essential for neuronal survival (Lee et al., 2012; Meyer et al., 2018).

However, before an oligodendrocyte can mature and participate in myelination, a complex differentiation process from




FIGURE 2

The transitional relationship between myelination and aging and the possible therapeutic advantage of cannabinoids | As emphasized throughout this review, the process of myelination is an intricate undertaking that involves all glial cells in the brain and is mainly driven by experience induced neuronal activity in adulthood and during aging. (A) The top panel of this figure shows some of the general contributions glial cells and neurons make to the overall process of myelination. With increasing age, a switch in cell state takes place in glial cells and neurons, ultimately leading to improper myelination and impairments in cognition. (B) The bottom panel showcases some of the benefits that cannabis (mainly focusing on THC and CBD) can have on myelination, as found in animal studies; with the majority looking at younger time points. Although evidence suggests improved myelination and cognition from cannabis use in aging animals, the paucity of studies focusing on this time point translates to relatively unknown overall effects of cannabis on myelination. This question mark indicates that topic, which is in need of increased research. Not only will this help fill in the gap of knowledge as to how cannabis impacts myelination across the lifespan, but will also better inform researchers on the effects of the endocannabinoid system on the aged brain. With this understanding, we are also better able to inform the public and health authorities on the impact cannabis use has on the senior population, who at the moment show increasing levels of consumption. Created with BioRender.com.

oligodendrocyte progenitor cell (OPC) must occur. These cells not only populate the developing brain, but remain present in the adult

brain, providing a reserve for the replacement of the mature oligodendrocyte population (Bergles and Richardson, 2016). On top of this, OPCs were shown to form direct contacts with neuronal synapses, to participate in synaptic engulfment, and to engage in axon pruning during development and adulthood, thus contributing to circuit formation and modulation in mice (Bergles et al., 2000; Auguste et al., 2022; Buchanan et al., 2022). Neuronal activity conveyed through these synapses relays information to OPCs that contributes to determining their proliferation, migration and differentiation, as well as oligodendrocyte survival, and myelination as a whole (Mitew et al., 2018; Moura et al., 2022).

1.2.1. Oligodendrocytes and OPCs in the aged brain

The degeneration and improper renewal of myelin in the aged brain is predictably associated with the oligodendrocyte population. Many of the extrinsic factors that impact oligodendrocyte function [e.g., pro-inflammatory cytokines, reactive oxygen species (ROS)] are the products of other glial cells and will be discussed in more detail subsequently. Changes in axonal signaling can also present challenges for continued myelination. As described, neuronal activity induces myelination—therefore, it is likely that altered activity due to neuronal dysfunction may lead to changes in activity-dependent myelination (Sams, 2021). Furthermore, mitochondrial dysfunction in the neurons of aged mice (12-month-old) resulted in decreased production of ATP and increased production of ROS, which may subsequently damage oligodendrocytes and OPCs over time, thereby preventing OPC differentiation and myelination (Stahon et al., 2016; Spaas et al., 2021). Although the axons remain functional and oligodendrocytes remain active in 12-month-old mice, neuronal viability will likely decline due to reduced ATP and increased production of free radicals, perhaps resulting in degeneration with increasing age (Stahon et al., 2016). Conversely, age-related degeneration of myelin, oligodendrocytes and the neuron-oligodendrocyte relationship also likely leads to a decrease in metabolic support provided by oligodendrocytes, further impacting neuronal function (Hill et al., 2018; Zhang X. et al., 2021).

In aging, oligodendrocytes and OPCs also accumulate signs of oxidative DNA (mitochondrial and nuclear) damage, a feature which is commonly found in neurodegenerative diseases like Alzheimer's disease and MS (Tse and Herrup, 2017). Oligodendrocytes and OPCs are particularly vulnerable to oxidative stress due to their extremely high metabolic demand needed for the endogenous production of myelin and their decreased ability to deal with free radicals (French et al., 2009; Giacci et al., 2018). For example, oligodendrocyte lineage cells were shown to have only half the glutathione—a major intracellular antioxidant—content compared to astrocytes in primary cell cultures from rats (Juurink et al., 1998). Importantly, synthesis of glutathione naturally declines during aging in mice, further rendering oligodendrocyte lineage cells susceptible to damage and dysfunction (Wang, 2003). Additionally, OPCs are particularly susceptible to damage by oxidizing agents because of a delay in the production or reduced activity of antioxidant enzymes (e.g., glutathione peroxidase), which increases with maturation (Back et al., 1998; Baud, 2004; Spaas et al., 2021). A recent review hypothesizes that this increased vulnerability to oxidative stress may inhibit OPC differentiation (Spaas et al., 2021).

A decline in newly formed mature oligodendrocytes has been observed in the aged brain (Soreq et al., 2017; Hill et al., 2018; Wang

et al., 2020; Rivera et al., 2021; Dimovasili et al., 2022). However, OPC density does not seem to decline with age when compared across the lifespan in mice, non-human primates or humans (Sim et al., 2002; Doucette et al., 2010; Yeung et al., 2014; Wang et al., 2020; Dimovasili et al., 2022). Although it is generally accepted that the OPC population remains stable in the aged brain, a more recent study found significantly reduced OPC (NG2⁺) density in the corpus callosum of 18-month-old mice (Rivera et al., 2021). Nonetheless, the inability of OPCs to differentiate into mature oligodendrocytes in the aged brain is not currently debated, and it likely contributes to the reduced capacity for remyelination and declining oligodendrocyte population (Sim et al., 2002; Neumann et al., 2019; Segel et al., 2019; Rivera et al., 2021; Zhang X. et al., 2021; Dimovasili et al., 2022). This reduced ability to differentiate may be a consequence of a markedly different proteome in aged OPCs (de la Fuente et al., 2020). For example, aldehyde dehydrogenase 1 family member A1 (ALDH1A1) and transcription factor 4 (TCF4) are both involved in OPC differentiation and have notably reduced expression levels in aged rats (>15-months-old; de la Fuente et al., 2020).

OPCs also display differing levels of ion channels and receptors depending on age and brain region (Spitzer et al., 2019). For example, the density of N-methyl-D-aspartate (NMDA) receptors on OPCs significantly declines with increasing age (>6-months-old) in multiple regions of the mouse brain, including the corpus callosum (Spitzer et al., 2019). This change likely affects activity-dependent myelination in the aged brain by significantly impacting the ability for OPCs to sense and act on glutamate released from active neurons (Gautier et al., 2015; Spitzer et al., 2019). Interestingly, this NMDA receptor-mediated activity-dependent myelination requires the simultaneous presence of glutamate and growth factors like brain derived neurotrophic factor (BDNF), hinting at the importance of surrounding glial cells (Lundgaard et al., 2013). As a note, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)/kainate receptors are important for the initial stages of remyelination after experimentally induced demyelination, where the binding of glutamate released from active axons within the lesion promotes proliferation, survival and differentiation of OPCs during development and adulthood in rodents (Gautier et al., 2015; Kougiountzidou et al., 2017). Although AMPA/kainate receptor levels remain relatively constant or increase with age, the impaired ability for neurons to form new synaptic contacts with OPCs, combined with a decline in NMDA receptors, may decrease OPC differentiation and activity-dependent myelination in the aged brain (Spitzer et al., 2019; Sams, 2021).

A general increase in ROS and free radicals in the aged brain and a phenomenon known as “niche stiffening” of the extracellular matrix (ECM) directly adjacent to OPCs can also prevent their differentiation (French et al., 2009; Segel et al., 2019). Niche stiffening is a dynamic process that physically alters the elasticity of tissue, which increases the rigidity of the ECM in the brain of aged mice (>14-months-old; Swift et al., 2013; Segel et al., 2019). The mechanosensitive ion channel PIEZO1 was found to be essential for the OPC detection of the elasticity of the ECM, and the knockdown of PIEZO1 resulted in increased proliferation and differentiation of OPCs in the aged CNS of mice (Segel et al., 2019). Furthermore, the introduction of aged OPCs into the ECM within the prefrontal cortex of neonatal rats rescued their proliferative and differentiating abilities, indicating the importance of the environment for these OPC functions (Segel et al., 2019).

Microglia and astrocytes may also influence OPC cell fate through their modulation of the ECM. For instance, primary microglial cells from aged (18–24-months-old) rats treated with an inflammatory stimulus [i.e., transforming growth factor- β (TGF- β)] creates a microglial-deposited ECM that promotes the differentiation of OPCs into astrocytes, thereby preventing oligodendrocyte differentiation and myelination (Baror et al., 2019). Furthermore, oligodendrocytes in aged mice (18-months-old) release factors that promote microglial-mediated survival of oligodendrocytes, but prevent OPC differentiation (Luan et al., 2021). Similarly, astrocytes were found to inhibit OPC differentiation and disrupt remyelination by releasing chondroitin sulfate proteoglycans into the ECM in primary mixed glial cell cultures from mice (Keough et al., 2016).

Lastly, as mentioned, neurotrophic factors (e.g., BDNF) released from glial cells impact the ability for OPCs to contribute to activity-dependent myelination. These are just a few examples of the influence surrounding glial cells have on OPCs, oligodendrocytes and the process of myelination. However, astrocytes and microglia are profoundly altered structurally and functionally in the aged brain, and these changes extend to the cells and structures they support (Figure 2A).

1.3. Astrocytes in myelination

Astrocytes are an extremely diverse group of glial cells that contribute to the blood–brain barrier (BBB), blood flow modulation, metabolite supply, and perform modulatory roles at the synapse involved in synaptic activity and plasticity (Sofroniew and Vinters, 2010). Furthermore, astrocytes are vital for myelination. The loss of astrocytes reduces the density of oligodendrocytes, initiates demyelination, and promotes myelin abnormalities (e.g., decompaction) in the white matter of developing and adolescent mice (7 days to ~1.5-months-old; Tognatta et al., 2020). These effects are likely partly due to local increases in extracellular glutamate causing excitotoxicity, and reduced trophic support [e.g., platelet-derived growth factor (PDGF)] from astrocytes (Tognatta et al., 2020). The uptake of glutamate from the extracellular space is an essential function performed by astrocytes that prevents excitotoxicity of neurons and glial cells (McDonald et al., 1998; Hassel et al., 2003; Matute et al., 2007; Goursaud et al., 2009; Mahmoud et al., 2019). Additionally, the controlled release of gliotransmitters (e.g., glutamate) from astrocytic hemichannels—membrane channels between cells and the extracellular space made up of connexin proteins—directly to synapses modulates synaptic transmission and plasticity, and was even shown to be essential for behavioral outputs including fear memory consolidation in rats (~2-months-old; Ye et al., 2003; Stehberg et al., 2012; Abudara et al., 2018).

Astrocytes also maintain appropriate levels of K^+ ions in the extracellular space and effectively disperse them throughout the pan-glial network that spans the entire brain (Beckner, 2020). The pan-glial network allows for the diffusion of ions and small metabolites between coupled cells connected through gap junctions, which are made up of adjoining hemichannels (Orthmann-Murphy et al., 2008; Stephan et al., 2021). The loss of astrocyte-oligodendrocyte gap junctions results in myelin pathology (e.g., vacuolation) and loss of astrocytes (Magnotti et al., 2011; Tress et al., 2012). Additionally, many human diseases characterized by demyelination (e.g., MS and

neuromyelitis optica) show early disruption of gap junctions between astrocytes and oligodendrocytes and a decline in connexin proteins (Markoullis et al., 2012; Masaki, 2015).

Astrocytes are also a main source of cholesterol, facilitate iron transport, and directly provide oligodendrocytes with metabolic support (*via* gap junctions) in the adult CNS, all contributing to the processes needed to synthesize myelin (Jurevics and Morell, 2002; Schulz et al., 2012; Saher and Stumpf, 2015; Camargo et al., 2017; Cheli et al., 2020). Lastly, a wide range of soluble factors [e.g., BDNF and chemokine (C-X-C motif) ligand 1 (CXCL1)] released by astrocytes can have myriad effects on myelination, as found in rodents (Tsai et al., 2002; Fulmer et al., 2014; Kiray et al., 2016).

1.3.1. Astrocytes and myelination in the aged brain

Due to the vast number of functions performed by astrocytes in the CNS, age-related dysfunction of these cells predictably has wide-ranging impacts on brain function. Firstly, observations in aged mice (20–24-months-old) reveal morphological changes in astrocytes and alterations in territorial domain that may result in reduced contacts between adjacent astrocytes, disconnecting them from the greater pan-glial network which is essential for many homeostatic functions (e.g., K^+ spatial buffering, metabolic and cholesterol supply to oligodendrocytes; Grosche et al., 2013; Popov et al., 2021; Verkhratsky et al., 2022). A steady reduction in astrocytic glutamate transporters, reduced capacity to buffer and disperse K^+ , and an overall decrease in their ability to sense synaptic activity (partly due to reduced density of ionotropic receptors) was also observed in aged mice (20–24-months-old), disrupting long-term potentiation of synapses in the hippocampus (Lalo et al., 2011; Popov et al., 2021). This is likely partly due to excess levels of glutamate in the extracellular space resulting in excitotoxicity, which damages neurons, oligodendrocytes, and myelin (Olney, 1971; Matute et al., 2007; Fu et al., 2009). Additionally, the increase in hemichannel activation due to an increase in pro-inflammatory cytokines [e.g., tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β)] in primary cell cultures and mice results in an increase in the release of various ions (i.e., K^+ , Ca^{2+}) and gliotransmitters (i.e., ATP, glutamate) into the extracellular space, further disrupting homeostasis and possibly contributing to neuronal death (Retamal et al., 2007; Froger et al., 2010; Karpuk et al., 2011; Orellana et al., 2011; Satarker et al., 2022). The cholesterol synthesis pathway is also significantly altered in astrocytes from aged mice (24-months-old), likely contributing to the observed decline of cholesterol in the aged brain and potentially hindering production of myelin (Boisvert et al., 2018; Palmer and Ousman, 2018).

An altered gene expression profile is further observed in aged astrocytes. Increases in genes associated with cytokine pathways, antigen presentation, the complement cascade, and reactivity [e.g., glial fibrillary acidic protein (*Gfap*)] were observed in astrocytes in the hippocampus, hypothalamus, visual cortex, striatum and cerebellum of aged mice (24-months-old; Boisvert et al., 2018; Clarke et al., 2018). GFAP is an intermediate filament protein commonly used as a marker for astrocytes, which significantly increases in pathological-like states (e.g., aging and MS), indicating heightened astrocyte reactivity in rodents and humans (Nichols et al., 1993; Saraste et al., 2021).

This elevated expression of GFAP has also been linked to astrocyte senescence (Salminen et al., 2011; Boisvert et al., 2018). Senescence refers to cells that enter into a distinct state characterized by

dysfunctional mitochondria, increased production of ROS, and an altered secretory profile as a consequence of DNA damage, telomere shortening and an altered environment (Sikora et al., 2021). The downstream effects of these pathways exacerbate inflammation and impair myelination *via* inhibition of the OPC cycle and loss of functional support for oligodendrocytes by astrocytes, as shown in primary cell cultures and in mice (Palmer and Ousman, 2018; Willis et al., 2020). Interestingly, astrocytes often do not transition into this more damaging/senescent-like state unless microglia induce the switch through the release of cytokines like IL-1 α and TNF (Herx et al., 2000; Liddelow et al., 2017; Clarke et al., 2018; Jha et al., 2019). There are numerous reasons for an altered microglial secretory profile in aging, but one specific to white matter regions could be an increase in myelin debris that cannot be efficiently cleared/metabolized by microglia, resulting in cellular stress (Safaiyan et al., 2016).

1.4. Microglia in myelination

Microglia—the resident immune cells of the CNS—perform vital functions in all stages of life. They act as gardeners, constantly surveying their surroundings looking for debris to clear, shaping and pruning synapses, maintaining appropriate glial and neuronal population sizes, modulating neuronal activity, and releasing various trophic factors to support growth and development of glial cells and neurons (Šimončičová et al., 2022). Microglia also play a substantial role in myelination, contributing to the developmental and experience-driven process of adaptive myelination (Kalafatakis and Karagozeos, 2021; Santos and Fields, 2021). In fact, microglia are present at higher densities in human white matter compared to gray matter, highlighting their importance in this environment (Mittelbronn et al., 2001; Askew et al., 2017). However, evidence is more contradictory for microglial density in mice. One study found increased density of microglia in the white matter of the forebrain, whereas an earlier study found increased density in gray matter from the entire mouse brain (Lawson et al., 1990; Savchenko et al., 2000). Therefore, it is important to keep in mind the region analyzed, as microglial density and function can greatly vary. Additionally, it is important to note that the microglial population is not homogenous, but instead exists as a continuum of states that contribute in divergent fashions to supporting brain development, activity, plasticity and integrity (Paolicelli et al., 2022).

Although microglia have a plethora of functions, three of their activities primarily contribute to myelination. (1) Microglia release a repertoire of soluble factors [e.g., insulin growth factor-1 (IGF-1), IL-1 β , TGF- β] that facilitate the promotion and prevention of myelination (Hsieh et al., 2004; Pang et al., 2007; Santos and Fields, 2021; McNamara et al., 2023). (2) Microglia phagocytose myelin debris, which is important as myelin debris can inhibit OPC differentiation, while efficient clearance of myelin debris allows for effective remyelination after experimental demyelination in rodents (Kotter, 2006; Neumann et al., 2008; Lampron et al., 2015). It was also reported that microglia are capable of removing incorrectly deposited myelin directly from the axon, contributing to the refinement of myelin sheaths, as shown in zebrafish and mice during development (Hughes and Appel, 2020; Djannatian et al., 2023). (3) Microglia dynamically contact active axons, guided by the nodal efflux of K⁺ ions. This interaction was associated with improved remyelination

after experimental demyelination in mice, and may be a way by which microglia prevent neuronal damage from hyperactivity (Madry et al., 2018; Ronzano et al., 2021). Microglia respond rapidly to hyperactive neurons and wrap their processes around axons to facilitate rapid repolarization, thus preventing excitotoxicity and maintaining neuronal viability in mice (Kato et al., 2016). Therefore, microglia are emerging as essential modulators of neuronal activity that substantially contribute to determining neuronal architecture and function (Badimon et al., 2020; Cserép et al., 2021).

1.4.1. Microglia and myelination in the aged brain

Microglia are not immune to the challenges of aging. Changes observed in aged mice (≥ 12 -months-old) include an upregulation of genes associated with the immune response, and a decrease in genes associated with environment probing and interactions with the ECM (Grabert et al., 2016; Angelova and Brown, 2019). Furthermore, an age-related metamorphosis in their secretory profile, resulting in increased pro-inflammatory markers is also a common characteristic of microglia, as observed in those sorted from the aged mouse brain (≥ 18 -months-old; Holling et al., 2004; Sierra et al., 2007; Norden and Godbout, 2013; Koellhoffer et al., 2017; Marschallinger et al., 2020). For instance, aged mice have microglia with increased levels of the nod-like receptor protein 3 (NLRP3) inflammasome (Youm et al., 2013). After inflammasome activation, an increase in the production of IL-1 β , IL-6, TNF- α and others are observed in the mouse brain (Youm et al., 2013; Hu et al., 2019). Many of these compounds are beneficial and required for proper myelination, however, a problem arises when this activation becomes chronic, as is the case in the aged brain (Tilstra et al., 2011). Prolonged activation of inflammatory pathways in astrocytes and microglia promote demyelination, while their inhibition has the potential to support remyelination in rodents (Jha et al., 2010; Raasch et al., 2011; Goldmann et al., 2013; Blank and Prinz, 2014).

Impairments in microglial phagocytosis of cellular debris is common in the aged mouse brain (≥ 20 -months-old) as well, perhaps contributing to the increase in inflammatory factors seen during aging (Ritzel et al., 2015; Safaiyan et al., 2016; Cantuti-Castelvetri et al., 2018; Marschallinger et al., 2020). Impaired phagocytosis could be partly attributed to the dysfunction of an overwhelmed clearance system (Safaiyan et al., 2016; Thériault and Rivest, 2016; Marschallinger et al., 2020). For instance, continuous increases in myelin debris cannot be accommodated by microglia in the long-term, which subsequently leads to dysfunctional lysosomal activity and a noticeable increase in insoluble lipofuscin-like granules—a marker of aging, dystrophy and possibly senescence—in mice (Safaiyan et al., 2016).

A specific population of microglia in the white matter of the aged brain (≥ 18 -months-old), white matter associated microglia (WAM), has an increased expression of triggering receptor expressed on myeloid cells 2 (TREM2)—a receptor important for phagocytosis, lipid metabolism and proper myelination (Poliani et al., 2015; Safaiyan et al., 2021). In aged humans (50–80-years-old), microglia in the white matter have increased expression of genes associated with lipid-metabolism as well [e.g., secreted phosphoprotein 1 (SPP1) and apolipoprotein E (APOE)], hinting at the possibility of the presence of WAMs in humans; although TREM2 did not significantly associate with these white matter microglial clusters (Sankowski et al., 2019; Safaiyan et al., 2021). Their function is likely similar to the

hypothesized function of WAMs in mice, which is thought to include myelin debris clearance and lipid metabolism (Safaiyan et al., 2021).

Interestingly, microglia isolated from whole brains of aged mice (≥ 21 -months-old) show significantly downregulated expression levels of TREM2 compared to younger mice (Hickman et al., 2013; Thomas et al., 2022). This could be due to a net decrease in microglial expression of TREM2, despite the increases seen in WAM. However, it has also been hypothesized that with time WAMs may become overwhelmed and enter into a senescent state, thereby reducing function and contributing to white matter degeneration and cognitive decline (Ahn et al., 2022). This senescent state—perhaps more common in WAM from rodents >18 -months of age—may have reduced expression of TREM2, contributing to the observed decline in expression with age.

Senescent microglia may have a reduced capacity to modulate neuronal activity because of genetic changes that result in reduced surveillance, migration, and sensitivity to endogenous ligands, and a heightened sensitivity to pathogens, as seen in 24-month-old mice (Hickman et al., 2013; Madry et al., 2018; Angelova and Brown, 2019). For example, due to the downregulation of genes for purinergic receptors (e.g., *P2ry12*) and potassium leak channels (e.g., *Thik-1*) in aged mice, the ability for microglia to sense hyperactive neurons and migrate toward them in order to facilitate rapid repolarization would be reduced, impairing their beneficial modulation (Hickman et al., 2013; Kato et al., 2016; Madry et al., 2018). This may be a contributing factor for neuronal hyperactivity and excitotoxicity found in the aged brain, which negatively affects memory (Bishop et al., 2010; Stargardt et al., 2015; Li et al., 2020).

The age-mediated alterations in microglial function that are evident in the aged brain are thought to arise from at least two factors. (1) Microglia become overwhelmed and cannot keep up with demand, and/or (2) shift into a less sensitive state trying to keep inflammatory signals to a minimum by reducing their reactivity to endogenous ligands (Hickman et al., 2013). Although some microglia may enter into a senescent state, many of these cells still have the ability to aid the aged brain and maintain/improve cognition. An interesting avenue to achieve this may be through the use of cannabis and the endocannabinoid system. Indeed, all glial cell types have receptors for cannabinoids, which have wide ranging effects on cellular function, indicating that cannabis may be beneficial for glial regulation of myelination (Figure 2B; Stella, 2010; Navarrete et al., 2014; Ilyasov et al., 2018; Martinez Ramirez et al., 2023).

1.5. The endocannabinoid system

The endocannabinoid system extends to most regions of the body. It encompasses naturally occurring endogenous (endo)cannabinoids, the enzymes needed for their formation and degradation, and cannabinoid receptors (Lu and Mackie, 2016). Although many receptors take part in endocannabinoid signaling [e.g., peroxisome proliferator-activated receptors (PPARs), transient receptor potential cation channels (TRPs)], the two main cannabinoid receptors are cannabinoid receptor type 1 (CB₁R) and cannabinoid receptor type 2 (CB₂R; Howlett, 2002).

CB₁Rs are widespread in the CNS and are the most prevalent G-protein coupled receptor in the mammalian brain (Herkenham et al., 1990; Marsicano and Lutz, 1999). In the hippocampus and

cortex, CB₁Rs have an especially high localization on inhibitory neurons, although their distribution and localization patterns differ throughout the human and rodent CNS (Glass et al., 1997; Marsicano and Lutz, 1999; Tsou et al., 1999; Fletcher-Jones et al., 2020). Interestingly, a significant number of CB₁Rs are not expressed on the cell surface, but instead localize to lysosomes and late endosomes, as shown in cell lines and primary cell cultures—possibly contributing to lysosomal integrity and function (Rozenfeld and Devi, 2008; Bilkei-Gorzo, 2012; Fletcher-Jones et al., 2020). Additionally, CB₁Rs also localize to mitochondria and can influence metabolism in neurons and glial cells, as observed in primary cell cultures and in mice (Bénard et al., 2012; Jimenez-Blasco et al., 2020).

However, the majority of neuronal CB₁Rs are found on pre-synaptic terminals, where their primary function is to suppress the release of neurotransmitters, altering the activation of post-synaptic channels and, therefore, modulating synaptic activity and plasticity (Mackie and Hille, 1992; Di Marzo et al., 2015; Zou and Kumar, 2018). The cannabinoid-mediated reduction in neurotransmitter release is achieved by the reduced influx of presynaptic Ca²⁺ due to the inhibition of voltage gated Ca²⁺ channels and of adenylyl cyclase, which downregulates cyclic adenosine monophosphate and protein kinase A, two cellular constituents involved in increasing the influx of Ca²⁺ (Castillo et al., 2012). CB₁R activation in pre-synaptic terminals is mainly facilitated through retrograde signaling of endocannabinoids released from the post-synapse (Kano et al., 2009; Castillo et al., 2012; Njoo et al., 2015).

CB₂Rs have much lower levels of expression in the CNS of humans and rodents (Lu and Mackie, 2016; Jordan and Xi, 2019). Interestingly, CB₂R mRNA expression can vastly increase during an inflammatory insult, with microglia from mice displaying as much as a 10-fold increase (Maresz et al., 2005). This finding indicated that the CB₂R likely plays a substantial role in CNS immune function, which has been subsequently supported in the literature (Turcotte et al., 2016; Komorowska-Müller and Schmöle, 2020). CBRs may attenuate pro-inflammatory cytokine secretion by interfering with the phosphorylation of mitogen activated protein kinases (MAPK), such as extracellular signal-regulated kinase (ERK), which is known to participate in pro-inflammatory pathways, as shown in microglial cell line cultures (Eljaschewitsch et al., 2006; Young and Denovan-Wright, 2022a,b). Furthermore, activation of CB₂Rs promotes IL-10 (an anti-inflammatory cytokine) secretion from primary microglia cells from mice by reducing the translocation of the transcription factor nuclear factor- κ B (NF- κ B) to the nucleus *via* reduced phosphorylation of I κ B Kinase- α (IKK α)—a subunit of the IKK complex that is essential for NF- κ B signaling—which subsequently prevents NF- κ B formation (Solt and May, 2008; Correa et al., 2010). Of note, NF- κ B-mediated inflammation is often *via* the inflammasome and is associated with many white matter associated diseases and is upregulated in the aged brain (Tilstra et al., 2011; Youm et al., 2013; Blank and Prinz, 2014; Rea et al., 2018). Evidence also suggests that CB₂Rs are present on post-synaptic terminals of neurons in rodents and non-human primates, although expression levels are relatively low and may depend on brain region (Brusco et al., 2008; Lanciego et al., 2011; Li and Kim, 2015; Stempel et al., 2016).

Importantly, astrocytes, oligodendrocytes, and OPCs possess CB₁Rs and CB₂Rs, highlighting the wide range of functions performed by this system (Navarrete et al., 2014; Ilyasov et al., 2018; Martinez Ramirez et al., 2023). The impact these receptors have with respect to

their activation by cannabinoids on individual cell-types will be discussed in more detail below (Figure 2B).

1.5.1. The endocannabinoid system in aging

The endocannabinoid system undergoes an unequivocal transition during aging across species (Bilkei-Gorzo, 2012; Bishay et al., 2013; Pascual et al., 2013; Di Marzo et al., 2015; Piyanova et al., 2015). The direction of this change relies on the region investigated and, therefore, its impact on brain function differs based on the affected region and the context in which it is examined.

A general trend in the literature suggests that CB₁R density decreases throughout the brain with aging, however, receptor function seems to differentially change depending on the region and cell type (Mato and Pazos, 2004; Bilkei-Gorzo, 2012; Di Marzo et al., 2015; Ginsburg and Hensler, 2022). Interestingly, G_{i/o}-coupled protein receptors decline in the aged brain as a whole (de Oliveira et al., 2019).

The CB₂R is less well characterized due to methodological difficulties and, therefore, the change in CB₂R expression with age is less well-known (Zhang et al., 2019). One study did not find any reductions in CB₂R density in any region analyzed from aged mice (22-months-old), while an earlier study found a significantly declined receptor density in synaptosomes, but not in overall membrane fractions from aged rats (24–28-months-old; Pascual et al., 2014; Hodges et al., 2020). This discrepancy could be a result of different rodent species and/or differential CB₂R expression based on cell type. It could be speculated that neuronal synaptic expression of CB₂Rs decline with age, whereas glial expression remains constant or potentially increases, which would warrant further investigation.

Sex differences are a common feature of the endocannabinoid system, although this depends on the type of measurement and regions analyzed (Laurikainen et al., 2019; Van Ryzin et al., 2019; De Meij et al., 2021; Levine et al., 2021; Vecchiarelli et al., 2022). For example, in the human brain, females exhibited increased binding of the CB₁R with age, whereas males did not show any change (Van Laere et al., 2008). Similarly, adult female CB₂R-knockout (KO) mice displayed larger alterations in synaptic markers compared to male mice, although both sexes exhibited deficits in social memory (Komorowska-Müller et al., 2021b).

Aging impacts the endocannabinoid system on multiple levels, and myelination relies on support from numerous cell types and is moderately guided by neuronal activity, two processes which are partly controlled by the endocannabinoid system. Therefore, any modification to endocannabinoid signaling will likely have an impact on myelination, one of the most important structural and functional aspects of the CNS.

1.6. Cannabis and the endocannabinoid system

Cannabinoids exert their influence over the endocannabinoid system mainly through CB₁Rs and CB₂Rs, which contribute to the sought after medicinal and recreational qualities of cannabis. The most common psychoactive cannabinoid, Δ-9-tetrahydrocannabinol (THC), has a relatively high affinity for the two cannabinoid

receptors (Pertwee, 2008). THC is generally considered to be a partial agonist for both CBRs, although its inhibitory effect on synapses can be comparable to that of a full agonist (Laaris et al., 2010). By contrast, cannabidiol (CBD)—the most common non-psychoactive cannabinoid—does not have a particularly high affinity for either CBR, but was shown to antagonize CBR agonists (Thomas et al., 2007). CBD is a negative allosteric modulator of CB₁Rs, and is suggested to act as an inverse agonist of CB₂Rs (Thomas et al., 2007; Laprairie et al., 2015). The anti-inflammatory effects attributed to CBD may be exerted through this inverse agonism of CB₂Rs (Thomas et al., 2007; Pertwee, 2008; Yu et al., 2020). However, it is important to note that both THC and CBD have many CBR-independent or indirect signaling mechanisms that also contribute to their overall outcomes (Pertwee, 2008; Stella, 2010). Interestingly, one indirect mechanism is the ability for CBD to inhibit fatty acid amide hydrolase (FAAH)—the enzyme required for the degradation of *N*-arachidonylethanolamine (anandamide; AEA), an endocannabinoid—which results in an increase in AEA (Watanabe et al., 1996; Bisogno et al., 2001; De Petrocellis et al., 2011). Of note, there is a large body of research looking at the potential benefits of inhibiting endocannabinoid metabolizing enzymes; with studies showing that increases in AEA have an immunomodulatory effect, and can potentially aid in MS, as shown in mice (Rossi et al., 2010; Vázquez et al., 2015a; Vecchiarelli et al., 2021).

The changes observed in the endocannabinoid system in the aging brain are mostly similar to those observed after chronic THC exposure (Yoo et al., 2020). The most noticeable effect observed after chronic THC exposure is the significant but reversible reduction in CB₁Rs, with cortical regions showing more extensive decreases in expression (Hirvonen et al., 2012; D'Souza et al., 2016; Augustin and Lovinger, 2022). However, it should be noted that these studies used exclusively male participants. Studies including females are lacking, which is a significant gap since sex differences with respect to the endocannabinoid system are well-described (Laurikainen et al., 2019; Levine et al., 2021).

Importantly, THC does not interact with the brain in equal measure, as found by Leishman et al. (2018). In this study, acute administration of THC [3 mg/kg; intraperitoneal injection (i.p)] differentially impacted the lipidome and transcriptome depending on the brain region and age of the subject, 2 h after administration. THC is distributed and metabolized in a region-specific manner, with highest levels in the hippocampus. Another interesting finding of this study was that adult mice (~4-months-old) displayed the largest changes after acute THC exposure compared to exposed ~1 and ~2-month-old mice, with a general downregulation of the endocannabinoid system. The effect of THC in the aged brain was not examined in this study, but the observed changes would likely be different than other time points.

Although THC and CBD are the main cannabinoids found in cannabis, it is important to note that 100s of different cannabinoids and other biologically active compounds exist in the plant, such as terpenes, including β-caryophyllene (Kopustinskiene et al., 2022). These compounds likely work synergistically to produce the effects of cannabis through multiple different signaling pathways, creating what is called the “entourage effect” (Ferber et al., 2020; Finlay et al., 2020). However, due to a lack of literature, the next sections will focus on THC, CBD, and some synthetic cannabinoids.

1.7. The impact of cannabis on microglia

One of the most essential microglial functions that promotes myelination is the clearance of myelin debris (Kotter, 2006; Lampron et al., 2015). The CB₂R has been shown to be important for phagocytosis in microglia. For instance, in CB₂R-KO primary microglia from mice, phagocytosis was significantly reduced in both steady-state conditions and following an inflammatory stimulus (i.e., TGF- β) compared to controls (Mecha et al., 2015). Similarly, activation of the CB₂R increased phagocytosis in primary cell cultures and improved the removal of amyloid- β_{40} peptides in a mouse model of Alzheimer's disease pathology (Ehrhart et al., 2005; Aso et al., 2016). Additionally, microglia from CB₂R-KO mice (18-months-old) had an age-dependent increase in lipofuscin granules compared to the control group, signifying reduced lysosomal degradation (Komorowska-Müller et al., 2021a). These findings indicate a significant role played by CB₂Rs in the proper removal and degradation of debris, and in the microglial response to environmental stimuli. However, other receptors likely also contribute, as CBD has been shown to promote phagocytosis through TRPs in primary microglial cell cultures from mice (Hassan et al., 2014; Yang et al., 2022). Briefly, TRPs are Ca²⁺-permeable channels known for their role in temperature sensation, but interestingly, have also been shown to play a role in the release of pro-inflammatory cytokines from microglia in mice exposed to immunogenic agents [i.e., lipopolysaccharide (LPS)] (Zhang Y. et al., 2021). The observed increase in phagocytosis due to CBD is thought to be related to the TRP-mediated increase in the influx of Ca²⁺ (Hassan et al., 2014).

Increased microglial phagocytosis of myelin debris following the application of 2-AG subcutaneously *via* an osmotic pump subsequently promoted remyelination in an adult mouse model of experimental demyelination (Mecha et al., 2019). This study also observed an altered secretory profile, with increases in IL-1 β , TNF- α , and IL-10 in the brain after 2-AG application. This is an example of an augmented immune response, where the benefits of pro-inflammatory cytokines in tandem with anti-inflammatory cytokines work synergistically to promote remyelination. However, microglia in the aged brain are responding to chronically elevated levels of pro-inflammatory factors resulting in reduced functional capacity.

Activation of the CB₂R by AEA in primary microglial cells from mice has been shown to reduce NF- κ B signaling and increase expression of IL-10 (Correa et al., 2010). The anti-inflammatory effect produced by IL-10 is partly through a negative feedback loop with astrocytes, where the binding of IL-10 causes the release of TGF- β from astrocytes, which subsequently attenuates pro-inflammatory cytokine production in primary microglia cells (Norden et al., 2014). However, astrocytes have reduced expression of IL-10 receptor-1 in aged mice (\geq 18-months-old), and fail to effectively diminish microglia-mediated inflammation (Norden et al., 2016; O'Neil et al., 2022). Therefore, compounds that can act directly on microglia to reduce pro-inflammatory cytokine production are of particular interest.

Selective CB₂R agonists (i.e., JWH-133; i.p.) reduce the release of pro-inflammatory cytokines from microglia in a mouse model of Alzheimer's disease pathology (Aso et al., 2013). Similarly, activation of CB₁Rs and CB₂Rs by synthetic cannabinoids [i.e., arachidonyl-2'-chloroethylamide (ACEA) and HU-308, respectively] reduced nitric oxide, TNF- α , IL-1 β and IL-6 release from spontaneous immortalized

microglia (SIM)-A9 cells in culture (Young and Denovan-Wright, 2022a).

CBD also has the ability to beneficially regulate the oxidative status in microglia by acting as an antioxidant, where it may directly scavenge ROS and/or inhibit the phosphorylation of upstream kinases needed for NF- κ B signaling in primary microglia cells cultured from mice (van den Berg et al., 2001; dos-Santos-Pereira et al., 2020; Atalay Ekiner et al., 2022). This results in reduced levels of IL-1 β and TNF- α independently of CB₁R, CB₂R or PPAR γ , as tested using receptor antagonists. These effects could also be partly regulated by increased levels of endocannabinoids or through TRP channels, since CBD has a relatively high affinity for TRP vanilloid receptor 1 (TRPV1), which has been shown to modulate cytokine production in microglia (Stampanoni Bassi et al., 2019). Furthermore, in a mouse (1-month-old) model of viral-induced demyelination, CBD (5 mg/kg; i.p.) attenuated morphological alterations in microglia, and reduced production of IL-1 β , chemokines, and vascular cell adhesion molecule-1 (VCAM-1) through adenosine A_{2A} receptors (Mecha et al., 2013). Interestingly, VCAM-1—a protein expressed by endothelial cells in the BBB that is involved in peripheral immune cell recruitment—expression increases in the aged brain, and the application of anti-VCAM-1 antibodies reduces microglial reactivity and improves memory and learning in aged (19-month-old) mice (Yousef et al., 2019). Overall, it is clear that CBD acts through a number of vastly different pathways that have overlapping effects on the brain.

Although the anti-inflammatory effects of the CB₂R are well-characterized, it is also important to note that the CB₂R seems essential for many environmental-induced immune responses in microglia. Primary microglia cell and organotypic hippocampal slice cultures generated from CB₂R-KO mice showed attenuation of the microglial immune response to toll-like receptor (TLR) ligands (e.g., TLR4/3/9), preventing the pro-inflammatory cascade usually associated with TLR ligands (Reusch et al., 2022). Therefore, although the CB₂R can function to suppress inflammation, it also contributes to its initiation. These studies highlight how complex the interaction between the endocannabinoid system and microglia is.

The CB₁R is also important for the inflammatory response, as a recent study found that inflammation was dependent on microglial CB₁Rs (De Meij et al., 2021). They found decreased pro-inflammatory cytokines in mice (2–5-months-old) with CB₁R-KO microglia exposed to LPS. However, it also increased sickness behavior in male, but not female mice (De Meij et al., 2021). Similarly, ablation of the CB₁R resulted in early age-related cognitive deficits in mice (Bilkei-Gorzo et al., 2005; Albayram et al., 2011). Therefore, the activation of CB₁Rs may be beneficial for preventing sickness behavior and age-related cognitive decline. This is true with respect to THC, but only with certain doses (Sarne, 2019). Ultra-low (0.002 mg/kg; i.p.) and low (3 mg/kg; i.p.) doses of THC resulted in improved cognitive function in old mice (24 and 18-month-old, respectively), whereas the same dose induced cognitive impairments in adult mice (2-months-old; Bilkei-Gorzo et al., 2017; Sarne et al., 2018). However, higher doses of THC have the reverse effect (Calabrese and Rubio-Casillas, 2018). This dose-dependent alteration was also observed in microglia in the 2-month-old mouse brain, where higher doses (20 mg/kg; i.p.) of THC resulted in the increased release of pro-inflammatory cytokines compared to lower doses (Cutando et al., 2013). In line with this, a recent study found that adolescent mice exposed to daily low-doses of

THC (5 mg/kg; i.p) resulted in the downregulation of genes responsible for the microglial response to an immune insult (e.g., IL-1 β , IL-6), which carried over into young adulthood (~1.5-months-old) but not maturity (~4-months-old; Lee et al., 2022). This blunting of the microglial response attenuated reactivity to LPS and, furthermore, altered behavior in mice, where exposed mice showed an inability to react appropriately to psychosocial stress (Lee et al., 2022). These impacts were mediated through the CB₁R, as tested using receptor antagonists. These studies highlight the differential impact cannabis can have on the developing brain and the aged brain, where low doses may be detrimental to young animals, yet beneficial to aged animals.

The reduced production of pro-inflammatory cytokines with THC administration may occur through the inhibition of NF- κ B signaling in microglia and astrocytes, as observed in cell culture experiments (Kozela et al., 2010; Rizzo et al., 2019). Interestingly, the overexpression of CB₁R in adult mice undergoing experimental demyelination resulted in delayed onset and reduced severity of symptoms, whereas CB₁R antagonism quickened symptom onset and increased the expression of inflammatory cytokines and NF- κ B proteins (Lou et al., 2016, 2018). Furthermore, an alteration in cell state and expression profile (i.e., increase in pro-inflammatory cytokines and nitric oxide) was observed in cultured BV-2 cells treated with a CB₁R antagonist (Lou et al., 2018).

It is clear that cannabinoids are involved in the microglial response to inflammation with respect to secretory profile and phagocytosis. This is the major way in which microglia may modulate myelination. However, it is important to note that there is still a paucity in *in vivo* experiments conducted with aged mice, emphasizing the need for increased research (Scipioni et al., 2022).

1.8. The impact of cannabis on astrocytes

The crosstalk that exists between microglia and astrocytes is essential for proper function and plasticity of the brain and maintenance of homeostasis (Jha et al., 2019; Matejuk and Ransohoff, 2020). However, in aging, certain aspects of this communication network become exacerbated. As discussed, the aged brain environment alters microglial state including their release of soluble factors. The concomitant impact aging has on astrocytes also induces a phenotypic switch that results in altered gene expression, perpetuation of inflammation, and recruitment of peripheral immune cells, which further exacerbates inflammation (Palmer and Ousman, 2018; Jha et al., 2019).

The ability for THC to inhibit NF- κ B and reduce the release of IL-1 β and TNF- α from microglia also extends to monocytes/macrophages and lymphocytes in human and rodent cell lines and primary cell cultures (Shivers et al., 1994; Kozela et al., 2010; Rizzo et al., 2019; Henriquez et al., 2020). This inhibition then translates into an observed reduction in the astrocytic release of IL-6 and monocyte chemoattractant protein-1 (MCP-1) in primary human cell cultures, thereby reducing inflammatory signaling and peripheral immune cell recruitment, respectively (Rizzo et al., 2019; Henriquez et al., 2020). Although IL-6 can be beneficial for many aspects of development including myelination, the chronically increased levels that are seen in aged humans and rodents are damaging and pro-inflammatory in nature (Ferrucci et al., 1999;

Godbout and Johnson, 2004; Kimura and Kishimoto, 2010; Ritzel et al., 2016; Porcher et al., 2021).

This protective effect offered by THC is thought to be facilitated by the activation of CB₂Rs, which is elevated in microglia and astrocytes within an inflammatory environment (Benito et al., 2008; Di Marzo et al., 2015; Cassano et al., 2017). However, other receptors may also play a role. Indeed, a recent study observed a marked inhibition of pro-inflammatory cytokines produced by IL-1 β -stimulated primary cell cultured human astrocytes when pre-/co-treated with WIN55,212-2—a synthetic cannabinoid that displays similar effects to THC (Compton et al., 1992; Fields et al., 2022). This effect was independent of the CB₁R and PPARs (Fields et al., 2022). Conversely, through PPAR γ , CBD reduced pro-inflammatory cytokine release, inhibited NF- κ B, and reduced GFAP expression in primary astrocyte cells stimulated with amyloid- β ₁₋₄₂ peptides, while also promoting neurogenesis in adult rats (Esposito et al., 2011). Furthermore, as discussed, CBD has the ability to diminish microglial cytokine production by scavenging ROS, diminishing NF- κ B activity, reducing VCAM-1 levels and increasing the availability of endocannabinoid ligands. This may also lessen the extent to which astrocytes participate in peripheral immune cell recruitment and inflammation (Mecha et al., 2013). Indeed, a recent study found an association between CBD administration, reduced phosphorylation of NF- κ B and reduced release of IL-6 from mouse primary cultured astrocytes stimulated with LPS (Wu et al., 2021).

However, cannabis includes both THC and CBD. Administration of Sativex[®]—an approved oromucosal spray (Health Canada and various European health agencies) containing THC (5 mg/kg) and CBD (5 mg/kg) for the treatment of symptoms associated with MS—reduced astrocyte reactivity and decreased the expression of pro-inflammatory cytokines released by microglia in a mouse model of MS (Feliú et al., 2015). Sativex[®] also preserved myelin morphology in mice exposed to virus-induced demyelination.

Inflammation can also be induced by disrupting the proper communication between astrocytes and neurons. The deletion of CB₁Rs from GABAergic neurons enhanced a phenotypic switch in astrocytes already associated with aging, including increased GFAP expression and amplified pro-inflammatory cytokine secretion in mice (Bilkei-Gorzo et al., 2018). Therefore, disruption of endocannabinoid signaling between neurons and astrocytes—which naturally occurs during aging—causes a deleterious transition in astrocytes that perpetuates cytokine-mediated damage. The application of cannabinoids may help since they directly act on astrocytes to diminish cytokine release (Sheng et al., 2005; Aguirre-Rueda et al., 2015; Rizzo et al., 2019; Fields et al., 2022).

Chronic inflammation was also identified as a major influencer of astrocytic gap junctions and hemichannels in rodents and humans (Bronzuoli et al., 2019; Peng et al., 2022). Typically, hemichannels remain mostly closed under “normal” conditions, but can be opened during pathological conditions; whereas the reverse is true for gap junctions (Peng et al., 2022). IL-1 β and TNF- α released from microglia can open astrocytic hemichannels and reduce coupling between astrocytes in primary cell/slice cultures and *in vivo* in mice (Même et al., 2006; Retamal et al., 2007; Abudara et al., 2015; Vázquez et al., 2015b). The application of synthetic cannabinoids (e.g., WIN55,212-2) were able to reduce the microglial release of these factors and directly act on primary astrocyte cells from mice to reduce hemichannel activation, preventing astrocytic uncoupling and maintaining gap

junctions (Froger et al., 2009; Gajardo-Gómez et al., 2017). Furthermore, direct activation of astrocytic CB₁Rs was found to be required for the observed decrease in hemichannel activation, as determined by CB₁R antagonism (Froger et al., 2009; Gajardo-Gómez et al., 2017). On the contrary, AEA (1 μM; topical application through a cortical cranial window) was shown to increase hemichannel activity in adult mice *in vivo*, resulting in a release of ATP that caused microglial process extension and migration toward the injury site (Vázquez et al., 2015b). Similarly, THC (5 mg/kg; i.p) has been shown to result in an increase in glutamate in the extracellular space by binding to astrocytic CB₁Rs in mice, contributing to long term depression at synapses and impairments in working memory (Navarrete and Araque, 2010; Han et al., 2012).

During acute inflammatory conditions, this increase in hemichannel activity may be beneficial for mounting an immune response to an insult, however, the prolonged release of many factors (i.e., glutamate) from astrocytes in chronic conditions may be associated with altered synaptic plasticity and memory impairments (Navarrete and Araque, 2010; Han et al., 2012; Vázquez et al., 2015b; Labra et al., 2018). Alternatively, if cannabinoids are able to reduce the opening of hemichannels during a chronic inflammatory event, it may subsequently prevent excitotoxicity caused by excess glutamate and reduce the release of pro-inflammatory cytokines, thus maintaining neuronal and astrocyte viability (Froger et al., 2009; Gajardo-Gómez et al., 2017). Indeed, a recent review highlights a potential signaling cascade involving NF-κB, p38 and nitric oxide in which cannabinoids prevent the release of glutamate from hemichannels (Labra et al., 2018). It can also be hypothesized that the conserved function of gap junctions would facilitate proper communication between astrocytes and oligodendrocytes, promoting myelination (Papaneophytou et al., 2019).

In summary, elevated levels of inflammatory factors partly initiated by microglia and other recruited immune cells causes a phenotypic switch in astrocytes that contributes to functionally perpetuating inflammation *via* the release of pro-inflammatory cytokines and recruitment of peripheral immune cells. Cannabinoids can inhibit the release of these compounds (i.e., TNF-α, IL-6, MCP-1) by reducing the activity of pro-inflammatory pathways (e.g., NF-κB) in immune cells and astrocytes and increasing the availability of endocannabinoids. The subsequent decrease in pro-inflammatory factors and direct action of (endo)cannabinoids may also modulate hemichannel activity, thereby contributing to changes in extracellular homeostasis, synaptic activity and plasticity, as well as glial support functions. Although these results from studies using younger animals can provide information on how cannabinoids impact astrocytes and what this could mean for the process of myelination, the unique environment present in the aged brain makes extrapolation conjectural.

1.9. The impact of cannabis on oligodendrocytes and OPCs

As discussed, oligodendrocytes and OPCs are both essential for proper myelination and maintenance of myelin. Chronic increases in ROS, pro-inflammatory cytokines and other damaging compounds released from immune cells and astrocytes have the ability to damage mature oligodendrocytes and OPCs, resulting in myelination impairments (Pang et al., 2003; Jurewicz et al., 2005; French et al., 2009; Peferoen et al., 2014; Guttenplan et al., 2021; Sams, 2021). The ability for cannabinoids to reduce the release of pro-inflammatory

cytokines from immune cells and astrocytes could therefore aid in the preservation of oligodendrocytes and OPCs. Furthermore, the antioxidant capacities offered by CBD could also protect these cells from oxidative stress.

However, in primary oligodendrocyte cell cultures from 12-day-old rats, CBD (100 nM–1 μM) resulted in mitochondrial dysfunction that led to increases in intracellular cytotoxic Ca²⁺ and ROS, which negatively impacted oligodendrocyte viability (Mato et al., 2010). Conversely, Mecha et al. (2012) found that CBD (1 μM) administered to inflammatory-induced primary oligodendrocyte cell cultures from the cortex of 2-day-old rats protected OPCs from oxidative stress and apoptosis. These studies found that these effects were independent of CB₁R, CB₂R, or PPARγ. The apparent discrepancy between studies is likely due to dosage and differences in age, region, and maturation of oligodendrocyte cells (Molina-Holgado et al., 2022).

Interestingly, CBD was also found to influence genes related to glycolysis—the major energy source for mature oligodendrocytes—and carbohydrate metabolism in oligodendrocytes (Rao et al., 2017; de Almeida et al., 2022). The data presented by de Almeida et al. (2022) suggests a slight downregulation of glycolysis in OPC and mature oligodendrocyte cell cultures (MO3.13). This finding is important because glycolysis produces lactate, an important energy metabolite transferred to neurons from oligodendrocytes, which has been shown to be essential for proper neuronal function in aged mice (12–24-months-old; Fünfschilling et al., 2012; Lee et al., 2012; Philips et al., 2021). Interestingly, the ablation of monocarboxylate transporter 1 (MCT1)—a lactate transporter—from OPCs resulted in hypomyelination and axonal degeneration in mature and older (18–24-months-old) mice, highlighting the understudied role played by OPCs in myelination among the adult brain (Philips et al., 2021). MCT1 expression naturally declines in the aging mouse brain, especially after 15-months of age (Ding et al., 2013; Philips et al., 2021). Therefore, further reductions in glycolysis due to CBD administration may prevent the beneficial support to neurons offered by oligodendrocytes and OPCs. Similarly, activation of mitochondrial CB₁Rs with THC resulted in reduced glycolytic activity and lactate production in primary cultured astrocytes (0–1-days-old), which was hypothesized to contribute to the subsequent impaired neuronal function and altered behavior (e.g., social interaction deficit) in young adult (8–12-weeks-old) mice (Jimenez-Blasco et al., 2020). This reduction in astrocytic lactate would also likely have an impact on the oligodendrocyte lineage cells. For example, global inhibition of lactate production prevented remyelination in the corpus callosum after experimental demyelination in mice (~3-months-old), while OPC-rich primary cell cultures displayed heightened differentiation when lactate was added to the glucose medium (Ichihara et al., 2017). Perhaps this increase in OPC differentiation due to the presence of lactate can aid in remyelination by replacing the oligodendrocyte pool, which would warrant further investigation.

Although the literature is split with respect to OPC number in the aged brain, their ability to differentiate is severely impaired in humans and rodents (Sim et al., 2002; Yeung et al., 2014; Segel et al., 2019; Luan et al., 2021; Rivera et al., 2021; Zhang X. et al., 2021). Therefore, the encouragement of differentiation and protection of OPC viability offered by certain cannabinoids is of particular relevance (Ilyasov et al., 2018; Molina-Holgado et al., 2022). 2-AG was shown to promote the differentiation of OPCs into mature myelinating oligodendrocytes in primary mixed glial cell cultures from rats through the activation of CB₁Rs and CB₂Rs (Gomez et al., 2010). OPCs also have the ability

to produce and release 2-AG themselves, indicating a potential autocrine and paracrine signaling mechanism that can further promote differentiation (Gomez et al., 2010, 2011). Pathways of note that have been shown to promote OPC differentiation through cannabinoid receptors are the phosphatidylinositol 3-kinase/Akt (PI3K/Akt) and mammalian target of rapamycin (mTOR), ERK/MAPK, and Rat sarcoma (Ras) homolog family member A/Rho-associated protein kinase (RhoA/ROCK; Molina-Holgado et al., 2002; Narayanan et al., 2009; Tyler et al., 2009; Gomez et al., 2010, 2011; Giacompo et al., 2017; Sánchez-de la Torre et al., 2022; Wang et al., 2022).

These findings have prompted further research investigating the effect of THC on OPC differentiation. A recent study found that the application of THC (3 mg/kg; i.p) to young mice (6-days-old) and organotypic cerebellar cultures promoted the differentiation of OPCs (Hueriga-Gómez et al., 2021). Furthermore, the same group found that THC (3 mg/kg; i.p) induced OPC differentiation and remyelination after experimentally-induced demyelination in the corpus callosum of adult (6–8-weeks-old) mice (Aguado et al., 2021). The modulation of OPC differentiation by THC is thought to be mediated by both CBRs, but mainly CB₁Rs, since antagonism of CB₁Rs prevented the beneficial effects observed. Furthermore, OPC(Ng2/Ai6)-CB₁R-KO mice displayed impaired OPC differentiation and myelination in the corpus callosum throughout the examined lifespan (≤2-months-old; Sánchez-de la Torre et al., 2022). The impaired differentiation of OPCs lacking CB₁Rs is thought to be partly due to an increase in RhoA/ROCK signaling. For example, THC (3 mg/kg, i.p) administered to WT mice resulted in reduced RhoA/ROCK proteins compared to vehicle exposed mice, leading to increased myelin-related proteins and enhanced OPC differentiation (Sánchez-de la Torre et al., 2022). However, OPC-CB₁R-KO mice had increased levels of RhoA/ROCK proteins observed with a lower density of mature oligodendrocytes and reduced immunofluorescence against myelin-related proteins, which did not significantly change with THC administration (Sánchez-de la Torre et al., 2022). These findings highlight the importance of the CB₁R in OPC differentiation, and its relationship with the RhoA/ROCK pathway. Furthermore, these findings are supported by previous studies that have shown the importance of RhoA/ROCK signaling in OPC differentiation (Baer et al., 2009; Pedraza et al., 2014). Similarly, the application of WIN55,212-2 (i.p) resulted in improved remyelination after experimentally-induced demyelination in mice (6–7-weeks-old) when administered at a dose of 0.5 mg/kg; whereas a dose of 1 mg/kg impaired remyelination (Tomas-Roig et al., 2020). The negative effects produced by 1 mg/kg of WIN55,212-2 are hypothesized to be due to stronger reductions in Ca²⁺ influx, resulting in reduced neuronal activity, which possibly hinders activity-dependent myelination (Tomas-Roig et al., 2020). Overall, it is evident that at the right dose cannabinoids can alter myelination and promote the maturation of oligodendrocyte lineage cells.

In summary, cannabinoids at the right doses have the ability to aid in remyelination and promote OPC differentiation after pathological insults. As outlined throughout this review, cannabinoids act on a plethora of different cell types and different targets within those cells, which directly and indirectly impact myelination. Although cannabinoid-induced alterations to microglia and astrocytes can influence OPC differentiation and oligodendrocyte function indirectly, it is also important to keep in mind that cannabinoids

directly act on OPCs to influence maturation and function. However, there are a lack of studies regarding the healthy aged brain. Although the aged brain does have increases in ROS, pro-inflammatory cytokines, and a general reduction in glial cell function, the extent of these changes and the dynamics at play are inherently different than those observed after the application of LPS or experimentally-induced demyelination at younger ages. Furthermore, findings from cell line and primary cell cultures need to be validated in *in vivo* experiments. Therefore, although studies outlined here point toward a role played by cannabinoids in myelination, more studies specifically looking at their impact in the healthy aged brain are required to confirm if these findings do indeed translate. Nonetheless, the evidence provided thus far suggests that cannabinoids may help promote myelination in the aged brain. Neuroimaging studies offer a different approach to visualize how cannabinoids impact brain communication and myelination, which allows elucidating large-scale changes that may result from the impact they have on glial cells and neuronal function.

1.10. The impact of cannabis on the human brain

Neuroimaging studies are valuable non-invasive techniques used to visualize large-scale changes in activity, connectivity, and structural alterations in the brain. DTI is a magnetic resonance imaging (MRI) technique that differentiates the degree and direction of water diffusion within an allotted space (Pierpaoli and Basser, 1996). Three main parameters are given by DTI for white matter: fractional anisotropy (FA), mean diffusivity (MD) and radial diffusivity (RD; Alexander et al., 2007). Generally, as described by Becker et al. (2015), increases in FA and decreases in MD and RD reflect increases in myelination.

The DTI literature encompassing cannabis use and myelin integrity is not consistent. Difficulties with respect to accurate reporting on usage and concentration of cannabis, and the ratio of cannabinoids—not to mention other confounding factors such as lifestyle and metabolism—make human cannabis studies increasingly difficult to perform. Furthermore, the DTI metrics that are used as proxies of myelin integrity do not specifically identify myelin abnormalities, but detect differences in water diffusion—a parameter that can be altered by axon packing density, axon caliber, and more confounding factors (Chang et al., 2017). However, it has been shown that FA can correlate with myelination quite accurately (Chang et al., 2017). With this in mind, some tentative directions can be ascertained.

Studies focusing on adolescent and young adults who heavily use cannabis have drawn cautious conclusions. Overall, reductions in FA and increases in RD and MD have been observed with chronic exposure to cannabis (Arnone et al., 2008; Ashtari et al., 2009; Gruber et al., 2014; Becker et al., 2015; Shollenbarger et al., 2015). However, Cousijn et al. (2022) did not find any significant differences between chronic, sporadic or control groups, while noting an association between reduced FA and the onset of regular cannabis use at younger ages (<18-years-old). This finding is generally supported in the literature, with many studies indicating that an earlier age of onset correlates with reduced myelin integrity and cognition (Gruber et al., 2012, 2014; Lisdahl et al., 2013; Orr et al., 2016).

Other studies with larger age ranges (~22–55-years of age) revealed similar results with respect to heavy cannabis use. Jakabek

et al. (2016) found reductions in FA in the forceps minor across all age ranges (18–55-years of age), but RD differed depending on age, with younger users having reduced RD and older users having higher RD. Similarly, Manza et al. (2020) reported compromised myelin integrity in heavy cannabis users compared to controls within the same age range. However, Filbey et al. (2014) observed an increase in FA and a decrease in RD, suggesting improvements in white matter integrity, although protracted use did result in a reversal of these findings.

On the contrary, a recent DTI study found that adults (47.85 ± 17.42-years of age) who use medical cannabis with moderate levels of CBD and low levels of THC have significantly increased FA and reduced MD in multiple white matter regions—notably the genu of the corpus callosum—after 6 months of use (Dahlgren et al., 2022). An important distinction between this study and the previously mentioned studies is that the participants were using cannabis for medicinal purposes, not recreationally. As stated by Dahlgren et al. (2022), the medicinal use of cannabis likely involves drastically different characteristics (i.e., age of onset, ratio of cannabinoids), potentially resulting in the increases in myelin integrity found in this study. Furthermore, the medicinal use of cannabis is likely to help treat symptoms associated with increased levels of inflammation, perhaps altering cannabinoid function compared to recreational users. These findings are supported by the numerous animal studies highlighted in this review that show that lower doses of cannabinoids administered less frequently are more beneficial than the reverse.

Despite indications that the brains response to cannabis substantially changes with age, to the best of our knowledge, no DTI study looking at the integrity of myelin has yet been performed in cannabis using seniors, creating a substantial gap in the literature. Through the use of functional MRI, functional connectivity—the statistical relationship that exists between different regions of the brain that are necessary for cognitive processes—can be quantified (Gaudet et al., 2020). Importantly, these functional connections are supported by, and significantly associated with white matter pathways (Hunt et al., 2016; Meier et al., 2016; Huntenburg et al., 2017; Vandewouw et al., 2021). Again, few studies are available for adults over the age of 60. However, one study found increased functional connectivity between the anterior cerebellum and the hippocampus, and with the posterior parahippocampal cortex in cannabis users aged 60–80-years-old (Watson et al., 2022). The authors suggest this finding may indicate a potential benefit of cannabis in the aged brain, although the extent and appearance of these potential benefits are still unclear.

Due to the paucity of studies conducted in older human adults, it is difficult to identify the impact cannabis has on myelination in this population using neuroimaging techniques at this time. Regardless, this evidence forms a basis for future exploration.

2. Conclusion

The studies examined in this review highlight the potential for cannabinoids to aid in myelination in the aged brain. At certain doses, cannabinoids have the ability to reduce the release of pro-inflammatory

cytokines from microglia and astrocytes, scavenge ROS, and promote OPC differentiation in a cell-autonomous manner, resulting in improved myelination after an inflammatory stimulus or demyelination. Furthermore, there is evidence suggesting that some cannabis use in the adult and older population may improve white matter integrity. However, the extreme lack of studies on this topic in the healthy aged brain currently prevents any definitive conclusions from being drawn. Therefore, future studies looking at the impact cannabinoids have on myelination in the aged brain and how this alters behavior and cognition should be performed. As a note, the myelination process in the peripheral nervous system is vastly different than in the CNS and was beyond the scope of this review, but cannabinoids may also influence this process. These studies will not only shed light onto how cannabinoids impact myelination, but will also add vital information as to how the endocannabinoid system contributes to modulating cognition in the aged brain.

Author contributions

CM: conceptualization, writing—original draft preparation, writing—review and editing, and visualizations. HV: conceptualization and writing—review and editing. M-ÉT: conceptualization, writing—review and editing, supervision, and funding acquisition. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by research grants from the Canadian Institutes of Health Research (CIHR) and Natural Sciences and Engineering Research Council of Canada (NSERC) awarded to M-ÉT. HV is the recipient of a CIHR postdoctoral fellowship and is a Michael Smith Health Research BC Research Trainee.

Acknowledgments

We acknowledge and respect the ɫəᑭ^wəŋən Peoples on whose traditional territory the University of Victoria (Victoria, BC, Canada) stands and the Songhees, Esquimalt and W̱SÁNEĆ Peoples whose historical relationships with the land continue to this day.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Aber, E. R., Griffey, C. J., Davies, T., Li, A. M., Yang, Y. J., Croce, K. R., et al. (2022). Oligodendroglial macroautophagy is essential for myelin sheath turnover to prevent neurodegeneration and death. *Cell Rep.* 41:111480. doi: 10.1016/j.celrep.2022.111480
- Abudara, V., Retamal, M. A., Del Rio, R., and Orellana, J. A. (2018). Synaptic functions of Hemichannels and Pannexons: a double-edged sword. *Front. Mol. Neurosci.* 11:435. doi: 10.3389/fnmol.2018.00435
- Abudara, V., Roux, L., Dallérac, G., Matias, I., Dulong, J., Mothet, J. P., et al. (2015). Activated microglia impairs neuroglial interaction by opening Cx43 hemichannels in hippocampal astrocytes: Astroglial Hemichannels impair Neuroglial interaction. *Glia* 63, 795–811. doi: 10.1002/glia.22785
- Aguado, T., Huerga-Gómez, A., Sánchez-de la Torre, A., Resel, E., Chara, J. C., Matute, C., et al. (2021). Δ^9 -tetrahydrocannabinol promotes functional remodeling in the mouse brain. *Br. J. Pharmacol.* 178, 4176–4192. doi: 10.1111/bph.15608
- Aguirre-Rueda, D., Guerra-Ojeda, S., Aldasoro, M., Iradi, A., Obrador, E., Mauricio, M. D., et al. (2015). WIN 55,212-2, agonist of cannabinoid receptors, prevents amyloid β 1-42 effects on astrocytes in primary culture. *PLoS One* 10:e0122843. doi: 10.1371/journal.pone.0122843
- Ahn, K., Lee, S.-J., and Mook-Jung, I. (2022). White matter-associated microglia: new players in brain aging and neurodegenerative diseases. *Ageing Res. Rev.* 75:101574. doi: 10.1016/j.arr.2022.101574
- Albayram, O., Alferink, J., Pitsch, J., Piyanova, A., Neitzert, K., Poppensieker, K., et al. (2011). Role of CB1 cannabinoid receptors on GABAergic neurons in brain aging. *Proc. Natl. Acad. Sci. U. S. A.* 108, 11256–11261. doi: 10.1073/pnas.1016442108
- Alexander, A. L., Lee, J. E., Lazar, M., and Field, A. S. (2007). Diffusion tensor imaging of the brain. *Neurotherapeutics* 4, 316–329. doi: 10.1016/j.nurt.2007.05.011
- Angelova, D. M., and Brown, D. R. (2019). Microglia and the aging brain: are senescent microglia the key to neurodegeneration? *J. Neurochem.* 151, 676–688. doi: 10.1111/jnc.14860
- Arancibia-Carcamo, I. L., and Attwell, D. (2014). The node of Ranvier in CNS pathology. *Acta Neuropathol.* 128, 161–175. doi: 10.1007/s00401-014-1305-z
- Arancibia-Carcamo, I. L., Ford, M. C., Cossell, L., Ishida, K., Tohyama, K., and Attwell, D. (2017). Node of Ranvier length as a potential regulator of myelinated axon conduction speed. *eLife* 6:e23329. doi: 10.7554/eLife.23329
- Arnone, D., Barrick, T. R., Chengappa, S., Mackay, C. E., Clark, C. A., and Abou-Saleh, M. T. (2008). Corpus callosum damage in heavy marijuana use: preliminary evidence from diffusion tensor tractography and tract-based spatial statistics. *NeuroImage* 41, 1067–1074. doi: 10.1016/j.neuroimage.2008.02.064
- Ashtari, M., Cervellione, K., Cottone, J., Ardekani, B. A., and Kumra, S. (2009). Diffusion abnormalities in adolescents and young adults with a history of heavy cannabis use. *J. Psychiatr. Res.* 43, 189–204. doi: 10.1016/j.jpsychires.2008.12.002
- Askew, K., Li, K., Olmos-Alonso, A., Garcia-Moreno, F., Liang, Y., Richardson, P., et al. (2017). Coupled proliferation and apoptosis maintain the rapid turnover of microglia in the adult brain. *Cell Rep.* 18, 391–405. doi: 10.1016/j.celrep.2016.12.041
- Aso, E., Andrés-Benito, P., Carmona, M., Maldonado, R., and Ferrer, I. (2016). Cannabinoid receptor 2 participates in amyloid- β processing in a mouse model of Alzheimer's disease but plays a minor role in the therapeutic properties of a cannabis-based medicine. *JAD* 51, 489–500. doi: 10.3233/JAD-150913
- Aso, E., Juvés, S., Maldonado, R., and Ferrer, I. (2013). CB2 cannabinoid receptor agonist ameliorates Alzheimer-like phenotype in A β PP/PS1 mice. *JAD* 35, 847–858. doi: 10.3233/JAD-130137
- Atalay Ekiner, S., Gęgotek, A., and Skrzydlewska, E. (2022). The molecular activity of cannabidiol in the regulation of Nrf2 system interacting with NF- κ B pathway under oxidative stress. *Redox Biol.* 57:102489. doi: 10.1016/j.redox.2022.102489
- Attia, H., Taha, M., and Abdellatif, A. (2019). Effects of aging on the myelination of the optic nerve in rats. *Int. J. Neurosci.* 129, 320–324. doi: 10.1080/00207454.2018.1529670
- Auguste, Y. S. S., Ferro, A., Kahng, J. A., Xavier, A. M., Dixon, J. R., Vrudhula, U., et al. (2022). Oligodendrocyte precursor cells engulf synapses during circuit remodeling in mice. *Nat. Neurosci.* 25, 1273–1278. doi: 10.1038/s41593-022-01170-x
- Augustin, S. M., and Lovinger, D. M. (2022). Synaptic changes induced by cannabinoid drugs and cannabis use disorder. *Neurobiol. Dis.* 167:105670. doi: 10.1016/j.nbd.2022.105670
- Back, S. A., Gan, X., Li, Y., Rosenberg, P. A., and Volpe, J. J. (1998). Maturation-dependent vulnerability of oligodendrocytes to oxidative stress-induced death caused by glutathione depletion. *J. Neurosci.* 18, 6241–6253. doi: 10.1523/JNEUROSCI.18-16-06241.1998
- Badimon, A., Strasburger, H. J., Ayata, P., Chen, X., Nair, A., Ikegami, A., et al. (2020). Negative feedback control of neuronal activity by microglia. *Nature* 586, 417–423. doi: 10.1038/s41586-020-2777-8
- Baer, A. S., Syed, Y. A., Kang, S. U., Mitteregger, D., Vig, R., ffrench-Constant, C., et al. (2009). Myelin-mediated inhibition of oligodendrocyte precursor differentiation can be overcome by pharmacological modulation of Fyn-RhoA and protein kinase C signalling. *Brain* 132, 465–481. doi: 10.1093/brain/awn334
- Bakiri, Y., Káradóttir, R., Cossell, L., and Attwell, D. (2011). Morphological and electrical properties of oligodendrocytes in the white matter of the corpus callosum and cerebellum: oligodendrocyte electrical properties. *J. Physiol.* 589, 559–573. doi: 10.1113/jphysiol.2010.201376
- Baror, R., Neumann, B., Segel, M., Chalut, K. J., Fancy, S. P. J., Schafer, D. P., et al. (2019). Transforming growth factor-beta renders ageing microglia inhibitory to oligodendrocyte generation by CNS progenitors. *Glia* 67, 1374–1384. doi: 10.1002/glia.23612
- Bartzokis, G. (2004). Age-related myelin breakdown: a developmental model of cognitive decline and Alzheimer's disease. *Neurobiol. Aging* 25, 5–18. doi: 10.1016/j.neurobiolaging.2003.03.001
- Bartzokis, G., Beckson, M., Lu, P. H., Nuechterlein, K. H., Edwards, N., and Mintz, J. (2001). Age-related changes in frontal and temporal lobe volumes in men: a magnetic resonance imaging study. *Arch. Gen. Psychiatry* 58:461. doi: 10.1001/archpsyc.58.5.461
- Baud, O. (2004). Glutathione peroxidase-catalase Cooperativity is required for resistance to hydrogen peroxide by mature rat oligodendrocytes. *J. Neurosci.* 24, 1531–1540. doi: 10.1523/JNEUROSCI.3989-03.2004
- Bechler, M. E., Byrne, L., and ffrench-Constant, C. (2015). CNS myelin sheath lengths are an intrinsic property of oligodendrocytes. *Curr. Biol.* 25, 2411–2416. doi: 10.1016/j.cub.2015.07.056
- Bechler, M. E., Swire, M., and ffrench-Constant, C. (2018). Intrinsic and adaptive myelination—a sequential mechanism for smart wiring in the brain: intrinsic and adaptive myelination mechanisms. *Devel. Neurobiol.* 78, 68–79. doi: 10.1002/dneu.22518
- Becker, M. P., Collins, P. F., Lim, K. O., Muetzel, R. L., and Luciana, M. (2015). Longitudinal changes in white matter microstructure after heavy cannabis use. *Dev. Cogn. Neurosci.* 16, 23–35. doi: 10.1016/j.dcn.2015.10.004
- Beckner, M. E. (2020). A roadmap for potassium buffering/dispersion via the glial network of the CNS. *Neurochem. Int.* 136:104727. doi: 10.1016/j.neuint.2020.104727
- Bénard, G., Massa, F., Puente, N., Lourenço, J., Bellocchio, L., Soria-Gómez, E., et al. (2012). Mitochondrial CB1 receptors regulate neuronal energy metabolism. *Nat. Neurosci.* 15, 558–564. doi: 10.1038/nn.3053
- Benito, C., Tolón, R. M., Pazos, M. R., Núñez, E., Castillo, A. I., and Romero, J. (2008). Cannabinoid CB₂ receptors in human brain inflammation: cannabinoid CB₂ receptors in human brain. *Br. J. Pharmacol.* 153, 277–285. doi: 10.1038/sj.bjp.0707505
- Bennett, I. J., and Madden, D. J. (2014). Disconnected aging: cerebral white matter integrity and age-related differences in cognition. *Neuroscience* 276, 187–205. doi: 10.1016/j.neuroscience.2013.11.026
- Bergles, D. E., and Richardson, W. D. (2016). Oligodendrocyte development and plasticity. *Cold Spring Harb. Perspect. Biol.* 8:a020453. doi: 10.1101/cshperspect.a020453
- Bergles, D. E., Roberts, J. D. B., Somogyi, P., and Jahr, C. E. (2000). Glutamatergic synapses on oligodendrocyte precursor cells in the hippocampus. *Nature* 405, 187–191. doi: 10.1038/35012083
- Bilkei-Gorzo, A. (2012). The endocannabinoid system in normal and pathological brain ageing. *Phil. Trans. R. Soc. B* 367, 3326–3341. doi: 10.1098/rstb.2011.0388
- Bilkei-Gorzo, A., Albayram, O., Ativie, F., Chasan, S., Zimmer, T., Bach, K., et al. (2018). Cannabinoid 1 receptor signaling on GABAergic neurons influences astrocytes in the ageing brain. *PLoS One* 13:e0202566. doi: 10.1371/journal.pone.0202566
- Bilkei-Gorzo, A., Albayram, O., Draffehn, A., Michel, K., Piyanova, A., Oppenheimer, H., et al. (2017). A chronic low dose of Δ^9 -tetrahydrocannabinol (THC) restores cognitive function in old mice. *Nat. Med.* 23, 782–787. doi: 10.1038/nm.4311
- Bilkei-Gorzo, A., Racz, I., Valverde, O., Otto, M., Michel, K., Sarstre, M., et al. (2005). Early age-related cognitive impairment in mice lacking cannabinoid CB1 receptors. *Proc. Natl. Acad. Sci. U. S. A.* 102, 15670–15675. doi: 10.1073/pnas.0504640102
- Bishay, P., Häussler, A., Lim, H.-Y., Oertel, B., Galve-Roperh, I., Ferreirós, N., et al. (2013). Anandamide deficiency and heightened neuropathic pain in aged mice. *Neuropharmacology* 71, 204–215. doi: 10.1016/j.neuropharm.2013.03.021
- Bishop, N. A., Lu, T., and Yankner, B. A. (2010). Neural mechanisms of ageing and cognitive decline. *Nature* 464, 529–535. doi: 10.1038/nature08983
- Bisogno, T., Hanuš, L., De Petrocellis, L., Tchilibon, S., Ponde, D. E., Brandi, I., et al. (2001). Molecular targets for cannabidiol and its synthetic analogues: effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide: Cannabidiol, VR1 receptors and anandamide inactivation. *Br. J. Pharmacol.* 134, 845–852. doi: 10.1038/sj.bjp.0704327
- Blank, T., and Prinz, M. (2014). NF- κ B signaling regulates myelination in the CNS. *Front. Mol. Neurosci.* 7:47. doi: 10.3389/fnmol.2014.00047
- Bobitt, J., Qualls, S. H., Schuchman, M., Wickersham, R., Lum, H. D., Arora, K., et al. (2019). Qualitative analysis of cannabis use among older adults in Colorado. *Drugs Aging* 36, 655–666. doi: 10.1007/s40266-019-00665-w
- Boisvert, M. M., Erikson, G. A., Shokhiev, M. N., and Allen, N. J. (2018). The aging astrocyte Transcriptome from multiple regions of the mouse brain. *Cell Rep.* 22, 269–285. doi: 10.1016/j.celrep.2017.12.039

- Boullerne, A. I. (2016). The history of myelin. *Exp. Neurol.* 283, 431–445. doi: 10.1016/j.expneurol.2016.06.005
- Bowley, M. P., Cabral, H., Rosene, D. L., and Peters, A. (2010). Age changes in myelinated nerve fibers of the cingulate bundle and corpus callosum in the rhesus monkey. *J. Comp. Neurol.* 518, 3046–3064. doi: 10.1002/cne.22379
- Brickman, A. M., Meier, I. B., Korgaonkar, M. S., Provenzano, F. A., Grieve, S. M., Siedlecki, K. L., et al. (2012). Testing the white matter retrogenesis hypothesis of cognitive aging. *Neurobiol. Aging* 33, 1699–1715. doi: 10.1016/j.neurobiolaging.2011.06.001
- Bronzuoli, M. R., Facchinetti, R., Valenza, M., Cassano, T., Steardo, L., and Scuderi, C. (2019). Astrocyte function is affected by aging and not Alzheimer's disease: a preliminary investigation in hippocampi of 3xTg-AD mice. *Front. Pharmacol.* 10:644. doi: 10.3389/fphar.2019.00644
- Brusco, A., Tagliaferro, P., Saez, T., and Onaivi, E. S. (2008). Postsynaptic localization of CB2 cannabinoid receptors in the rat hippocampus. *Synapse* 62, 944–949. doi: 10.1002/syn.20569
- Buchanan, J., Elabbady, L., Collman, F., Jorstad, N. L., Bakken, T. E., Ott, C., et al. (2022). Oligodendrocyte precursor cells ingest axons in the mouse neocortex. *Proc. Natl. Acad. Sci. U. S. A.* 119:e2202580119. doi: 10.1073/pnas.2202580119
- Buscham, T., Eichel, M., Siems, S., and Werner, H. (2019). Turning to myelin turnover. *Neural Regen. Res.* 14:2063. doi: 10.4103/1673-5374.262569
- Buyanova, I. S., and Arsalidou, M. (2021). Cerebral white matter myelination and relations to age, gender, and cognition: a selective review. *Front. Hum. Neurosci.* 15:662031. doi: 10.3389/fnhum.2021.662031
- Calabrese, E. J., and Rubio-Casillas, A. (2018). Biphasic effects of THC in memory and cognition. *Eur. J. Clin. Invest.* 48:e12920. doi: 10.1111/eci.12920
- Camargo, N., Goudriaan, A., van Deijk, A.-L. F., Otte, W. M., Brouwers, J. F., Lodder, H., et al. (2017). Oligodendroglial myelination requires astrocyte-derived lipids. *PLoS Biol.* 15:e1002605. doi: 10.1371/journal.pbio.1002605
- Cantuti-Castelvetri, L., Fitzner, D., Bosch-Queralt, M., Weil, M.-T., Su, M., Sen, P., et al. (2018). Defective cholesterol clearance limits remyelination in the aged central nervous system. *Science* 359, 684–688. doi: 10.1126/science.aan4183
- Cassano, T., Calcagnini, S., Pace, L., De Marco, F., Romano, A., and Gaetani, S. (2017). Cannabinoid receptor 2 signaling in neurodegenerative disorders: from pathogenesis to a promising therapeutic target. *Front. Neurosci.* 11:30. doi: 10.3389/fnins.2017.00030
- Castillo, P. E., Younts, T. J., Chávez, A. E., and Hashimoto, Y. (2012). Endocannabinoid signaling and synaptic function. *Neuron* 76, 70–81. doi: 10.1016/j.neuron.2012.09.020
- Chang, E. H., Argyelan, M., Aggarwal, M., Chandon, T.-S. S., Karlsgodt, K. H., Mori, S., et al. (2017). The role of myelination in measures of white matter integrity: combination of diffusion tensor imaging and two-photon microscopy of CLARITY intact brains. *NeuroImage* 147, 253–261. doi: 10.1016/j.neuroimage.2016.11.068
- Chapman, T. W., and Hill, R. A. (2020). Myelin plasticity in adulthood and aging. *Neurosci. Lett.* 715:134645. doi: 10.1016/j.neulet.2019.134645
- Cheli, V. T., Correale, J., Paez, P. M., and Pasquini, J. M. (2020). Iron metabolism in oligodendrocytes and astrocytes, implications for myelination and Remyelination. *ASN Neuro* 12:175909142096268. doi: 10.1177/1759091420962681
- Chong, S. Y. C., Rosenberg, S. S., Fancy, S. P. J., Zhao, C., Shen, Y.-A. A., Hahn, A. T., et al. (2012). Neurite outgrowth inhibitor Nogo-a establishes spatial segregation and extent of oligodendrocyte myelination. *Proc. Natl. Acad. Sci. U. S. A.* 109, 1299–1304. doi: 10.1073/pnas.1113540109
- Clarke, L. E., Liddel, S. A., Chakraborty, C., Münch, A. E., Heiman, M., and Barres, B. A. (2018). Normal aging induces A1-like astrocyte reactivity. *Proc. Natl. Acad. Sci. U. S. A.* 115, E1896–E1905. doi: 10.1073/pnas.1800165115
- Coelho, A., Fernandes, H. M., Magalhães, R., Moreira, P. S., Marques, P., Soares, J. M., et al. (2021). Signatures of white-matter microstructure degradation during aging and its association with cognitive status. *Sci. Rep.* 11:4517. doi: 10.1038/s41598-021-83983-7
- Compton, D. R., Gold, L. H., Ward, S. J., Balster, R. L., and Martin, B. R. (1992). Aminoalkylindole analogs: cannabimimetic activity of a class of compounds structurally distinct from delta 9-tetrahydrocannabinol. *J. Pharmacol. Exp. Ther.* 263, 1118–1126.
- Correa, F., Hernangómez, M., Mestre, L., Loria, F., Spagnolo, A., Docagne, F., et al. (2010). Anandamide enhances IL-10 production in activated microglia by targeting CB₂ receptors: roles of ERK1/2, JNK, and NF-κB: Anandamide enhances IL-10 production. *Glia* 58, 135–147. doi: 10.1002/glia.20907
- Cousijn, J., Toenders, Y. J., Velzen, L. S., and Kaag, A. M. (2022). The relation between cannabis use, dependence severity and white matter microstructure: a diffusion tensor imaging study. *Addict. Biol.* 27:e13081. doi: 10.1111/adb.13081
- Cox, S. R., Ritchie, S. J., Tucker-Drob, E. M., Liewald, D. C., Hagenaars, S. P., Davies, G., et al. (2016). Ageing and brain white matter structure in 3,513 UK biobank participants. *Nat. Commun.* 7:13629. doi: 10.1038/ncomms13629
- Cremers, L. G. M., de Groot, M., Hofman, A., Krestin, G. P., van der Lugt, A., Niessen, W. J., et al. (2016). Altered tract-specific white matter microstructure is related to poorer cognitive performance: the Rotterdam study. *Neurobiol. Aging* 39, 108–117. doi: 10.1016/j.neurobiolaging.2015.11.021
- Cserép, C., Pósfai, B., and Dénes, Á. (2021). Shaping neuronal fate: functional heterogeneity of direct microglia-neuron interactions. *Neuron* 109, 222–240. doi: 10.1016/j.neuron.2020.11.007
- Cutando, L., Busquets-García, A., Puighearnal, E., Gomis-González, M., Delgado-García, J. M., Gruart, A., et al. (2013). Microglial activation underlies cerebellar deficits produced by repeated cannabis exposure. *J. Clin. Invest.* 123, 2816–2831. doi: 10.1172/JCI67569
- D'Souza, D. C., Cortes-Briones, J. A., Ranganathan, M., Thurnauer, H., Creatura, G., Surti, T., et al. (2016). Rapid changes in cannabinoid 1 receptor availability in cannabis-dependent male subjects after abstinence from cannabis. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* 1, 60–67. doi: 10.1016/j.bpsc.2015.09.008
- Dahlgren, M. K., Gonenc, A., Sagar, K. A., Smith, R. T., Lambros, A. M., El-Abboud, C., et al. (2022). Increased white matter coherence following three and six months of medical cannabis treatment. *Cannabis Cannabinoid Res.* 2022:0097. doi: 10.1089/can.2022.0097
- de Almeida, V., Seabra, G., Reis-de-Oliveira, G., Zuccoli, G. S., Rumin, P., Fioramonte, M., et al. (2022). Cannabinoids modulate proliferation, differentiation, and migration signaling pathways in oligodendrocytes. *Eur. Arch. Psychiatry Clin. Neurosci.* 272, 1311–1323. doi: 10.1007/s00406-022-01425-5
- de la Fuente, A. G., Queiroz, R. M. L., Ghosh, T., McMurrin, C. E., Cubillos, J. F., Bergles, D. E., et al. (2020). Changes in the oligodendrocyte progenitor cell proteome with ageing. *Mol. Cell. Proteomics* 19, 1281–1302. doi: 10.1074/mcp.RA120.002102
- De Meij, J., Alfaneq, Z., Morel, L., Decoeur, F., Leyrolle, Q., Picard, K., et al. (2021). Microglial cannabinoid type 1 receptor regulates brain inflammation in a sex-specific manner. *Cannabis Cannabinoid Res.* 6, 488–507. doi: 10.1089/can.2020.0170
- de Oliveira, P. G., Ramos, M. L. S., Amaro, A. J., Dias, R. A., and Vieira, S. I. (2019). Gi/o-protein coupled receptors in the aging brain. *Front. Aging Neurosci.* 11:89. doi: 10.3389/fnagi.2019.00089
- De Petrocellis, L., Ligresti, A., Moriello, A. S., Allarà, M., Bisogno, T., Petrosino, S., et al. (2011). Effects of cannabinoids and cannabinoid-enriched cannabis extracts on TRP channels and endocannabinoid metabolic enzymes: novel pharmacology of minor plant cannabinoids. *Br. J. Pharmacol.* 163, 1479–1494. doi: 10.1111/j.1476-5381.2010.01166.x
- Di Marzo, V., Stella, N., and Zimmer, A. (2015). Endocannabinoid signalling and the deteriorating brain. *Nat. Rev. Neurosci.* 16, 30–42. doi: 10.1038/nrn3876
- Dimovasilis, C., Fair, A. E., Garza, I. R., Batterman, K. V., Mortazavi, F., Moore, T. L., et al. (2022). Aging compromises oligodendrocyte precursor cell maturation and efficient remyelination in the monkey brain. *GeroScience* 45, 249–264. doi: 10.1007/s11357-022-00621-4
- Ding, F., Yao, J., Rettberg, J. R., Chen, S., and Brinton, R. D. (2013). Early decline in glucose transport and metabolism precedes shift to Ketogenic system in female aging and Alzheimer's mouse brain: implication for bioenergetic intervention. *PLoS One* 8:e79977. doi: 10.1371/journal.pone.0079977
- Djannatian, M., Radha, S., Weikert, U., Safaiyan, S., Wrede, C., Deichsel, C., et al. (2023). Myelination generates aberrant ultrastructure that is resolved by microglia. *Journal of Cell Biology* 222:e202204010. doi: 10.1083/jcb.202204010
- dos-Santos-Pereira, M., Guimarães, F. S., del-Bel, E., Raisman-Vozari, R., and Michel, P. P. (2020). Cannabidiol prevents LPS-induced microglial inflammation by inhibiting ROS/NF-κB-dependent signaling and glucose consumption. *Glia* 68, 561–573. doi: 10.1002/glia.23738
- Doucette, J., Jiao, R., and Nazarali, A. J. (2010). Age-related and Cuprizone-induced changes in myelin and transcription factor gene expression and in oligodendrocyte cell densities in the rostral corpus callosum of mice. *Cell. Mol. Neurobiol.* 30, 607–629. doi: 10.1007/s10571-009-9486-z
- Duncan, I. D., Radcliff, A. B., Heidari, M., Kidd, G., August, B. K., and Wierenga, L. A. (2018). The adult oligodendrocyte can participate in remyelination. *Proc. Natl. Acad. Sci. U. S. A.* 115, E11807–E11816. doi: 10.1073/pnas.1808064115
- Ehrhart, J., Obregon, D., Mori, T., Hou, H., Sun, N., Bai, Y., et al. (2005). Stimulation of cannabinoid receptor 2 (CB2) suppresses microglial activation. *J. Neuroinflammation* 2:29. doi: 10.1186/1742-2094-2-29
- Eljaschewitsch, E., Witting, A., Mawrin, C., Lee, T., Schmidt, P. M., Wolf, S., et al. (2006). The Endocannabinoid Anandamide protects neurons during CNS inflammation by induction of MKP-1 in microglial cells. *Neuron* 49, 67–79. doi: 10.1016/j.neuron.2005.11.027
- Esposito, G., Scuderi, C., Valenza, M., Togna, G. I., Latina, V., De Filippis, D., et al. (2011). Cannabidiol reduces Aβ-induced Neuroinflammation and promotes hippocampal neurogenesis through PPARγ involvement. *PLoS One* 6:e28668. doi: 10.1371/journal.pone.0028668
- Faizy, T. D., Thaler, C., Broocks, G., Flottmann, F., Leischner, H., Knipf, H., et al. (2020). The myelin water fraction serves as a marker for age-related myelin alterations in the cerebral white matter – a multiparametric MRI aging study. *Front. Neurosci.* 14:136. doi: 10.3389/fnins.2020.00136
- Faria, O., Gonsalves, D. G., Nicholson, M., and Xiao, J. (2019). Activity-dependent central nervous system myelination throughout life. *J. Neurochem.* 148, 447–461. doi: 10.1111/jnc.14592

- Feliú, A., Moreno-Martet, M., Mecha, M., Carrillo-Salinas, F. J., de Lago, E., Fernández-Ruiz, J., et al. (2015). A Sativex®-like combination of phytocannabinoids as a disease-modifying therapy in a viral model of multiple sclerosis: Sativex® as a disease-modifying therapy in TMEV-IDD. *Br. J. Pharmacol.* 172, 3579–3595. doi: 10.1111/bph.13159
- Ferber, S. G., Namdar, D., Hen-Shoval, D., Eger, G., Koltai, H., Shoval, G., et al. (2020). The “entourage effect”: Terpenes coupled with cannabinoids for the treatment of mood disorders and anxiety disorders. *CN* 18, 87–96. doi: 10.2174/1570159X17666190903103923
- Ferrucci, L., Harris, T. B., Guralnik, J. M., Tracy, R. P., Corti, M.-C., Cohen, H. J., et al. (1999). Serum IL-6 level and the development of disability in older persons. *J. Am. Geriatr. Soc.* 47, 639–646. doi: 10.1111/j.1532-5415.1999.tb01583.x
- Fields, R. D. (2015). A new mechanism of nervous system plasticity: activity-dependent myelination. *Nat. Rev. Neurosci.* 16, 756–767. doi: 10.1038/nrn4023
- Fields, J. A., Swinton, M. K., Montilla-Perez, P., Ricciardelli, E., and Telese, F. (2022). The cannabinoid receptor agonist, WIN-55212-2, suppresses the activation of Proinflammatory genes induced by interleukin 1 Beta in human astrocytes. *Cannabis Cannabinoid Res.* 7, 78–92. doi: 10.1089/can.2020.0128
- Filbey, F. M., Aslan, S., Calhoun, V. D., Spence, J. S., Damaraju, E., Caprihan, A., et al. (2014). Long-term effects of marijuana use on the brain. *Proc. Natl. Acad. Sci. U. S. A.* 111, 16913–16918. doi: 10.1073/pnas.1415297111
- Finlay, D. B., Sircombe, K. J., Nimick, M., Jones, C., and Glass, M. (2020). Terpenoids from cannabis do not mediate an entourage effect by acting at cannabinoid receptors. *Front. Pharmacol.* 11:359. doi: 10.3389/fphar.2020.00359
- Fletcher-Jones, A., Hildick, K. L., Evans, A. J., Nakamura, Y., Henley, J. M., and Wilkinson, K. A. (2020). Protein Interactors and trafficking pathways that regulate the cannabinoid type 1 receptor (CB1R). *Front. Mol. Neurosci.* 13:108. doi: 10.3389/fnmol.2020.00108
- Ford, M. C., Alexandrova, O., Cossell, L., Stange-Marten, A., Sinclair, J., Kopp-Scheinplug, C., et al. (2015). Tuning of Ranvier node and internode properties in myelinated axons to adjust action potential timing. *Nat. Commun.* 6:8073. doi: 10.1038/ncomms9073
- French, H. M., Reid, M., Mamontov, P., Simmons, R. A., and Grinspan, J. B. (2009). Oxidative stress disrupts oligodendrocyte maturation. *J. Neurosci. Res.* 87, 3076–3087. doi: 10.1002/jnr.22139
- Froger, N., Orellana, J. A., Calvo, C.-F., Amigou, E., Kozoriz, M. G., Naus, C. C., et al. (2010). Inhibition of cytokine-induced connexin43 hemichannel activity in astrocytes is neuroprotective. *Mol. Cell. Neurosci.* 45, 37–46. doi: 10.1016/j.mcn.2010.05.007
- Froger, N., Orellana, J. A., Cohen-Salmon, M., Ezan, P., Amigou, E., SÁjcz, J. C., et al. (2009). Cannabinoids prevent the opposite regulation of astroglial connexin43 hemichannels and gap junction channels induced by pro-inflammatory treatments. *J. Neurochem.* 111, 1383–1397. doi: 10.1111/j.1471-4159.2009.06407.x
- Fu, Y., Sun, W., Shi, Y., Shi, R., and Cheng, J.-X. (2009). Glutamate Excitotoxicity inflicts Paranodal myelin splitting and retraction. *PLoS One* 4:e6705. doi: 10.1371/journal.pone.0006705
- Fulmer, C. G., VonDran, M. W., Stillman, A. A., Huang, Y., Hempstead, B. L., and Dreyfus, C. F. (2014). Astrocyte-derived BDNF supports myelin protein synthesis after Cuprizone-induced demyelination. *J. Neurosci.* 34, 8186–8196. doi: 10.1523/JNEUROSCI.4267-13.2014
- Fünfschilling, U., Supplie, L. M., Mahad, D., Boretius, S., Saab, A. S., Edgar, J., et al. (2012). Glycolytic oligodendrocytes maintain myelin and long-term axonal integrity. *Nature* 485, 517–521. doi: 10.1038/nature11007
- Furber, K. L., Lacombe, R. J. S., Caine, S., Thangaraj, M. P., Read, S., Rosendahl, S. M., et al. (2022). Biochemical alterations in white matter tracts of the aging mouse brain revealed by FTIR spectroscopy imaging. *Neurochem. Res.* 47, 795–810. doi: 10.1007/s11064-021-03491-y
- Gajardo-Gómez, R., Labra, V. C., Maturana, C. J., Shoji, K. F., Santibañez, C. A., Sáez, J. C., et al. (2017). Cannabinoids prevent the amyloid β -induced activation of astroglial hemichannels: a neuroprotective mechanism: CBs restore Neuroglial interaction. *Glia* 65, 122–137. doi: 10.1002/glia.23080
- Gaudet, I., Hüsser, A., Vannasing, P., and Gallagher, A. (2020). Functional brain connectivity of language functions in children revealed by EEG and MEG: a systematic review. *Front. Hum. Neurosci.* 14:62. doi: 10.3389/fnhum.2020.00062
- Gautier, H. O. B., Evans, K. A., Volbracht, K., James, R., Sitnikov, S., Lundgaard, L., et al. (2015). Neuronal activity regulates remyelination via glutamate signalling to oligodendrocyte progenitors. *Nat. Commun.* 6:8518. doi: 10.1038/ncomms9518
- Giacci, M. K., Bartlett, C. A., Smith, N. M., Iyer, K. S., Toomey, L. M., Jiang, H., et al. (2018). Oligodendroglia are particularly vulnerable to oxidative damage after Neurotrauma in vivo. *J. Neurosci.* 38, 6491–6504. doi: 10.1523/JNEUROSCI.1898-17.2018
- Giacoppo, S., Pollastro, F., Grassi, G., Bramanti, P., and Mazzoni, E. (2017). Target regulation of PI3K/Akt/mTOR pathway by cannabidiol in treatment of experimental multiple sclerosis. *Fitoterapia* 116, 77–84. doi: 10.1016/j.fitote.2016.11.010
- Gibson, E. M., Purger, D., Mount, C. W., Goldstein, A. K., Lin, G. L., Wood, L. S., et al. (2014). Neuronal activity promotes Oligodendrogenesis and adaptive myelination in the mammalian brain. *Science* 344:1252304. doi: 10.1126/science.1252304
- Ginsburg, B. C., and Hensler, J. G. (2022). Age-related changes in CB1 receptor expression and function and the behavioral effects of cannabinoid receptor ligands. *Pharmacol. Biochem. Behav.* 213:173339. doi: 10.1016/j.pbb.2022.173339
- Glass, M., Faull, R. L. M., and Dragunow, M. (1997). Cannabinoid receptors in the human brain: a detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience* 77, 299–318. doi: 10.1016/S0306-4522(96)00428-9
- Godbout, J. P., and Johnson, R. W. (2004). Interleukin-6 in the aging brain. *J. Neuroimmunol.* 147, 141–144. doi: 10.1016/j.jneuroim.2003.10.031
- Goldmann, T., Wieghofer, P., Müller, P. F., Wolf, Y., Varol, D., Yona, S., et al. (2013). A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nat. Neurosci.* 16, 1618–1626. doi: 10.1038/nn.3531
- Gomez, O., Arevalo-Martin, A., Garcia-Ovejero, D., Ortega-Gutierrez, S., Cisneros, J. A., Almazan, G., et al. (2010). The constitutive production of the endocannabinoid 2-arachidonoylglycerol participates in oligodendrocyte differentiation. *Glia* 58, 1913–1927. doi: 10.1002/glia.21061
- Gomez, O., Sanchez-Rodriguez, A., Le, M., Sanchez-Caro, C., Molina-Holgado, F., and Molina-Holgado, E. (2011). Cannabinoid receptor agonists modulate oligodendrocyte differentiation by activating PI3K/Akt and the mammalian target of rapamycin (mTOR) pathways: cannabinoids promote oligodendrocyte differentiation. *Br. J. Pharmacol.* 163, 1520–1532. doi: 10.1111/j.1476-5381.2011.01414.x
- Goursaud, S., Kozlova, E. N., Maloteaux, J.-M., and Hermans, E. (2009). Cultured astrocytes derived from corpus callosum or cortical grey matter show distinct glutamate handling properties. *J. Neurochem.* 108, 1442–1452. doi: 10.1111/j.1471-4159.2009.05889.x
- Grabert, K., Michoel, T., Karavolos, M. H., Clohisey, S., Baillie, J. K., Stevens, M. P., et al. (2016). Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nat. Neurosci.* 19, 504–516. doi: 10.1038/nn.4222
- Grosche, A., Grosche, J., Tackenberg, M., Scheller, D., Gerstner, G., Gumprecht, A., et al. (2013). Versatile and simple approach to determine astrocyte territories in mouse neocortex and hippocampus. *PLoS One* 8:e69143. doi: 10.1371/journal.pone.0069143
- Gruber, S. A., Dahlgren, M. K., Sagar, K. A., Gönenç, A., and Lukas, S. E. (2014). Worth the wait: effects of age of onset of marijuana use on white matter and impulsivity. *Psychopharmacology* 231, 1455–1465. doi: 10.1007/s00213-013-3326-z
- Gruber, S. A., Sagar, K. A., Dahlgren, M. K., Racine, M., and Lukas, S. E. (2012). Age of onset of marijuana use and executive function. *Psychol. Addict. Behav.* 26, 496–506. doi: 10.1037/a0026269
- Guttenplan, K. A., Weigel, M. K., Prakash, P., Wijewardhane, P. R., Hasel, P., Rufen-Blanchette, U., et al. (2021). Neurotoxic reactive astrocytes induce cell death via saturated lipids. *Nature* 599, 102–107. doi: 10.1038/s41586-021-03960-y
- Guttman, C. R. G., Jolesz, F. A., Kikinis, R., Killiany, R. J., Moss, M. B., Sandor, T., et al. (1998). White matter changes with normal aging. *Neurology* 50, 972–978. doi: 10.1212/WNL.50.4.972
- Han, J., Kesner, P., Metna-Laurent, M., Duan, T., Xu, L., Georges, F., et al. (2012). Acute cannabinoids impair working memory through Astroglial CB1 receptor modulation of hippocampal LTD. *Cells* 148, 1039–1050. doi: 10.1016/j.cell.2012.01.037
- Han, B. H., and Palamar, J. J. (2020). Trends in cannabis use among older adults in the United States, 2015–2018. *JAMA Intern. Med.* 180:609. doi: 10.1001/jamainternmed.2019.7517
- Hassan, S., Eldeeb, K., Millns, P. J., Bennett, A. J., Alexander, S. P. H., and Kendall, D. A. (2014). Cannabidiol enhances microglial phagocytosis via transient receptor potential (TRP) channel activation: Cannabidiol enhances microglial phagocytosis. *Br. J. Pharmacol.* 171, 2426–2439. doi: 10.1111/bph.12615
- Hassel, B., Boldingh, K. A., Narvesen, C., Iversen, E. G., and Skrede, K. K. (2003). Glutamate transport, glutamine synthetase and phosphate-activated glutaminase in rat CNS white matter. A quantitative study: glutamate uptake in white matter. *J. Neurochem.* 87, 230–237. doi: 10.1046/j.1471-4159.2003.01984.x
- Henriquez, J. E., Bach, A. P., Matos-Fernandez, K. M., Crawford, R. B., and Kaminski, N. E. (2020). Δ^9 -tetrahydrocannabinol (THC) impairs CD8+ T cell-mediated activation of astrocytes. *J. Neuroimmune Pharmacol.* 15, 863–874. doi: 10.1007/s11481-020-09912-z
- Herkenham, M., Lynn, A. B., Little, M. D., Johnson, M. R., Melvin, L. S., de Costa, B. R., et al. (1990). Cannabinoid receptor localization in brain. *Proc. Natl. Acad. Sci. U. S. A.* 87, 1932–1936. doi: 10.1073/pnas.87.5.1932
- Herx, L. M., Rivest, S., and Yong, V. W. (2000). Central nervous system-initiated inflammation and Neurotrophin in trauma: IL-1 β is required for the production of ciliary Neurotrophic factor. *J. Immunol.* 165, 2232–2239. doi: 10.4049/jimmunol.165.4.2232
- Hickman, S. E., Kingery, N. D., Ohsumi, T. K., Borowsky, M. L., Wang, L., Means, T. K., et al. (2013). The microglial sensome revealed by direct RNA sequencing. *Nat. Neurosci.* 16, 1896–1905. doi: 10.1038/nn.3554
- Hill, R. A., Li, A. M., and Grutzendler, J. (2018). Lifelong cortical myelin plasticity and age-related degeneration in the live mammalian brain. *Nat. Neurosci.* 21, 683–695. doi: 10.1038/s41593-018-0120-6

- Hines, J. H., Ravanelli, A. M., Schwandt, R., Scott, E. K., and Appel, B. (2015). Neuronal activity biases axon selection for myelination in vivo. *Nat. Neurosci.* 18, 683–689. doi: 10.1038/nn.3992
- Hinman, J. D., Peters, A., Cabral, H., Rosene, D. L., Hollander, W., Rasband, M. N., et al. (2006). Age-related molecular reorganization at the node of Ranvier. *J. Comp. Neurol.* 495, 351–362. doi: 10.1002/cne.20886
- Hirvonen, J., Goodwin, R. S., Li, C.-T., Terry, G. E., Zoghbi, S. S., Morse, C., et al. (2012). Reversible and regionally selective downregulation of brain cannabinoid CB1 receptors in chronic daily cannabis smokers. *Mol. Psychiatry* 17, 642–649. doi: 10.1038/mp.2011.82
- Hodges, E. L., Marshall, J. P., and Ashpole, N. M. (2020). Age-dependent hormesis-like effects of the synthetic cannabinoid CP55940 in C57BL/6 mice. *NPJ Aging Mech. Dis.* 6:7. doi: 10.1038/s41514-020-0045-7
- Holling, T. M., Schooten, E., and van Den Elsen, P. J. (2004). Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. *Hum. Immunol.* 65, 282–290. doi: 10.1016/j.humimm.2004.01.005
- Howlett, A. C. (2002). International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol. Rev.* 54, 161–202. doi: 10.1124/pr.54.2.161
- Hsieh, J., Aimone, J. B., Kaspar, B. K., Kuwabara, T., Nakashima, K., and Gage, F. H. (2004). IGF-I instructs multipotent adult neural progenitor cells to become oligodendrocytes. *J. Cell Biol.* 164, 111–122. doi: 10.1083/jcb.200308101
- Hu, M., Lin, Y., Zhang, B., Lu, D., Lu, Z., and Cai, W. (2019). Update of inflammasome activation in microglia/macrophage in aging and aging-related disease. *CNS Neurosci. Ther.* 25, 1299–1307. doi: 10.1111/cns.13262
- Huerga-Gómez, A., Aguado, T., Sánchez-de la Torre, A., Bernal-Chico, A., Matute, C., Mato, S., et al. (2021). Δ^9 -tetrahydrocannabinol promotes oligodendrocyte development and CNS myelination in vivo. *Glia* 69, 532–545. doi: 10.1002/glia.23911
- Hughes, A. N., and Appel, B. (2020). Microglia phagocytose myelin sheaths to modify developmental myelination. *Nat. Neurosci.* 23, 1055–1066. doi: 10.1038/s41593-020-0654-2
- Hughes, E. G., Orthmann-Murphy, J. L., Langseth, A. J., and Bergles, D. E. (2018). Myelin remodeling through experience-dependent oligodendrogenesis in the adult somatosensory cortex. *Nat. Neurosci.* 21, 696–706. doi: 10.1038/s41593-018-0121-5
- Hunt, B. A. E., Tevarie, P. K., Mouglin, O. E., Geades, N., Jones, D. K., Singh, K. D., et al. (2016). Relationships between cortical myeloarchitecture and electrophysiological networks. *Proc. Natl. Acad. Sci. U. S. A.* 113, 13510–13515. doi: 10.1073/pnas.1608587113
- Huntenburg, J. M., Bazin, P.-L., Goulas, A., Tardif, C. L., Villringer, A., and Margulies, D. S. (2017). A systematic relationship between functional connectivity and intracortical myelin in the human cerebral cortex. *Cereb. Cortex* 27, 981–997. doi: 10.1093/cercor/bhx030
- Huxley, A. F., and Stämpfli, R. (1949). Evidence for saltatory conduction in peripheral myelinated nerve fibres. *J. Physiol.* 108, 315–339. doi: 10.1113/jphysiol.1949.sp004335
- Ichihara, Y., Doi, T., Ryu, Y., Nagao, M., Sawada, Y., and Ogata, T. (2017). Oligodendrocyte progenitor cells directly utilize lactate for promoting cell cycling and differentiation: DIRECT EFFECTS OF LACTATE ON OPCs. *J. Cell. Physiol.* 232, 986–995. doi: 10.1002/jcp.25690
- Ilyasov, A. A., Milligan, C. E., Pharr, E. P., and Howlett, A. C. (2018). The Endocannabinoid system and oligodendrocytes in health and disease. *Front. Neurosci.* 12:733. doi: 10.3389/fnins.2018.00733
- Jakabek, D., Yücel, M., Lorenzetti, V., and Solowij, N. (2016). An MRI study of white matter tract integrity in regular cannabis users: effects of cannabis use and age. *Psychopharmacology* 233, 3627–3637. doi: 10.1007/s00213-016-4398-3
- Jha, M. K., Jo, M., Kim, J.-H., and Suk, K. (2019). Microglia-astrocyte crosstalk: an intimate molecular conversation. *Neuroscientist* 25, 227–240. doi: 10.1177/1073858418783959
- Jha, S., Srivastava, S. Y., Brickey, W. J., Iocca, H., Toews, A., Morrison, J. P., et al. (2010). The Inflammasome sensor, NLRP3, regulates CNS inflammation and demyelination via Caspase-1 and Interleukin-18. *J. Neurosci.* 30, 15811–15820. doi: 10.1523/JNEUROSCI.4088-10.2010
- Jimenez-Blasco, D., Busquets-Garcia, A., Hebert-Chatelain, E., Serrat, R., Vicente-Gutierrez, C., Ioannidou, C., et al. (2020). Glucose metabolism links astroglial mitochondria to cannabinoid effects. *Nature* 583, 603–608. doi: 10.1038/s41586-020-2470-y
- Jordan, C. J., and Xi, Z.-X. (2019). Progress in brain cannabinoid CB2 receptor research: from genes to behavior. *Neurosci. Biobehav. Rev.* 98, 208–220. doi: 10.1016/j.neubiorev.2018.12.026
- Jünemann, K., Marie, D., Worschech, F., Scholz, D. S., Grouiller, F., Kliegel, M., et al. (2022). Six months of piano training in healthy elderly stabilizes white matter microstructure in the fornix, compared to an active control group. *Front. Aging Neurosci.* 14:817889. doi: 10.3389/fnagi.2022.817889
- Jurevics, H., and Morell, P. (2002). Cholesterol for synthesis of myelin is made locally, not imported into brain. *J. Neurochem.* 64, 895–901. doi: 10.1046/j.1471-4159.1995.64020895.x
- Jurewicz, A., Matysiak, M., Tybor, K., Kilianek, L., Raine, C. S., and Selmaj, K. (2005). Tumour necrosis factor-induced death of adult human oligodendrocytes is mediated by apoptosis inducing factor. *Brain* 128, 2675–2688. doi: 10.1093/brain/awh627
- Juurink, B. H. J., Thorburne, S. K., and Hertz, L. (1998). Peroxide-scavenging deficit underlies oligodendrocyte susceptibility to oxidative stress. *Glia* 22, 371–378. doi: 10.1002/(SICI)1098-1136(199804)22:4<371::AID-GLIA6>3.0.CO;2-6
- Kalafatakis, I., and Karagogeos, D. (2021). Oligodendrocytes and microglia: key players in myelin development, Damage and Repair. *Biomolecules* 11:1058. doi: 10.3390/biom11071058
- Kano, M., Ohno-Shosaku, T., Hashimoto-dani, Y., Uchigashima, M., and Watanabe, M. (2009). Endocannabinoid-mediated control of synaptic transmission. *Physiol. Rev.* 89, 309–380. doi: 10.1152/physrev.00019.2008
- Karpuk, N., Burkovetskaya, M., Fritz, T., Angle, A., and Kielian, T. (2011). Neuroinflammation leads to region-dependent alterations in astrocyte gap junction communication and Hemichannel activity. *J. Neurosci.* 31, 414–425. doi: 10.1523/JNEUROSCI.5247-10.2011
- Kato, G., Inada, H., Wake, H., Akiyoshi, R., Miyamoto, A., Eto, K., et al. (2016). Microglial contact prevents excess depolarization and rescues neurons from Excitotoxicity. *eNeuro* 3:ENEURO.0004-16.2016. doi: 10.1523/ENEURO.0004-16.2016
- Keethakumar, A., Mehra, V. M., Khanlou, N., and Tamim, H. (2021). Cannabis use and patterns among middle and older aged Canadians prior to legalization: a sex-specific analysis of the Canadian tobacco, alcohol and drugs survey. *BMC Public Health* 21:26. doi: 10.1186/s12889-020-10074-z
- Keough, M. B., Rogers, J. A., Zhang, P., Jensen, S. K., Stephenson, E. L., Chen, T., et al. (2016). An inhibitor of chondroitin sulfate proteoglycan synthesis promotes central nervous system remyelination. *Nat. Commun.* 7:11312. doi: 10.1038/ncomms11312
- Kimura, A., and Kishimoto, T. (2010). IL-6: regulator of Treg/Th17 balance. *Eur. J. Immunol.* 40, 1830–1835. doi: 10.1002/eji.201040391
- Kiray, H., Lindsay, S. L., Hosseinzadeh, S., and Barnett, S. C. (2016). The multifaceted role of astrocytes in regulating myelination. *Exp. Neurol.* 283, 541–549. doi: 10.1016/j.expneurol.2016.03.009
- Koellhoffer, E., McCullough, L., and Ritzel, R. (2017). Old maids: aging and its impact on microglia function. *IJMS* 18:769. doi: 10.3390/ijms18040769
- Komorowska-Müller, J. A., Rana, T., Olabiyi, B. F., Zimmer, A., and Schmöle, A.-C. (2021a). Cannabinoid receptor 2 alters social memory and microglial activity in an age-dependent manner. *Molecules* 26:5984. doi: 10.3390/molecules26195984
- Komorowska-Müller, J. A., Ravichandran, K. A., Zimmer, A., and Schürmann, B. (2021b). Cannabinoid receptor 2 deletion influences social memory and synaptic architecture in the hippocampus. *Sci. Rep.* 11:16828. doi: 10.1038/s41598-021-96285-9
- Komorowska-Müller, J. A., and Schmöle, A.-C. (2020). CB2 receptor in microglia: the Guardian of self-control. *IJMS* 22:19. doi: 10.3390/ijms22010019
- Kopustinskiene, D. M., Masteikova, R., Lazauskas, R., and Bernatoniene, J. (2022). Cannabis sativa L. bioactive compounds and their protective role in oxidative stress and inflammation. *Antioxidants* 11:660. doi: 10.3390/antiox11040660
- Kotter, M. R. (2006). Myelin impairs CNS Remyelination by inhibiting oligodendrocyte precursor cell differentiation. *J. Neurosci.* 26, 328–332. doi: 10.1523/JNEUROSCI.2615-05.2006
- Kougoumtzidou, E., Shimizu, T., Hamilton, N. B., Tohyama, K., Sprengel, R., Monyer, H., et al. (2017). Signalling through AMPA receptors on oligodendrocyte precursors promotes myelination by enhancing oligodendrocyte survival. *eLife* 6:e28080. doi: 10.7554/eLife.28080
- Kozela, E., Pietr, M., Juknat, A., Rimmerman, N., Levy, R., and Vogel, Z. (2010). Cannabinoids Δ^9 -tetrahydrocannabinol and Cannabidiol differentially inhibit the lipopolysaccharide-activated NF- κ B and interferon- β /STAT Proinflammatory pathways in BV-2 microglial cells. *J. Biol. Chem.* 285, 1616–1626. doi: 10.1074/jbc.M109.069294
- Laaris, N., Good, C. H., and Lupica, C. R. (2010). Δ^9 -tetrahydrocannabinol is a full agonist at CB1 receptors on GABA neuron axon terminals in the hippocampus. *Neuropharmacology* 59, 121–127. doi: 10.1016/j.neuropharm.2010.04.013
- Labra, V. C., Santibáñez, C. A., Gajardo-Gómez, R., Díaz, E. F., Gómez, G. I., and Orellana, J. A. (2018). The Neuroglial dialog between cannabinoids and Hemichannels. *Front. Mol. Neurosci.* 11:79. doi: 10.3389/fnmol.2018.00079
- Lalo, U., Palygin, O., North, R. A., Verkhatsky, A., and Pankratov, Y. (2011). Age-dependent remodelling of ionotropic signalling in cortical astroglia: synaptic currents in aging astrocytes. *Aging Cell* 10, 392–402. doi: 10.1111/j.1474-9726.2011.00682.x
- Lampron, A., Laroche, A., Laflamme, N., Préfontaine, P., Plante, M.-M., Sánchez, M. G., et al. (2015). Inefficient clearance of myelin debris by microglia impairs remyelinating processes. *J. Exp. Med.* 212, 481–495. doi: 10.1084/jem.20141656
- Lanciego, J. L., Barroso-Chinea, P., Rico, A. J., Conte-Perales, L., Callén, L., Roda, E., et al. (2011). Expression of the mRNA coding the cannabinoid receptor 2 in the pallidum complex of *Macaca fascicularis*. *J. Psychopharmacol.* 25, 97–104. doi: 10.1177/0269881110367732
- Laprairie, R. B., Bagher, A. M., Kelly, M. E. M., and Denovan-Wright, E. M. (2015). Cannabidiol is a negative allosteric modulator of the cannabinoid CB₁ receptor: negative allosteric modulation of CB₁ by cannabidiol. *Br. J. Pharmacol.* 172, 4790–4805. doi: 10.1111/bph.13250
- Laurikainen, H., Tuominen, L., Tikka, M., Merisaari, H., Armio, R.-L., Sormunen, E., et al. (2019). Sex difference in brain CB1 receptor availability in man. *NeuroImage* 184, 834–842. doi: 10.1016/j.neuroimage.2018.10.013

- Lawson, L. J., Perry, V. H., Dri, P., and Gordon, S. (1990). Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience* 39, 151–170. doi: 10.1016/0306-4522(90)90229-W
- Lee, H.-L., Jung, K.-M., Fotio, Y., Squire, E., Palese, F., Lin, L., et al. (2022). Frequent low-dose Δ^9 -tetrahydrocannabinol in adolescence disrupts microglia homeostasis and disables responses to microbial infection and social stress in Young adulthood. *Biol. Psychiatry* 92, 845–860. doi: 10.1016/j.biopsych.2022.04.017
- Lee, Y., Morrison, B. M., Li, Y., Lengacher, S., Farah, M. H., Hoffman, P. N., et al. (2012). Oligodendroglia metabolically support axons and contribute to neurodegeneration. *Nature* 487, 443–448. doi: 10.1038/nature11314
- Leishman, E., Murphy, M., Mackie, K., and Bradshaw, H. B. (2018). Δ^9 -tetrahydrocannabinol changes the brain lipidome and transcriptome differentially in the adolescent and the adult. *Biochim. Biophys. Acta (BBA) – Mol. Cell Biol. Lipids* 1863, 479–492. doi: 10.1016/j.bbalip.2018.02.001
- Levine, A., Liktov-Busa, E., Lipinski, A. A., Couture, S., Balasubramanian, S., Aicher, S. A., et al. (2021). Sex differences in the expression of the endocannabinoid system within V1M cortex and PAG of Sprague Dawley rats. *Biol. Sex Differ.* 12:60. doi: 10.1186/s13293-021-00402-2
- Li, Y., and Kim, J. (2015). Neuronal expression of CB2 cannabinoid receptor mRNAs in the mouse hippocampus. *Neuroscience* 311, 253–267. doi: 10.1016/j.neuroscience.2015.10.041
- Li, Q., Marcu, D.-C., Palazzo, O., Turner, F., King, D., Spires-Jones, T. L., et al. (2020). High neural activity accelerates the decline of cognitive plasticity with age in *Caenorhabditis elegans*. *eLife* 9:e59711. doi: 10.7554/eLife.59711
- Liddel, S. A., Guttenplan, K. A., Clarke, L. E., Bennett, F. C., Bohlen, C. J., Schirmer, L., et al. (2017). Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 541, 481–487. doi: 10.1038/nature21029
- Lisdahl, K. M., Gilbert, E. R., Wright, N. E., and Shollenbarger, S. (2013). Dare to delay? The impacts of adolescent alcohol and marijuana use onset on cognition, brain structure, and function. *Front. Psych.* 4:53. doi: 10.3389/fpsy.2013.00053
- Liu, J., Dietz, K., DeLoyht, J. M., Pedre, X., Kelkar, D., Kaur, J., et al. (2012). Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nat. Neurosci.* 15, 1621–1623. doi: 10.1038/nn.3263
- Liu, H., Yang, Y., Xia, Y., Zhu, W., Leak, R. K., Wei, Z., et al. (2017). Aging of cerebral white matter. *Ageing Res. Rev.* 34, 64–76. doi: 10.1016/j.arr.2016.11.006
- Longoria, V., Parcel, H., Toma, B., Minhas, A., and Zeine, R. (2022). Neurological benefits, clinical challenges, and Neuropathologic promise of medical marijuana: a systematic review of cannabinoid effects in multiple sclerosis and experimental models of demyelination. *Biomedicine* 10:539. doi: 10.3390/biomedicines10030539
- Lou, Z.-Y., Cheng, J., Wang, X.-R., Zhao, Y.-F., Gan, J., Zhou, G.-Y., et al. (2018). The inhibition of CB1 receptor accelerates the onset and development of EAE possibly by regulating microglia/macrophages polarization. *Journal of Neuroimmunology* 317, 37–44. doi: 10.1016/j.jneuroim.2018.02.001
- Lu, H.-C., and Mackie, K. (2016). An introduction to the endogenous cannabinoid system. *Biol. Psychiatry* 79, 516–525. doi: 10.1016/j.biopsych.2015.07.028
- Lou, Z.-Y., Yu, W.-B., Chen, J., Li, L., Jiang, L.-S., Xiao, B.-G., et al. (2016). Neuroprotective Effect Is Driven Through the Upregulation of CB1 Receptor in Experimental Autoimmune Encephalomyelitis. *J Mol Neurosci* 58, 193–200. doi: 10.1007/s12031-015-0656-9
- Luan, W., Qi, X., Liang, F., Zhang, X., Jin, Z., Shi, L., et al. (2021). Microglia impede oligodendrocyte generation in aged brain. *JIR* 14, 6813–6831. doi: 10.2147/JIR.S338242
- Lundgaard, I., Luzhynskaya, A., Stockley, J. H., Wang, Z., Evans, K. A., Swire, M., et al. (2013). Neuregulin and BDNF induce a switch to NMDA receptor-dependent myelination by oligodendrocytes. *PLoS Biol.* 11:e1001743. doi: 10.1371/journal.pbio.1001743
- Mackie, K., and Hille, B. (1992). Cannabinoids inhibit N-type calcium channels in neuroblastoma-glioma cells. *Proc. Natl. Acad. Sci. U. S. A.* 89, 3825–3829. doi: 10.1073/pnas.89.9.3825
- Madry, C., Kyrargyri, V., Arancibia-Cárcamo, I. L., Jolivet, R., Kohsaka, S., Bryan, R. M., et al. (2018). Microglial ramification, surveillance, and interleukin-1 β release are regulated by the two-pore domain K⁺ channel THIK-1. *Neuron* 97, 299–312.e6. doi: 10.1016/j.neuron.2017.12.002
- Magnotti, L. M., Goodenough, D. A., and Paul, D. L. (2011). Deletion of oligodendrocyte Cx32 and astrocyte Cx43 causes white matter vacuolation, astrocyte loss and early mortality. *Glia* 59, 1064–1074. doi: 10.1002/glia.21179
- Mahmoud, S., Gharagozloo, M., Simard, C., and Gris, D. (2019). Astrocytes maintain glutamate homeostasis in the CNS by controlling the balance between glutamate uptake and release. *Cells* 8:184. doi: 10.3390/cells8020184
- Manza, P., Yuan, K., Shokri-Kojori, E., Tomasi, D., and Volkow, N. D. (2020). Brain structural changes in cannabis dependence: association with MAGL. *Mol. Psychiatry* 25, 3256–3266. doi: 10.1038/s41380-019-0577-z
- Maresz, K., Carrier, E. J., Ponomarev, E. D., Hillard, C. J., and Dittel, B. N. (2005). Modulation of the cannabinoid CB2 receptor in microglial cells in response to inflammatory stimuli. *J. Neurochem.* 95, 437–445. doi: 10.1111/j.1471-4159.2005.03380.x
- Markoullis, K., Sargiannidou, I., Schiza, N., Hadjisavvas, A., Roncaroli, F., Reynolds, R., et al. (2012). Gap junction pathology in multiple sclerosis lesions and normal-appearing white matter. *Acta Neuropathol.* 123, 873–886. doi: 10.1007/s00401-012-0978-4
- Marschallinger, J., Iram, T., Zardeneta, M., Lee, S. E., Lehallier, B., Haney, M. S., et al. (2020). Lipid-droplet-accumulating microglia represent a dysfunctional and proinflammatory state in the aging brain. *Nat. Neurosci.* 23, 194–208. doi: 10.1038/s41593-019-0566-1
- Marsicano, G., and Lutz, B. (1999). Expression of the cannabinoid receptor CB1 in distinct neuronal subpopulations in the adult mouse forebrain: CB1 expression in murine forebrain. *Eur. J. Neurosci.* 11, 4213–4225. doi: 10.1046/j.1460-9568.1999.00847.x
- Martinez Ramirez, C. E., Ruiz-Pérez, G., Stollenwerk, T. M., Behlke, C., Doherty, A., and Hillard, C. J. (2023). Endocannabinoid signaling in the central nervous system. *Glia* 71, 5–35. doi: 10.1002/glia.24280
- Masaki, K. (2015). Early disruption of glial communication via connexin gap junction in multiple sclerosis, Baló's disease and neuromyelitis optica: Connexin pathology in MS, BD and NMO. *Neuropathology* 35, 469–480. doi: 10.1111/neup.12211
- Matejuk, A., and Ransohoff, R. M. (2020). Crosstalk between astrocytes and microglia: an overview. *Front. Immunol.* 11:1416. doi: 10.3389/fimmu.2020.01416
- Mato, S., and Pazos, A. (2004). Influence of age, postmortem delay and freezing storage period on cannabinoid receptor density and functionality in human brain. *Neuropharmacology* 46, 716–726. doi: 10.1016/j.neuropharm.2003.11.004
- Mato, S., Victoria Sánchez-Gómez, M., and Matute, C. (2010). Cannabidiol induces intracellular calcium elevation and cytotoxicity in oligodendrocytes. *Glia* 58, 1739–1747. doi: 10.1002/glia.21044
- Mattson, M. P., and Magnus, T. (2006). Ageing and neuronal vulnerability. *Nat. Rev. Neurosci.* 7, 278–294. doi: 10.1038/nrn1886
- Matute, C., Alberdi, E., Domercq, M., Sánchez-Gómez, M.-V., Pérez-Samartin, A., Rodríguez-Antigüedad, A., et al. (2007). Excitotoxic damage to white matter. *J. Anat.* 210, 693–702. doi: 10.1111/j.1469-7580.2007.00733.x
- McDonald, J. W., Althomsons, S. P., Hyrc, K. L., Choi, D. W., and Goldberg, M. P. (1998). Oligodendrocytes from forebrain are highly vulnerable to AMPA/kainate receptor-mediated excitotoxicity. *Nat. Med.* 4, 291–297. doi: 10.1038/nm0398-291
- McNamara, N. B., Munro, D. A. D., Bestard-Cuche, N., Uyeda, A., Bogie, J. F. J., Hoffmann, A., et al. (2023). Microglia regulate central nervous system myelin growth and integrity. *Nature* 613, 120–129. doi: 10.1038/s41586-022-05534-y
- Mecha, M., Feliú, A., Carrillo-Salinas, F. J., Rueda-Zubiaurre, A., Ortega-Gutiérrez, S., de Sola, R. G., et al. (2015). Endocannabinoids drive the acquisition of an alternative phenotype in microglia. *Brain, Behavior, and Immunity* 49, 293–245. doi: 10.1016/j.bbi.2015.06.002
- Mecha, M., Feliú, A., Iñigo, P. M., Mestre, L., Carrillo-Salinas, F. J., and Guaza, C. (2013). Cannabidiol provides long-lasting protection against the deleterious effects of inflammation in a viral model of multiple sclerosis: a role for A2A receptors. *Neurobiol. Dis.* 59, 141–150. doi: 10.1016/j.nbd.2013.06.016
- Mecha, M., Torrao, A. S., Mestre, L., Carrillo-Salinas, F. J., Mechoulam, R., and Guaza, C. (2012). Cannabidiol protects oligodendrocyte progenitor cells from inflammation-induced apoptosis by attenuating endoplasmic reticulum stress. *Cell Death Dis.* 3:e331. doi: 10.1038/cddis.2012.71
- Mecha, M., Yanguas-Casás, N., Feliú, A., Mestre, L., Carrillo-Salinas, F., Azcoitia, I., et al. (2019). The endocannabinoid 2-AG enhances spontaneous remyelination by targeting microglia. *Brain Behav. Immun.* 77, 110–126. doi: 10.1016/j.bbi.2018.12.013
- Meier, J., Tewarie, P., Hillebrand, A., Douw, L., van Dijk, B. W., Stufflebeam, S. M., et al. (2016). A mapping between structural and functional brain networks. *Brain Connect.* 6, 298–311. doi: 10.1089/brain.2015.0408
- Même, W., Calvo, C., Froger, N., Ezan, P., Amigou, E., Koulakoff, A., et al. (2006). Proinflammatory cytokines released from microglia inhibit gap junctions in astrocytes: potentiation by β -amyloid. *FASEB J.* 20, 494–496. doi: 10.1096/fj.05.4297fj
- Meschkat, M., Steyer, A. M., Weil, M.-T., Kusch, K., Jahn, O., Piepkorn, L., et al. (2022). White matter integrity in mice requires continuous myelin synthesis at the inner tongue. *Nat. Commun.* 13:1163. doi: 10.1038/s41467-022-28720-y
- Meyer, N., Richter, N., Fan, Z., Siemonsmeier, G., Pivneva, T., Jordan, P., et al. (2018). Oligodendrocytes in the mouse corpus callosum maintain axonal function by delivery of glucose. *Cell Rep.* 22, 2383–2394. doi: 10.1016/j.celrep.2018.02.022
- Mitew, S., Gobius, I., Fenlon, L. R., McDougall, S. J., Hawkes, D., Xing, Y. L., et al. (2018). Pharmacogenetic stimulation of neuronal activity increases myelination in an axon-specific manner. *Nat. Commun.* 9:306. doi: 10.1038/s41467-017-02719-2
- Mittelbronn, M., Dietz, K., Schluessener, H. J., and Meyermann, R. (2001). Local distribution of microglia in the normal adult human central nervous system differs by up to one order of magnitude. *Acta Neuropathol.* 101, 249–255. doi: 10.1007/s004010000284
- Molina-Holgado, E., Esteban, P. F., Arevalo-Martin, Á., Moreno-Luna, R., Molina-Holgado, F., and Garcia-Ovejero, D. (2022). Endocannabinoid signaling in oligodendroglia. *Glia* 71, 91–102. doi: 10.1002/glia.24180
- Molina-Holgado, E., Vela, J. M., Arévalo-Martín, A., Almazán, G., Molina-Holgado, F., Borrell, J., et al. (2002). Cannabinoids promote oligodendrocyte progenitor survival:

- involvement of cannabinoid receptors and Phosphatidylinositol-3 kinase/Akt signaling. *J. Neurosci.* 22, 9742–9753. doi: 10.1523/JNEUROSCI.22-22-09742.2002
- Mount, C. W., and Monje, M. (2017). Wrapped to adapt: experience-dependent myelination. *Neuron* 95, 743–756. doi: 10.1016/j.neuron.2017.07.009
- Moura, D. M. S., Brennan, E. J., Brock, R., and Cocos, L. A. (2022). Neuron to oligodendrocyte precursor cell synapses: protagonists in oligodendrocyte development and myelination, and targets for therapeutics. *Front. Neurosci.* 15:779125. doi: 10.3389/fnins.2021.779125
- Narayanan, S. P., Flores, A. I., Wang, F., and Macklin, W. B. (2009). Akt signals through the mammalian target of Rapamycin pathway to regulate CNS myelination. *J. Neurosci.* 29, 6860–6870. doi: 10.1523/JNEUROSCI.0232-09.2009
- Navarrete, M., and Araque, A. (2010). Endocannabinoids potentiate synaptic transmission through stimulation of astrocytes. *Neuron* 68, 113–126. doi: 10.1016/j.neuron.2010.08.043
- Navarrete, M., Díez, A., and Araque, A. (2014). Astrocytes in endocannabinoid signalling. *Phil. Trans. R. Soc. B* 369:20130599. doi: 10.1098/rstb.2013.0599
- Neumann, B., Baror, R., Zhao, C., Segel, M., Dietmann, S., Rawji, K. S., et al. (2019). Metformin restores CNS Remyelination capacity by rejuvenating aged stem cells. *Cell Stem Cell* 25, 473–485.e8. doi: 10.1016/j.stem.2019.08.015
- Neumann, H., Kotter, M. R., and Franklin, R. J. M. (2008). Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain* 132, 288–295. doi: 10.1093/brain/awn109
- Nichols, N. R., Day, J. R., Laping, N. J., Johnson, S. A., and Finch, C. E. (1993). GFAP mRNA increases with age in rat and human brain. *Neurobiol. Aging* 14, 421–429. doi: 10.1016/0197-4580(93)90100-P
- Njoo, C., Agarwal, N., Lutz, B., and Kuner, R. (2015). The cannabinoid receptor CB1 interacts with the WAVE1 complex and plays a role in actin dynamics and structural plasticity in neurons. *PLoS Biol.* 13:e1002286. doi: 10.1371/journal.pbio.1002286
- Norden, D. M., Fenn, A. M., Dugan, A., and Godbout, J. P. (2014). TGF β produced by IL-10 redirected astrocytes attenuates microglial activation: IL-10 redirects immune activated astrocytes. *Glia* 62, 881–895. doi: 10.1002/glia.22647
- Norden, D. M., and Godbout, J. P. (2013). Review: microglia of the aged brain: primed to be activated and resistant to regulation: increased microglial reactivity with age. *Neuropathol. Appl. Neurobiol.* 39, 19–34. doi: 10.1111/j.1365-2990.2012.01306.x
- Norden, D. M., Trojanowski, P. J., Walker, F. R., and Godbout, J. P. (2016). Insensitivity of astrocytes to interleukin 10 signaling following peripheral immune challenge results in prolonged microglial activation in the aged brain. *Neurobiol. Aging* 44, 22–41. doi: 10.1016/j.neurobiolaging.2016.04.014
- O'Neil, S. M., Hans, E. E., Jiang, S., Wangler, L. M., and Godbout, J. P. (2022). Astrocyte immunosenescence and deficits in interleukin 10 signaling in the aged brain disrupt the regulation of microglia following innate immune activation. *Glia* 70, 913–934. doi: 10.1002/glia.24147
- O'Sullivan, M., Jones, D. K., Summers, P. E., Morris, R. G., Williams, S. C. R., and Markus, H. S. (2001). Evidence for cortical “disconnection” as a mechanism of age-related cognitive decline. *Neurology* 57, 632–638. doi: 10.1212/WNL.57.4.632
- Olney, J. W. (1971). Glutamate-induced neuronal necrosis in the infant mouse hypothalamus: an electron microscopic study. *J. Neuropathol. Exp. Neurol.* 30, 75–90. doi: 10.1097/00005072-197101000-00008
- Orellana, J. A., Froger, N., Ezan, P., Jiang, J. X., Bennett, M. V. L., Naus, C. C., et al. (2011). ATP and glutamate released via astroglial connexin 43 hemichannels mediate neuronal death through activation of pannexin 1 hemichannels: Astroglial hemichannels induce neuronal death. *J. Neurochem.* 118, 826–840. doi: 10.1111/j.1471-4159.2011.07210.x
- Orr, J. M., Paschall, C. J., and Banich, M. T. (2016). Recreational marijuana use impacts white matter integrity and subcortical (but not cortical) morphometry. *NeuroImage: Clinical* 12, 47–56. doi: 10.1016/j.nicl.2016.06.006
- Orthmann-Murphy, J. L., Abrams, C. K., and Scherer, S. S. (2008). Gap junctions couple astrocytes and oligodendrocytes. *J. Mol. Neurosci.* 35, 101–116. doi: 10.1007/s12031-007-9027-5
- Paes-Colli, Y., Aguiar, A. F. L., Isaac, A. R., Ferreira, B. K., Campos, R. M. P., Trindade, P. M. P., et al. (2022). Phytocannabinoids and cannabis-based products as alternative pharmacotherapy in neurodegenerative diseases: from hypothesis to clinical practice. *Front. Cell. Neurosci.* 16:917164. doi: 10.3389/fncel.2022.917164
- Pakkenberg, B., and Gundersen, H. J. (1997). Neocortical neuron number in humans: effect of sex and age. *J. Comp. Neurol.* 384, 312–320. doi: 10.1002/(SICI)1096-9861(19970728)384:2<312::AID-CNE10>3.0.CO;2-K
- Palmer, A. L., and Ousman, S. S. (2018). Astrocytes and aging. *Front. Aging Neurosci.* 10:337. doi: 10.3389/fnagi.2018.00337
- Pan, S., Mayoral, S. R., Choi, H. S., Chan, J. R., and Kheirbek, M. A. (2020). Preservation of a remote fear memory requires new myelin formation. *Nat. Neurosci.* 23, 487–499. doi: 10.1038/s41593-019-0582-1
- Pang, Y., Cai, Z., and Rhodes, P. G. (2003). Disturbance of oligodendrocyte development, hypomyelination and white matter injury in the neonatal rat brain after intracerebral injection of lipopolysaccharide. *Dev. Brain Res.* 140, 205–214. doi: 10.1016/S0165-3806(02)00606-5
- Pang, Y., Zheng, B., Fan, L.-W., Rhodes, P. G., and Cai, Z. (2007). IGF-1 protects oligodendrocyte progenitors against TNF α -induced damage by activation of PI3K/Akt and interruption of the mitochondrial apoptotic pathway. *Glia* 55, 1099–1107. doi: 10.1002/glia.20530
- Paolicelli, R. C., Sierra, A., Stevens, B., Tremblay, M.-E., Aguzzi, A., Ajami, B., et al. (2022). Microglia states and nomenclature: a field at its crossroads. *Neuron* 110, 3458–3483. doi: 10.1016/j.neuron.2022.10.020
- Papaneophytou, C., Georgiou, E., and Kleopa, K. A. (2019). The role of oligodendrocyte gap junctions in neuroinflammation. *Channels* 13, 247–263. doi: 10.1080/19336950.2019.1631107
- Papuć, E., and Rejdak, K. (2020). The role of myelin damage in Alzheimer's disease pathology. *Arch. Med. Sci.* 16, 345–351. doi: 10.5114/aoms.2018.76863
- Pascual, A. C., Gaveglione, V. L., Giusto, N. M., and Pasquaré, S. J. (2013). Aging modifies the enzymatic activities involved in 2-arachidonoylglycerol metabolism. *Biofactors* 39, 209–220. doi: 10.1002/biof.1055
- Pascual, A. C., Gaveglione, V. L., Giusto, N. M., and Pasquaré, S. J. (2014). Cannabinoid receptor-dependent metabolism of 2-arachidonoylglycerol during aging. *Exp. Gerontol.* 55, 134–142. doi: 10.1016/j.exger.2014.04.008
- Pedraza, C. E., Taylor, C., Pereira, A., Seng, M., Tham, C.-S., Izrael, M., et al. (2014). Induction of Oligodendrocyte Differentiation and In Vitro Myelination by Inhibition of Rho-Associated Kinase. *ASN Neuro.* 6:175909141453813. doi: 10.1177/1759091414538134
- Peferoen, L., Kipp, M., van der Valk, P., van Noort, J. M., and Amor, S. (2014). Oligodendrocyte-microglia cross-talk in the central nervous system. *Immunology* 141, 302–313. doi: 10.1111/imm.12163
- Peng, B., Xu, C., Wang, S., Zhang, Y., and Li, W. (2022). The role of Connexin Hemichannels in inflammatory diseases. *Biology* 11:237. doi: 10.3390/biology11020237
- Pertwee, R. G. (2008). The diverse CB $_1$ and CB $_2$ receptor pharmacology of three plant cannabinoids: Δ^9 -tetrahydrocannabinol, cannabidiol and Δ^9 -tetrahydrocannabivarin: Δ^9 -THC, CBD and Δ^9 -THCV. *Br. J. Pharmacol.* 153, 199–215. doi: 10.1038/sj.bjp.0707442
- Peters, A. (2009). The effects of normal aging on myelinated nerve fibers in monkey central nervous system. *Front. Neuroanat.* 3:11. doi: 10.3389/neuro.05.011.2009
- Phillips, T., Mironova, Y. A., Jouroukhin, Y., Chew, J., Vidensky, S., Farah, M. H., et al. (2021). MCT1 deletion in oligodendrocyte lineage cells causes late-onset Hypomyelination and axonal degeneration. *Cell Rep.* 34:108610. doi: 10.1016/j.celrep.2020.108610
- Phillips, K. A., Watson, C. M., Bearman, A., Knippenberg, A. R., Adams, J., Ross, C., et al. (2019). Age-related changes in myelin of axons of the corpus callosum and cognitive decline in common marmosets. *Am. J. Primatol.* 81:e22949. doi: 10.1002/ajp.22949
- Pierpaoli, C., and Basser, P. J. (1996). Toward a quantitative assessment of diffusion anisotropy. *Magn. Reson. Med.* 36, 893–906. doi: 10.1002/mrm.1910360612
- Piyanova, A., Lomazzo, E., Bindila, L., Lerner, R., Albayram, O., Ruhl, T., et al. (2015). Age-related changes in the endocannabinoid system in the mouse hippocampus. *Mech. Ageing Dev.* 150, 55–64. doi: 10.1016/j.mad.2015.08.005
- Poliani, P. L., Wang, Y., Fontana, E., Robinette, M. L., Yamanishi, Y., Gilfillan, S., et al. (2015). TREM2 sustains microglial expansion during aging and response to demyelination. *J. Clin. Invest.* 125, 2161–2170. doi: 10.1172/JCI77983
- Popov, A., Brazhe, A., Denisov, P., Sutyagina, O., Li, L., Lazareva, N., et al. (2021). Astrocyte dystrophy in ageing brain parallels impaired synaptic plasticity. *Aging Cell* 20:e13334. doi: 10.1111/acel.13334
- Porcher, L., Bruckmeier, S., Burbano, S. D., Finnell, J. E., Gorny, N., Klett, J., et al. (2021). Aging triggers an upregulation of a multitude of cytokines in the male and especially the female rodent hippocampus but more discrete changes in other brain regions. *J. Neuroinflammation* 18:219. doi: 10.1186/s12974-021-02252-6
- Raasch, J., Zeller, N., van Loo, G., Merkler, D., Mildner, A., Erny, D., et al. (2011). I κ B kinase 2 determines oligodendrocyte loss by non-cell-autonomous activation of NF- κ B in the central nervous system. *Brain* 134, 1184–1198. doi: 10.1093/brain/awq359
- Rao, V. T. S., Khan, D., Cui, Q.-L., Fuh, S.-C., Hossain, S., Almazan, G., et al. (2017). Distinct age and differentiation-state dependent metabolic profiles of oligodendrocytes under optimal and stress conditions. *PLoS One* 12:e0182372. doi: 10.1371/journal.pone.0182372
- Rea, I. M., Gibson, D. S., McGilligan, V., McNerlan, S. E., Alexander, H. D., and Ross, O. A. (2018). Age and age-related diseases: role of inflammation triggers and cytokines. *Front. Immunol.* 9:586. doi: 10.3389/fimmu.2018.00586
- Retamal, M. A., Froger, N., Palacios-Prado, N., Ezan, P., Saez, P. J., Saez, J. C., et al. (2007). Cx43 Hemichannels and gap junction channels in astrocytes are regulated oppositely by Proinflammatory cytokines released from activated microglia. *J. Neurosci.* 27, 13781–13792. doi: 10.1523/JNEUROSCI.2042-07.2007
- Reusch, N., Ravichandran, K. A., Olabiyi, B. F., Komorowska-Müller, J. A., Hansen, J. N., Ulas, T., et al. (2022). Cannabinoid receptor 2 is necessary to induce toll-like receptor-mediated microglial activation. *Glia* 70, 71–88. doi: 10.1002/glia.24089
- Ritzel, R. M., Crapser, J., Patel, A. R., Verma, R., Grenier, J. M., Chauhan, A., et al. (2016). Age-associated resident memory CD8 T cells in the central nervous system are

- primed To potentiate inflammation after ischemic brain injury. *J. I.* 196, 3318–3330. doi: 10.4049/jimmunol.1502021
- Ritzel, R. M., Patel, A. R., Pan, S., Crapser, J., Hammond, M., Jellison, E., et al. (2015). Age- and location-related changes in microglial function. *Neurobiol. Aging* 36, 2153–2163. doi: 10.1016/j.neurobiolaging.2015.02.016
- Rivera, A., Butt, A., and Azim, K. (2022). Resolving the age-related decline in central nervous system myelin turnover and drug discovery for oligodendroglial rejuvenation. *Neural Regen. Res.* 17:2677. doi: 10.4103/1673-5374.338995
- Rivera, A. D., Pieropan, F., Chacon-De-La-Rocha, I., Lecca, D., Abbracchio, M. P., Azim, K., et al. (2021). Functional genomic analyses highlight a shift in *Gpr17*-regulated cellular processes in oligodendrocyte progenitor cells and underlying myelin dysregulation in the aged mouse cerebrum. *Aging Cell* 20:e13335. doi: 10.1111/acel.13335
- Rizzo, M. D., Crawford, R. B., Bach, A., Sermet, S., Amalfitano, A., and Kaminski, N. E. (2019). Δ^9 -tetrahydrocannabinol suppresses monocyte-mediated astrocyte production of monocyte Chemoattractant protein 1 and Interleukin-6 in a toll-like receptor 7-stimulated human Coculture. *J. Pharmacol. Exp. Ther.* 371, 191–201. doi: 10.1124/jpet.119.260661
- Ronzano, R., Roux, T., Thetiot, M., Aigrot, M. S., Richard, L., Lejeune, F. X., et al. (2021). Microglia-neuron interaction at nodes of Ranvier depends on neuronal activity through potassium release and contributes to remyelination. *Nat. Commun.* 12:5219. doi: 10.1038/s41467-021-25486-7
- Rossi, S., Bernardi, G., and Centonze, D. (2010). The endocannabinoid system in the inflammatory and neurodegenerative processes of multiple sclerosis and of amyotrophic lateral sclerosis. *Exp. Neurol.* 224, 92–102. doi: 10.1016/j.expneurol.2010.03.030
- Rozenfeld, R., and Devi, L. A. (2008). Regulation of CB₁ cannabinoid receptor trafficking by the adaptor protein AP-3. *FASEB J.* 22, 2311–2322. doi: 10.1096/fj.07-102731
- Safaiyan, S., Besson-Girard, S., Kaya, T., Cantuti-Castelvetri, L., Liu, L., Ji, H., et al. (2021). White matter aging drives microglial diversity. *Neuron* 109, 1100–1117.e10. doi: 10.1016/j.neuron.2021.01.027
- Safaiyan, S., Kannaiyan, N., Snaidero, N., Brioschi, S., Biber, K., Yona, S., et al. (2016). Age-related myelin degradation burdens the clearance function of microglia during aging. *Nat. Neurosci.* 19, 995–998. doi: 10.1038/nn.4325
- Saher, G., and Stumpf, S. K. (2015). Cholesterol in myelin biogenesis and hypomyelinating disorders. *Biochim. Biophys. Acta* 1851, 1083–1094. doi: 10.1016/j.bbali.2015.02.010
- Salas-Wright, C. P., Vaughn, M. G., Cummings-Vaughn, L. A., Holzer, K. J., Nelson, E. J., AbiNader, M., et al. (2017). Trends and correlates of marijuana use among late middle-aged and older adults in the United States, 2002–2014. *Drug Alcohol Depend.* 171, 97–106. doi: 10.1016/j.drugalcdep.2016.11.031
- Salminen, A., Ojala, J., Kaarniranta, K., Haapasalo, A., Hiltunen, M., and Soyninen, H. (2011). Astrocytes in the aging brain express characteristics of senescence-associated secretory phenotype: astrocyte senescence in aging brain. *Eur. J. Neurosci.* 34, 3–11. doi: 10.1111/j.1460-9568.2011.07738.x
- Sams, E. C. (2021). Oligodendrocytes in the aging brain. *Neuronal Signaling* 5:NS20210008. doi: 10.1042/NS20210008
- Sánchez-de la Torre, A., Aguado, T., Hueriga-Gómez, A., Santamaría, S., Gentile, A., Chara, J. C., et al. (2022). Cannabinoid CB1 receptor gene inactivation in oligodendrocyte precursors disrupts oligodendrogenesis and myelination in mice. *Cell Death Dis.* 13:585. doi: 10.1038/s41419-022-05032-z
- Sankowski, R., Böttcher, C., Masuda, T., Geirsdottir, L., Sagar, S., Sindram, E., et al. (2019). Mapping microglia states in the human brain through the integration of high-dimensional techniques. *Nat. Neurosci.* 22, 2098–2110. doi: 10.1038/s41593-019-0532-y
- Santos, E. N., and Fields, R. D. (2021). Regulation of myelination by microglia. *Sci. Adv.* 7:eabk1131. doi: 10.1126/sciadv.abk1131
- Saraste, M., Bezukladova, S., Matilainen, M., Sucksdorff, M., Kuhle, J., Leppert, D., et al. (2021). Increased serum glial fibrillary acidic protein associates with microstructural white matter damage in multiple sclerosis. *Mult. Scler. Relat. Disord.* 50:102810. doi: 10.1016/j.msard.2021.102810
- Sarne, Y. (2019). Beneficial and deleterious effects of cannabinoids in the brain: the case of ultra-low dose THC. *Am. J. Drug Alcohol Abuse* 45, 551–562. doi: 10.1080/00952990.2019.1578366
- Sarne, Y., Toledano, R., Rachmany, L., Sasson, E., and Doron, R. (2018). Reversal of age-related cognitive impairments in mice by an extremely low dose of tetrahydrocannabinol. *Neurobiol. Aging* 61, 177–186. doi: 10.1016/j.neurobiolaging.2017.09.025
- Satarker, S., Bojja, S. L., Gurrum, P. C., Mudgal, J., Arora, D., and Nampoothiri, M. (2022). Astrocytic Glutamatergic transmission and its implications in neurodegenerative disorders. *Cells* 11:1139. doi: 10.3390/cells11071139
- Savchenko, V. L., McKanna, J. A., Nikonenko, I. R., and Skibo, G. G. (2000). Microglia and astrocytes in the adult rat brain: comparative immunocytochemical analysis demonstrates the efficacy of lipocortin 1 immunoreactivity. *Neuroscience* 96, 195–203. doi: 10.1016/S0306-4522(99)00538-2
- Scholz, J., Klein, M. C., Behrens, T. E. J., and Johansen-Berg, H. (2009). Training induces changes in white-matter architecture. *Nat. Neurosci.* 12, 1370–1371. doi: 10.1038/nn.2412
- Schulz, K., Kroner, A., and David, S. (2012). Iron efflux from astrocytes plays a role in Remyelination. *J. Neurosci.* 32, 4841–4847. doi: 10.1523/JNEUROSCI.5328-11.2012
- Scipioni, L., Ciaramellano, F., Carnicelli, V., Leuti, A., Lizzi, A. R., De Dominicis, N., et al. (2022). Microglial Endocannabinoid Signalling in AD. *Cells* 11:1237. doi: 10.3390/cells11071237
- Segel, M., Neumann, B., Hill, M. F. E., Weber, I. P., Viscomi, C., Zhao, C., et al. (2019). Niche stiffness underlies the ageing of central nervous system progenitor cells. *Nature* 573, 130–134. doi: 10.1038/s41586-019-1484-9
- Sheng, W. S., Hu, S., Min, X., Cabral, G. A., Lokensgard, J. R., and Peterson, P. K. (2005). Synthetic cannabinoid WIN55,212-2 inhibits generation of inflammatory mediators by IL-1 β -stimulated human astrocytes. *Glia* 49, 211–219. doi: 10.1002/glia.20108
- Shepherd, M. N., Pomicter, A. D., Velasco, C. S., Henderson, S. C., and Dupree, J. L. (2012). Paranodal reorganization results in the depletion of transverse bands in the aged central nervous system. *Neurobiol. Aging* 33, 203.e13–203.e24. doi: 10.1016/j.neurobiolaging.2010.08.001
- Shivers, S. C., Newton, C., Friedman, H., and Klein, T. W. (1994). Δ^9 -tetrahydrocannabinol (THC) modulates IL-1 bioactivity in human monocyte/macrophage cell lines. *Life Sci.* 54, 1281–1289. doi: 10.1016/0024-3205(94)00856-6
- Shollenbarger, S. G., Price, J., Wieser, J., and Lisdahl, K. (2015). Poorer frontolimbic white matter integrity is associated with chronic cannabis use, FAAH genotype, and increased depressive and apathy symptoms in adolescents and young adults. *NeuroImage: Clinical* 8, 117–125. doi: 10.1016/j.nicl.2015.03.024
- Sierra, A., Gottfried-Blackmore, A. C., McEwen, B. S., and Bulloch, K. (2007). Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia* 55, 412–424. doi: 10.1002/glia.20468
- Sikora, E., Bielak-Zmijewska, A., Dudkowska, M., Krzystyniak, A., Mosieniak, G., Wesierska, M., et al. (2021). Cellular senescence in brain aging. *Front. Aging Neurosci.* 13:646924. doi: 10.3389/fnagi.2021.646924
- Sim, F. J., Zhao, C., Penderis, J., and Franklin, R. J. M. (2002). The age-related decrease in CNS Remyelination efficiency is attributable to an impairment of both oligodendrocyte progenitor recruitment and differentiation. *J. Neurosci.* 22, 2451–2459. doi: 10.1523/JNEUROSCI.22-07-02451.2002
- Šimončičová, E., Gonçalves de Andrade, E., Vecchiarelli, H. A., Awogbindin, I. O., Delage, C. I., and Tremblay, M.-È. (2022). Present and future of microglial pharmacology. *Trends Pharmacol. Sci.* 43, 669–685. doi: 10.1016/j.tips.2021.11.006
- Sofroniew, M. V., and Vinters, H. V. (2010). Astrocytes: biology and pathology. *Acta Neuropathol.* 119, 7–35. doi: 10.1007/s00401-009-0619-8
- Solt, L. A., and May, M. J. (2008). The I κ B kinase complex: master regulator of NF- κ B signaling. *Immunol. Res.* 42, 3–18. doi: 10.1007/s12026-008-8025-1
- Soreq, L., Rose, J., Soreq, E., Hardy, J., Trabzuni, D., Cookson, M. R., et al. (2017). Major shifts in glial regional identity are a transcriptional Hallmark of human brain aging. *Cell Rep.* 18, 557–570. doi: 10.1016/j.celrep.2016.12.011
- Sowell, E. R., Peterson, B. S., Thompson, P. M., Welcome, S. E., Henkenius, A. L., and Toga, A. W. (2003). Mapping cortical change across the human life span. *Nat. Neurosci.* 6, 309–315. doi: 10.1038/nn1008
- Spaas, J., van Veggel, L., Schepers, M., Tiane, A., van Horsen, J., Wilson, D. M., et al. (2021). Oxidative stress and impaired oligodendrocyte precursor cell differentiation in neurological disorders. *Cell. Mol. Life Sci.* 78, 4615–4637. doi: 10.1007/s00118-021-03802-0
- Spitzer, S. O., Sitnikov, S., Kamen, Y., Evans, K. A., Kronenberg-Versteeg, D., Dietmann, S., et al. (2019). Oligodendrocyte progenitor cells become regionally diverse and heterogeneous with age. *Neuron* 101, 459–471.e5. doi: 10.1016/j.neuron.2018.12.020
- Stadelmann, C., Timmler, S., Barrantes-Freer, A., and Simons, M. (2019). Myelin in the central nervous system: structure, function, and pathology. *Physiol. Rev.* 99, 1381–1431. doi: 10.1152/physrev.00031.2018
- Stahon, K. E., Bastian, C., Griffith, S., Kidd, G. J., Brunet, S., and Balcen, S. (2016). Age-related changes in axonal and mitochondrial ultrastructure and function in white matter. *J. Neurosci.* 36, 9990–10001. doi: 10.1523/JNEUROSCI.1316-16.2016
- Stampanoni Bassi, M., Gentile, A., Iezzi, E., Zagaglia, S., Musella, A., Simonelli, I., et al. (2019). Transient receptor potential Vanilloid 1 modulates central inflammation in multiple sclerosis. *Front. Neurol.* 10:30. doi: 10.3389/fneur.2019.00030
- Stargardt, A., Swaab, D. F., and Bossers, K. (2015). The storm before the quiet: neuronal hyperactivity and A β in the presymptomatic stages of Alzheimer's disease. *Neurobiol. Aging* 36, 1–11. doi: 10.1016/j.neurobiolaging.2014.08.014
- Stehberg, J., Moraga-Amaro, R., Salazar, C., Becerra, A., Echeverría, C., Orellana, J. A., et al. (2012). Release of gliotransmitters through astroglial connexin 43 hemichannels is necessary for fear memory consolidation in the basolateral amygdala. *FASEB J.* 26, 3649–3657. doi: 10.1096/fj.11-198416
- Stella, N. (2010). Cannabinoid and cannabinoid-like receptors in microglia, astrocytes, and astrocytomas. *Glia* 58, 1017–1030. doi: 10.1002/glia.20983

- Stempel, A. V., Stumpf, A., Zhang, H.-Y., Özdoğan, T., Pannasch, U., Theis, A.-K., et al. (2016). Cannabinoid type 2 receptors mediate a cell type-specific plasticity in the hippocampus. *Neuron* 90, 795–809. doi: 10.1016/j.neuron.2016.03.034
- Stephan, J., Eitelmann, S., and Zhou, M. (2021). Approaches to study gap junctional coupling. *Front. Cell. Neurosci.* 15:640406. doi: 10.3389/fncel.2021.640406
- Swift, J., Ivanovska, I. L., Buxboim, A., Harada, T., Dingal, P. C. D. P., Pinter, J., et al. (2013). Nuclear Lamin-a scales with tissue stiffness and enhances matrix-directed differentiation. *Science* 341:1240104. doi: 10.1126/science.1240104
- Thériault, P., and Rivest, S. (2016). Microglia: senescence impairs clearance of myelin debris. *Curr. Biol.* 26, R772–R775. doi: 10.1016/j.cub.2016.06.066
- Thomas, A., Baillie, G. L., Phillips, A. M., Razdan, R. K., Ross, R. A., and Pertwee, R. G. (2007). Cannabidiol displays unexpectedly high potency as an antagonist of CB₁ and CB₂ receptor agonists *in vitro*: cannabinoid antagonism by cannabidiol. *Br. J. Pharmacol.* 150, 613–623. doi: 10.1038/sj.bjp.0707133
- Thomas, A. L., Lehn, M. A., Janssen, E. M., Hildeman, D. A., and Chougnet, C. A. (2022). Naturally-aged microglia exhibit phagocytic dysfunction accompanied by gene expression changes reflective of underlying neurodegenerative disease. *Sci. Rep.* 12:19471. doi: 10.1038/s41598-022-21920-y
- Tilstra, J. S., Clauson, C. L., Niedernhofer, L. J., and Robbins, P. D. (2011). NF- κ B in aging and disease. *Aging Dis.* 2, 449–465.
- Tognatta, R., Karl, M. T., Fyffe-Maricich, S. L., Popratiloff, A., Garrison, E. D., Schenck, J. K., et al. (2020). Astrocytes are required for oligodendrocyte survival and maintenance of myelin compaction and integrity. *Front. Cell. Neurosci.* 14:74. doi: 10.3389/fncel.2020.00074
- Tomas-Roig, J., Agbemenyah, H. Y., Celarain, N., Quintana, E., Ramió-Torrentà, L., and Havemann-Reinecke, U. (2020). Dose-dependent effect of cannabinoid WIN-55,212-2 on myelin repair following a demyelinating insult. *Sci. Rep.* 10:590. doi: 10.1038/s41598-019-57290-1
- Tress, O., Maglione, M., May, D., Pivneva, T., Richter, N., Seyfarth, J., et al. (2012). Panglial gap junctional communication is essential for maintenance of myelin in the CNS. *J. Neurosci.* 32, 7499–7518. doi: 10.1523/JNEUROSCI.0392-12.2012
- Tsai, H.-H., Frost, E., To, V., Robinson, S., French-Constant, C., Geertman, R., et al. (2002). The chemokine receptor CXCR2 controls positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. *Cells* 110, 373–383. doi: 10.1016/S0092-8674(02)00838-3
- Tse, K.-H., and Herrup, K. (2017). DNA damage in the oligodendrocyte lineage and its role in brain aging. *Mech. Ageing Dev.* 161, 37–50. doi: 10.1016/j.mad.2016.05.006
- Tsou, K., Mackie, K., Sañudo-Peña, M. C., and Walker, J. M. (1999). Cannabinoid CB1 receptors are localized primarily on cholecystokinin-containing GABAergic interneurons in the rat hippocampal formation. *Neuroscience* 93, 969–975. doi: 10.1016/S0306-4522(99)00086-X
- Turcotte, C., Blanchet, M.-R., Lavolette, M., and Flamand, N. (2016). The CB2 receptor and its role as a regulator of inflammation. *Cell. Mol. Life Sci.* 73, 4449–4470. doi: 10.1007/s00018-016-2300-4
- Tyler, W. A., Gangoli, N., Gokina, P., Kim, H. A., Covey, M., Levison, S. W., et al. (2009). Activation of the mammalian target of Rapamycin (mTOR) is essential for oligodendrocyte differentiation. *J. Neurosci.* 29, 6367–6378. doi: 10.1523/JNEUROSCI.0234-09.2009
- van den Berg, R., Haenen, G. R. M. M., van den Berg, H., and Bast, A. (2001). Transcription factor NF- κ B as a potential biomarker for oxidative stress. *Br. J. Nutr.* 86, S121–S127. doi: 10.1079/BJN2001340
- Van Laere, K., Goffin, K., Casteels, C., Dupont, P., Mortelmans, L., de Hoon, J., et al. (2008). Gender-dependent increases with healthy aging of the human cerebral cannabinoid-type 1 receptor binding using [18F]MK-9470 PET. *NeuroImage* 39, 1533–1541. doi: 10.1016/j.neuroimage.2007.10.053
- Van Ryzin, J. W., Marquardt, A. E., Argue, K. J., Vecchiarelli, H. A., Ashton, S. E., Arambula, S. E., et al. (2019). Microglial phagocytosis of newborn cells is induced by Endocannabinoids and sculpts sex differences in juvenile rat social play. *Neuron* 102, 435–449.e6. doi: 10.1016/j.neuron.2019.02.006
- Vandewouw, M. M., Hunt, B. A. E., Ziolkowski, J., and Taylor, M. J. (2021). The developing relations between networks of cortical myelin and neurophysiological connectivity. *NeuroImage* 237:118142. doi: 10.1016/j.neuroimage.2021.118142
- Vázquez, C., Tolón, R. M., Grande, M. T., Caraza, M., Moreno, M., Koester, E. C., et al. (2015a). Endocannabinoid regulation of amyloid-induced neuroinflammation. *Neurobiol. Aging* 36, 3008–3019. doi: 10.1016/j.neurobiolaging.2015.08.003
- Vázquez, C., Tolón, R. M., Pazos, M. R., Moreno, M., Koester, E. C., Cravatt, B. F., et al. (2015b). Endocannabinoids regulate the activity of astrocytic hemichannels and the microglial response against an injury: *in vivo* studies. *Neurobiol. Dis.* 79, 41–50. doi: 10.1016/j.nbd.2015.04.005
- Vecchiarelli, H. A., Aukema, R. J., Hume, C., Chiang, V., Morena, M., Keenan, C. M., et al. (2021). Genetic variants of fatty acid amide hydrolase modulate acute inflammatory responses to colitis in adult male mice. *Front. Cell. Neurosci.* 15:764706. doi: 10.3389/fncel.2021.764706
- Vecchiarelli, H. A., Morena, M., Lee, T. T. Y., Nastase, A. S., Aukema, R. J., Leitl, K. D., et al. (2022). Sex and stressor modality influence acute stress-induced dynamic changes in corticolimbic endocannabinoid levels in adult Sprague Dawley rats. *Neurobiol. Stress* 20:100470. doi: 10.1016/j.ynstr.2022.100470
- Verkhatsky, A., Lazareva, N., and Semyanov, A. (2022). Glial decline and loss of homeostatic support rather than inflammation defines cognitive aging. *Neural Regen. Res.* 17:565. doi: 10.4103/1673-5374.320979
- Vinke, E. J., de Groot, M., Venkatraghavan, V., Klein, S., Niessen, W. J., Ikram, M. A., et al. (2018). Trajectories of imaging markers in brain aging: the Rotterdam study. *Neurobiol. Aging* 71, 32–40. doi: 10.1016/j.neurobiolaging.2018.07.001
- Wake, H., Lee, P. R., and Fields, R. D. (2011). Control of local protein synthesis and initial events in myelination by action potentials. *Science* 333, 1647–1651. doi: 10.1126/science.1206998
- Wake, H., Ortiz, F. C., Woo, D. H., Lee, P. R., Angulo, M. C., and Fields, R. D. (2015). Nonsynaptic junctions on myelinating glia promote preferential myelination of electrically active axons. *Nat. Commun.* 6:7844. doi: 10.1038/ncomms8844
- Wang, H. (2003). Gender difference in glutathione metabolism during aging in mice. *Exp. Gerontol.* 38, 507–517. doi: 10.1016/S0531-5565(03)00036-6
- Wang, F., Ren, S.-Y., Chen, J.-F., Liu, K., Li, R.-X., Li, Z.-F., et al. (2020). Myelin degeneration and diminished myelin renewal contribute to age-related deficits in memory. *Nat. Neurosci.* 23, 481–486. doi: 10.1038/s41593-020-0588-8
- Wang, S., Wang, Y., and Zou, S. (2022). A glance at the molecules that regulate oligodendrocyte myelination. *CIMB* 44, 2194–2216. doi: 10.3390/cimb44050149
- Watanabe, K., Kayano, Y., Matsunaga, T., Yamamoto, I., and Yoshimura, H. (1996). Inhibition of Anandamide Amidase activity in mouse brain Microsomes by cannabinoids. *Biol. Pharm. Bull.* 19, 1109–1111. doi: 10.1248/bpb.19.1109
- Watson, K. K., Bryan, A. D., Thayer, R. E., Ellingson, J. M., Skrzyński, C. J., and Hutchison, K. E. (2022). Cannabis use and resting state functional connectivity in the aging brain. *Front. Aging Neurosci.* 14:804890. doi: 10.3389/fnagi.2022.804890
- Waxman, S. G. (1980). Determinants of conduction velocity in myelinated nerve fibers. *Muscle Nerve* 3, 141–150. doi: 10.1002/mus.880030207
- Westlye, L. T., Walhovd, K. B., Dale, A. M., Bjørnerud, A., Due-Tønnessen, P., Engvig, A., et al. (2010). Life-span changes of the human brain white matter: diffusion tensor imaging (DTI) and Volumetry. *Cereb. Cortex* 20, 2055–2068. doi: 10.1093/cercor/bhp280
- Whiting, P. E., Wolff, R. F., Deshpande, S., Di Nisio, M., Duffy, S., Hernandez, A. V., et al. (2015). Cannabinoids for medical use: a systematic review and meta-analysis. *JAMA* 313:2456. doi: 10.1001/jama.2015.6358
- Willis, C. M., Nicaise, A. M., Bongarzone, E. R., Givogri, M., Reiter, C. R., Heintz, O., et al. (2020). Astrocyte support for oligodendrocyte differentiation can be conveyed via extracellular vesicles but diminishes with age. *Sci. Rep.* 10:828. doi: 10.1038/s41598-020-57663-x
- Wong, R. (2018). A new strategic approach to successful aging and healthy aging. *Geriatrics* 3:86. doi: 10.3390/geriatrics3040086
- Wu, J., Chen, N., Liu, Y., Godlewski, G., Kaplan, H. J., Shrader, S. H., et al. (2021). Studies of involvement of G-protein coupled receptor-3 in cannabidiol effects on inflammatory responses of mouse primary astrocytes and microglia. *PLoS One* 16:e0251677. doi: 10.1371/journal.pone.0251677
- Wu, L. M. N., Williams, A., Delaney, A., Sherman, D. L., and Brophy, P. J. (2012). Increasing Internodal distance in myelinated nerves accelerates nerve conduction to a flat maximum. *Curr. Biol.* 22, 1957–1961. doi: 10.1016/j.cub.2012.08.025
- Yang, S., Du, Y., Zhao, X., Tang, Q., Su, W., Hu, Y., et al. (2022). Cannabidiol enhances microglial Beta-amyloid peptide phagocytosis and clearance via Vanilloid family type 2 channel activation. *IJMS* 23:5367. doi: 10.3390/ijms23105367
- Ye, Z.-C., Wyeth, M. S., Baltan-Tekkok, S., and Ransom, B. R. (2003). Functional hemichannels in astrocytes: a novel mechanism of glutamate release. *J. Neurosci.* 23, 3588–3596. doi: 10.1523/JNEUROSCI.23-09-03588.2003
- Yeung, M. S. Y., Zdunek, S., Bergmann, O., Bernard, S., Salehpour, M., Alkass, K., et al. (2014). Dynamics of oligodendrocyte generation and myelination in the human brain. *Cells* 159, 766–774. doi: 10.1016/j.cell.2014.10.011
- Yoo, H. B., DiMuzio, J., and Filbey, F. M. (2020). Interaction of cannabis use and aging: from molecule to mind. *J. Dual Diagn.* 16, 140–176. doi: 10.1080/15504263.2019.1665218
- Youn, Y.-H., Grant, R. W., McCabe, L. R., Albarado, D. C., Nguyen, K. Y., Ravussin, A., et al. (2013). Canonical Nlrp3 Inflammasome links systemic low-grade inflammation to functional decline in aging. *Cell Metab.* 18, 519–532. doi: 10.1016/j.cmet.2013.09.010
- Young, A. P., and Denovan-Wright, E. M. (2022a). Synthetic cannabinoids reduce the inflammatory activity of microglia and subsequently improve neuronal survival *in vitro*. *Brain Behav. Immun.* 105, 29–43. doi: 10.1016/j.bbi.2022.06.011
- Young, A. P., and Denovan-Wright, E. M. (2022b). The dynamic role of microglia and the Endocannabinoid system in Neuroinflammation. *Front. Pharmacol.* 12:806417. doi: 10.3389/fphar.2021.806417

- Young, K. M., Psachoulia, K., Tripathi, R. B., Dunn, S.-J., Cossell, L., Attwell, D., et al. (2013). Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. *Neuron* 77, 873–885. doi: 10.1016/j.neuron.2013.01.006
- Yousef, H., Czupalla, C. J., Lee, D., Chen, M. B., Burke, A. N., Zera, K. A., et al. (2019). Aged blood impairs hippocampal neural precursor activity and activates microglia via brain endothelial cell VCAM1. *Nat. Med.* 25, 988–1000. doi: 10.1038/s41591-019-0440-4
- Yu, Y., Li, L., Nguyen, D. T., Mustafa, S. M., Moore, B. M., and Jiang, J. (2020). Inverse Agonism of cannabinoid receptor type 2 confers anti-inflammatory and Neuroprotective effects following status epilepticus. *Mol. Neurobiol.* 57, 2830–2845. doi: 10.1007/s12035-020-01923-4
- Zhang, Y., Hou, B., Liang, P., Lu, X., Wu, Y., Zhang, X., et al. (2021). TRPV1 channel mediates NLRP3 inflammasome-dependent neuroinflammation in microglia. *Cell Death Dis.* 12:1159. doi: 10.1038/s41419-021-04450-9
- Zhang, X., Huang, N., Xiao, L., Wang, F., and Li, T. (2021). Replenishing the aged brains: targeting oligodendrocytes and myelination? *Front. Aging Neurosci.* 13:760200. doi: 10.3389/fnagi.2021.760200
- Zhang, H., Shen, H., Jordan, C. J., Liu, Q., Gardner, E. L., Bonci, A., et al. (2019). CB2 receptor antibody signal specificity: correlations with the use of partial CB2-knockout mice and anti-rat CB2 receptor antibodies. *Acta Pharmacol. Sin.* 40, 398–409. doi: 10.1038/s41401-018-0037-3
- Zou, S., and Kumar, U. (2018). Cannabinoid receptors and the Endocannabinoid system: signaling and function in the central nervous system. *IJMS* 19:833. doi: 10.3390/ijms19030833