



# Group I mGluR-dependent depotentiation in the lateral amygdala does not require the removal of calcium-permeable AMPA receptors

Kyungjoon Park<sup>1</sup>, Sukwoon Song<sup>1</sup>, Ingie Hong<sup>2</sup>, Beomjong Song<sup>3</sup>, Jeongyeon Kim<sup>4</sup>, Sungmo Park<sup>1</sup>, Junuk Lee<sup>1</sup>, Sangho Song<sup>4</sup>, Bobae An<sup>1</sup>, Jihye Kim<sup>1</sup>, C. Justin Lee<sup>4</sup>, Ki Soon Shin<sup>5</sup>, Sukwoo Choi<sup>1</sup> and Sukwon Lee<sup>1\*</sup>

<sup>1</sup> School of Biological Sciences, College of Natural Sciences, Seoul National University, Seoul, Korea (ROK)

<sup>2</sup> The Solomon H. Snyder Department of Neuroscience, Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD, USA

<sup>3</sup> Institute of Neuroscience, Technical University of Munich, Munich, Germany

<sup>4</sup> Center for Neural Science and Center for Connectomics, Korea Institute of Science and Technology, Seoul, Korea (ROK)

<sup>5</sup> Department of Biology, Department of Life and Nanopharmaceutical Sciences, Kyung Hee University, Seoul, Korea (ROK)

## Edited by:

Jeansok J. Kim, University of Washington, USA

## Reviewed by:

Hadley Bergstrom, National

Institutes of Health, USA

Roger Lee Clem, Icahn School of Medicine at Mount Sinai, USA

## \*Correspondence:

Sukwon Lee, School of Biological Sciences, College of Natural Sciences, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 151-742, Korea (ROK)  
e-mail: sukwon2@snu.ac.kr

There is conflicting evidence regarding whether calcium-permeable receptors are removed during group I mGluR-mediated synaptic depression. In support of this hypothesis, AMPAR rectification, a correlative index of the synaptic expression of GluA2-lacking calcium-permeable AMPARs (CP-AMPA), is known to decrease after the induction of several types of group I mGluR-mediated long-term depression (LTD), suggesting that a significant proportion of synaptic CP-AMPA is removed during synaptic depression. We have previously demonstrated that fear-conditioning-induced synaptic potentiation in the lateral amygdala is reversed by group 1 mGluR-mediated depotentiation. Here, we examined whether CP-AMPA is removed by mGluR1-mediated depotentiation of fear conditioning-induced synaptic potentiation. The synaptic expression of CP-AMPA was negligible before, increased significantly 12 h after, and returned to baseline 48 h after fear conditioning, as evidenced by the changes in the sensitivity of lateral amygdala synaptic responses to NASPM. Importantly, the sensitivity to NASPM was not altered after induction of depotentiation. Our findings, together with previous results, suggest that the removal of CP-AMPA is not required for the depotentiation of fear conditioning-induced synaptic potentiation at lateral amygdala synapses.

**Keywords:** calcium-permeable AMPA receptors, synaptic depotentiation, fear conditioning, lateral amygdala, long-term depression

## INTRODUCTION

Calcium-permeable AMPA receptors (CP-AMPA) are expressed transiently in excitatory neuron synapses but have been shown to be rapidly removed upon group I metabotropic glutamate receptor (mGluR) stimulation (Bellone and Lüscher, 2005, 2006; Clem and Haganir, 2010; McCutcheon et al., 2011; Loweth et al., 2013). Group 1 mGluR-induced synaptic depression also reduces the CP-AMPA-mediated current in GABAergic neurons (Kelly et al., 2009), suggesting that group 1 mGluRs may specifically target these CP-AMPA for synaptic removal. However, mGluR-LTD has been described in a wide variety of synaptic circuits that generally lack pre-existing CP-AMPA expression (for review, see Lüscher and Huber, 2010). Moreover, some types of mGluR-mediated synaptic depression require the phosphorylation/dephosphorylation of the AMPAR subunit GluA2 (Chung et al., 2003; Moulton et al., 2006), which CP-AMPA generally do not contain. Thus it is unclear whether group 1 mGluR-mediated synaptic

depression requires, or preferentially recruits, CP-AMPA removal.

The lateral amygdala (LA) is known to be critical for fear memory acquisition, extinction and renewal (LeDoux, 2000; Maren and Quirk, 2004; Kim et al., 2007; Myers and Davis, 2007; Knapska and Maren, 2009; Lee et al., 2013). The thalamic input synapses onto the LA (T-LA synapses) have been extensively studied as a site of learning-induced plasticity, and T-LA synaptic efficacy is tightly correlated with fear memory strength (McKernan and Shinnick-Gallagher, 1997; Rogan et al., 1997). Among several forms of synaptic plasticity at T-LA synapses, long-term depression (LTD) and/or depotentiation have been proposed as a cellular mechanism underlying extinction (or reconsolidation update) of fear memory (Lin et al., 2003, 2005; Kim et al., 2007; Clem and Haganir, 2010). LTD is a *de novo* decrease in synaptic efficacy, whereas depotentiation represents a net return of the potentiated synaptic efficacy to baseline (Collingridge et al., 2010). Although both

alterations result in a decrease in synaptic efficacy, the underlying mechanisms of these two types of plasticity may be different (Wagner and Alger, 1996; Kulla et al., 1999; Klausnitzer et al., 2004).

Several previous studies have reported that fear conditioning-induced synaptic potentiation *in vivo* at T-LA synapses can be depotentiated in brain slices prepared from conditioned animals (Kim et al., 2007; Clem and Hugarir, 2010, 2013). Our previous study has shown that depotentiation is blocked by intracellular dialysis of the GluA2<sub>3Y</sub> peptide, which prevents the internalization of GluA2-containing AMPARs (Ahmadian et al., 2004). Moreover, fear extinction reverses the conditioning-induced enhancements in the surface expression of synaptic GluA2 at LA synapses and occludes depotentiation, suggesting mutual mechanisms. Together, these findings suggest that depotentiation at T-LA synapses involves the internalization of GluA2-containing and, thus, calcium-impermeable AMPARs. However, conflicting evidence has also been presented primarily based on AMPAR rectification, an index of the synaptic expression of CP-AMPA, as GluA2-lacking CP-AMPA are removed during a type of LTD whose magnitude increases after fear conditioning (i.e., depotentiation-like plasticity) at T-LA synapses (Clem and Hugarir, 2013). One compromising factor in the latter study is that depotentiation could not be studied in isolation because the same stimuli also induces LTD before fear conditioning. Therefore, the specific subtypes of AMPARs involved in depotentiation of fear conditioning-induced synaptic potentiation are somewhat unclear.

In this study, we used the sensitivity of T-LA synaptic responses to 1-naphthylacetyl spermine (NASPM), a CP-AMPA antagonist, as an index of the synaptic expression of CP-AMPA to determine whether CP-AMPA are removed during depotentiation of fear conditioning-induced potentiation under conditions in which the depotentiation of fear conditioning-induced synaptic potentiation can be examined in isolation.

## MATERIALS AND METHODS

### ANIMALS AND AUDITORY FEAR CONDITIONING

All procedures were approved by the Institute of Laboratory Animal Resources of Seoul National University (Korea). Male Sprague-Dawley rats (4–5 weeks old) were maintained with free access to food and water under an inverted 12/12 h light/dark cycle (lights off at 09:00 h). Behavioral training was done during the dark portion of the light/dark cycle. For fear conditioning, rats were placed in a conditioning chamber and left undisturbed for 2 min. Then, a neutral tone (30 s, 2.8 kHz, 85 dB) coterminating with an electrical foot shock (1.0 mA, 1 s) was presented three times at an average interval of 100 s. After fear conditioning, rats were returned to their home cages until preparation of brain slices. Rats in naïve groups stayed in their home cages until brain slices were prepared.

### BRAIN SLICE PREPARATION

Sprague-Dawley rats (4–5 weeks old) were anesthetized with isoflurane and decapitated. Whole brains were isolated and placed in an ice-cold modified aCSF solution containing (in mM) 175 sucrose, 20 NaCl, 3.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 26 NaHCO<sub>3</sub>,

1.3 MgCl<sub>2</sub>, 11 D-(+)-glucose, and was gassed with 95% O<sub>2</sub>/5% CO<sub>2</sub>. Coronal slices (300 μm) including the LA were cut using a vibrating blade microtome (VT1200S, Leica Biosystems, Germany) and incubated in normal aCSF containing (in mM) 120 NaCl, 3.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 26 NaHCO<sub>3</sub>, 1.3 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 11 D-(+)-glucose, and was continuously bubbled at room temperature with 95% O<sub>2</sub>/5% CO<sub>2</sub>. Just before a given slice was transferred to the recording chamber, the cortex overlying the LA was cut away with a scalpel, so the addition of picrotoxin (100 μM; Abcam Plc., UK) would block cortical epileptic burst discharges from invading the LA.

### AFFERENT STIMULATION AND RECORDING CONDITIONS

We chose brain slices containing a well-isolated, sharply defined trunk (containing thalamic afferents) crossing the dorsolateral division of the LA where the somatosensory and auditory inputs converge. The sizes of the LA and the central amygdala were relatively constant in the utilized slices; when multiple trunks were observed, we used the closest trunk to the central nucleus of the amygdala. Unless otherwise noted, the thalamic afferents were stimulated at a frequency of 0.067 Hz using a concentric bipolar electrode (CBAEC75; FHC Inc., USA). The stimulation electrode was placed at the midpoint of the trunk between the internal capsule and the medial boundary of the LA. The regions and cells of interest for all recordings were located beneath the midpoint of the trunk spanning the LA horizontally.

### WHOLE-CELL PATCH-CLAMP RECORDINGS

Whole-cell recordings were made using an Axopatch 200A amplifier or Multiclamp 700A (Molecular Devices, Sunnyvale, CA, USA). For the whole-cell voltage-clamp recordings, the recordings were obtained using a Cs-based internal solution containing (in mM) 100 Cs-gluconate, 0.6 EGTA, 10 HEPES, 5 NaCl, 20 TEA, 4 Mg-ATP, 0.3 Na-GTP and 3 QX314, with the pH adjusted to 7.2 with CsOH and the osmolarity adjusted to approximately 297 mmol/kg with sucrose. The cells used were classified as principal neurons based on the pyramidal shape of their somata. We included picrotoxin (100 μM) in our recording solution to isolate excitatory synaptic transmission and block feed-forward GABAergic inputs to the principal neurons in the LA. The pipette resistances ranged from 3.5 to 4.5 Mohm. IR-DIC-enhanced visual guidance was used to select neurons that were 3–4 cell layers below the surface of the 300-μm-thick slices, which were held at 32 ± 1°C. The neurons were voltage-clamped at –70 mV except during paired pulse low-frequency stimulation (ppLFS-pairing), and the various solutions were delivered to the slices via gravity-driven superfusion at a flow rate of 1.4–1.5 ml/min. The pipette series resistance was monitored throughout each experiment, and the data were discarded if it changed by >20%. Whole-cell currents were filtered at 1 kHz, digitized at up to 20 kHz, and stored on a microcomputer (Clampex 9 software, Molecular Devices, Sunnyvale, CA, USA). EPSCs were monitored following stimulation at 0.067 Hz. One or two neurons were recorded per animal (a single neuron per slice). In the ppLFS-pairing protocol, stimulation for 3 min at 3 Hz was performed using paired pulses (50 ms interpulse interval) while the neuron was clamped at –50 mV, as described previously (Clem and Hugarir,

2013). Blockade of CP-AMPA receptors and mGluR1 was performed using NASPM (50  $\mu\text{M}$ ; Sigma-Aldrich) and LY367385 [(S)-(+)- $\alpha$ -amino-4-carboxy-2-methylbenzeneacetic acid] (100  $\mu\text{M}$ ; Tocris Bioscience). All EPSC amplitudes were normalized to an average of the baseline responses for the first 10 min and were expressed as percentages of the average baseline response. The percent inhibition by drugs (or vehicle) indicated difference in the average percentage of the responses between before and after drug (or vehicle) treatment. Therefore, the percent inhibition by NASPM in this study could be used to compare the amount of synaptic CP-AMPA receptors between before and after induction of LTD or depotentiation. The periods used to calculate these average responses were the 5 min immediately preceding drug treatment and the final 5 min after drug treatment, respectively. To prevent bias, we performed experiments in a blinded fashion. For improved visualization, the running averages of four data points were applied to the time-lapse experiments.


### STATISTICAL ANALYSIS

Between-group comparisons of the data were performed using either an unpaired *t*-test or one-way ANOVA with subsequent Newman-Keuls *post-hoc* comparison. A paired *t*-test was used to determine whether the post-treatment responses differed significantly from the baseline responses (Figures 1E, 3D). A *p*-value < 0.05 was considered to be statistically significant. The data from each neuron/slice were treated as independent samples. In all experiments using behaviorally trained rats, the data included samples from three or more animals.

## RESULTS

### SYNAPTIC EXPRESSION OF CP-AMPA RECEPTORS AT T-LA SYNAPSES AFTER FEAR CONDITIONING

We first characterized the synaptic expression of CP-AMPA receptors according to the sensitivity of T-LA synaptic responses to NASPM, a CP-AMPA receptor antagonist, at various time points (12 and 48 h) after fear conditioning in rats. We measured AMPAR-mediated EPSCs at T-LA synapses in the presence of D-AP5 (50  $\mu\text{M}$ ) via whole-cell voltage-clamp recording in acute brain slices prepared from conditioned (or naïve) rats. A negligible level of CP-AMPA receptors was found in slices prepared from naïve rats, but a significant level of CP-AMPA receptors was detected 12 h after conditioning. The enhanced level of CP-AMPA receptors returned to baseline in slices prepared 48 h after conditioning ( $F_{(2,13)} = 4.787$ ,  $p = 0.0319$ , one-way ANOVA; naïve,  $4.70 \pm 4.20\%$ ,  $n = 5$ ; 12 h,  $15.74 \pm 1.6\%$ ,  $n = 4$ ; 48 h,  $4.52 \pm 1.2\%$ ,  $n = 5$ ;  $p < 0.05$  for the 12 h group vs. the other groups, Newman-Keuls *post-hoc* test; Figures 1A–D), which was consistent with our previous study (Hong et al., 2013). Because NASPM has also been shown to inhibit kainate receptors (Koike et al., 1997; Cho et al., 2012), we examined whether UBP302 (10  $\mu\text{M}$ ), a specific antagonist of kainate receptors, blocked EPSCs at T-LA synapses when CP-AMPA receptor levels were elevated (12 h after conditioning). UBP302 displayed no significant effects on EPSCs at T-LA synapses in the presence of D-AP5 (50  $\mu\text{M}$ ) in slices prepared 12 h after fear conditioning ( $94.61 \pm 4.10\%$ ,  $n = 5$ ;  $p > 0.05$ ,



paired *t*-test; Figure 1E). In addition, we determined whether fear conditioning-induced synaptic potentiation was similar at those two time points (12 h vs. 48 h) after fear conditioning. There was no significant difference in the excitatory synaptic efficacy at T-LA synapses between the two groups (12 h after conditioning,  $11.25 \pm 1.57$  pA/ $\mu\text{A}$ ,  $n = 7$ ; 48 h after conditioning,  $11.84 \pm 1.36$  pA/ $\mu\text{A}$ ,  $n = 9$ ;  $p > 0.05$ , unpaired *t*-test; Figure 1F). Taken together, these results suggest that CP-AMPA receptors are transiently inserted into T-LA synapses after fear conditioning.


### CP-AMPA RECEPTORS ARE NOT REMOVED FROM T-LA SYNAPSES VIA DHPG-INDUCED DEPOTENTIATION


We have previously shown that DHPG, an agonist of group I mGluRs, induces the depotentiation of fear conditioning-induced

synaptic potentiation at T-LA synapses; that is, DHPG produces synaptic depression in slices prepared after fear conditioning but not in slices prepared from naïve rats or other controls (Kim et al., 2007; Hong et al., 2011). To examine whether CP-AMPA receptors are removed after induction of depotentiation, we monitored the changes in the sensitivity to NASPM after fear conditioning. Application of DHPG (100  $\mu$ M, 10 min) successfully induced synaptic depression in the presence of D-AP5 (50  $\mu$ M) in slices prepared 12 or 48 h after fear conditioning. After DHPG-induced depression had been stabilized, NASPM (or vehicle) was applied. NASPM treatment inhibited EPSCs significantly more than vehicle treatment when the CP-AMPA receptor levels were elevated (12 h after conditioning), but it did not exert a significant effect on EPSCs when CP-AMPA receptors were minimally expressed (48 h after conditioning) ( $F_{(2,16)} = 29.74$ ,  $p < 0.001$ , one-way ANOVA; vehicle group-12 h after conditioning,  $-0.42 \pm 1.04\%$ ,  $n = 5$ ; NASPM group-12 h after conditioning,  $18.59 \pm 2.15\%$ ,  $n = 6$ ; NASPM group-48 h after conditioning,  $1.78 \pm 2.11\%$ ,  $n = 6$ ;  $p < 0.0001$  for NASPM group-12 h after conditioning vs. the other groups, Newman-Keuls *post-hoc* test; **Figures 2A–D**). Thus, when CP-AMPA receptors are expressed at synapses, a significant proportion of CP-AMPA receptors appears to be retained even after the induction of depotentiation.

### CP-AMPA RECEPTORS ARE NOT REMOVED FROM T-LA SYNAPSES VIA ppLFS-INDUCED DEPOTENTIATION

Previous studies have shown that ppLFS also induces depotentiation or LTD (Kim et al., 2007; Hong et al., 2009). We performed a protocol of ppLFS-pairing which was used by Clem and Hugarir (2013). Unlike their findings that ppLFS-pairing induced LTD regardless of fear conditioning (either before or after conditioning), ppLFS-pairing produced no significant depression in slices from naïve rats ( $96.84 \pm 7.33\%$ ,  $n = 5$ ;  $p > 0.05$ , paired *t*-test; **Figure 3A**). Alternatively, ppLFS-pairing produced synaptic depression in slices prepared 12 h after conditioning, confirming the previously reported depotentiation of fear conditioning-induced synaptic potentiation (**Figures 3B,C**; Kim et al., 2007). After ppLFS-pairing-induced depression had been stabilized, NASPM (or vehicle) was applied. NASPM treatment inhibited EPSCs relative to vehicle treatment (vehicle,  $3.33 \pm 0.74\%$ ,  $n = 4$ ; NASPM,  $17.45 \pm 2.24\%$ ,  $n = 7$ ;  $p = 0.0013$ , unpaired *t*-test; **Figures 3B–D**). Therefore, similar to the results using DHPG, pre-existing CP-AMPA receptors appear to be largely retained after depotentiation. Because this particular result contradicts with the previous findings by Clem and Hugarir (2010, 2013), it was necessary to confirm that ppLFS-induced depression in the present study shared the same induction requirement with that in the two previous studies (i.e., mGluR1-dependency). Indeed, application of





**FIGURE 3 | Sensitivity to NASPM is maintained after ppLFS-pairing-induced depotentiation.** (A) ppLFS-pairing did not produce any significant changes in EPSCs. The slices were prepared from naïve rats. (B) pp-LFS-pairing produced synaptic depression, and vehicle treatment did not produce any further changes. The slices were prepared 12 h after conditioning. (C) ppLFS-pairing produced synaptic depression, and NASPM treatment produced further synaptic depression. The slices were prepared 12 h after conditioning.

(D) Average percent inhibition due to NASPM (or vehicle) treatment for the experiments shown in B (vehicle) and C (NASPM).

(E) ppLFS-pairing-induced depotentiation was blocked by the mGluR1 antagonist, LY367385. LY367385 (100  $\mu$ M) or vehicle (aCSF) was present during the entire recording period. The slices were prepared 12 h after conditioning. (F) A summary of NASPM-induced inhibition in all the groups in which the effects of NASPM treatment were evaluated.

\*\*  $p < 0.01$ . Scale bars: 50 pA and 10 ms.

the mGluR1 antagonist, LY367385, completely blocked ppLFS-pairing-induced depression (vehicle group,  $69.89 \pm 8.54\%$ ,  $n = 4$ ; LY367385 group,  $97.69 \pm 5.89\%$ ,  $n = 7$ ;  $p = 0.0223$ , unpaired  $t$ -test, Figure 3E).

Comparing the results from Figures 1–3 revealed that regardless of prior depotentiation, the primary factor that contributes to NASPM sensitivity at LA synapses is the duration after fear conditioning. NASPM-induced inhibition was minimal in naïve or 48 h, slices whereas in 12 h slices, NASPM-induced inhibition was prominent both before and after depotentiation, regardless of the depotentiation protocol (NASPM at baseline, NASPM after DHPG treatment and

NASPM after pp-LFS) when the CP-AMPA levels were elevated (12 h after conditioning) (NASPM at baseline (naïve rats),  $4.70 \pm 4.20\%$ ,  $n = 5$ ; NASPM at baseline (48 h after conditioning),  $4.52 \pm 1.24\%$ ,  $n = 5$ ; NASPM at baseline (12 h after conditioning),  $15.74 \pm 1.61\%$ ,  $n = 4$ ; NASPM after DHPG treatment (12 h after conditioning),  $18.59 \pm 2.15\%$ ,  $n = 6$ ; NASPM after ppLFS (12 h after conditioning),  $17.45 \pm 2.24\%$ ,  $n = 7$ ;  $F_{(4,26)} = 7.699$ ,  $p = 0.0005$ , one-way ANOVA;  $p > 0.05$  between the three groups prepared 12 h after conditioning,  $p < 0.05$  for these three groups vs. the other two groups, Newman-Keuls *post-hoc* test; Figure 3F). Thus, our data suggest that fear conditioning produces transient

insertion of CP-AMPA into T-LA synapses, but these CP-AMPA do not appear to be removed by depotentiation. The observation that depotentiation can be induced regardless of the expression of CP-AMPA also suggests that depotentiation does not require the pre-existence of CP-AMPA at synapses.

## DISCUSSION

Our findings suggest that removal of CP-AMPA from synapses does not contribute to the depotentiation of fear conditioning-induced synaptic potentiation. Under our experimental conditions, the sensitivity to NASPM is negligible before conditioning, increases 12 h after conditioning and returns to baseline 48 h after conditioning, suggesting that the synaptic expression of CP-AMPA is minimal before conditioning and increases transiently after conditioning. We have also determined whether NASPM selectively inhibits CP-AMPA, as it has also been shown to inhibit calcium-permeable kainate receptors. UBP302, a specific blocker of kainate receptors, does not inhibit T-LA EPSCs when CP-AMPA are elevated (but see Cho et al., 2012), ruling out the possibility that the sensitivity to NASPM is due to inhibition of kainate receptor-mediated currents. Importantly, the sensitivity to NASPM does not change after induction of depotentiation, even when the synaptic expression of CP-AMPA is elevated. This particular observation provides strong evidence that CP-AMPA are not removed from T-LA synapses during depotentiation of fear conditioning-induced synaptic potentiation.

Our results provide an example in which group 1 mGluR-mediated synaptic depression does not require or promote CP-AMPA removal. Instead, in T-LA synapses, it appears that mGluR1-dependent depotentiation leaves the pre-existing population of CP-AMPA largely intact. This result is in contrast with previous reports in which the pre-existing CP-AMPA content correlated to the extent of mGluR-LTD (Bellone and Lüscher, 2006), but our result is consistent with another report in which DHPG treatment induced synaptic depression regardless of prior CP-AMPA expression (McCutcheon et al., 2011). Together, these results shed light on the downstream molecular mechanisms of mGluR-mediated synaptic depression.

Two recent studies have reported the contribution of CP-AMPA to LTD at T-LA synapses (Clem and Hugarir, 2010, 2013). In these studies, the magnitude of LTD has been shown to increase after fear conditioning, and the increased portion of LTD might represent depotentiation of fear conditioning-induced synaptic potentiation, although other interpretations are also possible (e.g., metaplastic enhancements in the magnitude of LTD after conditioning). In addition, the basal expression of synaptic CP-AMPA is evident before fear conditioning, unlike our experimental conditions, which may involve a novel type of synaptic plasticity that is induced by calcium influx via pre-existing CP-AMPA. Therefore, although the most parsimonious explanation for the increased magnitude of LTD after fear conditioning is the reversal of fear conditioning-induced synaptic potentiation, it remains possible

that completely different types of synaptic plasticity (i.e., CP-AMPA-dependent LTD whose magnitude can be enhanced after fear conditioning) are involved in these two studies. Therefore, distinct mechanisms may underlie LTD and depotentiation of fear conditioning-induced synaptic potentiation at T-LA synapses; indeed, these two types of plasticity (depotentiation and LTD) are known to underlie fear extinction and reconsolidation update (a variant of extinction that produces fear memory erasure), respectively (Kim et al., 2007; Clem and Hugarir, 2010).

It is generally considered that in naïve animals (rats or mice), the synaptic expression of CP-AMPA is negligible in the LA (Mahanty and Sah, 1998; Polepalli et al., 2010). Thus, it may appear odd to detect a relatively large level of CP-AMPA under baseline conditions in the two previous studies (Clem and Hugarir, 2010, 2013). It will be interesting to determine which conditions affect the amount of synaptic CP-AMPA in the experimental subjects (see also Whitehead et al., 2013).

Our previous study (Kim et al., 2007) has shown that ppLFS-induced depotentiation (or LTD) is dependent on both mGluR and NMDAR activity at T-LA synapses. Similarly, Clem and Hugarir have also reported that ppLFS-induced LTD is dependent on both mGluR and NMDAR activity at T-LA synapses (Clem and Hugarir, 2010, 2013). In these three studies, ppLFS-induced LTD (or depotentiation) was completely blocked either by mGluR antagonists or by NMDAR antagonists unlike the case of the hippocampal LTD in which each antagonist partially blocked LTD (Oliet et al., 1997). Furthermore, Clem and Hugarir (2013) has shown that LFS-induced LTD is dependent on NMDARs, but not on mGluRs at T-LA synapses, and that LFS-induced LTD is produced via mechanisms that were completely different from those underlying ppLFS-induced LTD. Therefore, it is possible that ppLFS induces a unique form of LTD, which depends on both mGluRs and NMDARs, at T-LA synapses, and that these two types of receptors merge on the same intracellular signaling pathway.

There have been previous studies showing the existence of several forms of LTD in the lateral amygdala. Low-frequency or theta-frequency stimulation has been shown to produce LTD at naive synapses in the lateral amygdala (Heinbockel and Pape, 2000; Dalton et al., 2012; Clem and Hugarir, 2013). It remains to be elucidated whether these forms of LTD share similar mechanisms or not. Moreover, other new forms of LTD may coexist at lateral amygdala synapses and each of them may play a unique role in a distinct physiological process.

In summary, the present study has provided strong evidence that the removal of CP-AMPA from synapses does not contribute to the depotentiation of fear conditioning-induced synaptic potentiation. This conclusion is consistent with our and other previous studies showing that extinction (whose cellular substrate is the depotentiation of fear conditioning-induced synaptic potentiation) does not involve CP-AMPA removal (Clem and Hugarir, 2013; Lee et al., 2013). In contrast, removal of CP-AMPA via LTD at T-LA synapses has been proposed as a mechanism underlying reconsolidation update. Thus, the molecular mechanisms underlying LTD may differ from those underlying

depotential of fear conditioning-induced synaptic potentiation at T-LA synapses.

## ACKNOWLEDGMENTS

This work was supported by the National Research Foundation (NRF) of Korea Grant NRF-2011-0012876 funded by the Korean Ministry of Science and Technology (Sukwon Lee), by the NRF of Korea grant funded by the Korea government the Ministry of Education, Science and Technology (MEST) (No. 2011-0018209) (Sukwoo Choi), by the Original Technology Research Program for Brain Science through the NRF of Korea funded by MEST (No. 2011-0019226) (Sukwoo Choi), and by a grant from the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health and Welfare, Republic of Korea (grant number: HI12C1492) (Sukwoo Choi).

## REFERENCES

- Ahmadian, G., Ju, W., Liu, L., Wyszynski, M., Lee, S. H., Dunah, A. W., et al. (2004). Tyrosine phosphorylation of GluR2 is required for insulin-stimulated AMPA receptor endocytosis and LTD. *EMBO J.* 23, 1040–1050. doi: 10.1038/sj.emboj.7600126
- Bellone, C., and Lüscher, C. (2005). mGluRs induce a long-term depression in the ventral tegmental area that involves a switch of the subunit composition of AMPA receptors. *Eur. J. Neurosci.* 21, 1280–1288. doi: 10.1111/j.1460-9568.2005.03979.x
- Bellone, C., and Lüscher, C. (2006). Cocaine triggered AMPA receptor redistribution is reversed in vivo by mGluR-dependent long-term depression. *Nat. Neurosci.* 9, 636–641. doi: 10.1038/nn1682
- Cho, J. H., Bayazitov, I. T., Meloni, E. G., Myers, K. M., Carlezon, W. A. Jr., Zakharenko, S. S., et al. (2012). Coactivation of thalamic and cortical pathways induces input timing-dependent plasticity in amygdala. *Nat. Neurosci.* 15, 113–122. doi: 10.1038/nn.2993
- Chung, H. J., Steinberg, J. P., Haganir, R. L., and Linden, D. J. (2003). Requirement of AMPA receptor GluR2 phosphorylation for cerebellar long-term depression. *Science* 300, 1751–1755. doi: 10.1126/science.1082915
- Clem, R. L., and Haganir, R. L. (2010). Calcium-permeable AMPA receptor dynamics mediate fear memory erasure. *Science* 330, 1108–1112. doi: 10.1126/science.1195298
- Clem, R. L., and Haganir, R. L. (2013). Norepinephrine enhances a discrete form of long-term depression during fear memory storage. *J. Neurosci.* 33, 11825–11832. doi: 10.1523/JNEUROSCI.3317-12.2013
- Collingridge, G. L., Peineau, S., Howland, J. G., and Wang, Y. T. (2010). Long-term depression in the CNS. *Nat. Rev. Neurosci.* 11, 459–473. doi: 10.1038/nrn2867
- Dalton, G. L., Wu, D. C., Wang, Y. T., Floresco, S. B., and Phillips, A. G. (2012). NMDA GluN2A and GluN2B receptor splay separate roles in the induction of LTP and LTD in the amygdala and in the acquisition and extinction of conditioned fear. *Neuropharmacology* 62, 797–806. doi: 10.1016/j.neuropharm.2011.09.001
- Heinbockel, T., and Pape, H. C. (2000). Input-specific long-term depression in the lateral amygdala evoked by theta frequency stimulation. *J. Neurosci.* 20, RC68.
- Hong, I., Kim, J., Kim, J., Lee, S., Ko, H. G., Nader, K., et al. (2013). AMPA receptor exchange underlies transient memory destabilization on retrieval. *Proc. Natl. Acad. Sci. U S A* 110, 8218–8223. doi: 10.1073/pnas.1305235110
- Hong, I., Kim, J., Lee, J., Park, S., Song, B., Kim, J., et al. (2011). Reversible plasticity of fear memory-encoding amygdala synaptic circuits even after fear memory consolidation. *PLoS One* 6:e24260. doi: 10.1371/journal.pone.0024260
- Hong, I., Song, B., Lee, S., Kim, J., Kim, J., and Choi, S. (2009). Extinction of cued fear memory involves a distinct form of depotentiation at cortical input synapses onto the lateral amygdala. *Eur. J. Neurosci.* 30, 2089–2099. doi: 10.1111/j.1460-9568.2009.07004.x
- Kelly, L., Farrant, M., and Cull-Candy, S. G. (2009). Synaptic mGluR activation drives plasticity of calcium-permeable AMPA receptors. *Nat. Neurosci.* 12, 593–601. doi: 10.1038/nn.2309
- Kim, J., Lee, S., Park, K., Hong, I., Song, B., Son, G. H., et al. (2007). Amygdala depotentiation and fear extinction. *Proc. Natl. Acad. Sci. U S A* 104, 20955–20960. doi: 10.1073/pnas.0710548105
- Klausnitzer, J., Kulla, A., and Manahan-Vaughan, D. (2004). Role of the group III metabotropic glutamate receptor in LTP, depotentiation and LTD in the dentate gyrus of freely moving rats. *Neuropharmacology* 46, 160–170. doi: 10.1016/j.neuropharm.2003.09.019
- Knapska, E., and Maren, S. (2009). Reciprocal patterns of c-fos expression in the medial prefrontal cortex and amygdala after extinction and renewal of conditioned fear. *Learn. Mem.* 16, 486–493. doi: 10.1101/lm.14
- Koike, M., Iino, M., and Ozawa, S. (1997). Blocking effect of 1-naphthyl acetyl spermine on Ca(2+)-permeable AMPA receptors in cultured rat hippocampal neurons. *Neurosci. Res.* 29, 27–36. doi: 10.1016/s0168-0102(97)-9
- Kulla, A., Reymann, K. G., and Manahan-Vaughan, D. (1999). Time-dependent induction of depotentiation in the dentate gyrus of freely moving rats: involvement of group 2 metabotropic glutamate receptors. *Eur. J. Neurosci.* 11, 3864–3872. doi: 10.1046/j.1460-9568.1999.00807.x
- LeDoux, J. E. (2000). Emotion circuits in the brain. *Annu. Rev. Neurosci.* 23, 155–184. doi: 10.1146/annurev.neuro.23.1.155
- Lee, S., Song, B., Kim, J., Park, J., Hong, I., An, B., et al. (2013). GluA1 phosphorylation at serine 831 in the lateral amygdala is required for fear renewal. *Nat. Neurosci.* 16, 1436–1444. doi: 10.1038/nn.3491
- Lin, C.-H., Lee, C.-C., and Gean, P.-W. (2003). Involvement of a calcineurin cascade in amygdala depotentiation and quenching of fear memory. *Mol. Pharmacol.* 63, 44–52. doi: 10.1124/mol.63.1.44
- Lin, C.-H., Lee, C.-C., Huang, Y.-C., Wang, S.-J., and Gean, P.-W. (2005). Activation of group II metabotropic glutamate receptors induces depotentiation in amygdala slices and reduces fear-potentiated startle in rats. *Learn. Mem.* 12, 130–137. doi: 10.1101/lm.85304
- Loweth, J. A., Scheyer, A. F., Milovanovic, M., LaCrosse, A. L., Flores-Barrera, E., Werner, C. T., et al. (2013). Synaptic depression via mGluR1 positive allosteric modulation suppresses cue-induced cocaine craving. *Nat. Neurosci.* 17, 73–80. doi: 10.1038/nn.3590
- Lüscher, C., and Huber, K. M. (2010). Group 1 mGluR-dependent synaptic long-term depression: mechanisms and implications for circuitry and disease. *Neuron* 65, 445–459. doi: 10.1016/j.neuron.2010.01.016
- Mahanty, N. K., and Sah, P. (1998). Calcium-permeable AMPA receptors mediate long-term potentiation in interneurons in the amygdala. *Nature* 394, 683–687. doi: 10.1038/29312
- Maren, S., and Quirk, G. J. (2004). Neuronal signalling of fear memory. *Nat. Rev. Neurosci.* 5, 844–852. doi: 10.1038/nrn1535
- McCutcheon, J. E., Loweth, J. A., Ford, K. A., Marinelli, M., Wolf, M. E., and Tseng, K. Y. (2011). Group I mGluR activation reverses cocaine-induced accumulation of calcium-permeable AMPA receptors in nucleus accumbens synapses via a protein kinase C-dependent mechanism. *J. Neurosci.* 31, 14536–14541. doi: 10.1523/JNEUROSCI.3625-11.2011
- McKernan, M. G., and Shinnick-Gallagher, P. (1997). Fear conditioning induces a lasting potentiation of synaptic currents in vitro. *Nature* 390, 607–611. doi: 10.1038/37605
- Moult, P. R., Gladding, C. M., Sanderson, T. M., Fitzjohn, S. M., Bashir, Z. I., Molnar, E., et al. (2006). Tyrosine phosphatases regulate AMPA receptor trafficking during metabotropic glutamate receptor-mediated long-term depression. *J. Neurosci.* 26, 2544–2554. doi: 10.1523/jneurosci.4322-05.2006
- Myers, K. M., and Davis, M. (2007). Mechanisms of fear extinction. *Mol. Psychiatry* 12, 120–150. doi: 10.1038/sj.mp.4001939
- Oliet, S. H., Malenka, R. C., and Nicoll, R. A. (1997). Two distinct forms of long-term depression coexist in CA1 hippocampal pyramidal cells. *Neuron* 18, 969–982. doi: 10.1016/s0896-6273(00)80336-0
- Polepalli, J. S., Sullivan, R. K., Yanagawa, Y., and Sah, P. (2010). A specific class of interneuron mediates inhibitory plasticity in the lateral amygdala. *J. Neurosci.* 30, 14619–14629. doi: 10.1523/JNEUROSCI.3252-10.2010

- Rogan, M. T., Stäubli, U. V., and LeDoux, J. E. (1997). Fear conditioning induces associative long-term potentiation in the amygdala. *Nature* 390, 604–607. doi: 10.1038/37601
- Wagner, J. J., and Alger, B. E. (1996). Homosynaptic LTD and depotentiation: do they differ in name only? *Hippocampus* 6, 24–29. doi: 10.1002/(sici)1098-1063(1996)6:1<24::aid-hipo5>3.0.co;2-7
- Whitehead, G., Jo, J., Hogg, E. L., Piers, T., Kim, D. H., Seaton, G., et al. (2013). Acute stress causes rapid synaptic insertion of Ca<sup>2+</sup>-permeable AMPA receptors to facilitate long-term potentiation in the hippocampus. *Brain* 136(Pt. 12), 3753–3765. doi: 10.1093/brain/awt293

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 30 May 2014; accepted: 21 July 2014; published online: 08 August 2014.

Citation: Park K, Song S, Hong I, Song B, Kim J, Park S, Lee J, Song S, An B, Kim J, Lee CJ, Shin KS, Choi S and Lee S (2014) Group I mGluR-dependent depotentiation in the lateral amygdala does not require the removal of calcium-permeable AMPA receptors. *Front. Behav. Neurosci.* 8:269. doi: 10.3389/fnbeh.2014.00269

This article was submitted to the journal *Frontiers in Behavioral Neuroscience*.

Copyright © 2014 Park, Song, Hong, Song, Kim, Park, Lee, Song, An, Kim, Lee, Shin, Choi and Lee. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.