



# A new role for P2X<sub>4</sub> receptors as modulators of lung surfactant secretion

Pika Miklavc, Kristin E. Thompson and Manfred Frick\*

Institute of General Physiology, University of Ulm, Ulm, Germany

## Edited by:

Geoffrey Burnstock, University  
College Medical School, UK

## Reviewed by:

Jean-Pierre Timmermans, University  
of Antwerp (State University Centre  
Antwerp), Belgium

Ruth Dorothy Murrell-Lagnado,  
University of Cambridge, UK

## \*Correspondence:

Manfred Frick, Institute of General  
Physiology, University of Ulm,  
Albert-Einstein Allee 11, 89081 Ulm,  
Germany  
e-mail: manfred.frick@uni-ulm.de

In recent years, P2X receptors have attracted increasing attention as regulators of exocytosis and cellular secretion. In various cell types, P2X receptors have been found to stimulate vesicle exocytosis directly via Ca<sup>2+</sup> influx and elevation of the intracellular Ca<sup>2+</sup> concentration. Recently, a new role for P2X<sub>4</sub> receptors as regulators of secretion emerged. Exocytosis of lamellar bodies (LBs), large storage organelles for lung surfactant, results in a local, fusion-activated Ca<sup>2+</sup> entry (FACE) in alveolar type II epithelial cells. FACE is mediated via P2X<sub>4</sub> receptors that are located on the limiting membrane of LBs and inserted into the plasma membrane upon exocytosis of LBs. The localized Ca<sup>2+</sup> influx at the site of vesicle fusion promotes fusion pore expansion and facilitates surfactant release. In addition, this inward-rectifying cation current across P2X<sub>4</sub> receptors mediates fluid resorption from lung alveoli. It is hypothesized that the concomitant reduction in the alveolar lining fluid facilitates insertion of surfactant into the air-liquid interphase thereby "activating" it. These findings constitute a novel role for P2X<sub>4</sub> receptors in regulating vesicle content secretion as modulators of the secretory output during the exocytic post-fusion phase.


**Keywords: P2X<sub>4</sub> receptor, lamellar body, alveolar epithelial cell, exocytosis, calcium, cellular secretion, pulmonary surfactant**

## INTRODUCTION

In recent years, P2X receptors have attracted increasing attention as regulators of exocytosis and cellular secretion in a wide variety of organs including the lungs (Burnstock et al., 2012). P2X receptors are membrane cation channels that are activated by extracellular adenosine triphosphate (ATP), the molecular and functional properties of which have been reviewed in detail elsewhere (Surprenant, 1996; North, 2002; Khakh and North, 2006; Burnstock and Kennedy, 2011; Kaczmarek-Hajek et al., 2012). ATP has been known to stimulate cellular secretion for several decades (Rodriguez Candela and Garcia-Fernandez, 1963; Diamant and Kruger, 1967). One of the earliest indications for involvement of P2X receptors in stimulating secretion came from the studies of Cockcroft and Gomperts (1979a,b, 1980). They found that ATP triggers degranulation and histamine release in mast cells via activation of P<sub>2Z</sub> (Cockcroft and Gomperts, 1980), which later turned out to be P2X<sub>7</sub> (Surprenant et al., 1996). Since the first cloning of P2X receptor subunits in 1994 (Brake et al., 1994; Valera et al., 1994), P2X receptors have been found to stimulate and modulate various cellular secretion pathways, including fluid secretion in exocrine glands and epithelia (Novak, 2011), secretion of cytokines via release of plasma-derived microvesicles (Solini et al., 1999; MacKenzie et al., 2001) or exosomes (Qu et al., 2007; Qu and Dubyak, 2009).

Moreover, several members of the P2X family have been implicated in regulating exocytosis of secretory organelles in a variety of cell types (Gu and MacDermott, 1997; Ulmann et al., 2008; Jacques-Silva et al., 2010; Gutierrez-Martin et al., 2011; Huang

et al., 2011). Substantial evidence suggests that P2X receptor activation stimulates exocytosis directly via influx of Ca<sup>2+</sup> from the extracellular space and elevation of the cytoplasmic Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>c</sub>; Kim et al., 2004; Shigetomi and Kato, 2004; Jacques-Silva et al., 2010; Hayoz et al., 2012). It is well established that a series of Ca<sup>2+</sup>-dependent steps during the exocytic pre-fusion stage is essential for fusion of exocytic vesicles with the plasma membrane (Burgoyne and Morgan, 1998; Sudhof, 2004; Neher and Sakaba, 2008). Ca<sup>2+</sup> can either enter through P2X receptor pores themselves or through voltage-gated Ca<sup>2+</sup> channels, which are activated as a consequence of the P2X receptor-mediated membrane depolarization (Khakh and North, 2006). In line with these findings, several studies proposed a role for P2X<sub>4</sub> receptors in exocytosis that is mediated via an increase in the intracellular Ca<sup>2+</sup> concentration. P2X<sub>4</sub> receptors have a relatively slow desensitization (5–10 s) and a high Ca<sup>2+</sup> permeability, Ca<sup>2+</sup> contributes 8% of the whole current in human P2X<sub>4</sub> (Wang et al., 1996; Garcia-Guzman et al., 1997; North, 2002; Egan and Khakh, 2004). Hence, activation of P2X<sub>4</sub> receptors can generate sufficient increases in [Ca<sup>2+</sup>]<sub>c</sub> to stimulate regulated exocytosis. Indeed, insulin secretion from pancreatic islets (Ohtani et al., 2011) and exocytic response in parotid acinar cells (Bhattacharya et al., 2012) following stimulation with ATP were augmented in the presence of ivermectin, a selective potentiator of P2X<sub>4</sub> receptor currents (Khakh et al., 1999). P2X<sub>4</sub> activation was also found to modulate glutamate and gamma-aminobutyric acid (GABA) release in hypothalamic neurons (Vavra et al., 2011) and brain-derived neurotrophic factor (BDNF) in microglial cells (Trang et al., 2009).



In all of these systems, activation of P2X receptors adjusts the secretory output predominantly by modulating the number of vesicles that fuse with the plasma membrane. Depending on the cell type and the shape of the Ca<sup>2+</sup> signal, the rise in [Ca<sup>2+</sup>]<sub>c</sub> triggers fusion of secretory vesicles with the plasma membrane, but also affects maturation and trafficking of secretory vesicles to the plasma membrane (Neher and Sakaba, 2008; Dolensek et al., 2011; Gutierrez-Martin et al., 2011).

### VESICULAR P2X<sub>4</sub> RECEPTORS PROMOTE SURFACTANT SECRETION VIA FACE – “FUSION-ACTIVATED Ca<sup>2+</sup>-ENTRY”

Apart from regulating secretion via adjusting the number of fusing organelles the amount and composition of the secretory output is – at least for exocytosis of large secretory granules and secretion of bulky vesicle contents – modulated following fusion of the vesicle with the plasma membrane during the so-called exocytic “post-fusion” phase. Recent evidence also suggests a role for P2X<sub>4</sub> receptors therein. It has been demonstrated that activation of P2X<sub>4</sub> receptors following vesicle–plasma membrane fusion modulates the secretion and activation of pulmonary surfactant (Miklavc et al., 2011; Dietl et al., 2012; Thompson et al., 2013).

Pulmonary surfactant is secreted via exocytosis of lamellar bodies (LBs), large lysosome-related storage organelles in alveolar type II (AII) epithelial cells. Surfactant is stored in LBs as densely packed membranous structures that do not readily diffuse out of fused LBs following opening of the exocytic fusion pore. Rather, surfactant is so insoluble, that it may remain entrapped within the fused vesicle for many minutes and the slowly expanding fusion pore acts as a mechanical barrier for the release (Dietl et al., 2001; Haller et al., 2001; Singer et al., 2003; Dietl and Haller, 2005; Miklavc et al., 2012).

Miklavc et al. (2010) initially discovered that exocytosis of LBs results in localized Ca<sup>2+</sup> influx at the site of vesicle fusion which they termed “FACE” for “fusion-activated Ca<sup>2+</sup>-entry”. Subsequently, they found that FACE is mediated via activation of P2X<sub>4</sub> receptors expressed on the limiting membranes of LBs (Miklavc et al., 2011). Upon exocytosis of LBs, the P2X<sub>4</sub> receptor is readily part of the apical membrane as soon as membrane fusion is

completed (Miklavc et al., 2009). Activation of P2X<sub>4</sub> in the presence of extracellular ATP then results in a transient, non-selective, inward-rectifying, cation current at the site of the fused vesicle (Miklavc et al., 2011; Thompson et al., 2013) (Figure 1). The relatively high Ca<sup>2+</sup> permeability of P2X<sub>4</sub> receptors (North, 2002) causes a local, transient rise of [Ca<sup>2+</sup>]<sub>c</sub> around the fused vesicle which promotes fusion pore expansion (Miklavc et al., 2011). In AII cells, vesicle content (i.e., surfactant) release is tightly regulated via Ca<sup>2+</sup>-dependent fusion pore expansion (Haller et al., 2001) and it has been demonstrated that FACE via P2X<sub>4</sub> receptors on LBs directly facilitates surfactant release in the alveolus (Miklavc et al., 2011).

Ca<sup>2+</sup> channels localized in the membranes of the secretory vesicles that respond to changes in the membrane potential or extracellular agonists upon fusion are ideally suited for generating a localized rise in Ca<sup>2+</sup> and selectively affect the individual fused vesicle. Yet, so far such mechanisms have only been known in invertebrates (Smith et al., 2000; Yao et al., 2009; Miklavc and Frick, 2011) and P2X<sub>4</sub> receptors on LBs resemble the first analog mechanism in mammals. It will be interesting to see whether a similar role for P2X receptors is present in other secretory cells. Similar to LBs in AII cells, many different cell types harbor secretory lysosomes or lysosome-related organelles to store for secretory products that are released via exocytosis of these organelles (Dell’Angelica et al., 2000; Blott and Griffiths, 2002; Luzio et al., 2007). Many of these contain rather bulky, macromolecular vesicle contents and release is modulated via the exocytic post-fusion phase (Thorn, 2009). In addition, it is well established that P2X receptors, in particular P2X<sub>4</sub>, are predominantly located within lysosomal compartments and inserted into the cell surface upon exocytosis (Qureshi et al., 2007; Toyomitsu et al., 2012).

### VESICULAR P2X<sub>4</sub> RECEPTORS FACILITATE “ACTIVATION” OF SURFACTANT

Following release into the alveolar hypophase surfactant maintains its compact organization, constituting lamellar body-like particles (LBPs; Haller et al., 2004). To gain its vital function of reducing the surface tension within alveoli, it needs to be inserted

into the air–liquid interface. Freshly released LBPs disintegrate when they contact an air–liquid interface, leading to instantaneous spreading and insertion of surfactant material at this interface (Dietl and Haller, 2005). Thompson et al. (2013) demonstrated that, in addition to facilitating fusion pore dilation, FACE via P2X<sub>4</sub> also drives fluid resorption from the alveolar lumen. The P2X<sub>4</sub> mediated inward-rectifying cation current on the apical side results in vectorial ion transport across ATII cells, which in turn promotes apical to basolateral fluid transport (Thompson et al., 2013) (**Figure 1**). FACE-dependent transepithelial fluid resorption is a rather transient process which requires the presence of luminal ATP or other P2X<sub>4</sub> agonists and hence it is unlikely that it is a major contributor to regulation of alveolar liquid homeostasis under physiological conditions (Folkesson and Matthay, 2006). However, it has been suggested that this localized alveolar fluid resorption results in temporary thinning of the alveolar hypophase which in turn promotes contact between LBPs and the interphase and facilitates adsorption of newly released surfactant into the air–liquid interface (Thompson et al., 2013). Hence, activation of P2X<sub>4</sub> and FACE (which in order to embrace the true nature of FACE should now be referred to as “fusion-activated cation entry”) facilitates surfactant release via fusion pore opening and contributes to “activation” or “functionalising” of surfactant. Such a temporal and local coordination of surfactant secretion and reduction of alveolar lining fluid could constitute a powerful mechanism for fine-tuning surfactant replenishment – the integrators being vesicular P2X<sub>4</sub> receptors and extracellular ATP.

### ATP AS INTEGRATOR FOR SURFACTANT SECRETION AND “ACTIVATION”

It is intriguing that extracellular ATP plays multiple functions for surfactant secretion in the alveolus. Apart from P2X<sub>4</sub> receptors, ATII cells also express P2Y<sub>2</sub> receptors (Garcia-Verdugo et al., 2008; Burnstock et al., 2012) and activation thereof is one of the most potent stimuli for LB exocytosis and surfactant secretion (Rice and Singleton, 1987; Frick et al., 2001; Andreeva et al., 2007; Dietl et al., 2010). Hence, ATP is integrating the entire secretion process from stimulating LB exocytosis to facilitating surfactant release and “activating” surfactant during the post-fusion phase.

Despite this importance of ATP for lung function, the origins of ATP in the alveoli are still elusive. It has been reported that ATP is present in the pulmonary hypophase (Patel et al., 2005), however, the estimated concentration under resting conditions is in the low nM range (Bove et al., 2010), well below the EC<sub>50</sub> values for P2X<sub>4</sub> activation (North, 2002) or P2Y<sub>2</sub> activation (Lazarowski et al., 1995; Brunschweiler and Muller, 2006).

Cell stretch during deep inflation is considered the most potent if not only physiologically relevant stimulus for surfactant secretion (Wirtz and Dobbs, 2000; Dietl et al., 2004, 2010; Frick et al., 2004) and stretch-induced ATP release from alveolar epithelial cells (Patel et al., 2005; Mishra et al., 2011) could represent a key regulatory element (Dietl et al., 2010). Several possible pathways for ATP release have been described in the respiratory epithelia. ATP can either be released into the hypophase via regulated exocytosis from secretory cells (Kreda et al., 2010; Okada et al., 2011), or in a conductive way via

pannexin hemichannels (Ransford et al., 2009; Seminario-Vidal et al., 2011) or P2X<sub>7</sub> receptors (Mishra et al., 2011). In particular, local ATP release within individual alveoli may provide an ideal mechanism to gradually adapt local surfactant secretion to local demands. The alveolar epithelium consists of only two cell types; besides surfactant secreting ATII cells, flat alveolar type I (ATI) cells cover most of the alveolar surface. In contrast to primary ATII cells that only express P2X<sub>4</sub> receptors (Miklavc et al., 2011) ATI cells express P2X<sub>4</sub> and P2X<sub>7</sub> receptors (Weinhold et al., 2010; Burnstock et al., 2012). P2X<sub>7</sub> knock-out mice fail to increase surfactant secretion in response to hyperventilation and substantial evidence suggests that ATP release via P2X<sub>7</sub> receptors on ATI cells maintains alveolar surfactant homeostasis in response to increased alveolar distension by stimulating P2Y<sub>2</sub> receptors on ATII cells (Mishra et al., 2011) and, in light of our recent findings, possible activation of P2X<sub>4</sub> (Miklavc et al., 2011; Thompson et al., 2013). In addition to responding to mechanical distension of alveoli, alveolar epithelial cells also respond to increased tension forces at the air–liquid interphase with exocytic release of ATP (e.g., upon local depletion of surfactant or when coming in close proximity to the air–liquid interphase following a decrease in alveolar hypophase height; Ramsingh et al., 2011).

Whether ATII cells also release ATP, to act in an autocrine feedback loop, is still unknown. Many secretory vesicles, including lysosome-related organelles, have been found to contain significant amounts of ATP (Bodin and Burnstock, 2001; Praetorius and Leipziger, 2009; Lazarowski et al., 2011) and it has been reported that ATP is released from ATII-like A549 cells, likely via exocytosis (Tatur et al., 2008; Ramsingh et al., 2011). It is tempting to speculate that LBs contain ATP and hence provide the ligand for the P2X<sub>4</sub> receptors themselves. In such a scenario, the high degree of pH sensitivity of this receptor (Clarke et al., 2000; Zsembergy et al., 2003; Coddou et al., 2011) could prevent intravesicular activation of the receptor in the presence of vesicular ATP (pH of LB is <6.1; Chander et al., 1986).

Also, under pathophysiological conditions resulting from many chronic lung diseases, release of purine nucleotides from respiratory epithelia is significantly increased (Adriaensen and Timmermans, 2004; Lommatzsch et al., 2010). It has been demonstrated that trauma-induced damage of the alveolus leads to substantial ATP release and that extracellular ATP is a key player to rescue alveolar function following damage, including regulation of surfactant secretion (Riteau et al., 2010; Belete et al., 2011). In addition, several studies have demonstrated up-regulation of P2X receptors in various cell types during pathological conditions including inflammation, tumor growth, and injury (Burnstock and Kennedy, 2011) and it has been hypothesized that chronic extracellular ATP may be responsible (Geisler et al., 2013). Such a mechanism could be particularly relevant in the lung, and P2X receptors may play an even greater role in many pathological conditions with chronically increased extracellular ATP levels. Initial evidence came from studies indicating that smoke-induced lung inflammation leads to increased levels of ATP in broncho-alveolar fluid and up-regulation of P2X<sub>7</sub> expression (Lommatzsch et al., 2010; Lucattelli et al., 2011). A similar role for P2X<sub>4</sub> receptors under pathophysiological conditions is still to be confirmed. However,

it is becoming increasingly evident that purinergic signaling is taking center stage in regulating secretion of pulmonary surfactant and adapting it to local demands under physiological and diseased conditions. P2X<sub>4</sub> receptors on LBs provide an ideal mechanism for fine-tuning surfactant secretion via ATP levels in the alveolar hypophase.

Despite recent advances in our understanding how purinergic signaling in the alveolus, in particular activation of vesicular P2X<sub>4</sub> receptors, modulates LB exocytosis, surfactant secretion and activation of released surfactant, several important questions still

remain to be answered: What is the physiological relevance of such a complex regulatory mechanism? What are the sources of ATP under physiological and more importantly pathophysiological conditions? And – extending the scope from the lung – is purinergic signaling a general mechanism for secretion of large, macromolecular vesicle contents or is it unique to LB exocytosis and surfactant secretion? The answers to these questions warrant further research and certainly promise an increased understanding of the role of P2X receptors in regulating exocytosis and cellular secretion.

## REFERENCES

- Adriaensen, D., and Timmermans, J. P. (2004). Purinergic signalling in the lung: important in asthma and COPD? *Curr. Opin. Pharmacol.* 4, 207–214. doi: 10.1016/j.coph.2004.01.010
- Andreeva, A. V., Kutuzov, M. A., and Voyno-Yasenetskaya, T. A. (2007). Regulation of surfactant secretion in alveolar type II cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 293, L259–L271. doi: 10.1152/ajplung.00112.2007
- Belete, H. A., Hubmayr, R. D., Wang, S., and Singh, R. D. (2011). The role of purinergic signaling on deformation induced injury and repair responses of alveolar epithelial cells. *PLoS ONE* 6:e27469. doi: 10.1371/journal.pone.0027469
- Bhattacharya, S., Verrill, D. S., Carbone, K. M., Brown, S., Yule, D. I., and Giovannucci, D. R. (2012). Distinct contributions by ionotropic purinoceptor subtypes to ATP-evoked calcium signals in mouse parotid acinar cells. *J. Physiol.* 590, 2721–2737. doi: 10.1113/jphysiol.2012.228148
- Blott, E. J., and Griffiths, G. M. (2002). Secretory lysosomes. *Nat. Rev. Mol. Cell Biol.* 3, 122–131. doi: 10.1038/nrm732
- Bodin, P., and Burnstock, G. (2001). Purinergic signalling: ATP release. *Neurochem. Res.* 26, 959–969. doi: 10.1023/A:1012388618693
- Bove, P. F., Grubb, B. R., Okada, S. F., Ribeiro, C. M., Rogers, T. D., Randell, S. H., et al. (2010). Human alveolar type II cells secrete and absorb liquid in response to local nucleotide signaling. *J. Biol. Chem.* 285, 34939–34949. doi: 10.1074/jbc.M110.162933
- Brake, A. J., Wagenbach, M. J., and Julius, D. (1994). New structural motif for ligand-gated ion channels defined by an ionotropic ATP receptor. *Nature* 371, 519–523. doi: 10.1038/371519a0
- Brunschweiler, A., and Muller, C. E. (2006). P2 receptors activated by uracil nucleotides – an update. *Curr. Med. Chem.* 13, 289–312. doi: 10.2174/092986706775476052
- Burgoyne, R. D., and Morgan, A. (1998). Calcium sensors in regulated exocytosis. *Cell Calcium* 24, 367–376. doi: 10.1016/S0143-4160(98)90060-4
- Burnstock, G., Brouns, I., Adriaensen, D., and Timmermans, J. P. (2012). Purinergic signaling in the airways. *Pharmacol. Rev.* 64, 834–868. doi: 10.1124/pr.111.005389
- Burnstock, G., and Kennedy, C. (2011). P2X receptors in health and disease. *Adv. Pharmacol.* 61, 333–372. doi: 10.1016/B978-0-12-385526-8.00011-4
- Chander, A., Johnson, R. G., Reicherter, J., and Fisher, A. B. (1986). Lung lamellar bodies maintain an acidic internal pH. *J. Biol. Chem.* 261, 6126–6131.
- Clarke, C. E., Benham, C. D., Bridges, A., George, A. R., and Meadows, H. J. (2000). Mutation of histidine 286 of the human P2X<sub>4</sub> purinoceptor removes extracellular pH sensitivity. *J. Physiol.* 523(Pt 3), 697–703. doi: 10.1111/j.1469-7793.2000.00697.x
- Cockcroft, S., and Gomperts, B. D. (1979a). Activation and inhibition of calcium-dependent histamine secretion by ATP ions applied to rat mast cells. *J. Physiol.* 296, 229–243.
- Cockcroft, S., and Gomperts, B. D. (1979b). ATP induces nucleotide permeability in rat mast cells. *Nature* 279, 541–542. doi: 10.1038/279541a0
- Cockcroft, S., and Gomperts, B. D. (1980). The ATP<sub>4</sub>-receptor of rat mast cells. *Biochem. J.* 188, 789–798.
- Coddou, C., Yan, Z., Obsil, T., Huidobro-Toro, J. P., and Stojilkovic, S. S. (2011). Activation and regulation of purinergic P2X receptor channels. *Pharmacol. Rev.* 63, 641–683. doi: 10.1124/pr.110.003129
- Dell'Angelica, E. C., Mullins, C., Caplan, S., and Bonifacino, J. S. (2000). Lysosome-related organelles. *FASEB J.* 14, 1265–1278. doi: 10.1096/fj.14.10.1265
- Diamant, B., and Kruger, P. G. (1967). Histamine release from isolated rat peritoneal mast cells induced by adenosine-5'-triphosphate. *Acta Physiol. Scand.* 71, 291–302. doi: 10.1111/j.1748-1716.1967.tb03736.x
- Dietl, P., Frick, M., Mair, N., Bertocchi, C., and Haller, T. (2004). Pulmonary consequences of a deep breath revisited. *Biol. Neonate* 85, 299–304. doi: 10.1159/000078176
- Dietl, P., and Haller, T. (2005). Exocytosis of lung surfactant: from the secretory vesicle to the air-liquid interface. *Annu. Rev. Physiol.* 67, 595–621. doi: 10.1146/annurev.physiol.67.040403.102553
- Dietl, P., Haller, T., and Frick, M. (2012). Spatio-temporal aspects, pathways and actions of Ca<sup>2+</sup> in surfactant secreting pulmonary alveolar type II pneumocytes. *Cell Calcium* 52, 296–302. doi: 10.1016/j.ceca.2012.04.010
- Dietl, P., Haller, T., Mair, N., and Frick, M. (2001). Mechanisms of surfactant exocytosis in alveolar type II cells *in vitro* and *in vivo*. *News Physiol. Sci.* 16, 239–243.
- Dietl, P., Liss, B., Felder, E., Miklavc, P., and Wirtz, H. (2010). Lamellar body exocytosis by cell stretch or purinergic stimulation: possible physiological roles, messengers and mechanisms. *Cell. Physiol. Biochem.* 25, 1–12. doi: 10.1159/000272046
- Dolensek, J., Skelin, M., and Rupnik, M. S. (2011). Calcium dependencies of regulated exocytosis in different endocrine cells. *Physiol. Res.* 60(Suppl. 1), S29–S38.
- Egan, T. M., and Khakh, B. S. (2004). Contribution of calcium ions to P2X channel responses. *J. Neurosci.* 24, 3413–3420. doi: 10.1523/JNEUROSCI.5429-03.2004
- Folkesson, H. G., and Matthay, M. A. (2006). Alveolar epithelial ion and fluid transport: recent progress. *Am. J. Respir. Cell Mol. Biol.* 35, 10–19. doi: 10.1165/rcmb.2006-0080SF
- Frick, M., Bertocchi, C., Jennings, P., Haller, T., Mair, N., Singer, W., et al. (2004). Ca<sup>2+</sup> entry is essential for cell strain-induced lamellar body fusion in isolated rat type II pneumocytes. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 286, L210–L220. doi: 10.1152/ajplung.00332.2003
- Frick, M., Eschertzhuber, S., Haller, T., Mair, N., and Dietl, P. (2001). Secretion in alveolar type II cells at the interface of constitutive and regulated exocytosis. *Am. J. Respir. Cell Mol. Biol.* 25, 306–315. doi: 10.1165/ajrcmb.25.3.4493
- Garcia-Guzman, M., Soto, F., Gomez-Hernandez, J. M., Lund, P. E., and Stuhmer, W. (1997). Characterization of recombinant human P2X<sub>4</sub> receptor reveals pharmacological differences to the rat homologue. *Mol. Pharmacol.* 51, 109–118.
- Garcia-Verdugo, I., Ravasio, A., De Paco, E. G., Synguelakis, M., Ivanova, N., Kanellopoulos, J., et al. (2008). Long-term exposure to LPS enhances the rate of stimulated exocytosis and surfactant secretion in alveolar type II cells and upregulates P2Y<sub>2</sub> receptor expression. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 295, L708–L717. doi: 10.1152/ajplung.00536.2007
- Geisler, J. C., Corbin, K. L., Li, Q., Feranchak, A. P., Nunemaker, C. S., and Li, C. (2013). Vesicular nucleotide transporter-mediated ATP release regulates insulin secretion. *Endocrinology* 154, 675–684. doi: 10.1210/en.2012-1818
- Gu, J. G., and MacDermott, A. B. (1997). Activation of ATP P2X receptors elicits glutamate release from sensory neuron synapses. *Nature* 389, 749–753. doi: 10.1038/39639
- Gutierrez-Martin, Y., Bustillo, D., Gomez-Villafuertes, R., Sanchez-Nogueiro, J., Torregrosa-Hetland, C., Binz, T., et al. (2011). P2X<sub>7</sub> receptors trigger ATP exocytosis and modify secretory vesicle dynamics in neuroblastoma cells. *J. Biol. Chem.* 286, 11370–11381. doi: 10.1074/jbc.M110.139410
- Haller, T., Dietl, P., Pfaller, K., Frick, M., Mair, N., Paulmichl, M., et al. (2001). Fusion pore expansion is a slow, discontinuous, and Ca<sup>2+</sup>-dependent process regulating secretion from alveolar type II cells. *J. Cell Biol.* 155, 279–289. doi: 10.1083/jcb.200102106
- Haller, T., Dietl, P., Stockner, H., Frick, M., Mair, N., Tinhofer,

- I., et al. (2004). Tracing surfactant transformation from cellular release to insertion into an air-liquid interface. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 286, L1009–L1015. doi: 10.1152/ajplung.00342.2003
- Hayoz, S., Jia, C., and Hegg, C. (2012). Mechanisms of constitutive and ATP-evoked ATP release in neonatal mouse olfactory epithelium. *BMC Neurosci.* 13:53. doi: 10.1186/1471-2202-13-53
- Huang, Y. A., Stone, L. M., Pereira, E., Yang, R., Kinnamon, J. C., Dvoryanchikov, G., et al. (2011). Knocking out P2X receptors reduces transmitter secretion in taste buds. *J. Neurosci.* 31, 13654–13661. doi: 10.1523/JNEUROSCI.3356-11.2011
- Jacques-Silva, M. C., Correa-Medina, M., Cabrera, O., Rodriguez-Diaz, R., Makeeva, N., Fachado, A., et al. (2010). ATP-gated P2X<sub>3</sub> receptors constitute a positive autocrine signal for insulin release in the human pancreatic beta cell. *Proc. Natl. Acad. Sci. U.S.A.* 107, 6465–6470. doi: 10.1073/pnas.0908935107
- Kaczmarek-Hajek, K., Lorinczi, E., Hausmann, R., and Nicke, A. (2012). Molecular and functional properties of P2X receptors – recent progress and persisting challenges. *Purinergic Signal.* 8, 375–417. doi: 10.1007/s11302-012-9314-7
- Khakh, B. S., and North, R. A. (2006). P2X receptors as cell-surface ATP sensors in health and disease. *Nature* 442, 527–532. doi: 10.1038/nature04886
- Khakh, B. S., Proctor, W. R., Dunwiddie, T. V., Labarca, C., and Lester, H. A. (1999). Allosteric control of gating and kinetics at P2X<sub>4</sub> receptor channels. *J. Neurosci.* 19, 7289–7299.
- Kim, J. H., Nam, J. H., Kim, M. H., Koh, D. S., Choi, S. J., Kim, S. J., et al. (2004). Purinergic receptors coupled to intracellular Ca<sup>2+</sup> signals and exocytosis in rat prostate neuroendocrine cells. *J. Biol. Chem.* 279, 27345–27356. doi: 10.1074/jbc.M313575200
- Kreda, S. M., Seminario-Vidal, L., Van Heusden, C. A., O'Neal, W., Jones, L., Boucher, R. C., et al. (2010). Receptor-promoted exocytosis of airway epithelial mucin granules containing a spectrum of adenine nucleotides. *J. Physiol.* 588, 2255–2267. doi: 10.1113/jphysiol.2009.186643
- Lazarowski, E. R., Sesma, J. I., Seminario-Vidal, L., and Kreda, S. M. (2011). Molecular mechanisms of purine and pyrimidine nucleotide release. *Adv. Pharmacol.* 61, 221–261. doi: 10.1016/B978-0-12-385526-8.00008-4
- Lazarowski, E. R., Watt, W. C., Stutts, M. J., Boucher, R. C., and Harden, T. K. (1995). Pharmacological selectivity of the cloned human P2U-purinoceptor: potent activation by diadenosine tetraphosphate. *Br. J. Pharmacol.* 116, 1619–1627. doi: 10.1111/j.1476-5381.1995.tb16382.x
- Lommatzsch, M., Cicko, S., Muller, T., Lucattelli, M., Bratke, K., Stoll, P., et al. (2010). Extracellular adenosine triphosphate and chronic obstructive pulmonary disease. *Am. J. Respir. Crit. Care Med.* 181, 928–934. doi: 10.1164/rccm.200910-1506OC
- Lucattelli, M., Cicko, S., Muller, T., Lommatzsch, M., De Cunto, G., Cardini, S., et al. (2011). P2X<sub>7</sub> receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am. J. Respir. Cell Mol. Biol.* 44, 423–429. doi: 10.1165/rcmb.2010-0038OC
- Luzio, J. P., Pryor, P. R., and Bright, N. A. (2007). Lysosomes: fusion and function. *Nat. Rev. Mol. Cell Biol.* 8, 622–632. doi: 10.1038/nrm2217
- MacKenzie, A., Wilson, H. L., Kiss-Toth, E., Dower, S. K., North, R. A., and Surprenant, A. (2001). Rapid secretion of interleukin-1beta by microvesicle shedding. *Immunity* 15, 825–835. doi: 10.1016/S1074-7613(01)00229-1
- Miklavc, P., Albrecht, S., Wittekindt, O. H., Schullian, P., Haller, T., and Dietl, P. (2009). Existence of exocytotic hemifusion intermediates with a lifetime of up to seconds in type II pneumocytes. *Biochem. J.* 424, 7–14. doi: 10.1042/BJ20091094
- Miklavc, P., and Frick, M. (2011). Vesicular calcium channels as regulators of the exocytotic post-fusion phase. *Commun. Integr. Biol.* 4, 796–798.
- Miklavc, P., Frick, M., Wittekindt, O. H., Haller, T., and Dietl, P. (2010). Fusion-activated Ca<sup>2+</sup> entry: an “active zone” of elevated Ca<sup>2+</sup> during the post-fusion stage of lamellar body exocytosis in rat type II pneumocytes. *PLoS ONE* 5:e10982. doi: 10.1371/journal.pone.0010982
- Miklavc, P., Hecht, E., Hobi, N., Wittekindt, O. H., Dietl, P., Kranz, C., et al. (2012). Actin coating and compression of fused secretory vesicles are essential for surfactant secretion—a role for Rho, formins and myosin II. *J. Cell Sci.* 125, 2765–2774. doi: 10.1242/jcs.105262
- Miklavc, P., Mair, N., Wittekindt, O. H., Haller, T., Dietl, P., Felder, E., et al. (2011). Fusion-activated Ca<sup>2+</sup> entry via vesicular P2X<sub>4</sub> receptors promotes fusion pore opening and exocytotic content release in pneumocytes. *Proc. Natl. Acad. Sci. U.S.A.* 108, 14503–14508. doi: 10.1073/pnas.1101039108
- Mishra, A., Chintagari, N. R., Guo, Y., Weng, T., Su, L., and Liu, L. (2011). Purinergic P2X<sub>7</sub> receptor regulates lung surfactant secretion in a paracrine manner. *J. Cell Sci.* 124, 657–668. doi: 10.1242/jcs.066977
- Neher, E., and Sakaba, T. (2008). Multiple roles of calcium ions in the regulation of neurotransmitter release. *Neuron* 59, 861–872. doi: 10.1016/j.neuron.2008.08.019
- North, R. A. (2002). Molecular physiology of P2X receptors. *Physiol. Rev.* 82, 1013–1067.
- Novak, I. (2011). Purinergic signalling in epithelial ion transport: regulation of secretion and absorption. *Acta Physiol. (Oxf.)* 202, 501–522. doi: 10.1111/j.1748-1716.2010.02225.x
- Ohtani, M., Ohura, K., and Oka, T. (2011). Involvement of P2X receptors in the regulation of insulin secretion, proliferation and survival in mouse pancreatic beta-cells. *Cell. Physiol. Biochem.* 28, 355–366. doi: 10.1159/000331752
- Okada, S. F., Zhang, L., Kreda, S. M., Abdullah, L. H., Davis, C. W., Pickles, R. J., et al. (2011). Coupled nucleotide and mucin hypersecretion from goblet-cell metaplastic human airway epithelium. *Am. J. Respir. Cell Mol. Biol.* 45, 253–260. doi: 10.1165/rcmb.2010-0253OC
- Patel, A. S., Reigada, D., Mitchell, C. H., Bates, S. R., Margulies, S. S., and Koval, M. (2005). Paracrine stimulation of surfactant secretion by extracellular ATP in response to mechanical deformation. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 289, L489–L496. doi: 10.1152/ajplung.00074.2005
- Praetorius, H. A., and Leipziger, J. (2009). ATP release from non-excitable cells. *Purinergic Signal.* 5, 433–446. doi: 10.1007/s11302-009-9146-2
- Qu, Y., and Dubyak, G. R. (2009). P2X<sub>7</sub> receptors regulate multiple types of membrane trafficking responses and non-classical secretion pathways. *Purinergic Signal.* 5, 163–173. doi: 10.1007/s11302-009-9132-8
- Qu, Y., Franchi, L., Nunez, G., and Dubyak, G. R. (2007). Nonclassical IL-1 beta secretion stimulated by P2X<sub>7</sub> receptors is dependent on inflammasome activation and correlated with exosome release in murine macrophages. *J. Immunol.* 179, 1913–1925.
- Qureshi, O. S., Paramasivam, A., Yu, J. C., and Murrell-Lagnado, R. D. (2007). Regulation of P2X<sub>4</sub> receptors by lysosomal targeting, glycan protection and exocytosis. *J. Cell Sci.* 120, 3838–3849. doi: 10.1242/jcs.010348
- Ramsingh, R., Grygorczyk, A., Solecki, A., Cherkaoui, L. S., Berthiaume, Y., and Grygorczyk, R. (2011). Cell deformation at the air-liquid interface induces Ca<sup>2+</sup>-dependent ATP release from lung epithelial cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 300, L587–L595. doi: 10.1152/ajplung.00345.2010
- Ransford, G. A., Fregien, N., Qiu, F., Dahl, G., Conner, G. E., and Salathe, M. (2009). Pannexin 1 contributes to ATP release in airway epithelia. *Am. J. Respir. Cell Mol. Biol.* 41, 525–534. doi: 10.1165/rcmb.2008-0367OC
- Rice, W. R., and Singleton, F. M. (1987). P2Y-purinoceptor regulation of surfactant secretion from rat isolated alveolar type II cells is associated with mobilization of intracellular calcium. *Br. J. Pharmacol.* 91, 833–838. doi: 10.1111/j.1476-5381.1987.tb11282.x
- Riteau, N., Gasse, P., Fauconnier, L., Gombault, A., Couegnat, M., Fick, L., et al. (2010). Extracellular ATP is a danger signal activating P2X<sub>7</sub> receptor in lung inflammation and fibrosis. *Am. J. Respir. Crit. Care Med.* 182, 774–783. doi: 10.1164/rccm.201003-0359OC
- Rodriguez Candela, J. L., and Garcia-Fernandez, M. C. (1963). Stimulation of secretion of insulin by adenosine-triphosphate. *Nature* 197, 1210. doi: 10.1038/1971210a0
- Seminario-Vidal, L., Okada, S. F., Sesma, J. I., Kreda, S. M., Van Heusden, C. A., Zhu, Y., et al. (2011). Rho signaling regulates pannexin 1-mediated ATP release from airway epithelia. *J. Biol. Chem.* 286, 26277–26286. doi: 10.1074/jbc.M111.260562
- Shigetomi, E., and Kato, F. (2004). Action potential-independent release of glutamate by Ca<sup>2+</sup> entry through presynaptic P2X receptors elicits postsynaptic firing in the brainstem autonomic network. *J. Neurosci.* 24, 3125–3135. doi: 10.1523/JNEUROSCI.0090-04.2004
- Singer, W., Frick, M., Haller, T., Bernet, S., Ritsch-Martel, M., and Dietl, P. (2003). Mechanical forces impeding exocytotic surfactant release revealed by optical tweezers. *Biophys. J.* 84, 1344–1351. doi: 10.1016/S0006-3495(03)74950-9
- Smith, R. M., Baibakov, B., Ikebuchi, Y., White, B. H., Lambert, N. A.,

- Kaczmarek, L. K., et al. (2000). Exocytotic insertion of calcium channels constrains compensatory endocytosis to sites of exocytosis. *J. Cell Biol.* 148, 755–767. doi: 10.1083/jcb.148.4.755
- Solini, A., Chiozzi, P., Morelli, A., Fellin, R., and Di Virgilio, F. (1999). Human primary fibroblasts *in vitro* express a purinergic P2X<sub>7</sub> receptor coupled to ion fluxes, microvesicle formation and IL-6 release. *J. Cell Sci.* 112(Pt 3), 297–305.
- Sudhof, T. C. (2004). The synaptic vesicle cycle. *Annu. Rev. Neurosci.* 27, 509–547. doi: 10.1146/annurev.neuro.26.041002.131412
- Surprenant, A. (1996). Functional properties of native and cloned P2X receptors. *Ciba Found. Symp.* 198, 208–219; discussion 219–222.
- Surprenant, A., Rassendren, F., Kawashima, E., North, R. A., and Buell, G. (1996). The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X<sub>7</sub>). *Science* 272, 735–738. doi: 10.1126/science.272.5262.735
- Tatur, S., Kreda, S., Lazarowski, E., and Grygorczyk, R. (2008). Calcium-dependent release of adenosine and uridine nucleotides from A549 cells. *Purinergic Signal.* 4, 139–146. doi: 10.1007/s11302-007-9059-x
- Thompson, K. E., Korbmacher, J. P., Hecht, E., Hobi, N., Wittekindt, O. H., Dietl, P., et al. (2013). Fusion-activated cation entry (FACE) via P2X<sub>4</sub> couples surfactant secretion and alveolar fluid transport. *FASEB J.* 27, 1772–1783. doi: 10.1096/fj.12-220533
- Thorn, P. (2009). New insights into the control of secretion. *Commun. Integr. Biol.* 2, 315–317. doi: 10.4161/cib.2.4.8262
- Toyomitsu, E., Tsuda, M., Yamashita, T., Tozaki-Saitoh, H., Tanaka, Y., and Inoue, K. (2012). CCL<sub>2</sub> promotes P2X<sub>4</sub> receptor trafficking to the cell surface of microglia. *Purinergic Signal.* 8, 301–310. doi: 10.1007/s11302-011-9288-x
- Trang, T., Beggs, S., Wan, X., and Salter, M. W. (2009). P2X<sub>4</sub>-receptor-mediated synthesis and release of brain-derived neurotrophic factor in microglia is dependent on calcium and p38-mitogen-activated protein kinase activation. *J. Neurosci.* 29, 3518–3528. doi: 10.1523/JNEUROSCI.5714-08.2009
- Ulmann, L., Hatcher, J. P., Hughes, J. P., Chaumont, S., Green, P. J., Conquet, F., et al. (2008). Up-regulation of P2X<sub>4</sub> receptors in spinal microglia after peripheral nerve injury mediates BDNF release and neuropathic pain. *J. Neurosci.* 28, 11263–11268. doi: 10.1523/JNEUROSCI.2308-08.2008
- Valera, S., Hussy, N., Evans, R. J., Adami, N., North, R. A., Surprenant, A., et al. (1994). A new class of ligand-gated ion channel defined by P2x receptor for extracellular ATP. *Nature* 371, 516–519. doi: 10.1038/371516a0
- Vavra, V., Bhattacharya, A., and Zemkova, H. (2011). Facilitation of glutamate and GABA release by P2X receptor activation in supraoptic neurons from freshly isolated rat brain slices. *Neuroscience* 188, 1–12. doi: 10.1016/j.neuroscience.2011.04.067
- Wang, C. Z., Namba, N., Gonoj, T., Inagaki, N., and Seino, S. (1996). Cloning and pharmacological characterization of a fourth P2X receptor subtype widely expressed in brain and peripheral tissues including various endocrine tissues. *Biochem. Biophys. Res. Commun.* 220, 196–202. doi: 10.1006/bbrc.1996.0380
- Weinhold, K., Krause-Buchholz, U., Rodel, G., Kasper, M., and Barth, K. (2010). Interaction and interrelation of P2X<sub>7</sub> and P2X<sub>4</sub> receptor complexes in mouse lung epithelial cells. *Cell. Mol. Life Sci.* 67, 2631–2642. doi: 10.1007/s00018-010-0355-1
- Wirtz, H. R., and Dobbs, L. G. (2000). The effects of mechanical forces on lung functions. *Respir. Physiol.* 119, 1–17. doi: 10.1016/S0034-5687(99)00092-4
- Yao, C. K., Lin, Y. Q., Ly, C. V., Ohyama, T., Haueter, C. M., Moiseenkova-Bell, V. Y., et al. (2009). A synaptic vesicle-associated Ca<sup>2+</sup> channel promotes endocytosis and couples exocytosis to endocytosis. *Cell* 138, 947–960. doi: 10.1016/j.cell.2009.06.033
- Zsembery, A., Boyce, A. T., Liang, L., Peti-Peterdi, J., Bell, P. D., and Schwiebert, E. M. (2003). Sustained calcium entry through P2X nucleotide receptor channels in human airway epithelial cells. *J. Biol. Chem.* 278, 13398–13408. doi: 10.1074/jbc.M212277200

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 05 August 2013; paper pending published: 02 September 2013; accepted: 12 September 2013; published online: 08 October 2013.

Citation: Miklavc P, Thompson KE and Frick M (2013) A new role for P2X<sub>4</sub> receptors as modulators of lung surfactant secretion. *Front. Cell. Neurosci.* 7:171. doi: 10.3389/fncel.2013.00171

This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2013 Miklavc, Thompson and Frick. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.