



Hypoxia-Inducible Factor-1 α Target Genes Contribute to Retinal Neuroprotection

Lin Cheng^{1*}, Honghua Yu^{1,2}, Naihong Yan³, Kunbei Lai¹ and Mengqing Xiang^{1,4*}

¹ State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China, ² Department of Ophthalmology, General Hospital of Guangzhou Military Command of PLA, Guangzhou, China, ³ Department of Ophthalmology and Ophthalmic Laboratories, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, Chengdu, China, ⁴ Center for Advanced Biotechnology and Medicine and Department of Pediatrics, Rutgers University-Robert Wood Johnson Medical School, Piscataway, NJ, USA

Hypoxia-inducible factor (HIF) is a transcription factor that facilitates cellular adaptation to hypoxia and ischemia. Long-standing evidence suggests that one isotype of HIF, HIF-1 α , is involved in the pathogenesis of various solid tumors and cardiac diseases. However, the role of HIF-1 α in retina remains poorly understood. HIF-1 α has been recognized as neuroprotective in cerebral ischemia in the past two decades. Additionally, an increasing number of studies has shown that HIF-1 α and its target genes contribute to retinal neuroprotection. This review will focus on recent advances in the studies of HIF-1 α and its target genes that contribute to retinal neuroprotection. A thorough understanding of the function of HIF-1 α and its target genes may lead to identification of novel therapeutic targets for treating degenerative retinal diseases including glaucoma, age-related macular degeneration, diabetic retinopathy, and retinal vein occlusions.

OPEN ACCESS

Edited by:

Francesco Moccia,
University of Pavia, Italy

Reviewed by:

Teresa Pasqua,
University of Calabria, Italy
Claudio Bucolo,
University of Catania, Italy

*Correspondence:

Lin Cheng
kjade.cheng@gmail.com
Mengqing Xiang
xiangmq3@mail.sysu.edu.cn

Received: 10 July 2016

Accepted: 23 January 2017

Published: 27 February 2017

Citation:

Cheng L, Yu H, Yan N, Lai K and Xiang M (2017) Hypoxia-Inducible Factor-1 α Target Genes Contribute to Retinal Neuroprotection. *Front. Cell. Neurosci.* 11:20. doi: 10.3389/fncel.2017.00020

Keywords: HIF-1 α , hypoxia preconditioning, retina, neuroprotection, retinal degeneration


INTRODUCTION


Hypoxia or ischemia stress is generally harmful for the organism but may become beneficial under certain circumstances. Reduced oxygenation leads to induction of a number of hypoxia-responsive genes. The known hypoxia-inducible factor (HIF) targets such as erythropoietin (EPO), vascular endothelial growth factor (VEGF), heme oxygenase-1 (HO-1), adrenomedullin (ADM), glucose transporter-1 (Glut-1), etc., exert neuroprotective effects on the central nervous system (CNS; Harten et al., 2010). For instance, hypoxia preconditioning has been shown to increase

Abbreviations: HIF, Hypoxia-inducible factor; EPO, erythropoietin; VEGF, vascular endothelial growth factor; HO-1, heme oxygenase-1; ADM, adrenomedullin; Glut-1, glucose transporter-1; CNS, central nervous system; AMD, age-related macular degeneration; DR, diabetic retinopathy; RVO, retinal vein occlusion; bHLH, helix-loop-helix; PHD, prolyl-hydroxylases; VHL E3, von Hippel-Lindau E3 ubiquitin ligase; ARNT, aryl hydrocarbon receptor nuclear translocator; HREs, HIF-responsive elements; EPO-R, EPO receptors; RPE, retinal pigment epithelium; ARE/EpRE, antioxidant/electrophile response elements; eNOS, endothelial nitric oxide synthase; BRB, blood-retinal barrier; qPCR, quantitative polymerase chain reaction; RGCs, retinal ganglion cells; rhEPO, recombinant human EPO; AAVs, adeno-associated viruses; IOP, intraocular pressure; BDNF, brain-derived neurotrophic factor; IGF, insulin-like growth factor; GDNF, glial cell line-derived neurotrophic factor; rMSCs, rat mesenchymal stem cells; SI, sodium iodate; Nrf2, erythroid 2-related factor; GCL, ganglion cell layer; INL, inner nuclear layer; CNV, choroid neovascularization; rd1, retinal degeneration 1; ONL, outer nuclear layer; rds, retinal degeneration slow; TNF- α , tumor necrosis factor- α ; IL-1 β , interleukin-1 β ; PDT, photodynamic therapy; PDR, proliferative diabetic retinopathy; DME, diabetic macular edema; ROS, reactive oxygen species; siRNAs, small interfering RNAs; bFGF, basic fibroblast growth factor; iNOS, inducible nitric oxide synthase.

resistance of harmful insults by the brain (Kitagawa et al., 1990; Glazier et al., 1994; Gidday, 2006). Unlike in the brain, however, knowledge about the neuroprotective effects of HIF targets in the retina is limited. The roles of HIF targets in retinal neurogenesis and function are not well-defined and there is no consensus on their actual functions in the retina. Retinal degeneration diseases are the leading causes of blindness worldwide and some of them are closely related to ischemia (Song et al., 2003; Zheng et al., 2007; Fulton et al., 2009). Retinal hypoxia is known as the common pathogenic condition leading to vision loss, including in diseases like glaucoma, age-related macular

degeneration (AMD), diabetic retinopathy (DR), retinal vein occlusion (RVO), and some retinal degeneration secondary to stroke or Alzheimer's disease (Blanks et al., 1989; Katz et al., 1989). One isotype of HIF, HIF-1 α , is abundantly expressed in the retina (Figure 1B; Zhu et al., 2013). Under normoxia, HIF-1 α protein is rapidly degraded and its expression is hardly detectable in the adult retinal tissue (Jewell et al., 2001). Under hypoxia stress, however, HIF-1 α expression is upregulated and the protein is translocated to the nucleus to stimulate the expression of its downstream targets (Figure 2). These downstream targets such as EPO and VEGF, which are produced in a relatively low level





to stabilize the normal development or integrity of retina, are increased substantially under hypoxia to accommodate these alterations. When overexpressed, these target proteins may cause some ocular diseases. On the other hand, the HIF-1 α target genes such as EPO, HO-1, and Glut-1, have the properties to promote neurogenesis themselves. Therefore, unraveling the neuroprotective network of HIF-1 α target genes in the retina is important for understanding the mechanisms underlying ocular diseases.

HIF-1 α AND ITS TARGET GENES ARE INVOLVED IN THE PATHOPHYSIOLOGICAL ROLES OF RETINA

In humans, HIF family members include HIF-1, HIF-2, and HIF-3. They have different roles in hypoxia. HIF-1 triggers the expression of several genes to promote survival in hypoxia. HIF-2 regulates other functions such as erythropoietin production in normoxia or hypoxia (Eckardt and Kurtz, 2005) and

high-altitude adaptation (Yi et al., 2010; Hanaoka et al., 2012). HIF-3 is considered to be a negative regulator of hypoxia-inducible gene expression (Augstein et al., 2011; Heikkilä et al., 2011; Ando et al., 2013). The structures of HIFs and HIF-1 α expression in normal human nervous tissues are shown in **Figure 1**. HIF is composed of an alpha (α) and a beta (β) subunit. The α subunit belongs to the helix-loop-helix (bHLH) family of transcription factors. Under normoxia, it is hydroxylated at specific proline residues by HIF prolyl-hydroxylases (PHD), allowing its recognition and ubiquitination by the von Hippel-Lindau E3 ubiquitin ligase (VHL E3), which labels it for rapid degradation by the proteasome (Maxwell et al., 1999). The β subunit is a constitutively-expressed aryl hydrocarbon receptor nuclear translocator (ARNT). HIF α and β are constantly being produced and degraded at aerobic conditions. While under hypoxemia in the retina, only HIF-1 α persists and the HIF-1 α / β complex is translocated to the nucleus, where it binds to HIF-responsive elements (HREs) and induces the transcription of its target genes involved in oxygen delivery and energy metabolism (**Figure 2**). The target genes regulated by HIF-1 α are mainly EPO, VEGF, HO-1, ADM, and Glut-1.

EPO is the first identified gene related to hypoxia which in turn led to the discovery of HIF (Semenza and Wang, 1992). It is a glycoprotein hormone predominately secreted by kidney and liver to stimulate bone marrow erythrocyte production. EPO and EPO receptors (EPO-R) are widely expressed in different organs including brain, spinal cord, retina (mainly retinal pigment epithelium, RPE), heart, gut, kidney, muscle, lung, and testis (Erbayraktar et al., 2003; Gassmann et al., 2003; Hernandez et al., 2006). EPO is the dominant HIF-1 α target and acts as an initial factor to initiate the hypoxia condition as well as works independently without VEGF to promote neovascularization (Jaquet et al., 2002; Watanabe et al., 2005). VEGF is a HIF-1 α responsive cytokine and is the primary contributor to mitosis and the development of abnormal vessel growth and angiogenesis. In a physiological setting, endogenous VEGF, VEGF-A in particular, is required for RPE stability as well as maintenance and function of adult retina neuronal cells, especially the survival of Müller cells and photoreceptors (Nishijima et al., 2007; Saint-Geniez et al., 2008). VEGF-associated complications are extensively described in retinal angiogenesis diseases. VEGF is highly expressed under hypoxia; its expression leads to retinal vessel leakage and retinal endothelial cell proliferation and migration, thereby resulting in neovascularization. Anti-VEGF therapy (bevacizumab or ranibizumab) is largely prescribed for treatment of angiogenesis and vessel permeability in wet AMD, DR and RVO (Moravski et al., 2003; Ray et al., 2004; Nagpal et al., 2007). HO-1, one target product of HIF-1 α , is an antioxidative stress transcription factor and is regulated by antioxidant/electrophile response elements (ARE/EpRE) that scavenge heme to iron, carbon monoxide and biliverdin. HO-1 is also involved in the vasculature and angiogenesis of tumors, wounds and so on. ADM is a small, secreted multifunctional peptide expressed in various organs including the retina (Lopez and Martinez, 2002; Blom et al., 2012). ADM and its receptor are localized in neural tissues of the embryo (Montuenga et al., 1997) and are present throughout the retina of the adult mouse (Blom et al., 2012).

Lack of a functional ADM gene resulted in embryonic lethality (Caron and Smithies, 2001; Shindo et al., 2001; Iesato et al., 2013). ADM heterozygous knockout mice exhibited reduced expression of VEGF and endothelial nitric oxide synthase (eNOS) during retinal angiogenesis, and ADM blockade reduced pathological retinal angiogenesis (Iesato et al., 2013). ADM is also involved in some inflammatory and proliferative ocular diseases whose pathogenesis is ischemia (Takahashi, 2001; Udono-Fujimori et al., 2003). Its expression is upregulated in response to hypoxic preconditioning in the retina (Udono et al., 2001; Thiersch et al., 2008; Zhu et al., 2008; Ponnaluri et al., 2011). It works together with VEGF to promote angiogenic effect in ischemia (Iimuro et al., 2004). Glucose is imperative for normal functioning of retinal neurons and glial cells as they have high-energy demands. Glucose entry into the retina is a critical part for retinal normoxia. Glucose transporter-1 (Glut-1), also known as solute carrier family 2, is an integral membrane glycoprotein and a sole glucose transporter between blood and retina (blood-retinal barrier, BRB). The aerobic glycolysis in the retina is reflected by the increased blood flow metabolite of Glut-1. Glut-1 is also one of the several adaptive metabolites including glucoytic enzymes and pyruvate dehydrogenase kinase, to allow retina to better respond to the ischemia condition. In cultured Müller glial cells, quantitative polymerase chain reaction (qPCR) revealed that Glut-1 mRNA expression was 5.85- and 116-fold greater than that of Glut-3 and -4, respectively (Hosoya et al., 2008), suggesting that Glut-1 is a dominant transporter in regulating glucose utilization.

The expression of HIF-1 α target genes and the associated effects in the retina are presented in **Table 1**. Their neuroprotective role in retina, especially in the models of glaucoma, wet AMD and DR, is highlighted and discussed.

HIF-1 α AND ITS TARGET GENES IN MODELS OF RETINAL GANGLION CELL DEGENERATION

Retinal ganglion cells (RGCs), located near the inner surface of the retina, receive visual information from photoreceptors and convey visual signals to the brain via their axons. Damage to the optic nerve and RGCs lead to permanent loss of vision. RGC degeneration models mimic the diseases of glaucoma, optic neuritis, traumatic optic neuropathy injury, and etc. HIF-1 α was detected in the retina and optic nerve head of human glaucomatous eyes compared with the control eyes, and its expression was closely concordant with the location of visual field defects recorded in these eyes (Tezel and Wax, 2004). These observations provide direct evidence that tissue hypoxia is present in the retina and optic nerve head of human glaucomatous eyes, and hypoxic signaling is a component of the pathogenic mechanisms of glaucomatous neurodegeneration.

The dominant HIF-1 α target EPO is well-defined in protecting against ischemic brain injury in the CNS (Marti et al., 2000; Ehrenreich et al., 2002; Kretz et al., 2005). In the retina, endogenous EPO protects neuroretinal function in ischemic retinopathy. Böcker-Meffert et al. first reported that EPO

TABLE 1 | Effects of HIF-1 α target genes on promoting retinal neuroprotection.

HIF-1 α targets	Expression in the retina	Approximate molecular weight (in human)	Effects	References
EPO	\uparrow EPO in RPE than in neuroretina, \uparrow intravitreal EPO in diabetic patients than nondiabetic patients	34KD	Polycythemia, anti-apoptosis, neurotrophic, angiogenic effect	Garcia-Ramirez et al., 2008
VEGF-A	RPE, retinal capillaries, etc.	Monomer: 21KD; dimer: 42KD	Stimulate vasculogenesis and endothelial cell growth, promote permeabilization of blood vessels and cell migration, and inhibit apoptosis	Shima et al., 1995; Saint-Geniez et al., 2008
HO-1	Müller cells, RPE	32KD	Catalyze the degradation of heme, and therefore produce biliverdin, iron and carbon monoxide. Respond to oxidative stress, hypoxia, heavy metals, cytokines, and so on. Promote vasculature and angiogenesis	Choi and Alam, 1996; Kikuchi et al., 2005; Morse and Choi, 2005
ADM	Photoreceptor outer segments, ONL, Müller and amacrine cell somata in the INL, and some somata in the GCL	12167KD~6028.73KD, varies	Vasodilator, upregulate angiogenesis, increase tolerance to oxidative stress and hypoxic injury	Blom et al., 2012
Glut-1	RPE, choroidal, iridial and pars planus, BRB, Müller cell, lens, iris, photoreceptors	55KD	Facilitate the transport of glucose across BRB	Nihira et al., 1995
bFGF	Immature photoreceptors during postnatal development, photoreceptor outer segment/interphotoreceptor matrix complex, INL in adult retina	17.4KD	Wound healing, endothelial cell mitogen, mediate angiogenesis	Gao and Hollyfield, 1995
iNOS	Ganglion cells, INL and glial cells in diabetic eyes, microglia of the developing quail retina	150-160KD, divided into nNOS and eNOS	Synthesis of nitric oxide (NO), NO is a messenger involved in vasodilatation, neurotransmission, antimicrobial and anti-tumor activities	Abu El-Asrar et al., 2004; Sierra et al., 2014

RPE, retinal pigment epithelium; ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion cell layer; BRB, blood-retinal barrier; nNOS, neuronal nitric oxide synthase; eNOS, endothelial nitric oxide synthase.

\uparrow , increased.

could promote neural outgrowth of axotomized RGCs in a dose-dependent manner (Böcker-Meffert et al., 2002). Systemic administration of recombinant human EPO (rhEPO) before or immediately after retinal ischemia was found to not only reduce histopathological damage and promote functional recovery in electroretinography, but also decrease apoptotic neurons (Junk et al., 2002). Sullivan et al. showed that systemic administration of adeno-associated viruses (AAVs)-mediated mutant form of EPO (EPOR76E) preserved RGCs and visual function in DBA/2J glaucomatous mice. The rescued RGCs retained their axonal projections within the optic nerve and the hematocrit did not exceed normal limits (Sullivan et al., 2011). Retrobulbar administration of EPO also protected RGCs from acute elevated intraocular pressure (IOP; Zhong et al., 2008) and axotomy (Kilic et al., 2005). The survival rate of the RGCs 14 days after axotomy lesion was markedly increased from about 20% to more than 60% (Kilic et al., 2005), which surpasses the neuroprotection efficacy of classical neurotrophins, such as brain-derived neurotrophic factor (BDNF; Klocker et al., 2000), insulin-like growth factor-1 (IGF-1; Kermer et al., 2000) or glial cell line-derived neurotrophic factor (GDNF; Kilic et al., 2004). Consistent with these notions, systemic applied recombinant EPO protected EPO-deficient

mice against profound retinal ischemia damage (Mowat et al., 2012) in RGCs (Junk et al., 2002). And subretinal transplantation of EPO gene-modified rat mesenchymal stem cells (rMSCs) worked better than rMSCs alone for sodium iodate (SI)-induced retinal degeneration in rats (Guan et al., 2013). Taken together, the neuroprotective role of EPO in RGCs is clear and is mainly accomplished by its anti-apoptosis (Chung et al., 2009), neurotrophic, and angiogenic effects.

VEGF, another HIF-1 α target, was depicted to have protective roles for non-vascular cells in various models (Yang and Cepko, 1996; Coultas et al., 2005; Maharaj and D'Amore, 2007; Emerich et al., 2010). For instance, some isotypes of VEGF displayed neuroprotective properties *in vitro* (Jin et al., 2000). *In vivo* studies showed that endogenous VEGF-A165b isoform exerted neuroprotective effects in response to glutamatergic excitotoxicity on RGCs during rat retinal ischemia-reperfusion injury, through activation of VEGFR-2 and MEK1/2 pathways and inhibition of caspase-3 (Beazley-Long et al., 2013). VEGF-A promoted glaucomatous RGC survival via VEGFR-2, thereby highlighting the potential risks associated with using VEGF-A antagonists in ocular diseases (Foxton et al., 2013). Anti-VEGF reduced the potential benefit of VEGF in neuroprotection and

exacerbated RGC apoptosis (Nishijima et al., 2007). Similar to EPO, VEGF protects axotomized RGCs from degeneration through ERK1/2 and Akt pathways (Böcker-Meffert et al., 2002; Kilic et al., 2006). Ischemic preconditioning prior to ischemia-reperfusion injury showed increased VEGF-A levels. And the VEGF-A levels substantially decreased in a dose-dependent manner following RGC apoptosis. Despite the overwhelming evidence indicating a role for VEGF in neuroretinal protection, a few groups did not observe any damage to retinal photoreceptors or RGCs after blocking VEGF or its receptors (Miki et al., 2010; Sobaci et al., 2013; Demirel et al., 2015). These discrepancies suggest a possibility that the neuroprotective function exerted by VEGF may be conditional and context-dependent. Therefore, further studies to clarify the neurotrophic role of VEGF in RGCs is warranted.

Oxidative stress is considered to be an early event of various retinal diseases (Liu et al., 2007). And indeed, HO-1 exerts a protective effect on ocular diseases. Previous findings suggested that functional human HO-1 could prevent RGC death in the adult rat retina after pressure-induced ischemia (Hegazy et al., 2000). The nuclear factor erythroid 2-related factor (Nrf2)/HO-1-antioxidant pathway was activated in ischemia-reperfusion-induced rat retinal damage, and this activation led to better preservation of ganglion cell layer (GCL) and inner nuclear layer (INL; Varga et al., 2013; He et al., 2014). Nrf2 and HO-1 were also increased in attenuated GCL damage following limb remote ischemic conditioning of the retina (Zhang X. et al., 2014). Overexpression of HO-1 in RGCs facilitated the survival of some RGCs for up to 7 days after optic nerve crush (Himori et al., 2014). AAV-mediated HO-1 gene transfer (AAV-HO-1) into the vitreous promoted RGC survival following ischemia/reperfusion-induced damage, while inhibiting HO-1 counteracted the effect in rats (Peng et al., 2008). Stimulation of Nrf2/HO-1 axis could be of interest for the treatment of retinal degeneration such as glaucoma, AMD, and DR which will be discussed below (Foresti et al., 2013, 2015). Furthermore, increased ADM was found in the vitreous humor of glaucoma (Evereklioglu et al., 2002), DR (Ito et al., 2003; Shaw et al., 2006; Weng et al., 2006), uveitis, and vitreoretinal patients (Udono et al., 2002), and in the plasma of retinitis pigmentosa individuals (Vingolo et al., 2005). Glucose deprivation may result in an increased ability to protect RGCs from glutamate-induced excitotoxicity (Toft-Kehler et al., 2014), but the function of Glut-1 was not tested in the RGC degeneration models.

The efficacy of HIF-1 α and its target genes in protecting against RGC degeneration may open new perspectives for the treatment of glaucoma, optic neuritis, and traumatic optic neuropathy injuries. Developing new treatment targeting on HIF-1 α target genes for RGC degeneration deserves concerted efforts in the near future.

HIF-1 α AND ITS TARGET GENES IN MODELS OF PHOTORECEPTOR DEGENERATION AND WET AMD

The pathogenesis of AMD is not well-known. The wet AMD, also known as neovascular or exudative AMD, causes central

vision loss due to RPE atrophy, neovascularization, and death of photoreceptors. Oxidative stress was implicated in the senescence of RPE cells and the pathogenesis of AMD. Development of choroid neovascularization (CNV), which is mediated mainly by HIF-1 α and VEGF, is the most significant threat for AMD; therefore, inhibiting HIF-1 α and VEGF is often used for treating AMD. However, it was found that focused activation of HIF transcription factors in normoxic photoreceptors resulted in a transient protection of rods against light damage (Lange et al., 2011). Pyruvate application, which stabilized HIF-1 α and HIF-2 α , protected the mouse retina against white light damage (Ren et al., 2011).

Unlike HIF-1 α , the effect of EPO on protecting photoreceptors and RPE cells is uncontroversial. EPO is generally regarded as a potent neuroprotective factor for photoreceptors (Becerra and Amaral, 2002). In the adult mammalian retina, systemically applied EPO was protective against light-induced photoreceptor apoptosis (Becerra and Amaral, 2002; Grimm et al., 2002). It stabilized the retinal vasculature and inhibited the development of focal vascular lesions in photoreceptor degeneration (Shen et al., 2014). Subretinal delivery of EPO protected against light-induced and genetic photoreceptor degeneration (Zhang et al., 2008; Rex et al., 2009; Colella et al., 2011; Busch et al., 2014) through the ERK1/2 and Akt pathway (Shen et al., 2010). An earlier report revealed that by either transgenic or systemic applications, EPO protected against apoptotic cell death during acute, light-induced photoreceptor cell death but not in genetically based retinal degeneration such as in retinal degeneration 1 (rd1) or dominant retinitis pigmentosa mouse models (Grimm et al., 2004). Findings from a study demonstrated that subretinal injection of EPO AAVs resulted in approximately an 8 μ m thicker outer nuclear layer (ONL) in retinas of retinal degeneration slow (rds) mice compared to the control group (Hines-Beard et al., 2013). Nevertheless, one may argue that the neuroprotective effect of EPO is short-lived and the ONL may degenerate without a repeated hypoxia preconditioning (Grimm et al., 2005). For RPE cells, EPO, and EPO-R are highly enriched in light-induced apoptotic RPE cells (Chung et al., 2009). EPO protected RPE from oxidative damage by decreasing inflammatory cytokines of tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β), and inhibiting caspase-3-like activity (Gawad et al., 2009; Wang et al., 2009).

Anti-VEGF agents are widely used in the treatment of choroidal neovascularization for AMD. However, long-term VEGF inhibition could have deleterious effects on the remaining healthy retina given the well-documented role of VEGF in normal development of the retinal vasculature (Hernandez and Simo, 2012). Non-isoform-specific inhibition of VEGF for treating angiogenesis can constrain the neuroprotective role of some isoforms of VEGF and damage retinal and sensory neurons. Inhibiting VEGF following photodynamic therapy (PDT) which causes elevated VEGF resulted in photoreceptor apoptosis, suggesting that VEGF after PDT is neuroprotective (Suzuki et al., 2011). Depletion of VEGF-A specifically in neural progenitor cells resulted in thinner retina and aberrant cortex development (Haigh et al., 2003).

Expression of HO-1 in RPE is essential for the survival of photoreceptors (Satarug et al., 2008) and increases following repetitive hypoxia preconditioning in the retina (Zhu et al., 2007). HO-1 is also a sensitive marker for light-induced insult in the retina (Kutty et al., 1995). An increased expression of HO-1 is thought to be a cytoprotectant against light-induced oxidative damage in the retina (Kutty et al., 1995). HO-1 overexpression protected photoreceptors from cellular damage caused by intense light exposure (Sun et al., 2007) and protected RPE cells against the toxic effect induced by high glucose and oxidative/nitrosative stress (Castilho et al., 2012). RPE cells derived from neovascular AMD patients displayed much higher HO-1 and HO-2 antigen levels compared to those from younger individuals, suggesting that HO has protective mechanisms against oxidation (Frank et al., 1999).

ADM, another HIF-1 α target, is the first neurotransmitter/neuromodulator found in the RPE (Udono et al., 2000, 2002). It can promote hypoxia vasodilation in retinal arteries (Maenhaut et al., 2007). Additionally, ADM suppressed the migration, proliferation and tube formation of human RPE cells under hypoxia (Chen et al., 2014). Administration of ADM in mice inhibited macrophage migration from RPE to prevent choroidal neovascularization (Yuda et al., 2012). These results were in agreement with a neuroprotective role for ADM in ischemic brain injury (Willis et al., 1996; Dogan et al., 1997; Xia et al., 2004, 2006; Miyashita et al., 2006; Igarashi et al., 2014; Zhang H. et al., 2014).

The above evidence suggests that HIF-1 α and its target genes can act as a retinal shield against photoreceptor degeneration and wet AMD injuries in the retina. Further studies are required to clarify the role of these transcription factors in retinal pathology to determine whether they can be exploited as potential therapeutic targets in treating photoreceptor degeneration and wet AMD.

HIF-1 α AND ITS TARGET GENES IN MODELS OF DIABETIC RETINOPATHY

Elevated HIF-1 α and VEGF are generally accepted to be harmful for diabetic retinopathy, despite some studies found that anti-VEGF in DR rats had detrimental effects on neuronal cells (RGC, amacrine and bipolar cells) which are present in the inner layers of the retina (Park et al., 2014). The situation is different for the HIF-1 α targets such as EPO, HO-1, ADM, and Glut-1. Strikingly high levels of intravitreal EPO have been found in both proliferative diabetic retinopathy (PDR) and diabetic macular edema (DME) patients (Hernandez et al., 2006). EPO-R is upregulated in DR neurosensory retina in response to diabetic stress (Junk et al., 2002; Zhang et al., 2008). The upregulation of EPO/EPO-R leads to a maintenance-survival mechanism thereby adapting to insulin increase in early DR. Intravitreal injection of EPO induced downregulation of EPO-R, VEGF, and

VEGF receptor in streptozocin-induced DR rats (Mitsuhashi et al., 2013). In DR, EPO-derived peptide and EPO helix-B domain significantly protected against neuroglial and vascular degeneration without altering hematocrit or exacerbating neovascularization or thrombosis in diabetic rats (McVicar et al., 2011). Administering suberythropoietic EPO is also vaso- and neuro-protective in experimental early DR both *in vivo* and *in vitro* (Zhang et al., 2008; Wang et al., 2011; Mitsuhashi et al., 2013).

HO-1 was found to be increased in the retina of db/db DR mice and cultured retinal explants (He et al., 2013). It was reported that HIF-1 α -mediated, long-lasting HO-1 elevation contributed to long-term retinal ischemia tolerance (Zhu et al., 2007). Induction of HO-1 by compounds or oxidative stress reduced damage in the retina (Mandal et al., 2009; Kim et al., 2012; Chao et al., 2013; Rappoport et al., 2013; Koskela et al., 2014; Lee et al., 2014) and decreased retinal microvascular complications in DR (Gerald et al., 2008) through the IRS1/PI3K/Akt2 pathway (Cukiernik et al., 2003). Overexpression of HO-1 in the retina restored visual function in diabetic rat models (Gerald et al., 2008; Shyong et al., 2008) and systemic hypoxia mice (Zhu et al., 2007). Furthermore, diminished HO-1 expression in RPE cells was found in diabetic patients (da Silva et al., 1997). HO-1 also exerts anti-inflammatory, anti-apoptotic and anti-proliferative effects through Nrf2/ERK-related signaling (Peng et al., 2011; Fan et al., 2012). It could drive the resolution of inflammation by reducing macrophage infiltration in the ischemia-reperfusion injured retina (Sun et al., 2010). Other *in vitro* experiments supported a similar anti-inflammatory role for HO-1 (Willis et al., 1996). Inhibition of upregulated HO-1 in Müller cells resulted in increased infiltration of inflammatory cells and destruction of the retina in ischemia-reperfusion injury rats (Ulyanova et al., 2001; Arai-Gaun et al., 2004). Inhibition of HO-1 also enhanced reactive oxygen species (ROS) production and the toxic effect (Castilho et al., 2012). These findings have revealed that HO-1 is cytoprotective by helping the cells to adapt to the oxidative stress environment.

For ADM, its plasma level was lower in juvenile type 1 diabetes patients without retinopathy than that in healthy subjects (Semeran et al., 2013). Glut-1, as indicator of glucose transportation in the retina, played a big role in the pathophysiology of DR. In the human and monkey, Glut-1 is diffusely expressed throughout the whole retina, including the BRB cells, RPE, vascular endothelium, and rod and cone inner and outer segments (Nihira et al., 1995). Elevated levels of Glut-1 increased the ability of retina to utilize glucose for normal metabolic activity (Schubert, 2005). Knockdown or blockade of Glut-1 by small interfering RNAs (siRNAs) significantly reduced retinal glucose in diabetic mice (Lu et al., 2013). In contrast, some studies suggested that inhibiting Glut-1 could prevent expression of early biomarkers of DR (Lu et al., 2013) and that Glut-1 was downregulated in oxidative RPEs (Fernandes et al., 2011). Expression of Glut-1 in the microvasculature does not account for the glucose accumulation in diabetic retina (Puchowicz et al., 2004). Therefore, the

TABLE 2 | Selected reports of HIF-1 α target genes involved in retinal neuroprotection.

HIF-1 α targets	Interventions	Observed objects	Findings	Conclusions	References
EPO	Systemic administration of recombinant human EPO (rhEPO) Treat with EPO and VEGF <i>In vitro</i> culture under glycosylated insult Damage from ROS	Transient global retinal ischemia induced by raising IOP Retinal explants from postnatal rats Chemical-induced insults in primary retinal neurons Oxidant-treated cultured human RPE cells	\uparrow EPO-R in the retina. EPO with soluble EPO-R exacerbated ischemic injury. \downarrow histopathological damage, \uparrow functional recovery in ERG EPO and VEGF improved neurite outgrowth of RGCs EPO was shown to be antiapoptotic, \uparrow Bcl-XL and p-BAD, \downarrow Bax EPO \uparrow RPE cells viability, \downarrow inflammatory cytokines TNF- α and IL-1 β , \downarrow cell DNA fragmentation and membrane phosphatidylserine exposure, \downarrow ROS, \downarrow caspase-3 \downarrow p-STAT-5, Bcl-XL in RGCs, \uparrow p-ERK-1/2 and p-Akt, \downarrow caspase-3	Exogenous EPO has an antiapoptotic mechanism of action; EPO is as a neuroprotective agent in acute neuronal ischemic injury EPO and VEGF have a significant and specific biological effect on neurite regrowth of axotomized RGCs; EPO and VEGF have a neuroprotective and neuroregenerative role on RGCs in ischemic retina EPO/EPO-R acts through ERK-1/2 and Akt pathways EPO protects against oxidative injury-induced cell death and mitochondrial dysfunction in RPE cells through modulation of p-Akt1 and mitochondrial membrane potential	Junk et al., 2002 Böcker-Meffert et al., 2002 Shen et al., 2010 Wang et al., 2009
	Optic nerve transection EPO single intravitreal injection	<i>In vivo</i> retrograde degeneration of RGCs in mouse line tg21 that constitutively expresses human EPO Diabetic Sprague-Dawley rats	\uparrow p-STAT-5, Bcl-XL in RGCs, \uparrow p-ERK-1/2 and p-Akt, \downarrow caspase-3 \uparrow EPO-R in the neurosensory retina, improvement of photoreceptor survival, but endogenous EPO in neurosensory retina was unchanged, activation of the ERK but not the STAT-5 pathway \downarrow caspase-3, \uparrow Bcl-XL, anti-apoptosis of photoreceptors; \uparrow p-JAK2, p-Akt and p-ERK-1/2 by 400ng EPO treatment \downarrow caspase-1, \uparrow EPO-R in photoreceptors; stabilized the retinal vasculature, \downarrow photoreceptor apoptosis, \uparrow CD34 ⁺ cells into the retina	Predict clinical implementation of recombinant human EPO not only in patients with acute ischemic stroke but also with more delayed degenerative neurological diseases EPO/EPO-R is a maintenance-survival mechanism of retinal neurons; responds to the insults of early diabetes other than ischemia; Intravitreally injection of EPO in early diabetes may prevent retinal cell death and protect the BRB function	Kilic et al., 2005 Zhang et al., 2008
	EPO intravitreal injection Acute hypoxia-induced EPO; systematic administration of EPO	Photoreceptor degeneration in retinal detachment rat Photoreceptor degeneration by light and surgery in mice or rats	Photoreceptor degeneration in retinal detachment rat Photoreceptor degeneration by light and surgery in mice or rats	Intravitreal injection of 400ng EPO is safe and photoreceptor-protective; EPO may activate PI3K/Akt and MAPK/ERK-1/2 pathways EPO protects photoreceptors through \downarrow p75NTR-pro-NT3 signaling, \uparrow production and mobilization of bone marrow derived cells	Xie et al., 2012a,b Grimm et al., 2002; Shen et al., 2014
VEGF	Intravitreal anti-VEGF-A antibody injection Axotomy	Streptozotocin-induced diabetic rat retina A transgenic mouse line that constitutively expresses human VEGF	\uparrow RGC death, novel apoptosis in amacrine and bipolar cells, \downarrow p-Akt RGCs of VEGF- transgenic mice were protected against delayed degeneration after axotomy; \uparrow p-ERK-1/2, \uparrow p-Akt, \downarrow p38, \downarrow caspase-3	The p-Akt pathway, which plays a neuroprotective role via VEGF, was significantly affected by VEGF inhibition; \downarrow VEGF may have detrimental effects on neuronal cells VEGF exerts neuroprotection by dual activation of ERK-1/2 and Akt pathways	Park et al., 2014 Kilic et al., 2006

(Continued)

TABLE 2 | Continued

HIF-1 α targets	Interventions	Observed objects	Findings	Conclusions	References
	VEGF-A, VEGF-A165b treatment	Rat glaucoma model or ischemia-reperfusion injury in rats or <i>in vitro</i> retina culture or RGC insults in rat retinal ischemia-reperfusion injury	Dose-dependent \downarrow in retinal neuron apoptosis, \uparrow VEGFR-2; VEGF-A acts directly on RGC to promote survival, VEGFR-2 signaling via the pathway of phosphoinositide-3-kinase/Akt; VEGF-A protects RGC via VEGFR-2; neuroprotective through activation of VEGFR-2 and MEK-1/2, not via p38 activation, \downarrow caspase-3	Antagonism of VEGF-A function presents a risk to neuronal survival; VEGF-A165b may be therapeutically useful for pathologies that involve neuronal damage, non-isoform-specific inhibition of VEGF-A may be damaging to retinal and sensory neurons	Nishijima et al., 2007; Beazley-Long et al., 2013; Foxton et al., 2013
	siRNA-based \downarrow VEGF	Müller cells and photoreceptors	\downarrow INL and ONL thickness, \downarrow retinal function in ERG	Endogenous VEGF is required for visual function	Saint-Geniez et al., 2008
HO-1	\uparrow HO-1 in photoreceptors by AAV gene subretinal injection	Light-injured Sprague-Dawley rats	Partially preserved retina structure and attenuated apoptosis in photoreceptors, \downarrow c-fos, \downarrow p53, \uparrow p38, \uparrow bcl-2, \uparrow c-FLIP	The anti-apoptotic mechanisms of HO-1 may be related to \uparrow p38, bcl-2 and c-FLIP and \downarrow c-fos and p53	Sun et al., 2007
	Insulin-induced HO-1 treatment	Bovine retinal endothelial cells (BREC) and pericyte cells (BRPC) from fresh calf eyes	Insulin-induced HO-1 through PI3-kinase/Akt pathway without affecting ERK and p38 MAPK; insulin regulated HO-1 expression via IRS1 and Akt2 pathways; \downarrow INF- α B, \downarrow caspase-8 and apoptosis via the IRS1/PI3K/Akt2/HO-1 pathway	Insulin activates HO-1 expression via IRS1/PI3K/Akt2 signaling	Geraldes et al., 2008
ADM	Culture under normoxic or hypoxic conditions	RPE cells	Hypoxia \uparrow ADM in all three human RPE cell lines, ADM treatment \downarrow the hypoxia-induced cell number decrease	ADM induced by hypoxia protects cell damage in RPE cells	Udono et al., 2001
	Oxygen-induced retinopathy	Heterozygous KO mouse of ADM [ADM(+/-)] and its receptors, inducible endothelial cell-specific RAMP2 KO mouse line [DI-E-RAMP2(-/-)]	\downarrow VEGF and eNOS in ADM(+/-) retinas, DI-E-RAMP2(-/-) showed abnormal retinal vascular patterns in the early stages of development, ADM enhanced the proliferation and migration of RPE cells, intravitreal injection of anti-ADM antibody \downarrow pathological retinal angiogenesis	The ADM-RAMP2 system is crucially involved in retinal angiogenesis; ADM and its receptor system are potential therapeutic targets for controlling pathological retinal angiogenesis	Iesato et al., 2013
Glut-1	Knockdown Glut-1 by siRNA and systemic administration of Glut-1 inhibitor	Diabetic mice	\downarrow Retinal glucose by \downarrow Glut-1; Glut-1 inhibitors reduced glucose and glycohemoglobin levels in RBC, prevented early biomarkers of DR including superoxide radicals, chaperone protein β 2 crystallin and VEGF	Anti-Glut-1 treatment is a promising therapeutic target for preventing DR	Lu et al., 2013

(Continued)

TABLE 2 | Continued

HIF-1 α targets	Interventions	Observed objects	Findings	Conclusions	References
	Diabetics induced by streptozotocin	Streptozotocin-induced diabetic rats vs. nondiabetic rats (normoglycemic and acute-hyperglycemic)	Retinal glucose influx in the diabetic rats was lower than in the nondiabetic acute-hyperglycemic group, but not in the normoglycemic group. \uparrow glucose in the diabetic retina than the nondiabetic retina	The accumulation of glucose in the diabetic retina cannot be explained by increased endothelial-glucose uptake (Glut-1)	Puchowicz et al., 2004
bFGF	Genetic degeneration of photoreceptors	Photoreceptor degeneration rd mice	\uparrow bFGF in the outer retina during photoreceptor degeneration, weakly present in some cells in the INL	Neuronal degeneration is accompanied by \uparrow bFGF in degenerating neurons prior to cell death	Gao and Hollyfield, 1995
iNOS	Ischemia preconditioning	WT mice vs. WT mice treated with NOS inhibitor before preconditioning vs. iNOS/eNOS/mNOS KO mice	Ischemic tolerance was not achieved in the retinas of NOS KO mice; NOS inhibitor to WT mice blocked the development of ischemic tolerance	NO derived from both eNOS and nNOS is a required molecular signal in the adaptive response to ischemic preconditioning in the retina	Zhu et al., 2006

ERG, electroretinography; Bcl-XL, B-cell lymphoma-extra large; p, phosphorylation; BAD, Bcl2-associated agonist of cell death; Bax, Bcl2-associated x protein; ERK, extracellular signal-regulated kinase; Akt, Akt serine/threonine kinase; STAT, signal transducer and activator of transcription; JAK2, Janus kinase 2; PI3K, phosphoinositide 3-kinase; MAPK or MEK, mitogen-activated protein kinase; p75NTR-pro-NT3, p75 neurotrophin receptor-pro-mature neurotrophin-3; c-fos, fos proto-oncogene; bcl-2, B-cell lymphoma-2; c-FLIP cellular FLICE (FADD-like IL-1 β -converting enzyme)-inhibitory protein; IRS1, insulin receptor substrate 1; NF- κ B, nuclear factor- κ B; Di-E-RAMP2(-/-), drug-inducible vascular endothelial cell-specific RAMP2 knockout (-/-) mice; RAMP2, receptor activity modifying protein 2; RBC, red blood cells; WT, wildtype; KO, knockout; iNOS, inducible nitric oxide synthase; eNOS, endothelial nitric oxide synthase; nNOS, neuronal nitric oxide synthase; \uparrow , promote or increase or induce; \downarrow , inhibit or decrease or suppress. rd mouse is a spontaneous mouse mutant in which photoreceptors degenerate shortly after birth.

neuroprotective function of Glut-1 in the retina needs to be further investigated.

DISCUSSION

Some other HIF-1 α targets such as basic fibroblast growth factor (bFGF; Cuevas et al., 1998; Yu et al., 2004; Valter et al., 2005), inducible nitric oxide synthase (iNOS; Sakamoto et al., 2006; Zhu et al., 2006; Sun et al., 2010), etc. which are less reported in retinal neuroprotection are not discussed here. The selected reports of HIF-1 α target genes on retinal neuroprotection are summarized in Table 2.

HIF-1 α together with its target genes are essential for retinal development, vasculature stability, proper retinal function, and vision maintenance. Hypoxia pre-conditioning mimics the physiopathological responses of the retina to provide protection for neuronal cells. PHD inhibitors stabilize HIF transcription factors, especially HIF-1 α , to mediate genetic adaptation to hypoxia. Under hypoxia, the neuroprotective effect of HIF-1 α targets might be multimodal, including promotion of oxygen and glucose supply, neovascularization, antioxidization, anti-inflammation, anti-apoptosis, and neurotrophs.

Nevertheless, the HIF-1 α target gene therapy may be a double-edged sword (Grimm and Willmann, 2012). High-dose toxicity and longtime exposure may over-activate HIF-1 α and bring about severe consequences in terms of side effects. Several lines of evidence have suggested that HIF-1 α overexpression can contribute to retinal pathologies, including retinopathy of prematurity, DR, AMD, glaucoma, and high altitude retinopathy (Arjamaa and Nikinmaa, 2006; Willmann et al., 2013). It may also promote neovascularization, oncogenesis, and metastasis. New method seeking to manipulate the HIF-1 α target genes to treat various ocular diseases as well as to reduce the side effects are warranted for new therapeutic avenues. Take the dominant HIF-1 α target EPO as an example. Accumulating evidence makes EPO particularly attractive in preventing retinal degeneration in the early stages of retinal diseases. However, before applying EPO for retinal degenerative diseases, two critical points need to be taken into account. The first is the EPO form. Longtime usage and overdose of EPO lead to chronic polycythemia, which is a lethal side effect (Gassmann et al., 2003). Engineering an effective package for neuroprotective EPO and attenuating erythropoietic and angiogenic activity is the key for such a therapy. Secondly, timing and dose are crucial for EPO treatment of retinal diseases. It has been shown that intraocular delivery of EPO has a lower efficiency than when administered systemically. This suggests that hormone systemic effects contribute to progressive neuroprotection (Colella et al., 2011). Furthermore, high levels of EPO is not protective due to its bell-shaped dose curve (Hines-Beard et al., 2013), and the neuroprotective dose is higher than that required for erythropoiesis (Coleman and Brines, 2004). Early retinal EPO replenishment improves retinal vascular stability, but elevated EPO levels during the proliferation stage contribute to neovascularization and ocular diseases (Coleman and Brines, 2004; Chen et al., 2008). Therefore, early administration of EPO at a proper dose for retinal

degeneration is important, for it can activate endogenous EPO-R in the affected tissue through stimulation of potent ischemic preconditioning. The treating course and strategy should be tailored according to patients' responses. Continuous delivery of EPO may lead to elevated local EPO and cause insults. Overall, timing, dose and route of administration are the important factors to be considered for balancing favorable vs. detrimental effects during EPO retinal therapy. Similar problems may be seen in exploiting other HIF-1 α targets for treating retinal degeneration.

HIF-1 α being a "master switch" for regulating all oxygen-dependent retinal diseases is critical in construction of new therapeutic avenues for treating retractable retinal insults. Additional preclinical experiments need to be conducted in order to elucidate the physiological and neuroprotective role of HIF-1 α targets in the retina. A more refined understanding of the complex functions of HIF-1 α and target genes will help to manipulate these genes and define new and more-specific targets for retinal degenerative diseases.

REFERENCES

- Abu El-Asrar, A. M., Meersschaert, A., Dralands, L., Missotten, L., and Geboes, K. (2004). Inducible nitric oxide synthase and vascular endothelial growth factor are colocalized in the retinas of human subjects with diabetes. *Eye* 18, 306–313. doi: 10.1038/sj.eye.6700642
- Ando, H., Natsume, A., Iwami, K., Ohka, F., Kuchimaru, T., Kizaka-Kondoh, S., et al. (2013). A hypoxia-inducible factor (HIF)-3 α splicing variant, HIF-3 α 4 impairs angiogenesis in hypervascular malignant meningiomas with epigenetically silenced HIF-3 α 4. *Biochem. Biophys. Res. Commun.* 433, 139–144. doi: 10.1016/j.bbrc.2013.02.044
- Arai-Gaun, S., Katai, N., Kikuchi, T., Kurokawa, T., Ohta, K., and Yoshimura, N. (2004). Heme oxygenase-1 induced in muller cells plays a protective role in retinal ischemia-reperfusion injury in rats. *Invest. Ophthalmol. Vis. Sci.* 45, 4226–4232. doi: 10.1167/jovs.04-0450
- Arjamaa, O., and Nikinmaa, M. (2006). Oxygen-dependent diseases in the retina: role of hypoxia-inducible factors. *Exp. Eye Res.* 83, 473–483. doi: 10.1016/j.exer.2006.01.016
- Augstein, A., Poitz, D. M., Braun-Dullaeus, R. C., Strasser, R. H., and Schmeisser, A. (2011). Cell-specific and hypoxia-dependent regulation of human HIF-3 α : inhibition of the expression of HIF target genes in vascular cells. *Cell. Mol. Life Sci.* 68, 2627–2642. doi: 10.1007/s00018-010-0575-4
- Beazley-Long, N., Hua, J., Jehle, T., Hulse, R. P., Dersch, R., Lehring, C., et al. (2013). VEGF-A165b is an endogenous neuroprotective splice isoform of vascular endothelial growth factor A *in vivo* and *in vitro*. *Am. J. Pathol.* 183, 918–929. doi: 10.1016/j.ajpath.2013.05.031
- Becerra, S. P., and Amaral, J. (2002). Erythropoietin—an endogenous retinal survival factor. *N. Engl. J. Med.* 347, 1968–1970. doi: 10.1056/NEJMcibr022629
- Blanks, J. C., Hinton, D. R., Sadun, A. A., and Miller, C. A. (1989). Retinal ganglion cell degeneration in Alzheimer's disease. *Brain Res.* 501, 364–372. doi: 10.1016/0006-8993(89)90653-7
- Blom, J., Giove, T. J., Pong, W. W., Blute, T. A., and Eldred, W. D. (2012). Evidence for a functional adrenomedullin signaling pathway in the mouse retina. *Mol. Vis.* 18, 1339–1353.
- Böcker-Meffert, S., Rosenstiel, P., Röhl, C., Warneke, N., Held-Feindt, J., Sievers, J., et al. (2002). Erythropoietin and VEGF promote neural outgrowth from retinal explants in postnatal rats. *Invest. Ophthalmol. Vis. Sci.* 43, 2021–2026.
- Busch, S., Kannt, A., Kolibabka, M., Schlotterer, A., Wang, Q., Lin, J., et al. (2014). Systemic treatment with erythropoietin protects the neurovascular unit in a rat model of retinal neurodegeneration. *PLoS ONE* 9:e102013. doi: 10.1371/journal.pone.0102013

AUTHOR CONTRIBUTIONS

LC and MX conceived the review. LC wrote the manuscript. HY, NY, and KL joined discussions and revised the paper. MX made critical revision and language editing. The manuscript has been reviewed and approved by all named authors.

ACKNOWLEDGMENTS

We thank Dr. Sumit Bhattacharya at Department of Neurosciences, University of Toledo for language editing. This work was supported in part by the Fundamental Research Funds of the State Key Laboratory of Ophthalmology in Sun Yat-sen University, National Basic Research Program (973 Program) of China (2015CB964600), National Natural Science Foundation of China (81670862, 81603200), New Jersey Health Foundation, China Postdoctoral Science Foundation (2015M580758), and Natural Science Foundation of Guangdong Province (2016A030310201).

- Caron, K. M., and Smithies, O. (2001). Extreme hydrops fetalis and cardiovascular abnormalities in mice lacking a functional Adrenomedullin gene. *Proc. Natl. Acad. Sci. U.S.A.* 98, 615–619. doi: 10.1073/pnas.98.2.615
- Castilho, Á., Aveira, C. A., Leal, E. C., Simões, N. F., Fernandes, C. R., Meirinhos, R. I., et al. (2012). Heme oxygenase-1 protects retinal endothelial cells against high glucose- and oxidative/nitrosative stress-induced toxicity. *PLoS ONE* 7:e42428. doi: 10.1371/journal.pone.0042428
- Chao, H. M., Chuang, M. J., Liu, J. H., Liu, X. Q., Ho, L. K., Pan, W. H., et al. (2013). Baicalein protects against retinal ischemia by antioxidation, antiapoptosis, downregulation of HIF-1 α , VEGF, and MMP-9 and upregulation of HO-1. *J. Ocul. Pharmacol. Ther.* 29, 539–549. doi: 10.1089/jop.2012.0179
- Chen, J., Connor, K. M., Aderman, C. M., and Smith, L. E. (2008). Erythropoietin deficiency decreases vascular stability in mice. *J. Clin. Invest.* 118, 526–533. doi: 10.1172/jci33813
- Chen, Z., Liu, G., Xiao, Y., and Lu, P. (2014). Adrenomedullin22-52 suppresses high-glucose-induced migration, proliferation, and tube formation of human retinal endothelial cells. *Mol. Vis.* 20, 259–269.
- Choi, A. M., and Alam, J. (1996). Heme oxygenase-1: function, regulation, and implication of a novel stress-inducible protein in oxidant-induced lung injury. *Am. J. Respir. Cell Mol. Biol.* 15, 9–19. doi: 10.1165/ajrcmb.15.1.8679227
- Chung, H., Lee, H., Lamoke, F., Hrushesky, W. J., Wood, P. A., and Jahng, W. J. (2009). Neuroprotective role of erythropoietin by antiapoptosis in the retina. *J. Neurosci. Res.* 87, 2365–2374. doi: 10.1002/jnr.22046
- Colella, P., Iodice, C., Di Vicino, U., Annunziata, I., Surace, E. M., and Auricchio, A. (2011). Non-erythropoietic erythropoietin derivatives protect from light-induced and genetic photoreceptor degeneration. *Hum. Mol. Genet.* 20, 2251–2262. doi: 10.1093/hmg/ddr115
- Coleman, T., and Brines, M. (2004). Science review: recombinant human erythropoietin in critical illness: a role beyond anemia? *Crit. Care* 8, 337–341. doi: 10.1186/cc2897
- Coultas, L., Chawengsaksophak, K., and Rossant, J. (2005). Endothelial cells and VEGF in vascular development. *Nature* 438, 937–945. doi: 10.1038/nature04479
- Cuevas, P., Carceller, F., Redondo-Horcajo, M., Lozano, R. M., and Giménez-Gallego, G. (1998). Systemic administration of acidic fibroblast growth factor ameliorates the ischemic injury of the retina in rats. *Neurosci. Lett.* 255, 1–4. doi: 10.1016/S0304-3940(98)00672-7
- Cukiernik, M., Mukherjee, S., Downey, D., and Chakabarti, S. (2003). Heme oxygenase in the retina in diabetes. *Curr. Eye Res.* 27, 301–308. doi: 10.1076/ceyr.27.5.301.17227

- da Silva, J. L., Stoltz, R. A., Dunn, M. W., Abraham, N. G., and Shibahara, S. (1997). Diminished heme oxygenase-1 mRNA expression in RPE cells from diabetic donors as quantitated by competitive RT/PCR. *Curr. Eye Res.* 16, 380–386. doi: 10.1076/ceyr.16.4.380.10695
- Demirel, S., Batioglu, F., Özmert, E., and Erenler, F. (2015). The effect of multiple injections of ranibizumab on retinal nerve fiber layer thickness in patients with age-related macular degeneration. *Curr. Eye Res.* 40, 87–92. doi: 10.3109/02713683.2014.917190
- Dogan, A., Suzuki, Y., Koketsu, N., Osuka, K., Saito, K., Takayasu, M., et al. (1997). Intravenous infusion of adrenomedullin and increase in regional cerebral blood flow and prevention of ischemic brain injury after middle cerebral artery occlusion in rats. *J. Cereb. Blood Flow Metab.* 17, 19–25. doi: 10.1097/00004647-199701000-00004
- Eckardt, K. U., and Kurtz, A. (2005). Regulation of erythropoietin production. *Eur. J. Clin. Invest.* 35(Suppl. 3), 13–19. doi: 10.1111/j.1365-2362.2005.01525.x
- Ehrenreich, H., Hasselblatt, M., Dembowski, C., Cepek, L., Lewczuk, P., Stiefel, M., et al. (2002). Erythropoietin therapy for acute stroke is both safe and beneficial. *Mol. Med.* 8, 495–505.
- Emerich, D. F., Silva, E., Ali, O., Mooney, D., Bell, W., Yu, S. J., et al. (2010). Injectable VEGF hydrogels produce near complete neurological and anatomical protection following cerebral ischemia in rats. *Cell Transplant.* 19, 1063–1071. doi: 10.3727/096368910X498278
- Erbayraktar, S., Yilmaz, O., Gökmen, N., and Brines, M. (2003). Erythropoietin is a multifunctional tissue-protective cytokine. *Curr. Hematol. Rep.* 2, 465–470.
- Evereklioglu, C., Doganay, S., Er, H., and Yurekli, M. (2002). Aqueous humor adrenomedullin levels differ in patients with different types of glaucoma. *Jpn. J. Ophthalmol.* 46, 203–208. doi: 10.1016/S0021-5155(01)00501-9
- Fan, J., Xu, G., Jiang, T., and Qin, Y. (2012). Pharmacologic induction of heme oxygenase-1 plays a protective role in diabetic retinopathy in rats. *Invest. Ophthalmol. Vis. Sci.* 53, 6541–6556. doi: 10.1167/iovs.11-9241
- Fernandes, R., Hosoya, K., and Pereira, P. (2011). Reactive oxygen species downregulate glucose transport system in retinal endothelial cells. *Am. J. Physiol. Cell Physiol.* 300, C927–C936. doi: 10.1152/ajpcell.00140.2010
- Foresti, R., Bains, S. K., Pitchumony, T. S., de Castro Bras, L. E., Drago, F., Dubois-Randé, J. L., et al. (2013). Small molecule activators of the Nrf2-HO-1 antioxidant axis modulate heme metabolism and inflammation in BV2 microglia cells. *Pharmacol. Res.* 76, 132–148. doi: 10.1016/j.phrs.2013.07.010
- Foresti, R., Bucolo, C., Platania, C. M., Drago, F., Dubois-Randé, J. L., and Motterlini, R. (2015). Nrf2 activators modulate oxidative stress responses and bioenergetic profiles of human retinal epithelial cells cultured in normal or high glucose conditions. *Pharmacol. Res.* 99, 296–307. doi: 10.1016/j.phrs.2015.07.006
- Foxton, R. H., Finkelstein, A., Vijay, S., Dahlmann-Noor, A., Khaw, P. T., Morgan, J. E., et al. (2013). VEGF-A is necessary and sufficient for retinal neuroprotection in models of experimental glaucoma. *Am. J. Pathol.* 182, 1379–1390. doi: 10.1016/j.ajpath.2012.12.032
- Frank, R. N., Amin, R. H., and Puklin, J. E. (1999). Antioxidant enzymes in the macular retinal pigment epithelium of eyes with neovascular age-related macular degeneration. *Am. J. Ophthalmol.* 127, 694–709. doi: 10.1016/S0002-9394(99)00032-X
- Fulton, A. B., Akula, J. D., Mocko, J. A., Hansen, R. M., Benador, I. Y., Beck, S. C., et al. (2009). Retinal degenerative and hypoxic ischemic disease. *Doc. Ophthalmol.* 118, 55–61. doi: 10.1007/s10633-008-9127-8
- Gao, H., and Hollyfield, J. G. (1995). Basic fibroblast growth factor in retinal development: differential levels of bFGF expression and content in normal and retinal degeneration (rd) mutant mice. *Dev. Biol.* 169, 168–184. doi: 10.1006/dbio.1995.1135
- Garcia-Ramirez, M., Hernandez, C., and Simo, R. (2008). Expression of erythropoietin and its receptor in the human retina: a comparative study of diabetic and nondiabetic subjects. *Diabetes Care* 31, 1189–1194. doi: 10.2337/dc07-2075
- Gassmann, M., Heinicke, K., Soliz, J., and Ogunshola, O. O. (2003). Non-erythroid functions of erythropoietin. *Adv. Exp. Med. Biol.* 543, 323–330. doi: 10.1007/978-1-4419-8997-0_22
- Gawad, A. E., Schlichting, L., Strauss, O., and Zeitz, O. (2009). Antiapoptotic properties of erythropoietin: novel strategies for protection of retinal pigment epithelial cells. *Eye* 23, 2245–2250. doi: 10.1038/eye.2008.398
- Geraldes, P., Yagi, K., Ohshiro, Y., He, Z., Maeno, Y., Yamamoto-Hiraoka, J., et al. (2008). Selective regulation of heme oxygenase-1 expression and function by insulin through IRS1/phosphoinositide 3-kinase/Akt-2 pathway. *J. Biol. Chem.* 283, 34327–34336. doi: 10.1074/jbc.M807036200
- Gidday, J. M. (2006). Cerebral preconditioning and ischaemic tolerance. *Nat. Rev. Neurosci.* 7, 437–448. doi: 10.1038/nrn1927
- Glazier, S. S., O'Rourke, D. M., Graham, D. I., and Welsh, F. A. (1994). Induction of ischemic tolerance following brief focal ischemia in rat brain. *J. Cereb. Blood Flow Metab.* 14, 545–553. doi: 10.1038/jcbfm.1994.68
- Grimm, C., Hermann, D. M., Bogdanova, A., Hotop, S., Kilic, U., Wenzel, A., et al. (2005). Neuroprotection by hypoxic preconditioning: HIF-1 and erythropoietin protect from retinal degeneration. *Semin. Cell Dev. Biol.* 16, 531–538. doi: 10.1016/j.semcdb.2005.03.004
- Grimm, C., Wenzel, A., Groszer, M., Mayser, H., Seeliger, M., Samardzija, M., et al. (2002). HIF-1-induced erythropoietin in the hypoxic retina protects against light-induced retinal degeneration. *Nat. Med.* 8, 718–724. doi: 10.1038/nm723
- Grimm, C., Wenzel, A., Stanescu, D., Samardzija, M., Hotop, S., Groszer, M., et al. (2004). Constitutive overexpression of human erythropoietin protects the mouse retina against induced but not inherited retinal degeneration. *J. Neurosci.* 24, 5651–5658. doi: 10.1523/JNEUROSCI.1288-04.2004
- Grimm, C., and Willmann, G. (2012). Hypoxia in the eye: a two-sided coin. *High Alt. Med. Biol.* 13, 169–175. doi: 10.1089/ham.2012.1031
- Guan, Y., Cui, L., Qu, Z., Lu, L., Wang, F., Wu, Y., et al. (2013). Subretinal transplantation of rat MSCs and erythropoietin gene modified rat MSCs for protecting and rescuing degenerative retina in rats. *Curr. Mol. Med.* 13, 1419–1431. doi: 10.2174/15665240113139990071
- Haigh, J. J., Morelli, P. I., Gerhardt, H., Haigh, K., Tsien, J., Damert, A., et al. (2003). Cortical and retinal defects caused by dosage-dependent reductions in VEGF-A paracrine signaling. *Dev. Biol.* 262, 225–241. doi: 10.1016/S0012-1606(03)00356-7
- Hanaoka, M., Droma, Y., Basnyat, B., Ito, M., Kobayashi, N., Katsuyama, Y., et al. (2012). Genetic variants in EPAS1 contribute to adaptation to high-altitude hypoxia in Sherpas. *PLoS ONE* 7:e50566. doi: 10.1371/journal.pone.0050566
- Harten, S. K., Ashcroft, M., and Maxwell, P. H. (2010). Prolyl hydroxylase domain inhibitors: a route to HIF activation and neuroprotection. *Antioxid. Redox Signal.* 12, 459–480. doi: 10.1089/ars.2009.2870
- He, M., Pan, H., Chang, R. C., So, K. F., Brecha, N. C., and Pu, M. (2014). Activation of the Nrf2/HO-1 antioxidant pathway contributes to the protective effects of *Lycium barbarum* polysaccharides in the rodent retina after ischemia-reperfusion-induced damage. *PLoS ONE* 9:e84800. doi: 10.1371/journal.pone.0084800
- He, M., Pan, H., Xiao, C., and Pu, M. (2013). Roles for redox signaling by NADPH oxidase in hyperglycemia-induced heme oxygenase-1 expression in the diabetic retina. *Invest. Ophthalmol. Vis. Sci.* 54, 4092–4101. doi: 10.1167/iovs.13-12004
- Hegazy, K. A., Dunn, M. W., and Sharma, S. C. (2000). Functional human heme oxygenase has a neuroprotective effect on adult rat ganglion cells after pressure-induced ischemia. *Neuroreport* 11, 1185–1189. doi: 10.1097/00001756-200004270-00008
- Heikkilä, M., Pasanen, A., Kivirikko, K. I., and Myllyharju, J. (2011). Roles of the human hypoxia-inducible factor (HIF)-3 α variants in the hypoxia response. *Cell. Mol. Life Sci.* 68, 3885–3901. doi: 10.1007/s00018-011-0679-5
- Hernandez, C., Fonollosa, A., Garcia-Ramirez, M., Higuera, M., Catalan, R., Miralles, A., et al. (2006). Erythropoietin is expressed in the human retina and it is highly elevated in the vitreous fluid of patients with diabetic macular edema. *Diabetes Care* 29, 2028–2033. doi: 10.2337/dc06-0556
- Hernandez, C., and Simo, R. (2012). Neuroprotection in diabetic retinopathy. *Curr. Diab. Rep.* 12, 329–337. doi: 10.1007/s11892-012-0284-5
- Himori, N., Maruyama, K., Yamamoto, K., Yasuda, M., Ryu, M., Omodaka, K., et al. (2014). Critical neuroprotective roles of heme oxygenase-1 induction against axonal injury-induced retinal ganglion cell death. *J. Neurosci. Res.* 92, 1134–1142. doi: 10.1002/jnr.23398
- Hines-Beard, J., Desai, S., Haag, R., Esumi, N., D'Surney, L., Parker, S., et al. (2013). Identification of a therapeutic dose of continuously delivered erythropoietin in the eye using an inducible promoter system. *Curr. Gene Ther.* 13, 275–281. doi: 10.2174/15665232113139990024
- Hosoya, K., Nakamura, G., Akanuma, S., Tomi, M., and Tachikawa, M. (2008). Dehydroascorbic acid uptake and intracellular ascorbic acid accumulation

- in cultured Muller glial cells (TR-MUL). *Neurochem. Int.* 52, 1351–1357. doi: 10.1016/j.neuint.2008.02.001
- Iesato, Y., Toriyama, Y., Sakurai, T., Kamiyoshi, A., Ichikawa-Shindo, Y., Kawate, H., et al. (2013). Adrenomedullin-RAMP2 system is crucially involved in retinal angiogenesis. *Am. J. Pathol.* 182, 2380–2390. doi: 10.1016/j.ajpath.2013.02.015
- Igarashi, K., Sakurai, T., Kamiyoshi, A., Ichikawa-Shindo, Y., Kawate, H., Yamauchi, A., et al. (2014). Pathophysiological roles of adrenomedullin-RAMP2 system in acute and chronic cerebral ischemia. *Peptides* 62, 21–31. doi: 10.1016/j.peptides.2014.08.013
- Imuro, S., Shindo, T., Moriyama, N., Amaki, T., Niu, P., Takeda, N., et al. (2004). Angiogenic effects of adrenomedullin in ischemia and tumor growth. *Circ. Res.* 95, 415–423. doi: 10.1161/01.RES.0000138018.61065.d1
- Ito, S., Fujisawa, K., Sakamoto, T., and Ishibashi, T. (2003). Elevated adrenomedullin in the vitreous of patients with diabetic retinopathy. *Ophthalmologica* 217, 53–57. doi: 10.1159/000068244
- Jaquet, K., Krause, K., Tawakol-Khodai, M., Geidel, S., and Kuck, K. H. (2002). Erythropoietin and VEGF exhibit equal angiogenic potential. *Microvasc. Res.* 64, 326–333. doi: 10.1006/mvre.2002.2426
- Jewell, U. R., Kvietikova, I., Scheid, A., Bauer, C., Wenger, R. H., and Gassmann, M. (2001). Induction of HIF-1 α in response to hypoxia is instantaneous. *FASEB J.* 15, 1312–1314.
- Jin, K. L., Mao, X. O., and Greenberg, D. A. (2000). Vascular endothelial growth factor: direct neuroprotective effect in *in vitro* ischemia. *Proc. Natl. Acad. Sci. U.S.A.* 97, 10242–10247. doi: 10.1073/pnas.97.18.10242
- Junk, A. K., Mammis, A., Savitz, S. I., Singh, M., Roth, S., Malhotra, S., et al. (2002). Erythropoietin administration protects retinal neurons from acute ischemia-reperfusion injury. *Proc. Natl. Acad. Sci. U.S.A.* 99, 10659–10664. doi: 10.1073/pnas.152321399
- Katz, B., Rimmer, S., Iragui, V., and Katzman, R. (1989). Abnormal pattern electroretinogram in Alzheimer's disease: evidence for retinal ganglion cell degeneration? *Ann. Neurol.* 26, 221–225.
- Kermer, P., Klocker, N., Labes, M., and Bahr, M. (2000). Insulin-like growth factor-I protects axotomized rat retinal ganglion cells from secondary death via PI3-K-dependent Akt phosphorylation and inhibition of caspase-3 *in vivo*. *J. Neurosci.* 20, 2–8.
- Kikuchi, G., Yoshida, T., and Noguchi, M. (2005). Heme oxygenase and heme degradation. *Biochem. Biophys. Res. Commun.* 338, 558–567. doi: 10.1016/j.bbrc.2005.08.020
- Kilic, U., Kilic, E., Dietz, G. P., and Bahr, M. (2004). The TAT protein transduction domain enhances the neuroprotective effect of glial-cell-line-derived neurotrophic factor after optic nerve transection. *Neurodegener. Dis.* 1, 44–49. doi: 10.1159/000076669
- Kilic, U., Kilic, E., Jarve, A., Guo, Z., Spudich, A., Bieber, K., et al. (2006). Human vascular endothelial growth factor protects axotomized retinal ganglion cells *in vivo* by activating ERK-1/2 and Akt pathways. *J. Neurosci.* 26, 12439–12446. doi: 10.1523/JNEUROSCI.0434-06.2006
- Kilic, U., Kilic, E., Soliz, J., Bassetti, C. I., Gassmann, M., and Hermann, D. M. (2005). Erythropoietin protects from axotomy-induced degeneration of retinal ganglion cells by activating ERK-1/2. *FASEB J.* 19, 249–251. doi: 10.1096/fj.04-2493fje
- Kim, K. J., Kim, K. S., Kim, N. R., and Chin, H. S. (2012). Effects of simvastatin on the expression of heme oxygenase-1 in human RPE cells. *Invest. Ophthalmol. Vis. Sci.* 53, 6456–6464. doi: 10.1167/iovs.12-9658
- Kitagawa, K., Matsumoto, M., Tagaya, M., Hata, R., Ueda, H., Niinobe, M., et al. (1990). 'Ischemic tolerance' phenomenon found in the brain. *Brain Res.* 528, 21–24. doi: 10.1016/0006-8993(90)90189-I
- Klocker, N., Kermer, P., Weishaupt, J. H., Labes, M., Ankerhold, R., and Bahr, M. (2000). Brain-derived neurotrophic factor-mediated neuroprotection of adult rat retinal ganglion cells *in vivo* does not exclusively depend on phosphatidylinositol-3'-kinase/protein kinase B signaling. *J. Neurosci.* 20, 6962–6967.
- Koskela, A., Reinisalo, M., Hyttinen, J. M., Kaarniranta, K., and Karjalainen, R. O. (2014). Pinosylvin-mediated protection against oxidative stress in human retinal pigment epithelial cells. *Mol. Vis.* 20, 760–769.
- Kretz, A., Happold, C. J., Marticke, J. K., and Isenmann, S. (2005). Erythropoietin promotes regeneration of adult CNS neurons via Jak2/Stat3 and PI3K/AKT pathway activation. *Mol. Cell. Neurosci.* 29, 569–579. doi: 10.1016/j.mcn.2005.04.009
- Kutty, R. K., Kutty, G., Wiggert, B., Chader, G. J., Darrow, R. M., and Organisciak, D. T. (1995). Induction of heme oxygenase 1 in the retina by intense visible light: suppression by the antioxidant dimethylthiourea. *Proc. Natl. Acad. Sci. U.S.A.* 92, 1177–1181. doi: 10.1073/pnas.92.4.1177
- Lange, C., Heynen, S. R., Tanimoto, N., Thiersch, M., Le, Y. Z., Meneau, I., et al. (2011). Normoxic activation of hypoxia-inducible factors in photoreceptors provides transient protection against light-induced retinal degeneration. *Invest. Ophthalmol. Vis. Sci.* 52, 5872–5880. doi: 10.1167/iovs.11-7204
- Lee, D., Kim, K. Y., Shim, M. S., Kim, S. Y., Ellisman, M. H., Weinreb, R. N., et al. (2014). Coenzyme Q10 ameliorates oxidative stress and prevents mitochondrial alteration in ischemic retinal injury. *Apoptosis* 19, 603–614. doi: 10.1007/s10495-013-0956-x
- Liu, Q., Ju, W. K., Crowston, J. G., Xie, F., Perry, G., Smith, M. A., et al. (2007). Oxidative stress is an early event in hydrostatic pressure induced retinal ganglion cell damage. *Invest. Ophthalmol. Vis. Sci.* 48, 4580–4589. doi: 10.1167/iovs.07-0170
- Lopez, J., and Martinez, A. (2002). Cell and molecular biology of the multifunctional peptide, adrenomedullin. *Int. Rev. Cytol.* 221, 1–92. doi: 10.1016/S0074-7696(02)21010-4
- Lu, L., Seidel, C. P., Iwase, T., Stevens, R. K., Gong, Y. Y., Wang, X., et al. (2013). Suppression of GLUT1; a new strategy to prevent diabetic complications. *J. Cell. Physiol.* 228, 251–257. doi: 10.1002/jcp.24133
- Maenhaut, N., Boussery, K., Delaey, C., and Van de Voorde, J. (2007). Control of retinal arterial tone by a paracrine retinal relaxing factor. *Microcirculation* 14, 39–48. doi: 10.1080/10739680601072131
- Maharaj, A. S., and D'Amore, P. A. (2007). Roles for VEGF in the adult. *Microvasc. Res.* 74, 100–113. doi: 10.1016/j.mvr.2007.03.004
- Mandal, M. N., Patlolla, J. M., Zheng, L., Agbaga, M. P., Tran, J. T., Wicker, L., et al. (2009). Curcumin protects retinal cells from light- and oxidant stress-induced cell death. *Free Radic. Biol. Med.* 46, 672–679. doi: 10.1016/j.freeradbiomed.2008.12.006
- Marti, H. J., Bernaudin, M., Bellail, A., Schoch, H., Euler, M., Petit, E., et al. (2000). Hypoxia-induced vascular endothelial growth factor expression precedes neovascularization after cerebral ischemia. *Am. J. Pathol.* 156, 965–976. doi: 10.1016/S0002-9440(10)64964-4
- Maxwell, P. H., Wiesener, M. S., Chang, G. W., Clifford, S. C., Vaux, E. C., Cockman, M. E., et al. (1999). The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 399, 271–275.
- McVicar, C. M., Hamilton, R., Colhoun, L. M., Gardiner, T. A., Brines, M., Cerami, A., et al. (2011). Intervention with an erythropoietin-derived peptide protects against neurological and vascular degeneration during diabetic retinopathy. *Diabetes* 60, 2995–3005. doi: 10.2337/db11-0026
- Miki, A., Miki, K., Ueno, S., Wersinger, D. M., Berlinicke, C., Shaw, G. C., et al. (2010). Prolonged blockade of VEGF receptors does not damage retinal photoreceptors or ganglion cells. *J. Cell. Physiol.* 224, 262–272. doi: 10.1002/jcp.22129
- Mitsuhashi, J., Morikawa, S., Shimizu, K., Ezaki, T., Yasuda, Y., and Hori, S. (2013). Intravitreal injection of erythropoietin protects against retinal vascular regression at the early stage of diabetic retinopathy in streptozotocin-induced diabetic rats. *Exp. Eye Res.* 106, 64–73. doi: 10.1016/j.exer.2012.11.001
- Miyashita, K., Itoh, H., Arai, H., Suganami, T., Sawada, N., Fukunaga, Y., et al. (2006). The neuroprotective and vasculo-neuro-regenerative roles of adrenomedullin in ischemic brain and its therapeutic potential. *Endocrinology* 147, 1642–1653. doi: 10.1210/en.2005-1038
- Montuenga, L. M., Martinez, A., Miller, M. J., Unsworth, E. J., and Cuttitta, F. (1997). Expression of adrenomedullin and its receptor during embryogenesis suggests autocrine or paracrine modes of action. *Endocrinology* 138, 440–451. doi: 10.1210/endo.138.1.4881
- Moravski, C. J., Skinner, S. L., Stubbs, A. J., Sarlos, S., Kelly, D. J., Cooper, M. E., et al. (2003). The renin-angiotensin system influences ocular endothelial cell proliferation in diabetes: transgenic and interventional studies. *Am. J. Pathol.* 162, 151–160. doi: 10.1016/S0002-9440(10)63806-0
- Morse, D., and Choi, A. M. (2005). Heme oxygenase-1: from bench to bedside. *Am. J. Respir. Crit. Care Med.* 172, 660–670. doi: 10.1164/rccm.200404-465SO
- Mowat, F. M., Gonzalez, F., Luhmann, U. F., Lange, C. A., Duran, Y., Smith, A. J., et al. (2012). Endogenous erythropoietin protects neuroretinal

- function in ischemic retinopathy. *Am. J. Pathol.* 180, 1726–1739. doi: 10.1016/j.ajpath.2011.12.033
- Nagpal, M., Nagpal, K., and Nagpal, P. N. (2007). A comparative debate on the various anti-vascular endothelial growth factor drugs: pegaptanib sodium (Macugen), ranibizumab (Lucentis) and bevacizumab (Avastin). *Indian J. Ophthalmol.* 55, 437–439. doi: 10.4103/0301-4738.36478
- Nihira, M., Anderson, K., Gorin, F. A., and Burns, M. S. (1995). Primate rod and cone photoreceptors may differ in glucose accessibility. *Invest. Ophthalmol. Vis. Sci.* 36, 1259–1270.
- Nishijima, K., Ng, Y. S., Zhong, L., Bradley, J., Schubert, W., Jo, N., et al. (2007). Vascular endothelial growth factor-A is a survival factor for retinal neurons and a critical neuroprotectant during the adaptive response to ischemic injury. *Am. J. Pathol.* 171, 53–67. doi: 10.2353/ajpath.2007.061237
- Park, H. Y., Kim, J. H., and Park, C. K. (2014). Neuronal cell death in the inner retina and the influence of vascular endothelial growth factor inhibition in a diabetic rat model. *Am. J. Pathol.* 184, 1752–1762. doi: 10.1016/j.ajpath.2014.02.016
- Peng, P. H., Chao, H. M., Juan, S. H., Chen, C. F., Liu, J. H., and Ko, M. L. (2011). Pharmacological preconditioning by low dose cobalt protoporphyrin induces heme oxygenase-1 overexpression and alleviates retinal ischemia-reperfusion injury in rats. *Curr. Eye Res.* 36, 238–246. doi: 10.3109/02713683.2010.539760
- Peng, P. H., Ko, M. L., Chen, C. F., and Juan, S. H. (2008). Haem oxygenase-1 gene transfer protects retinal ganglion cells from ischaemia/reperfusion injury. *Clin. Sci.* 115, 335–342. doi: 10.1042/CS20070384
- Ponnaluri, V. K., Vadlapatla, R. K., Vavilala, D. T., Pal, D., Mitra, A. K., and Mukherji, M. (2011). Hypoxia induced expression of histone lysine demethylases: implications in oxygen-dependent retinal neovascular diseases. *Biochem. Biophys. Res. Commun.* 415, 373–377. doi: 10.1016/j.bbrc.2011.10.075
- Puchowicz, M. A., Xu, K., Magness, D., Miller, C., Lust, W. D., Kern, T. S., et al. (2004). Comparison of glucose influx and blood flow in retina and brain of diabetic rats. *J. Cereb. Blood Flow Metab.* 24, 449–457. doi: 10.1097/00004647-200404000-00010
- Rappoport, D., Morzaev, D., Weiss, S., Vieyra, M., Nicholson, J. D., Leiba, H., et al. (2013). Effect of intravitreal injection of bevacizumab on optic nerve head leakage and retinal ganglion cell survival in a mouse model of optic nerve crush. *Invest. Ophthalmol. Vis. Sci.* 54, 8160–8171. doi: 10.1167/iovs.13-12771
- Ray, D., Mishra, M., Ralph, S., Read, I., Davies, R., and Brenchley, P. (2004). Association of the VEGF gene with proliferative diabetic retinopathy but not proteinuria in diabetes. *Diabetes* 53, 861–864. doi: 10.2337/diabetes.53.3.861
- Ren, H., Liu, N. Y., Song, X. F., Ma, Y. S., and Zhai, X. Y. (2011). A novel specific application of pyruvate protects the mouse retina against white light damage: differential stabilization of HIF-1 α and HIF-2 α . *Invest. Ophthalmol. Vis. Sci.* 52, 3112–3118. doi: 10.1167/iovs.10-5605
- Rex, T. S., Wong, Y., Kodali, K., and Merry, S. (2009). Neuroprotection of photoreceptors by direct delivery of erythropoietin to the retina of the retinal degeneration slow mouse. *Exp. Eye Res.* 89, 735–740. doi: 10.1016/j.exer.2009.06.017
- Saint-Geniez, M., Maharaj, A. S., Walshe, T. E., Tucker, B. A., Sekiyama, E., Kurihara, T., et al. (2008). Endogenous VEGF is required for visual function: evidence for a survival role on muller cells and photoreceptors. *PLoS ONE* 3:e3554. doi: 10.1371/journal.pone.0003554
- Sakamoto, K., Yonoki, Y., Kubota, Y., Kuwagata, M., Saito, M., Nakahara, T., et al. (2006). Inducible nitric oxide synthase inhibitors abolished histological protection by late ischemic preconditioning in rat retina. *Exp. Eye Res.* 82, 512–518. doi: 10.1016/j.exer.2005.08.011
- Satarug, S., Wisedpanichkij, R., Takeda, K., Li, B., Na-Bangchang, K., Moore, M. R., et al. (2008). Prostaglandin D2 induces heme oxygenase-1 mRNA expression through the DP2 receptor. *Biochem. Biophys. Res. Commun.* 377, 878–883. doi: 10.1016/j.bbrc.2008.10.094
- Schubert, D. (2005). Glucose metabolism and Alzheimer's disease. *Ageing Res. Rev.* 4, 240–257. doi: 10.1016/j.arr.2005.02.003
- Semenza, G. L., and Wang, G. L. (1992). A nuclear factor induced by hypoxia via *de novo* protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol. Cell. Biol.* 12, 5447–5454. doi: 10.1128/MCB.12.12.5447
- Semeran, K., Pawlowski, P., Lisowski, L., Szczepaniak, I., Wojtowicz, J., Lawicki, S., et al. (2013). Plasma levels of IL-17, VEGF, and adrenomedullin and S-cone dysfunction of the retina in children and adolescents without signs of retinopathy and with varied duration of diabetes. *Mediators Inflamm.* 2013:274726. doi: 10.1155/2013/274726
- Shaw, S. G., Boden, J. P., Biecker, E., Reichen, J., and Rothen, B. (2006). Endothelin antagonism prevents diabetic retinopathy in NOD mice: a potential role of the angiogenic factor adrenomedullin. *Exp. Biol. Med.* 231, 1101–1105.
- Shen, J., Wu, Y., Xu, J. Y., Zhang, J., Sinclair, S. H., Yanoff, M., et al. (2010). ERK- and Akt-dependent neuroprotection by erythropoietin (EPO) against glyoxal-AGEs via modulation of Bcl-xL, Bax, and BAD. *Invest. Ophthalmol. Vis. Sci.* 51, 35–46. doi: 10.1167/iovs.09-3544
- Shen, W., Chung, S. H., Irhimeh, M. R., Li, S., Lee, S. R., and Gillies, M. C. (2014). Systemic administration of erythropoietin inhibits retinopathy in RCS rats. *PLoS ONE* 9:e104759. doi: 10.1371/journal.pone.0104759
- Shima, D. T., Adamis, A. P., Ferrara, N., Yeo, K. T., Yeo, T. K., Allende, R., et al. (1995). Hypoxic induction of endothelial cell growth factors in retinal cells: identification and characterization of vascular endothelial growth factor (VEGF) as the mitogen. *Mol. Med.* 1, 182–193.
- Shindo, T., Kurihara, Y., Nishimatsu, H., Moriyama, N., Kakoki, M., Wang, Y., et al. (2001). Vascular abnormalities and elevated blood pressure in mice lacking adrenomedullin gene. *Circulation* 104, 1964–1971. doi: 10.1161/hc4101.097111
- Shyong, M. P., Lee, F. L., Hen, W. H., Kuo, P. C., Wu, A. C., Cheng, H. C., et al. (2008). Viral delivery of heme oxygenase-1 attenuates photoreceptor apoptosis in an experimental model of retinal detachment. *Vision Res.* 48, 2394–2402. doi: 10.1016/j.visres.2008.07.017
- Sierra, A., Navascues, J., Cuadros, M. A., Calvente, R., Martin-Oliva, D., Ferrer-Martin, R. M., et al. (2014). Expression of inducible nitric oxide synthase (iNOS) in microglia of the developing quail retina. *PLoS ONE* 9:e106048. doi: 10.1371/journal.pone.0106048
- Sobaci, G., Gungor, R., and Ozge, G. (2013). Effects of multiple intravitreal anti-VEGF injections on retinal nerve fiber layer and intraocular pressure: a comparative clinical study. *Int. J. Ophthalmol.* 6, 211–215. doi: 10.3980/j.issn.2222-3959.2013.02.20
- Song, S. K., Sun, S. W., Ju, W. K., Lin, S. J., Cross, A. H., and Neufeld, A. H. (2003). Diffusion tensor imaging detects and differentiates axon and myelin degeneration in mouse optic nerve after retinal ischemia. *Neuroimage* 20, 1714–1722. doi: 10.1016/j.neuroimage.2003.07.005
- Sullivan, T. A., Geisert, E. E., Hines-Beard, J., and Rex, T. S. (2011). Systemic adeno-associated virus-mediated gene therapy preserves retinal ganglion cells and visual function in DBA/2J glaucomatous mice. *Hum. Gene Ther.* 22, 1191–1200. doi: 10.1089/hum.2011.052
- Sun, M. H., Pang, J. H., Chen, S. L., Han, W. H., Ho, T. C., Chen, K. J., et al. (2010). Retinal protection from acute glaucoma-induced ischemia-reperfusion injury through pharmacologic induction of heme oxygenase-1. *Invest. Ophthalmol. Vis. Sci.* 51, 4798–4808. doi: 10.1167/iovs.09-4086
- Sun, M. H., Pang, J. H., Chen, S. L., Kuo, P. C., Chen, K. J., Kao, L. Y., et al. (2007). Photoreceptor protection against light damage by AAV-mediated overexpression of heme oxygenase-1. *Invest. Ophthalmol. Vis. Sci.* 48, 5699–5707. doi: 10.1167/iovs.07-0340
- Suzuki, M., Ozawa, Y., Kubota, S., Hirasawa, M., Miyake, S., Noda, K., et al. (2011). Neuroprotective response after photodynamic therapy: role of vascular endothelial growth factor. *J. Neuroinflammation* 8:176. doi: 10.1186/1742-2094-8-176
- Takahashi, K. (2001). Adrenomedullin from a pheochromocytoma to the eye: implications of the adrenomedullin research for endocrinology in the 21st century. *Tohoku J. Exp. Med.* 193, 79–114. doi: 10.1620/tjem.193.79
- Tezel, G., and Wax, M. B. (2004). Hypoxia-inducible factor 1 α in the glaucomatous retina and optic nerve head. *Arch. Ophthalmol.* 122, 1348–1356. doi: 10.1001/archophth.122.9.1348
- Thiersch, M., Raffelsberger, W., Frigg, R., Samardzija, M., Wenzel, A., Poch, O., et al. (2008). Analysis of the retinal gene expression profile after hypoxic preconditioning identifies candidate genes for neuroprotection. *BMC Genomics* 9:73. doi: 10.1186/1471-2164-9-73
- Toft-Kehler, A. K., Skytt, D. M., Poulsen, K. A., Braendstrup, C. T., Gegelashvili, G., Waagepetersen, H., et al. (2014). Limited energy supply in Muller cells alters glutamate uptake. *Neurochem. Res.* 39, 941–949. doi: 10.1007/s11064-014-1289-z
- Udono, T., Takahashi, K., Abe, T., Shibahara, S., and Tamai, M. (2002). Elevated immunoreactive-adrenomedullin levels in the aqueous humor of

- patients with uveitis and vitreoretinal disorders. *Peptides* 23, 1865–1868. doi: 10.1016/S0196-9781(02)00145-6
- Udono, T., Takahashi, K., Nakayama, M., Murakami, O., Durlu, Y. K., Tamai, M., et al. (2000). Adrenomedullin in cultured human retinal pigment epithelial cells. *Invest. Ophthalmol. Vis. Sci.* 41, 1962–1970.
- Udono, T., Takahashi, K., Nakayama, M., Yoshinoya, A., Totsune, K., Murakami, O., et al. (2001). Induction of adrenomedullin by hypoxia in cultured retinal pigment epithelial cells. *Invest. Ophthalmol. Vis. Sci.* 42, 1080–1086.
- Udono-Fujimori, R., Udono, T., Totsune, K., Tamai, M., Shibahara, S., and Takahashi, K. (2003). Adrenomedullin in the eye. *Regul. Pept.* 112, 95–101. doi: 10.1016/S0167-0115(03)00027-2
- Ulyanova, T., Szel, A., Kuty, R. K., Wiggert, B., Caffè, A. R., Chader, G. J., et al. (2001). Oxidative stress induces heme oxygenase-1 immunoreactivity in Muller cells of mouse retina in organ culture. *Invest. Ophthalmol. Vis. Sci.* 42, 1370–1374.
- Valter, K., Bisti, S., Gargini, C., Di Loreto, S., Maccarone, R., Cervetto, L., et al. (2005). Time course of neurotrophic factor upregulation and retinal protection against light-induced damage after optic nerve section. *Invest. Ophthalmol. Vis. Sci.* 46, 1748–1754. doi: 10.1167/iovs.04-0657
- Varga, B., Gesztelyi, R., Bombicz, M., Haines, D., Szabo, A. M., Kemeny-Beke, A., et al. (2013). Protective effect of alpha-melanocyte-stimulating hormone (alpha-MSH) on the recovery of ischemia/reperfusion (I/R)-induced retinal damage in a rat model. *J. Mol. Neurosci.* 50, 558–570. doi: 10.1007/s12031-013-9998-3
- Vingolo, E. M., Lupo, S., Domanico, D., Cotesta, D., Petramala, L., Grenga, R., et al. (2005). Adrenomedullin plasma concentrations in patients with retinitis pigmentosa. *Clin. Biochem.* 38, 735–738. doi: 10.1016/j.clinbiochem.2005.04.007
- Wang, Q., Gorbey, S., Pfister, F., Hoger, S., Dorn-Beineke, A., Krugel, K., et al. (2011). Long-term treatment with suberythropoietic Epo is vaso- and neuroprotective in experimental diabetic retinopathy. *Cell. Physiol. Biochem.* 27, 769–782. doi: 10.1159/000330085
- Wang, Z. Y., Shen, L. J., Tu, L., Hu, D. N., Liu, G. Y., Zhou, Z. L., et al. (2009). Erythropoietin protects retinal pigment epithelial cells from oxidative damage. *Free Radic. Biol. Med.* 46, 1032–1041. doi: 10.1016/j.freeradbiomed.2008.11.027
- Watanabe, D., Suzuma, K., Matsui, S., Kurimoto, M., Kiryu, J., Kita, M., et al. (2005). Erythropoietin as a retinal angiogenic factor in proliferative diabetic retinopathy. *N. Engl. J. Med.* 353, 782–792. doi: 10.1056/NEJMoa041773
- Weng, Y., Shen, F., and Zhang, X. M. (2006). Study on increasing expression of adrenomedullin in diabetic rats' retinal pigment epithelium cells. *Fen Zi Xi Bao Sheng Wu Xue Bao* 39, 373–377.
- Willis, D., Moore, A. R., Frederick, R., and Willoughby, D. A. (1996). Heme oxygenase: a novel target for the modulation of the inflammatory response. *Nat. Med.* 2, 87–90. doi: 10.1038/nm0196-87
- Willmann, G., Fischer, M. D., Schatz, A., Schommer, K., and Gekeler, F. (2013). Retinal vessel leakage at high altitude. *JAMA* 309, 2210–2212. doi: 10.1001/jama.2013.5550
- Xia, C. F., Yin, H., Borlongan, C. V., Chao, J., and Chao, L. (2004). Adrenomedullin gene delivery protects against cerebral ischemic injury by promoting astrocyte migration and survival. *Hum. Gene Ther.* 15, 1243–1254. doi: 10.1089/hum.2004.15.1243
- Xia, C. F., Yin, H., Borlongan, C. V., Chao, J., and Chao, L. (2006). Posts ischemic infusion of adrenomedullin protects against ischemic stroke by inhibiting apoptosis and promoting angiogenesis. *Exp. Neurol.* 197, 521–530. doi: 10.1016/j.expneurol.2005.10.027
- Xie, Z., Chen, F., Wu, X., Zhuang, C., Zhu, J., Wang, J., et al. (2012a). Effects of supplemental erythropoietin on its receptor expression and signal transduction pathways in rat model of retinal detachment. *Curr. Eye Res.* 37, 138–144. doi: 10.3109/02713683.2011.647225
- Xie, Z., Chen, F., Wu, X., Zhuang, C., Zhu, J., Wang, J., et al. (2012b). Safety and efficacy of intravitreal injection of recombinant erythropoietin for protection of photoreceptor cells in a rat model of retinal detachment. *Eye* 26, 144–152. doi: 10.1038/eye.2011.254
- Yang, X., and Cepko, C. L. (1996). Flk-1, a receptor for vascular endothelial growth factor (VEGF), is expressed by retinal progenitor cells. *J. Neurosci.* 16, 6089–6099.
- Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z. X., Pool, J. E., et al. (2010). Sequencing of 50 human exomes reveals adaptation to high altitude. *Science* 329, 75–78. doi: 10.1126/science.1190371
- Yu, D. Y., Cringle, S., Valter, K., Walsh, N., Lee, D., and Stone, J. (2004). Photoreceptor death, trophic factor expression, retinal oxygen status, and photoreceptor function in the P23H rat. *Invest. Ophthalmol. Vis. Sci.* 45, 2013–2019. doi: 10.1167/iovs.03-0845
- Yuda, K., Takahashi, H., Inoue, T., Ueta, T., Iriyama, A., Kadonosono, K., et al. (2012). Adrenomedullin inhibits choroidal neovascularization via CCL2 in the retinal pigment epithelium. *Am. J. Pathol.* 181, 1464–1472. doi: 10.1016/j.ajpath.2012.06.028
- Zhang, H., Tang, B., Yin, C. G., Chen, Y., Meng, Q. L., Jiang, L., et al. (2014). Plasma adrenomedullin levels are associated with long-term outcomes of acute ischemic stroke. *Peptides* 52, 44–48. doi: 10.1016/j.peptides.2013.11.025
- Zhang, J., Wu, Y., Jin, Y., Ji, F., Sinclair, S. H., Luo, Y., et al. (2008). Intravitreal injection of erythropoietin protects both retinal vascular and neuronal cells in early diabetes. *Invest. Ophthalmol. Vis. Sci.* 49, 732–742. doi: 10.1167/iovs.07-0721
- Zhang, X., Jizhang, Y., Xu, X., Kwicien, T. D., Li, N., Zhang, Y., et al. (2014). Protective effects of remote ischemic conditioning against ischemia/reperfusion-induced retinal injury in rats. *Vis. Neurosci.* 31, 245–252. doi: 10.1017/S0952523814000121
- Zheng, L., Gong, B., Hatala, D. A., and Kern, T. S. (2007). Retinal ischemia and reperfusion causes capillary degeneration: similarities to diabetes. *Invest. Ophthalmol. Vis. Sci.* 48, 361–367. doi: 10.1167/iovs.06-0510
- Zhong, Y. S., Liu, X. H., Cheng, Y., and Min, Y. J. (2008). Erythropoietin with retrobulbar administration protects retinal ganglion cells from acute elevated intraocular pressure in rats. *J. Ocul. Pharmacol. Ther.* 24, 453–459. doi: 10.1089/jop.2008.0021
- Zhu, Y., Ohlemiller, K. K., McMahan, B. K., Park, T. S., and Gidday, J. M. (2006). Constitutive nitric oxide synthase activity is required to trigger ischemic tolerance in mouse retina. *Exp. Eye Res.* 82, 153–163. doi: 10.1016/j.exer.2005.06.005
- Zhu, Y., Zhang, L., and Gidday, J. M. (2008). Deferoxamine preconditioning promotes long-lasting retinal ischemic tolerance. *J. Ocul. Pharmacol. Ther.* 24, 527–535. doi: 10.1089/jop.2008.0082
- Zhu, Y., Zhang, L., and Gidday, J. M. (2013). Role of hypoxia-inducible factor-1 α in preconditioning-induced protection of retinal ganglion cells in glaucoma. *Mol. Vis.* 19, 2360–2372.
- Zhu, Y., Zhang, Y., Ojwang, B. A., Brantley, M. A. Jr., and Gidday, J. M. (2007). Long-term tolerance to retinal ischemia by repetitive hypoxic preconditioning: role of HIF-1 α and heme oxygenase-1. *Invest. Ophthalmol. Vis. Sci.* 48, 1735–1743. doi: 10.1167/iovs.06-1037

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Cheng, Yu, Yan, Lai and Xiang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.