



Multiple Kinases Involved in the Nicotinic Modulation of Gamma Oscillations in the Rat Hippocampal CA3 Area

JianGang Wang^{1,2†}, XiaoLong He^{3†}, Fangli Guo^{1,4}, XiangLin Cheng⁵, Yali Wang^{1,4}, XiaoFang Wang¹, ZhiWei Feng¹, Martin Vreugdenhil^{6,7} and ChengBiao Lu^{1,3,4*}

¹Key Laboratory for the Brain Research of Henan Province, Xinxiang Medical University, Xinxiang, China, ²Department of Pathophysiology, Xinxiang Medical University, Xinxiang, China, ³Key Laboratory of Neuronal Oscillation and Disease, Yantze University Medical School, JingZhou, China, ⁴Department of Neurobiology and Physiology, Xinxiang Medical University, Xinxiang, China, ⁵Department of Laboratory Medicine, Yantze University Affiliated Hospital, JingZhou, China, ⁶Department of Psychology, Xinxiang Medical University, Xinxiang, China, ⁷School of Life Sciences, Birmingham City University, Birmingham, UK

OPEN ACCESS

Edited by:

Tycho M. Hoogland,
Erasmus MC, Netherlands

Reviewed by:

Richardson N. Leão,
Brain Institute, Brazil
Fiona Elizabeth Nicole LeBeau,
Newcastle University, UK

*Correspondence:

ChengBiao Lu
johnlu9000@hotmail.com

[†]These authors have contributed
equally to this work.

Received: 07 September 2016

Accepted: 15 February 2017

Published: 06 March 2017

Citation:

Wang J, He X, Guo F, Cheng X,
Wang Y, Wang X, Feng Z,
Vreugdenhil M and Lu C
(2017) Multiple Kinases Involved in
the Nicotinic Modulation of Gamma
Oscillations in the Rat Hippocampal
CA3 Area.
Front. Cell. Neurosci. 11:57.
doi: 10.3389/fncel.2017.00057

Neuronal synchronization at gamma band frequency (20–80 Hz, γ oscillations) is closely associated with higher brain function, such as learning, memory and attention. Nicotinic acetylcholine receptors (nAChRs) are highly expressed in the hippocampus, and modulate hippocampal γ oscillations, but the intracellular mechanism underlying such modulation remains elusive. We explored multiple kinases by which nicotine can modulate γ oscillations induced by kainate in rat hippocampal area CA3 *in vitro*. We found that inhibitors of cyclic AMP dependent kinase (protein kinase A, PKA), protein kinase C (PKC), N-methyl-D-aspartate receptor (NMDA) receptors, Phosphoinositide 3-kinase (PI3K) and extracellular signal-related kinases (ERK), each individually could prevent the γ oscillation-enhancing effect of 1 μ M nicotine, whereas none of them affected baseline γ oscillation strength. Inhibition of the serine/threonine kinase Akt increased baseline γ oscillations and partially blocked its nicotinic enhancement. We propose that the PKA-NMDAR-PI3K-ERK pathway modifies cellular properties required for the nicotinic enhancement of γ oscillations, dependent on a PKC-ERK mediated pathway. These signaling pathways provide clues for restoring γ oscillations in pathological conditions affecting cognition. The suppression of γ oscillations at 100 μ M nicotine was only dependent on PKA-NMDAR activation and may be due to very high intracellular calcium levels.

Keywords: hippocampus, nAChR, PKA, PKC, ERK1/2, PI3 kinase, Akt, γ oscillations

INTRODUCTION

Hippocampal oscillations reflect the coordinated, rhythmic activity emerging from mutually connected interneuron and principal cell populations. Fast network oscillations in the gamma frequency band (30–80 Hz, γ) are associated with higher brain functions, such as visual attention and memory (Fries et al., 2001; Lutzenberger et al., 2002; Jensen et al., 2007). γ oscillations provide the ms-precision timing matrix necessary for effective information processing and the maintenance of behaviorally relevant information in working memory (Roux et al., 2012). γ oscillations are impaired in neuropsychological disorders such as schizophrenia and Alzheimer's disease which may explain the deficits of working memory in these disorders (Cho et al., 2006; Leicht et al., 2010; Klein et al., 2016).

Nicotine modulates neuronal network activity via nicotinic acetylcholine receptor (nAChR) activation. Nicotine induced hippocampal theta oscillations (Lu and Henderson, 2010) and increased the auditory-evoked or the stimulation-evoked hippocampal γ oscillations (Song et al., 2005; Featherstone et al., 2012), which may well explain the improvement of cognitive function in both normal subjects and patients with schizophrenia (Phillips et al., 2007). Recently, we found that nicotine at a low concentration increases hippocampal γ oscillations induced by kainate *in vitro*, while at a high concentration, nicotine decreases γ activity (Wang et al., 2015). However, the intracellular mechanism involved in nicotinic modulation of γ oscillation is unclear.

nAChR are ligand-gated cation selective channels that are widely distributed in the brain and classified as non- $\alpha 7$ and $\alpha 7$ nAChR subtypes, based on the calcium permeability of these channels, with $\alpha 7$ nAChR having relative high calcium permeability (Hu et al., 2002). These nAChRs may be selectively expressed in specific neuronal populations, and modulate neuronal function. For example, the selective expression of $\alpha 2$ nAChR in a subset of hippocampus area CA1 interneurons (Oriens-Lacunosum Moleculare cells) modulates information flow and long-term potentiation (LTP) in the CA1 region (Leão et al., 2012).

Nicotine activates cAMP-dependent protein kinase A (PKA) and extracellular signal-related kinase (ERK 1/2) via $\alpha 7$ nAChR activation in hippocampal neurons (Dajas-Bailador et al., 2002). The cAMP/PKA-mitogen-activated protein kinase (MAPK) pathway is a signal-transduction cascade which is neuron-specific (Waltereit and Weller, 2003). In addition, nicotine may also activate protein kinase C (PKC) to increase the density of non- $\alpha 7$ nAChR ($\alpha 4\beta 2$ nAChR) in the cell membrane via PKC-dependent phosphorylation of the $\alpha 4$ subunit (Wecker et al., 2010). Furthermore, the nicotine-induced phosphorylation of PKC may be an upstream event for the increased phosphorylation of ERK1/2 (Sugano et al., 2006). Thus, the PKC-ERK signaling cascade may also contribute to nicotinic modulation of γ oscillations.

Regulation of neuronal N-methyl-D-aspartate receptors (NMDARs) by protein kinases such as PKA and PKC is critical in synaptic transmission and plasticity (Lan et al., 2001; Skeberdis et al., 2006). NMDAR activation mediates the phosphorylation of ERK and Akt/PKB (a serine and threonine protein kinase and an effector of a phosphatidylinositol 3-kinase (PI3K); Perkinson et al., 2002; Crossthwaite et al., 2004). Interestingly, NMDAR inhibition abolished the effect of nicotine on γ oscillations (Wang et al., 2015). Therefore, nicotine may modulate γ oscillations through NMDAR-mediated Ca^{2+} influx, activating MAPK.

Nicotinic modulation of synaptic plasticity is associated with the selective activation of some kinases (PKA, ERK and PI3K), but not of other kinases (tyrosine kinase, Src, Ca^{2+} -calmodulin-dependent protein kinase II or p38 MAPK; Welsby et al., 2009). Therefore, in order to establish by what signaling pathways nicotine can modulate γ oscillations, we tested the

role of multiple kinases, including PKA, PKC, ERK, PI3K and Akt in addition to the role of the NMDAR in nicotinic modulation of γ oscillations. We found that nicotinic modulation of γ oscillation requires both PKA and NMDAR activity and is partially blocked by PKC, ERK1/2 and PI3K inhibitors, suggesting that the NMDAR-PKA signaling pathway plays a critical role whereas PKC, ERK 1/2 and PI3K play a less critical role in acute nicotinic modulation of hippocampal γ oscillations.

MATERIALS AND METHODS

Animals

All procedures were carried out in accordance with the UK Animals (Scientific Procedures) Act 1986 and with Animal Ethics and Administrative Council of Henan province, P.R. China, and all efforts were made to minimize animal suffering and to reduce the number of animals used. Electrophysiological studies were performed on hippocampal slices prepared from Wistar rats (male, 3–6 week-old).

Animals were anesthetized by intraperitoneal injection of Sagatal (sodium pentobarbitone, 100 mg kg^{-1} , Rhône Mérieux Ltd, Harlow, UK). When all pedal reflexes were abolished, the animals were perfused intracardially with chilled (5°C), sucrose-based solution contained (in mM): 225 sucrose, 3 KCl, 1.25 NaH_2PO_4 , 24 NaHCO_3 , 6 MgSO_4 , 0.5 CaCl_2 and 10 glucose (305 mOsm l^{-1}). Horizontal slices (450 μm) of the brain were cut in the sucrose-based solution, using a Leica VT1000S vibratome (Leica Microsystems UK, Milton Keynes, UK).

Electrophysiological Recording, Data Acquisition and Analysis

For extracellular field potential recordings, two hippocampal slices were transferred to an interface recording chamber. The slices were maintained at a temperature of 33°C at the interface between artificial cerebrospinal fluid (ACSF) and warm humidified carbogen gas (95% O_2 –5% CO_2). The ACSF contained (in mM): 126 NaCl, 3 KCl, 1.25 NaH_2PO_4 , 24 NaHCO_3 , 2 MgSO_4 , 2 CaCl_2 and 10 Glucose (pH 7.4, 310 mOsm l^{-1}). The slices were allowed to equilibrate in this medium for 1 h prior to recording. The local field potential was recorded from stratum pyramidale of area CA3c, using glass microelectrodes containing ACSF (resistance 2–5 $\text{M}\Omega$). Field potentials were amplified with an Axoprobe 1A amplifier (Axon Instruments, Union City, CA, USA) and band-pass filtered between 0.5 Hz and 2 kHz using a Neurolog NL125 (Digitimer Ltd, Welwyn Garden City, UK). Electrical interference from the mains supply was eliminated with 50 Hz noise eliminators (HumBug; Digitimer Ltd). Data were then digitized at a sample rate of 5–10 kHz using a CED 1401 plus ADC board (Cambridge Electronic Design, Cambridge, UK).

Data were analyzed off-line using Spike 2 software (Cambridge Electronic Design). Power spectra were generated using a fast Fourier transform algorithm to provide a quantitative measure of the frequency components in a 60-s recording epoch,

where power, a quantitative measure of the oscillation strength, was plotted as function of frequency. The parameters used for measuring the oscillatory activity in the slice were peak frequency and γ power, calculated as the summated power in 20–60 Hz range.

All statistical tests were performed using SigmaStat software (SPSS Inc., San Francisco, CA, USA). Results are expressed as mean \pm standard error of mean. Statistical significance for comparison between two groups or between three groups was determined using tests described in the text or in the figure legends, as appropriate. As γ power varies strongly between slices, the effect of drugs was expressed as a change, normalized to pre-drug values and the statistical comparisons were performed on normalized changes. Measures were considered statistically significant if $P < 0.05$.

Drugs used for Electrophysiology

All standard reagents, except where indicated, were obtained either from Sigma-Aldrich (UK). D-(–)-2-amino-5-phosphonopentanoic acid (D-AP5), bicuculline methochloride and 2,3,-dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide (NBQX), 1,4-Diamino-2,3-dicyano-1,4-bis(o-aminophenylmercapto) butadiene (U0126), wortmannin and H-89 were purchased from Tocris Cookson Ltd (Bristol, UK). Gö6983 was purchased from Santa Cruz Biotechnology (Dallas, Tx, USA). Triciribine (TCBN) was purchased from Sigma-Aldrich (Dorset, UK). Stock solutions, at 1000 times the working concentration, were made up in water or in dimethylsulphoxide, and were stored in individual aliquots at -45°C . Working solutions were prepared freshly on the day of the experiment.

RESULTS

Nicotinic Modulation of Hippocampal γ Oscillations

Oscillatory activity was induced in the slices by addition of 200 nM kainate to the ACSF and allowed to develop and stabilize for 1 h before further drugs were applied. We have previously reported that the persistent γ oscillation (20–60 Hz) induced by kainate in rat hippocampal CA3 area can be either enhanced or reduced by nicotine, depending on the concentration of nicotine used. This was confirmed in this study by acute perfusion of nicotine at 1 μM or 100 μM (Figure 1). At 1 μM nicotine caused a $83.4 \pm 21.6\%$ increase ($P < 0.01$, paired t test, $n = 13$) and at 100 μM a $54.9 \pm 5.2\%$ decrease ($P < 0.001$, paired t test, $n = 12$; Figure 1C).

Kainate-induced oscillation peak frequency was 28.4 ± 0.5 Hz, $n = 23$ and was not significantly affected by 1 μM nicotine (27.3 ± 0.8 Hz, $P > 0.05$, paired t test, $n = 13$) or 100 μM nicotine (26.9 ± 0.7 Hz, $P > 0.05$, paired t test, $n = 12$).

The Effect of PKA Inhibition on the Nicotinic Modulation of γ Oscillations

To test whether PKA is involved in the modulation of γ oscillations by nicotine, we perfused the slices with the PKA inhibitor H89 (10 μM for 40 min), followed by addition of


nicotine (1 μM for 30 min or 100 μM for 30 min in two different sets of experiments). H89 alone had no effect on γ power ($101.6 \pm 3.8\%$ of baseline, one way repeated measures (RM) ANOVA, $P > 0.05$, $n = 12$). In the presence of H89, 1 μM nicotine caused no significant change in γ power ($107.3 \pm 5.6\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. H89, $n = 12$, Figure 2). Further application of 100 μM nicotine caused no significant change in γ power neither ($95.8 \pm 5.9\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. H89, $n = 8$, Figure 2). The complete block of the nicotinic modulation of γ oscillations by inhibition of PKA inhibition, suggests that PKA is required for the nicotinic modulation of γ oscillations.

The Effect of PKC Inhibition on the Nicotinic Modulation of γ Oscillations

Nicotine activates PKC via non- $\alpha 7$ nAChR activation (Wecker et al., 2010). To test whether PKC is involved in the nicotinic modulation of γ oscillations, we perfused slices with the conventional PKC inhibitor Gö6983 (10 μM), followed by nicotine application. Gö6983 had no effect on γ power ($95.6 \pm 5.8\%$ of baseline, one way RM ANOVA, $P > 0.05$, $n = 5$). In the presence of Gö6983 nicotine caused no significant change in γ power at 1 μM ($100.6 \pm 4.3\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. Gö6983, $n = 5$, Figure 3), but caused a significant decrease in γ power at 100 μM ($41.2 \pm 5.6\%$ of baseline, one way RM ANOVA, $P < 0.001$, vs. Gö6983, $n = 5$, Figure 3), which was not different from the effect of 100 μM nicotine alone (t test, $P > 0.05$). These results suggest that PKC is only involved in nicotinic enhancement of γ oscillations, not in the suppression at a high nicotine concentration.

The Effect of NMDA Receptor Antagonism on the Nicotinic Modulation of γ Oscillations

Nicotine enhances NMDA receptor function through activation of nAChR (Aramakis and Metherate, 1998; Yang et al., 2013). We further tested the role of NMDAR activation on nicotinic modulation of γ oscillations. Application of the NMDA receptor antagonist D-AP5 (1 μM) alone had no effect on γ power ($109.7 \pm 4.3\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. baseline γ , $n = 8$), but blocked the nicotinic modulation of γ oscillations (example in Figures 4A,B). In the presence of D-APV, nicotine (1 μM) caused no significant change in γ power ($106.3 \pm 5.9\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. D-AP5, $n = 8$, Figure 4E), which was smaller than the effect of nicotine alone (Student t -test, $P < 0.001$). Similarly, in a different set of experiments D-AP5 reduced the suppression of γ oscillations by 100 μM nicotine (example in Figures 4C,D), D-AP5 (1 μM) had no effect on control γ power ($112.9 \pm 5.3\%$ of baseline, one way RM ANOVA, $P > 0.05$, vs. baseline γ , $n = 8$), but reduced the suppression of γ power by 100 μM nicotine (104.7 ± 10.3 of baseline, one way RM ANOVA, $P > 0.05$, vs. D-AP5, $n = 8$; Figure 4E). In comparison with the effect of 100 μM nicotine alone, the effect of D-AP5 + nicotine on γ power ($11.07 \pm 7.7\%$ decrease) was significantly smaller (Student t -test, $P < 0.001$, vs. nicotine alone). The prevention of the nicotinic



modulation of γ oscillations by D-AP5, suggests that NMDA receptor activation is required for the nicotinic modulation of γ oscillations.

The Effect of ERK Inhibition on the Nicotinic Modulation of γ Oscillations

NMDAR activation mediates the phosphorylation of ERK (Perkinton et al., 2002), which suggests that ERK is a downstream molecule of NMDAR. ERK1 and ERK2 are two closely related MAPK. Previous studies indicate that nicotine induced phosphorylation of ERK via $\alpha 7$ nAChR in hippocampal neurons (Dajas-Bailador et al., 2002). We thus examined whether ERK1/2 activation is required for the nicotinic modulation of

γ oscillations. U0126, a selective MEK kinase inhibitor, known to block ERK1/2 phosphorylation, was used to test the role of ERK1/2 in the nicotinic modulation of γ oscillations. U0126 (2.5 μ M) alone had no effect on γ oscillations (Figure 5) and subsequent application of nicotine (1 μ M) caused no significant change in γ power ($101.8 \pm 0.9\%$ of baseline, $P > 0.05$ vs. U0126, $n = 8$, Figure 5). In the presence of U0126, 100 μ M nicotine reduced γ power by $19.3 \pm 3.0\%$ ($P < 0.001$, vs. U0126, $n = 8$, Figure 5), which was less than the effect of 100 μ M nicotine alone ($49 \pm 3.7\%$ decrease, Student t -test, $P < 0.001$). These results suggest that ERK1/2 is involved in the nicotinic modulation of γ oscillations.




FIGURE 2 | The effect of protein kinase A (PKA) inhibitor on the nicotinic modulation of γ oscillations. (A,C) The extracellular field potential recordings from the CA3 stratum pyramidale, for the kainate alone (control, trace taken at a time point that γ oscillation was stable for 60 min after kainate), in the presence of PKA inhibitor, 10 μ M H89 (40 min after H89), and after additional application of 1 μ M nicotine (30 min after nicotine; **A**) or 100 μ M nicotine (30 min after nicotine; **C**). **(B,D)** The power spectra corresponding to **(A,C)**. **(E)** γ power as % of baseline γ power for control, in the presence of H89, and with additional application of 1 μ M nicotine ($n = 12$) or 100 μ M nicotine ($n = 8$).




FIGURE 3 | The effect of protein kinase C (PKC) inhibitor on the nicotinic modulation of γ oscillations. (A,C) The extracellular field recordings from CA3 pyramidal for the kainate alone, in the presence of PKC inhibitor, 10 μ M Gö6983 (40 min after Gö6983), and after additional application of 1 μ M nicotine (30 min after nicotine; **A**) or 100 μ M nicotine (30 min after nicotine; **C**). **(B,D)** The power spectra corresponding to **(A,C)**. **(E)** γ power as % of baseline γ power for the kainate alone, Gö6983 and with additional application of 1 μ M nicotine ($n = 5$) or 100 μ M nicotine ($n = 5$) ($***P < 0.001$, one way RM ANOVA, compared with Gö6983 alone).





FIGURE 4 | The effect of N-methyl-D-aspartate receptor (NMDA) receptor antagonist on the nicotinic modulation of γ oscillations. (A,C) The extracellular field recordings from CA3 pyramidal for the kainate alone, NMDA receptor antagonist, 1 μ M D-AP5 (40 min after D-AP5), and after additional application of 1 μ M nicotine (30 min after nicotine; **A**) or 100 μ M nicotine (30 min after nicotine; **C**). **(B,D)** The power spectra corresponding to **(A,C)**. **(E)** γ power as % of baseline γ power for control, D-AP5, and with additional application of 1 μ M nicotine ($n = 8$) or 100 μ M nicotine ($n = 8$).

The Effects of PI3 Kinase Inhibitor on Nicotinic Modulation of γ Oscillation

PI3K is a central mediator of NMDA receptor signaling to Erk1/2 (Perkinton et al., 2002) and $\alpha 7$ nAChR activation reduced

amyloid β neurotoxicity through PI3K activation (Kihara et al., 2001). We therefore tested whether PI3K is involved in the effect of nicotine on γ oscillations, using wortmannin, a cell-permeable, irreversible inhibitor of PI3K. Wortmannin (0.2 μ M)



alone had no effects on γ oscillations ($104.5 \pm 9.3\%$ of baseline, $P > 0.05$, $n = 8$). Subsequent application of nicotine (1 μM) caused no significant change in γ power ($102.1 \pm 5.3\%$ of

baseline, $P > 0.05$, vs. wortmannin, $n = 8$). In the presence of wortmannin the 100 μM nicotine-induced reduction of γ was $13.2 \pm 2.8\%$ ($P < 0.05$, vs. wortmannin, $n = 8$,

Figure 6), which was smaller than that of 100 μ M nicotine alone ($49 \pm 3.7\%$ decrease, Student *t*-test, $P < 0.01$). These results suggest that PI3K is involved in the nicotinic modulation of γ oscillations.

The Effects of Akt Inhibitor on Nicotinic Modulation of γ Oscillation

NMDAR activation also mediates the phosphorylation of Akt/PKB in addition to that of ERK (Perkinton et al., 2002). We then tested whether Akt, a serine and threonine protein kinase and an effector of a PI3K, affects nicotinic modulation of γ oscillations, using Triciribine (TCBN), a selective Akt inhibitor. TCBN (5 μ M) caused a small but significant increase in γ power ($118.1 \pm 6.5\%$ of baseline, $P < 0.05$ vs. baseline, $n = 9$). Subsequent application of nicotine (1 μ M) caused a $23.3 \pm 11.3\%$ increase in γ power ($P < 0.01$, vs. TCBN, $n = 9$, **Figure 7**), which is smaller than that in the absence of TCBN (Student *t*-test, $P < 0.05$). In the presence of TCBN, 100 μ M nicotine caused a $36.7 \pm 7.7\%$ decrease in γ power ($P < 0.01$, vs. TCBN, $n = 9$, **Figure 7**), which was not different from the effect of nicotine alone (Student *t* test, $P > 0.05$). These results indicate that Akt activation is involved in the nicotinic modulation of γ oscillations.

DISCUSSION

In this study, we found that activation of a PKA/PKC-NMDAR-ERK/PI3K signaling pathway is required for nicotinic enhancement of γ oscillations, with a partial involvement of Akt. PKA and NMDAR activation are also required for inhibition of γ oscillations by high concentration of nicotine, with a partial involvement of the downstream signaling kinase ERK or PI3K. None of above-mentioned kinases except for Akt is directly involved in the generation of γ oscillations.

Signaling Pathways Potentially Involved in the γ Oscillation Enhancing Effect of Low Concentrations of Nicotine

Our data are consistent with the nicotinic enhancement of auditory evoked γ oscillations in mouse hippocampus via $\alpha 4\beta 2$ nAChR activation at low micromolar concentrations (Phillips et al., 2007; Featherstone et al., 2012). The nicotinic γ oscillation enhancement may be related to the activation of NMDAR and subsequent calcium influx, as D-AP5, in line with previous observations (Wang et al., 2015), completely blocked the effects of nicotine. Calcium influx through the NMDAR is the start of many signaling pathways that alter neuronal and network functions (Perkel et al., 1993; Sobczyk and Svoboda, 2007). The lack of effect of the NMDAR antagonist on γ oscillations replicates our previous findings (Wang et al., 2015; Chen et al., 2016) and that of others (Cunningham et al., 2003; Fisahn et al., 2004). It indicates that under normal slice conditions there is little NMDAR activation. How can NMDAR inhibition have such an effect on the nicotinic enhancement of γ oscillations? One explanation

is that nAChR receptor activation causes an enhancement of NMDAR activation to levels that give substantial calcium influx and calcium-dependent signaling pathways that subsequently affect properties that determine γ oscillation strength. Indeed, it has been reported that $\alpha 7$ nAChRs form a protein complex with the NMDAR through a protein-protein interaction, and activation of $\alpha 7$ nAChR upregulates NMDAR-mediated whole cell currents (Li et al., 2013). With both PI3K and ERK1/2 activity required for the γ oscillation-enhancing effect of nicotine, one potential downstream signaling pathway has been identified. ERK1/2 is known to be one of the downstream molecules of NMDAR activation and PI3K is a central mediator of NMDAR signaling to ERK1/2 (Perkinton et al., 2002; Crossthwaite et al., 2004; Skeberdis et al., 2006).

Stimulation of $\alpha 7$ nAChR activates PI3K in cultured cortical neurons (Kihara et al., 2001) and activation of nAChRs engages calcium-dependent signaling pathways including PI3K (Liu et al., 2008). The ERK activation may be a common pathway for both nAChR and NMDA receptor activation, as stimulation of $\alpha 7$ -nAChRs activates calcium-dependent intracellular signaling molecules including ERK1/2 (Chang and Berg, 2001; Hu et al., 2002; Gubbins et al., 2010). Inhibition of PKA blocked the nicotine-induced ERK1/2 activity in hippocampal neurons in cultures (Dajas-Bailador et al., 2002), thus PKA could be an upstream kinase of ERK1/2 (**Figure 8**). Although our study is not exhaustive we have identified the PI3K-ERK1/2 pathway as essential for the NMDAR-mediated nicotinic enhancement of γ oscillations. Akt is a putative downstream molecule of PI3K and a downstream molecule of NMDAR activation, since it was reported that in striatal neurons NMDAR-dependent Akt activation was mediated by PI3K (Crossthwaite et al., 2004). Akt was only partially involved in the γ oscillation enhancement. The Akt inhibitor caused an increase in γ oscillations suggesting that Akt activation is suppressing γ oscillations *in vitro*, which is in agreement with our recent observation (Wang et al., 2016).

How can NMDAR-mediated activation of the PI3K-ERK/Akt signaling pathway affect γ oscillation strength? γ oscillations emerge from the rhythmic IPSCs generated by interneurons, activated by pyramidal cells in a feedback manner that, in turn, synchronize the firing of neurons (Mann and Paulsen, 2005). If acting on interneurons the PI3K-ERK/Akt pathway could affect γ oscillations by enhancing intrinsic excitability and GABA release. Although the signaling pathway involved is unknown, $\alpha 7$ nAChR activation modulates hippocampal interneurons and enhances GABA release (Hulo and Muller, 2001; Wanaverbecq et al., 2007), which would increase γ oscillations. If acting on pyramidal neurons the PI3K-ERK/Akt pathway could affect γ oscillations by increasing intrinsic excitability and glutamate release.

PKA activation was also required for the nicotinic enhancement of γ oscillations and can form the link between nAChR activation and NMDAR-mediated signaling pathways responsible for the enhancement. PKA can be activated by nAChR activation (Dajas-Bailador et al., 2002) and can, in turn, phosphorylate NMDARs, causing an increase in Ca^{2+} permeability (Skeberdis et al., 2006). It is, however, remarkable






FIGURE 6 | The effect of Phosphoinositide 3-kinase (PI3K) inhibitor on the nicotinic modulation of γ oscillations. (A,C) The extracellular field recordings from CA3 pyramidal for the kainate alone, in the presence of the PI3K inhibitor 200 nM wortmannin (40 min after nicotine), and after additional application of 1 μ M nicotine (30 min after nicotine; **A**) or 100 μ M nicotine (30 min after nicotine; **C**). **(B,D)** The power spectra corresponding to **(A,C)**. **(E)** γ power as % of baseline γ power for the kainate alone, wortmannin, and with additional application of 1 μ M nicotine ($n = 8$) or 100 μ M nicotine ($n = 8$; * $P < 0.05$, one way RM ANOVA, compared with wortmannin alone).



that this NMDAR-mediated calcium influx is required while nAChR are Ca^{2+} permeable themselves (Didier et al., 1995). Previously we have shown that for the full nicotinic γ oscillation enhancement both $\alpha_4\beta_2$ and α_7 type nAChRs are required, but mostly the $\alpha_4\beta_2$ type (Wang et al., 2015), which is less Ca^{2+} permeable than the α_7 type (Fucile, 2004). Furthermore, the location of the limited calcium influx through nAChRs may be

different from the location of the NMDAR-mediated calcium influx required to activate the signaling pathway responsible for the modulation of γ oscillations.

PKC activation was also required for the nicotinic enhancement of γ oscillations. PKC activation can increase NMDA channel opening and insert more NMDA channels into the plasma membrane (Lan et al., 2001), thus increasing



NMDAR-mediated calcium influx. However, since both PKA and PKC activation is required, it is unlikely that PKC and PKA activation converge on NMDAR modulation, because then the effects would be additive. PKC activates ERK1/2 (Cox et al., 2004), so could also converge with the PKA-NMDAR-PI3K pathway to activate ERK1/2 in a way that requires both. Alternatively, it is possible that PKC activation act directly on one or more of the γ oscillation strength determining factors (intrinsic excitability and neurotransmitter release), in co-dependence with the PKA-NMDA-IP3K-ERK/Akt pathway (Figure 8).

Impaired γ oscillations have been associated with cognitive deficits typical to schizophrenia (Cho et al., 2006) and Alzheimer's disease (Klein et al., 2016). Nicotine use is increased in schizophrenia patients, where it has cognitive benefits (Boggs et al., 2014). Also nicotine increases visual attention and working

memory in healthy volunteers (Ernst et al., 2001). It is possible that this is due to nicotinic enhancement of γ oscillations, which was significant at concentrations as low as 0.1 μM (Wang et al., 2015), comparable to plasma levels achieved by cigarette smoking. Further elucidating the pathways involved may well lead to identifying therapeutic targets that can increase γ oscillations effectively and safely.

A limitation of this study is that we have not defined the specific cell subtypes that are involved in the nicotinic enhancement of γ oscillations. It appears that expression of nAChR subtypes in hippocampus is highly selective, which determines the precise modulation of specific synapse function (Leão et al., 2012). Thus, defining the specific cell type which may contribute to the nicotinic enhancement of γ oscillations has to be the focus of future studies.

Signaling Pathways Potentially Involved in the γ Oscillation Suppressing Effect of High Concentrations of Nicotine

Both blocking NMDARs and PKA inhibition completely prevented the suppression of γ oscillations at a high concentration of nicotine, whereas the downstream molecules in the pathway (ERK, PI3K) only partially contributed to the suppression, and PKC and Akt were not involved. These results suggest that nicotinic enhancement or inhibition of γ oscillations involves both shared and different mechanisms.

The suppressing effect of high concentrations of nicotine was stronger in the presence of nAChR antagonists (Wang et al., 2015), which seems to indicate an effect of nicotine on the NMDAR bypassing the nAChR. One possibility is that the intensity of NMDAR activation and subsequent calcium influx may differentiate between enhancement (low calcium influx) and suppression (high calcium influx). Because γ oscillations are a high energy demanding neuronal activity (Kann et al., 2011), a large calcium influx may overwhelmingly overcome the role of downstream kinases such as ERK, PI3K or Akt, causing the collapse in γ oscillations via mitochondrial membrane depolarization (Kann et al., 2011; Lu et al., 2012). A reverse correlation between intracellular calcium and γ oscillations was observed in aged hippocampal neurons (Lu et al., 2011, 2012). The physiological role of the γ oscillation-suppressing effect of high concentrations of nicotine is limited, because plasma nicotine levels will normally not exceed 1 μM .

Possible Cellular Mechanisms of Nicotinic Modulations of γ Oscillations

Our results reveal that activation of a signaling pathway of PKA/PKC-NMDA-ERK/PI3K contributes to the nicotinic modulation of γ oscillations. Nicotine may increase NMDAR activation through PKA- and PKC-dependent phosphorylation, which leads to the Ca^{2+} influx and activation of the downstream kinases such as ERK, PI3K and Akt and subsequently modulation of γ oscillations. The nicotinic modulation of γ oscillations may depend on the intensity of NMDAR activation, a weak or strong activation of NMDAR by a low or high concentration of nicotine enhances or inhibits γ , respectively (Figure 8).

AUTHOR CONTRIBUTIONS

CL designed the experiment; JW, FG, XH, XC, YW and XW performed the experiments; CL, JW, ZF, XH and MV wrote the manuscript; CL, JW, XH, XC, YW and XW analyzed the data. All authors reviewed the manuscript.

REFERENCES

Aramakis, V. B., and Metherate, R. (1998). Nicotine selectively enhances NMDA receptor-mediated synaptic transmission during postnatal development in sensory neocortex. *J. Neurosci.* 18, 8485–8495.

Boggs, D. L., Carlson, J., Cortes-Briones, J., Krystal, J. H., and D'Souza, D. C. (2014). Going up in smoke? A review of nAChRs-based treatment strategies for improving cognition in schizophrenia. *Curr. Pharm. Des.* 20, 5077–5092. doi: 10.2174/1381612819666131216121019

Chang, K. T., and Berg, D. K. (2001). Voltage-gated channels block nicotinic regulation of CREB phosphorylation and gene expression in neurons. *Neuron* 32, 855–865. doi: 10.1016/s0896-6273(01)00516-5

Chen, K., Nakauchi, S., Su, H., Tanimoto, S., and Sumikawa, K. (2016). Early postnatal nicotine exposure disrupts the $\alpha 2^*$ nicotinic acetylcholine receptor-mediated control of oriens-lacunosum moleculare cells during adolescence in rats. *Neuropharmacology* 101, 57–67. doi: 10.1016/j.neuropharm.2015.09.022

Cho, R. Y., Konecky, R. O., and Carter, C. S. (2006). Impairments in frontal cortical γ synchrony and cognitive control in schizophrenia. *Proc. Natl. Acad. Sci. U S A* 103, 19878–19883. doi: 10.1073/pnas.0609440103

Cox, S., Harvey, B. K., Sanchez, J. F., Wang, J. Y., and Wang, Y. (2004). Mediation of BMP7 neuroprotection by MAPK and PKC IN rat primary cortical cultures. *Brain Res.* 1010, 55–61. doi: 10.1016/j.brainres.2004.02.068

Crossthwaite, A. J., Valli, H., and Williams, R. J. (2004). Inhibiting Src family tyrosine kinase activity blocks glutamate signalling to ERK1/2 and Akt/PKB but not JNK in cultured striatal neurones. *J. Neurochem.* 88, 1127–1139. doi: 10.1046/j.1471-4159.2004.02257.x

Cunningham, M. O., Davies, C. H., Buhl, E. H., Kopell, N., and Whittington, M. A. (2003). γ oscillations induced by kainate receptor activation in the entorhinal cortex *in vitro*. *J. Neurosci.* 23, 9761–9769.

Dajas-Bailador, F. A., Soliakov, L., and Wonnacott, S. (2002). Nicotine activates the extracellular signal-regulated kinase 1/2 via the $\alpha 7$ nicotinic acetylcholine receptor and protein kinase A, in SH-SY5Y cells and hippocampal neurones. *J. Neurochem.* 80, 520–530. doi: 10.1046/j.0022-3042.2001.00725.x

Didier, M., Berman, S. A., Lindstrom, J., and Bursztajn, S. (1995). Characterization of nicotinic acetylcholine receptors expressed in primary cultures of cerebellar granule cells. *Mol. Brain Res.* 30, 17–28. doi: 10.1016/0169-328x(94)00266-h

Ernst, M., Heishman, S. J., Spurgeon, L., and London, E. D. (2001). Smoking history and nicotine effects on cognitive performance. *Neuropsychopharmacology* 25, 313–319. doi: 10.1016/s0893-133x(01)00257-3

Featherstone, R. E., Phillips, J. M., Thieu, T., Ehrlichman, R. S., Halene, T. B., Leiser, S. C., et al. (2012). Nicotine receptor subtype-specific effects on auditory evoked oscillations and potentials. *PLoS One* 7:e39775. doi: 10.1371/journal.pone.0039775

Fisahn, A., Contractor, A., Traub, R. D., Buhl, E. H., Heinemann, S. F., and McBain, C. J. (2004). Distinct roles for the kainate receptor subunits GluR5 and GluR6 in kainate-induced hippocampal γ oscillations. *J. Neurosci.* 24, 9658–9668. doi: 10.1523/JNEUROSCI.2973-04.2004

Fries, P., Reynolds, J. H., Rorie, A. E., and Desimone, R. (2001). Modulation of oscillatory neuronal synchronization by selective visual attention. *Science* 291, 1560–1563. doi: 10.1126/science.1055465

Fucile, S. (2004). Ca^{2+} permeability of nicotinic acetylcholine receptors. *Cell Calcium* 35, 1–8. doi: 10.1016/j.ceca.2003.08.006

Gubbins, E. J., Gopalakrishnan, M., and Li, J. (2010). $\alpha 7$ nAChR-mediated activation of MAP kinase pathways in PC12 cells. *Brain Res.* 1328, 1–11. doi: 10.1016/j.brainres.2010.02.083

Hu, M., Liu, Q. S., Chang, K. T., and Berg, D. K. (2002). Nicotinic regulation of CREB activation in hippocampal neurons by glutamatergic and nonglutamatergic pathways. *Mol. Cell. Neurosci.* 21, 616–625. doi: 10.1006/mcne.2002.1202

ACKNOWLEDGMENTS

This study was supported by the National Natural Science Foundation of China (NSFC, Grant Nos. 81271422; 31070938) and by the Project of TaiHang Scholar of Xinxiang Medical University.

Hulo, S., and Muller, D. (2001). Tetrodotoxin-sensitive enhancement of inhibition in CA1 pyramidal neurones by nicotine. *Neuroreport* 12, 1351–1354. doi: 10.1097/00001756-200105250-00012

Jensen, O., Kaiser, J., and Lachaux, J. P. (2007). Human γ -frequency oscillations associated with attention and memory. *Trends Neurosci.* 30, 317–324. doi: 10.1016/j.tins.2007.05.001

Kann, O., Huchzermeyer, C., Kovács, R., Wirtz, S., and Schuelke, M. (2011). γ oscillations in the hippocampus require high complex I gene expression and strong functional performance of mitochondria. *Brain* 134, 345–358. doi: 10.1093/brain/awq333

Kihara, T., Shimohama, S., Sawada, H., Honda, K., Nakamizo, T., Shibasaki, H., et al. (2001). $\alpha 7$ nicotinic receptor transduces signals to phosphatidylinositol 3-kinase to block A β -amyloid-induced neurotoxicity. *J. Biol. Chem.* 276, 13541–13546. doi: 10.1074/jbc.M008035200

Klein, A. S., Donoso, J. R., Kempter, R., Schmitz, D., and Beed, P. (2016). Early cortical changes in γ oscillations in Alzheimer's disease. *Front. Syst. Neurosci.* 10:83. doi: 10.3389/fnsys.2016.00083

Lan, J. Y., Skeberdis, V. A., Jover, T., Grooms, S. Y., Lin, Y., Aranedo, R. C., et al. (2001). Protein kinase C modulates NMDA receptor trafficking and gating. *Nat. Neurosci.* 4, 382–390. doi: 10.1038/86028

Leão, R. N., Mikulovic, S., Leão, K. E., Munguba, H., Gezelius, H., Enjin, A., et al. (2012). OLM interneurons differentially modulate CA3 and entorhinal inputs to hippocampal CA1 neurons. *Nat. Neurosci.* 15, 1524–1530. doi: 10.1038/nn.3235

Leicht, G., Kirsch, V., Giegling, I., Karch, S., Hantschk, I., Möller, H. J., et al. (2010). Reduced early auditory evoked γ -band response in patients with schizophrenia. *Biol. Psychiatry* 67, 224–231. doi: 10.1016/j.biopsych.2009.07.033

Li, S., Nai, Q., Lipina, T. V., Roder, J. C., and Liu, F. (2013). $\alpha 7$ nAChR/NMDAR coupling affects NMDAR function and object recognition. *Mol. Brain* 6:58. doi: 10.1186/1756-6606-6-58

Liu, Z., Conroy, W. G., Stawicki, T. M., Nai, Q., Neff, R. A., and Berg, D. K. (2008). EphB receptors co-distribute with a nicotinic receptor subtype and regulate nicotinic downstream signaling in neurons. *Mol. Cell. Neurosci.* 38, 236–244. doi: 10.1016/j.mcn.2008.02.013

Lu, C. B., Hamilton, J. B., Powell, A. D., Toescu, E. C., and Vreugdenhil, M. (2011). Effect of ageing on CA3 interneuron sAHP and γ oscillations is activity-dependent. *Neurobiol. Aging* 32, 956–965. doi: 10.1016/j.neurobiolaging.2009.05.006

Lu, C. B., and Henderson, Z. (2010). Nicotine induction of theta frequency oscillations in rodent hippocampus *in vitro*. *Neuroscience* 166, 84–93. doi: 10.1016/j.neuroscience.2009.11.072

Lu, C. B., Vreugdenhil, M., and Toescu, E. C. (2012). The effect of aging-associated impaired mitochondrial status on kainate-evoked hippocampal γ oscillations. *Neurobiol. Aging* 33, 2692–2703. doi: 10.1016/j.neurobiolaging.2012.01.001

Lutzenberger, W., Ripper, B., Busse, L., Birbaumer, N., and Kaiser, J. (2002). Dynamics of γ -band activity during an audiospatial working memory task in humans. *J. Neurosci.* 22, 5630–5638.

Mann, E. O., and Paulsen, O. (2005). Mechanisms underlying γ ('40 Hz') network oscillations in the hippocampus—a mini-review. *Prog. Biophys. Mol. Biol.* 87, 67–76. doi: 10.1016/j.pbiomolbio.2004.06.004

Perkel, D. J., Petrozzino, J. J., Nicoll, R. A., and Connor, J. A. (1993). The role of Ca^{2+} entry via synaptically activated NMDA receptors in the induction of long-term potentiation. *Neuron* 11, 817–823. doi: 10.1016/0896-6273(93)90111-4

Perkinton, M. S., Ip, J. K., Wood, G. L., Crossthwaite, A. J., and Williams, R. J. (2002). Phosphatidylinositol 3-kinase is a central mediator of NMDA receptor signalling to MAP kinase (Erk1/2), Akt/PKB and CREB in striatal neurones. *J. Neurochem.* 80, 239–254. doi: 10.1046/j.0022-3042.2001.00699.x

- Phillips, J. M., Ehrlichman, R. S., and Siegel, S. J. (2007). Mecamylamine blocks nicotine-induced enhancement of the P20 auditory event-related potential and evoked γ . *Neuroscience* 144, 1314–1323. doi: 10.1016/j.neuroscience.2006.11.003
- Roux, F., Wibral, M., Mohr, H. M., Singer, W., and Uhlhaas, P. J. (2012). γ -band activity in human prefrontal cortex codes for the number of relevant items maintained in working memory. *J. Neurosci.* 32, 12411–12420. doi: 10.1523/JNEUROSCI.0421-12.2012
- Skeberdis, V. A., Chevaleyre, V., Lau, C. G., Goldberg, J. H., Pettit, D. L., Suadicani, S. O., et al. (2006). Protein kinase A regulates calcium permeability of NMDA receptors. *Nat. Neurosci.* 9, 501–510. doi: 10.1038/nn1664
- Sobczyk, A., and Svoboda, K. (2007). Activity-dependent plasticity of the NMDA-receptor fractional Ca^{2+} current. *Neuron* 53, 17–24. doi: 10.1016/j.neuron.2006.11.016
- Song, C., Murray, T. A., Kimura, R., Wakui, M., Ellsworth, K., Javedan, S. P., et al. (2005). Role of $\alpha 7$ -nicotinic acetylcholine receptors in tetanic stimulation-induced γ oscillations in rat hippocampal slices. *Neuropharmacology* 48, 869–880. doi: 10.1016/j.neuropharm.2005.01.003
- Sugano, T., Yanagita, T., Yokoo, H., Satoh, S., Kobayashi, H., and Wada, A. (2006). Enhancement of insulin-induced PI3K/Akt/GSK-3 β and ERK signaling by neuronal nicotinic receptor/PKC- α /ERK pathway: up-regulation of IRS-1/-2 mRNA and protein in adrenal chromaffin cells. *J. Neurochem.* 98, 20–33. doi: 10.1111/j.1471-4159.2006.03846.x
- Waltereit, R., and Weller, M. (2003). Signaling from cAMP/PKA to MAPK and synaptic plasticity. *Mol. Neurobiol.* 27, 99–106. doi: 10.1385/mn:27:1:99
- Wanaverbecq, N., Semyanov, A., Pavlov, I., Walker, M. C., and Kullmann, D. M. (2007). Cholinergic axons modulate GABAergic signaling among hippocampal interneurons via postsynaptic $\alpha 7$ nicotinic receptors. *J. Neurosci.* 27, 5683–5693. doi: 10.1523/JNEUROSCI.1732-07.2007
- Wang, Y., Wang, Z., Wang, J., Wang, Y., Henderson, Z., Wang, X., et al. (2015). The modulation of nicotinic acetylcholine receptors on the neuronal network oscillations in rat hippocampal CA3 area. *Sci. Rep.* 5:9493. doi: 10.1038/srep09493
- Wang, J., Zhao, J., Liu, Z., Guo, F., Wang, Y., Wang, X., et al. (2016). Acute ethanol inhibition of γ oscillations is mediated by Akt and GSK3 β . *Front. Cell. Neurosci.* 10:189. doi: 10.3389/fncel.2016.00189
- Wecker, L., Pollock, V. V., Pacheco, M. A., and Pastoor, T. (2010). Nicotine-induced up regulation of $\alpha 4\beta 2$ neuronal nicotinic receptors is mediated by the protein kinase C-dependent phosphorylation of $\alpha 4$ subunits. *Neuroscience* 171, 12–22. doi: 10.1016/j.neuroscience.2010.09.005
- Welsby, P. J., Rowan, M. J., and Anwyl, R. (2009). Intracellular mechanisms underlying the nicotinic enhancement of LTP in the rat dentate gyrus. *Eur. J. Neurosci.* 29, 65–75. doi: 10.1111/j.1460-9568.2008.06562.x
- Yang, Y., Paspalas, C. D., Jin, L. E., Picciotto, M. R., Arnsten, A. F., and Wang, M. (2013). Nicotinic $\alpha 7$ receptors enhance NMDA cognitive circuits in dorsolateral prefrontal cortex. *Proc. Natl. Acad. Sci. U S A* 110, 12078–12083. doi: 10.1073/pnas.1307849110

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Wang, He, Guo, Cheng, Wang, Wang, Feng, Vreugdenhil and Lu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.