



# Mechanisms of Long Non-Coding RNAs in the Assembly and Plasticity of Neural Circuitry

Andi Wang<sup>1†</sup>, Junbao Wang<sup>1†</sup>, Ying Liu<sup>2\*</sup> and Yan Zhou<sup>1,2\*</sup>

<sup>1</sup> Hubei Key Laboratory of Cell Homeostasis, College of Life Sciences, Wuhan University, Wuhan, China, <sup>2</sup> Medical Research Institute, Wuhan University, Wuhan, China

The mechanisms underlying development processes and functional dynamics of neural circuits are far from understood. Long non-coding RNAs (lncRNAs) have emerged as essential players in defining identities of neural cells, and in modulating neural activities. In this review, we summarized latest advances concerning roles and mechanisms of lncRNAs in assembly, maintenance and plasticity of neural circuitry, as well as lncRNAs' implications in neurological disorders. We also discussed technical advances and challenges in studying functions and mechanisms of lncRNAs in neural circuitry. Finally, we proposed that lncRNA studies would advance our understanding on how neural circuits develop and function in physiology and disease conditions.

## OPEN ACCESS

### Edited by:

Minmin Luo,  
Tsinghua University, China

### Reviewed by:

Nicoletta Berardi,  
Consiglio Nazionale Delle Ricerche  
(CNR), Italy  
Xiaosong Gu,  
Nantong University, China  
Chuan Zhou,  
Institute of Zoology (CAS), China

### \*Correspondence:

Ying Liu  
y.liu@whu.edu.cn  
Yan Zhou  
yan.zhou@whu.edu.cn

<sup>†</sup> These authors have contributed  
equally to this work.

**Received:** 02 May 2017

**Accepted:** 28 September 2017

**Published:** 23 October 2017

### Citation:

Wang A, Wang J, Liu Y and Zhou Y  
(2017) Mechanisms of Long  
Non-Coding RNAs in the Assembly  
and Plasticity of Neural Circuitry.  
*Front. Neural Circuits* 11:76.  
doi: 10.3389/fncir.2017.00076

**Keywords:** long non-coding RNA, neural circuit, cell fates, synaptogenesis, synaptic plasticity, CRISPR-Cas9

## INTRODUCTION

The human brain confers on us the abilities of perceptions, thoughts, emotions, actions, and memories. Over many years, the neuroscientists have strived to understand the molecular, cellular, circuit and behavior-level mechanisms that underlie these processes. Around a century ago, Santiago Ramon y Cajal proposed the neuron doctrine postulating that the relationship between nerve cells was not continuous, but contiguous. Cajal, in his *Theory of Dynamic Polarization*, described how information, in the form of electrical signals, travels within individual neurons, from their dendrites to their cell bodies and finally to their axons. We now know cognition, emotion, memory, and action are generated by circuits and networks of thousands to millions of interconnected neural cells, mostly neurons. Neural circuits are both anatomical and functional entities, composed of a series of interconnected neurons and glial cells with diverse properties and functions. However, it remains largely elusive how specific types of neural cells assemble the neural circuits in different brain regions and how specific neural circuits perform their signal processing functions during cognitive processes and behaviors. This requires detailed knowledge on the construction of neural circuits at the single-cell resolution and on the spatiotemporal pattern of neuronal activity (Poo et al., 2016). The United States BRAIN (Brain Research through Advancing Innovative Neurotechnologies) initiative was launched in 2013, which was intent to “accelerate the development and application of innovative technologies to construct a dynamic picture of brain function that integrates neuronal and circuit activity over time and space.” To achieve this goal “requires an integrated view of its component cell types, their local and long range synaptic connections, their electrical and chemical activity over time, and the functional consequences of that activity at the levels of circuits, the brain, and behavior” (NIH, 2014).

The process of neuronal specification, migration, and circuit formation is enormously complex in time and space during development, with multiple levels of regulation. Deciphering this

process requires both a high-throughput neuronal subclass identification and an integrative approach that considers dynamic, multilayered transcriptional regulation (Molyneaux et al., 2015). However, current regulatory models are limited to a number of regulators, mostly transcription factors, which account for a limited subset of key nodes within a broader regulatory network that is believed to be far more complex (Greig et al., 2013). The transient expression, flexible structures, and dynamic localization of RNA molecules enable fine-tuning genome arrangement, scaffolding and transcription functions, thus precisely regulating gene expression in a time and site-specific manner. Recent work indeed points to the critical role of long non-coding RNAs (lncRNAs) in transcriptional and post-transcriptional regulation of gene expression, the formation of complexes with epigenetic regulatory machinery, and chromosomal architecture organization (Rinn and Chang, 2012; Quinn and Chang, 2016). Therefore, lncRNAs participate in numerous physiological and pathological processes including maintenance of pluripotency, lineage specification, organogenesis, tumorigenesis, and metabolism (Wang et al., 2011; Ramos et al., 2013; Li and Chang, 2014; Yang et al., 2014; Wu et al., 2015). Although recent reviews have covered many aspects of lncRNAs in the assembly and plasticity of neural circuits, this field is fast-growing with new evidence reinforcing the notion that lncRNAs are pivotal in cell fate determination and in modulating neural activity (Ng et al., 2013a; Aprea and Calegari, 2015; Briggs et al., 2015). In this review, we highlighted roles and mechanisms of lncRNAs in assembly, maintenance, plasticity and abnormality of neural circuitry. Given the *cis*- and *trans*- regulatory mechanisms by lncRNAs and/or their embedding DNA elements, along with far more uncharacterized lncRNAs than protein-coding genes, strategies and technologies in studying lncRNAs were also discussed. Finally, we speculate that findings in lncRNA studies would deepen our understanding on neural circuitry composition and functional dynamics in physiology and disease conditions.

## **lncRNAs ARE ABUNDANT IN BRAIN AND DISPLAY SPATIOTEMPORAL SPECIFICITY**

Current data from the ENCODE consortium suggest that up to 75% of the human genome may be transcribed (Djebali et al., 2012), but only about 1–2% of the human genome seems to encode protein (Birney et al., 2007; Church et al., 2009). Thus, most transcripts are non-protein-coding RNA (ncRNA) transcripts (Chodroff et al., 2010). lncRNAs are usually defined as non-protein coding transcripts longer than 200 nucleotides (nt) to exclude small regulatory RNAs such as short interfering RNAs (siRNAs), microRNAs (miRNAs), small nucleolar RNAs (snoRNAs), Piwi-interacting RNAs (piRNAs), and other short RNAs. Occasionally, functional short peptides can be derived from lncRNAs (Matsumoto et al., 2017). Until now, the NONCODE database have annotated 101,700 lncRNA genes in the human genome (Zhao et al., 2016). Remarkably, 40% of lncRNAs are expressed predominantly in the brain (Derrien et al., 2012). While many lncRNA genes overlap protein-coding genes

in sense or antisense directions, others resides in genomic regions previously termed “gene deserts,” between protein-coding genes (intergenic) (Carninci et al., 2005; Cheng et al., 2005; Katayama et al., 2005; Kapranov et al., 2007a,b; Qureshi et al., 2010). In a recent study, using FANTOM5 (Functional Annotation of Mammalian cDNA) cap analysis of gene expression (CAGE) data, Hon et al. generated an atlas of nearly 30,000 human lncRNA genes with typical 5' ends and expression profiles across 1,829 human primary cell types and tissues. Interestingly, most intergenic lncRNAs (lincRNAs) originate from enhancers rather than from promoters. Incorporating genetic and expression data implicates around 20,000 potentially functional lncRNAs in multiple diseases and in transcriptional regulation (Hon et al., 2017).

Some lncRNAs have distinct molecular biogenesis features compared to protein-coding transcripts. Using native elongating transcript sequencing (mNET-seq), Schlackow et al. found human lincRNAs and protein-coding pre-mRNAs are transcribed by different Pol II phospho-CTD (the C-terminal domain) isoforms. lincRNAs are rarely spliced, mainly non-polyadenylated, and are stabilized in the nucleoplasm (Schlackow et al., 2017). lncRNA conservation includes four dimensions: the sequence, structure, function, and expression from syntenic genome loci (Diederichs, 2014). In fact, lncRNA exons are significantly more conserved than neutrally evolving sequences, albeit at lower levels than protein-coding genes (Derrien et al., 2012). Interestingly, lncRNA promoters are more conserved than their exons, and nearly as conserved as promoters of protein-coding gene (Guttman et al., 2009).

lncRNAs are generally expressed at lower levels than protein-coding transcripts, and exhibit more cell- and tissue-specific expression patterns. Moreover, lncRNA expression is vigorously regulated during neural development (Mercer et al., 2010; Belgard et al., 2011; Aprea et al., 2013; Molyneaux et al., 2015) and upon neural activity (Lipovich et al., 2012; Barry et al., 2014), suggesting their specific functional roles. Analyzing *in situ* hybridization data from ABA (the Allen Brain Atlas), numerous lncRNAs are found to be expressed in the adult mouse brain and most of them were present in specific neuroanatomical structures or cell types such as particular cortical regions or the hippocampus (Mercer et al., 2008). Similarly to the expression of fate-determining protein-coding genes, these region-specific and dynamic expression patterns of lncRNAs could be orchestrated by *cis*-regulatory elements (enhancers), chromatin status, and cell-type-specific or activity-dependent transcription factors (TFs) (Ramos et al., 2013).

## **lncRNAs CONTROL NEURAL CELL FATES VIA CIS- AND TRANS- REGULATORY MECHANISMS**

Production of neurons and glial cells during neural development is an intricate but highly stereotyped process that necessitates accurate spatiotemporal controlling of neural stem/progenitor cells (NSPCs) self-renewal and differentiation (Zhou, 2012). The mature mammalian neocortex, for example, is a multi-layered

structure and the layers-specific projection neurons are generated sequentially by cortical neural stem/precursor cells (NSPCs) lying in the ventricular zone/subventricular zone (VZ/SVZ) over developmental time. Intriguingly, cortical NPCs acquire restrictions in fate potential progressively over developmental time, which are largely cell-intrinsic (McConnell, 1995; Desai and McConnell, 2000; Shen et al., 2006; Gaspard et al., 2008). In contrast, cortical interneurons, which usually make local and inhibitory connections, are produced mostly from precursors in the ventral telencephalon and cortical hem and undergo tangential migration into the cortex (Anderson et al., 2002; Wonders and Anderson, 2006). Notably, most neurons are not directly derived from bipolar radial glial neural precursor cells (RGCs) but are from multipolar intermediate progenitor cells, which are direct progenies of RGCs and may undergo a few rounds of replication prior to differentiation. This so-called indirect cortical neurogenesis is more prevalent in primates than in rodents (Qian et al., 2000; Franco and Muller, 2013; Greig et al., 2013). Cortical neurogenesis is followed by gliogenesis, which occurs mostly after birth in mice.

### Cis-Acting lncRNAs

Recent studies unveil that lncRNAs have essential regulatory roles balancing NSPC self-renewal and differentiation (Figures 1A–I; Table 1). Initially, researchers were interested in lncRNAs transcribed from loci overlapping with or adjacent to transcription factor (TF) genes known to be essential for NSPC fate choices. The rationale is based on the facts that lncRNAs can regulate transcription locally (*in cis*) (Wang et al., 2008; Guil and Esteller, 2012; Dimitrova et al., 2014; Engreitz et al., 2016; Luo et al., 2016). One of such examples, *utNgn1*, is a noncoding RNA transcribed from an enhancer region (3.8–7.8 kb upstream of the transcription start site) of mouse *Neurogenin1* (*Neurog1*), a key TF that promotes neuronal fate specification (Figure 1G). Expression pattern of *utNgn1* is highly correlated with that of *Neurog1* mRNA. Moreover, *utNgn1* is required for the expression of *Neurog1* during neuronal differentiation of cortical NPCs (Onoguchi et al., 2012).


Interestingly, many such lncRNAs can simultaneously target distal genes by associating with *cis*-elements in the genome, TFs and epigenetic modifiers. lncRNA *Evf2* (also known as *Dlx6os1* or *Dlx6as*) is transcribed from the ultra-conserved intergenic region between the *Dlx5* and *Dlx6* genes, encoding two members of the DLX homeodomain-containing protein family essential for interneuron development (Feng et al., 2006). Disruption of mouse *Evf2* transcription results in decreased numbers of GABAergic interneurons in early postnatal hippocampus and dentate gyrus, and reduced synaptic inhibition in CA1 layer of the adult hippocampus (Bond et al., 2009). Mechanistically, *Evf2* controls the expression of interneuron lineage genes, including *Gad1*, *Dlx5*, and *Dlx6*, by both *cis*- and *trans*-acting mechanisms. *Evf2* guides methyl-CpG-binding protein MECP2 and the transcription factor DLX to regulatory elements in the *Dlx5/6* intergenic region, thus regulating *Dlx5/6* expression by modulating the opposing interactions between DLX and MECP2, and by modulating *Dlx5/6* ultra-conserved enhancer site-specific methylation (Figure 1H; Berghoff et al., 2013). *Paupar* (*Pax6*

*Upstream Antisense RNA*) is a single-exon lncRNA transcribed from 8.5 kb upstream of the *Pax6* gene in mouse, which is evolutionarily conserved in term of genomic organization and sequence. Knockdown of *Paupar* induces neural differentiation of Neuro-2a neuroblastoma cells. *Paupar* acts locally to regulate *Pax6* expression in a transcript-dependent manner. Interestingly, *Paupar* also functions *in trans* to control distal neural gene expression on a large scale. *Paupar* transcript physically associates with PAX6 protein and localizes at promoters of *Sox2*, *Nanog*, and *Hes1* to regulate cell-cycle progression and differentiation of Neuro-2a cells (Figure 1B; Vance et al., 2014).

*Linc-Brn1b* is transcribed from a 6.8 kb genomic locus about 10 kb downstream of the *Brn1* (*Pou3f3*), a well-studied TF gene involved in cortical development. *Linc-Brn1b*'s expression is restricted in germinal zones (VZ/SVZ) of the early developing brain but becomes prominent in the cortical plate neurons in late cortical neurogenesis, indicating its role in regulating neuronal differentiation. Deletion of the *linc-Brn1b* locus leads to significant decrease in *Brn1* expression. Moreover, *linc-Brn1b*-null cerebral cortices displayed an expression signature indicative of decreased cellular proliferation and increased neuronal differentiation. Consistently, the embryonic cortices of *linc-Brn1b*-null mice showed decreased numbers of intermediate progenitors and upper layer (II-IV) projection neurons, accompanied by an expansion of deep layer neurons. Furthermore, *linc-Brn1b*-null mice exhibit reduced barrel size and number in the somatosensory cortex. All these suggest *linc-Brn1b* specifies cortical NPC fate and regulate area patterning and layer formation. However, it's elusive the cortical defects found in *linc-Brn1b*-null mice are due to depletion of *linc-Brn1b* transcript or the deletion of its embedding *cis* element (Sauvageau et al., 2013). *Dali*, another lncRNA transcribed downstream of *Brn1* (*Pou3f3*) locus, also controls neuronal differentiation partly via its positive regulation of *Brn1* expression. Depletion of *Dali* in Neuro-2a neuroblastoma cell inhibits its neuronal differentiation induced by retinoic acid. Intriguingly, similar to *Paupar*, *Dali* also interacts directly with the protein product of its neighboring genes, BRN1, to regulate expression of a large set of neural differentiation genes *in trans*. Moreover, *Dali* associates with DNMT1, a DNA methyltransferase, to regulate DNA methylation status of promoter CpG islands (Figure 1E; Chalei et al., 2014).

### Trans-Acting lncRNAs

Recently, several lncRNAs are reported to mainly function *in trans* by directing TFs or chromatin modifiers to important loci, thus regulating expression of distal genes that are essential for cell fate specifications. *Pnky* is a conserved nuclear lncRNA predominantly expressed in NSPCs of both the embryonic and postnatal brain. Depletion of *Pnky* promotes neuronal lineage specification and augments the population of transit-amplifying cell, leading to increased neurogenesis. RNA pull-down assay identified PTBP1, an RNA-splicing factor that potentiates neural development and neuronal reprogramming (Keppetipola et al., 2012; Xue et al., 2013), as the binding partner of *Pnky*. In NSPCs, *Pnky* and PTBP1 promote neurogenesis by regulating



**FIGURE 1 |** lncRNAs regulate different aspects of neural circuitry assembly and function (left) via *cis*- and *trans*- mechanisms (right). (**A–G**) lncRNAs can control neurogenesis of neural stem/progenitor cells via regulating expressions of proximal genes (**B, E, G**) and/or distal genes by associating with fate-determining transcription factors (**A–C, E**), acting as competing endogenous RNAs (**D**) and regulating alternative splicing (**F**). (**H**) *Evt2* controls expression of *Dlx5/Dlx6* *in cis* and *Gad1* *in trans* to regulate GABAergic interneuron specification. (**I**) *lncOL1* interacts with SUZ12 to repress a gene program that maintains oligodendrocyte progenitor state, thereby promoting OL myelination. (**J–L**) lncRNAs regulate neurite outgrowth and synaptogenesis via *cis*-regulation (**J**), alternative splicing (**K**), and translational control (**L**).

**TABLE 1** | Examples of lncRNA activities in neural circuitry assembly and function.

Process	lncRNA	Cell/tissue distribution	Biological function/phenotypes	Molecular mechanisms: Cis and/or trans regulation	Protein partner
Neural stem cell proliferation	<i>Pnky</i>	Mouse and human NSPCs	Regulates neuronal differentiation of embryonic and adult NSPCs	<i>Trans</i> : <i>Pnky</i> and PTBP1 regulate the expression and alternative splicing of an overlapping set of transcripts to promote neurogenesis	PTBP1
	<i>LncND</i>	Radial glia NSPCs	Promotes expansion of radial glial NSPCs	<i>Trans</i> : <i>LncND</i> compete with the 3'UTR of <i>NOTCH1</i> or <i>NOTCH2</i> for the binding of miR-143-3p to promote Notch signal pathway	/
	<i>Paupar</i>	Neuro-2a neuroblastoma cells*	Knockdown of <i>Paupar</i> induces neural differentiation of Neuro-2a neuroblastoma cells	<i>Cis</i> : <i>Paupar</i> regulates <i>Pax6</i> expression locally. <i>Trans</i> : <i>Paupar</i> also associates with PAX6 protein and localizes at promoters of <i>SOX2</i> , <i>NANOG</i> , and <i>HES1</i>	PAX6
	<i>utNgn1</i>	Mouse NSPCs	Promotes neuronal differentiation	<i>Cis</i> : <i>utNgn1</i> is transcribed from an enhancer region of <i>Neurog1</i> and positively regulates its expression	/
Neuronal differentiation	<i>RMST</i>	Midbrain dopaminergic neuronal precursor cells	Promotes neuronal differentiation	<i>Trans</i> : <i>RMST</i> interacts with <i>SOX2</i> to regulate neurogenic genes including <i>ASCL1</i> and <i>DLX1</i>	<i>SOX2</i> , hnRNPA2/B1
	<i>Tuna</i>	Mouse ESCs; mouse and zebrafish CNS	Regulates pluripotency and neural differentiation of ESCs	<i>Trans</i> : <i>TUNA</i> formed a complex with three pluripotency related RNA-binding proteins, PTBP1, hnRNP-K, and NCL	PTBP1, HNRNP-K, NCL
	<i>Linc-Brn1b</i>	Developing mouse neocortex; SVZ and upper cortical layers	Specifies cortical NSPC fate and regulate area patterning and layer formation of mouse neocortex	<i>Cis</i> : Deletion of the <i>linc-Brn1b</i> locus leads to significant decrease in <i>Brn1</i> expression	/
	<i>NBAT-1</i>	SH-SY5Y cells*	Loss of <i>NBAT-1</i> increases cellular proliferation and invasion	<i>Trans</i> : <i>NBAT-1</i> interacts with <i>EZH2</i> to suppress expression of <i>SOX9</i> , <i>OSMR</i> , and <i>VCAN</i>	<i>EZH2</i>
	<i>Miat/Gomafu</i>	NSPCs, neurons	Controls retinal development. Dysregulated in schizophrenia	<i>Trans</i> : <i>Gomafu/Miat</i> regulates splicing of neuronal genes, including <i>DISC1</i> , <i>ERBB4</i> , and <i>WNT7B</i> , probably via association with splicing factors SF1, SRSF1, and QKI	QKI, SRSF1, SF1
	<i>Dali</i>	Neuro-2a neuroblastoma cells*	Depletion of <i>Dali</i> in Neuro-2a neuroblastoma cell inhibits its neuronal differentiation induced by retinoic acid	<i>Cis</i> : <i>Dali</i> maintains <i>Brn1</i> expression. <i>Trans</i> : <i>Dali</i> interacts with the DNMT1 to regulates DNA methylation status of CpG island-associated promoters; interacts with <i>BRN1</i> to regulate expression of neural differentiation genes	<i>BRN1</i> , <i>DNMT1</i>
	<i>Kcna2-as</i>	DRG sensory neurons	Upregulation of <i>Kcna2-as</i> decreased <i>Kcna2</i> mRNA, reduced total voltage-gated potassium current, increased excitability in DRG neurons, ultimately leading to neuropathic pain symptoms	<i>Cis</i> : <i>Kcna2-as</i> silences <i>Kcna2</i> expression	/
Neurites outgrowth and synaptogenesis	<i>BC1/BC200</i>	Dendrites and somata of neurons	Regulates synaptic excitability	<i>Trans</i> : <i>BC1</i> controls protein translation in synaptodendritic microdomains	<i>FMRP</i> , <i>elf4A</i> , <i>PABP</i>
	<i>BDNF-AS</i>	DRG sensory neurons	Depletion of <i>BDNF-AS</i> promotes neuronal outgrowth and adult neurogenesis	<i>Cis</i> : <i>BDNF-AS</i> negatively regulates <i>BDNF</i> expression by recruiting <i>EZH2</i> , a <i>PRC2</i> core component	<i>EZH2</i>
	<i>Malat1</i>	Differentiated projection neurons	Promotes dendrite maturation and synaptogenesis in cultured hippocampal neurons	<i>Trans</i> : <i>Malat1</i> associates with SR family splicing factors to controls expression of synaptic molecules including <i>Nlgn1</i> and <i>SynCAM1</i>	<i>SF2/ASF</i> , <i>SC35</i>
	<i>Evl2</i>	Postmitotic interneurons	Ensures proper formation of GABA-dependent neuronal circuitry	<i>Cis</i> and <i>trans</i> : <i>Evl2</i> associates with <i>DLX1/2</i> and <i>MECP2</i> at the regulatory elements in the <i>Dlx5/6</i> intergenic region to control <i>Dlx5</i> , <i>Dlx6</i> and <i>Gad1</i> expression	<i>DLX1</i> , <i>DLX2</i> , <i>BRG1</i> , <i>MECP2</i>
Glial cells	<i>Six3OS</i>	Retinal progenitor cells	Regulates retinal cell specification, neuron and oligodendrocyte differentiation	<i>Trans</i> : <i>Six3OS</i> binds to <i>EZH2</i> and <i>EYA</i> family members to regulate expression of <i>SIX3</i> target genes	<i>EZH2</i> , <i>EYA1</i> , <i>EYA3</i> , <i>EYA4</i>
	<i>LncOL1</i>	Mature oligodendrocytes (OLs)	Promotes oligodendrocyte myelination	<i>Trans</i> : <i>LncOL1</i> interacts with <i>SUZ12</i> to suppress a gene program that maintains OL progenitor state, thereby promoting OL myelination.	<i>SUZ12</i>

\*Research subject; ESCs, embryonic stem cells; NSPCs, neural stem/progenitor cells; V-SVZ, ventricular-subventricular zone; DRG, dorsal root ganglion.

the expression and alternative splicing of an overlapping set of transcripts (Ramos et al., 2015).

Rhabdomyosarcoma 2-associated transcript (*RMST*), a lncRNA with prominent expression in midbrain dopaminergic neuronal precursors, is required for neuronal differentiation of human ESCs (Uhde et al., 2010; Ng et al., 2012). *RMST* is negatively regulated by the transcription factor REST and upregulated during neuronal differentiation. *RMST* interacts with SOX2 to co-activate a large pool of neurogenic genes such as *ASCL1*, *NEUGOG2*, *HEY2* and *DLX1* to promote neuronal differentiation (Figure 1C; Ng et al., 2013b). *TUNA* (Tcl1 Upstream Neuron-Associated lincRNA, or megamind), a highly conserved lincRNA that show specific expression in developing CNS of zebrafish and mice, was required for pluripotency maintenance and proper neural differentiation of mouse embryonic stem cells. *TUNA* formed a complex with three pluripotency related RNA-binding proteins (RBPs), hnRNP-K, PTBP1, and NCL, at the promoters of *Nanog*, *Sox2*, and *Fgf4* to regulate gene expression. Furthermore, disruption of *TUNA* expression in zebrafish caused impaired locomotor function (Figure 1A; Lin et al., 2014).

## Competitive Endogenous RNAs

Some lncRNAs contains multiple complementary sites for microRNAs (miRNAs). These competitive endogenous RNAs (ceRNAs) act as molecular sponges for miRNAs through their miRNA binding sites (also known as miRNA response elements, MRE), thereby de-repressing all target genes of the respective miRNA family (Cesana et al., 2011; Salmena et al., 2011; Tay et al., 2014). Human *lncND* (neurodevelopment) contains 16 MREs for miR-143-3p and is highly expressed in the progenitor zone (VZ/SVZ) in developing human neocortex where it co-localizes with NSPC markers such as *PAX6*. *lncND* positively regulates the expression of *NOTCH1* and *NOTCH2*, two receptors genes essential for NSPC self-renewal, by competing the binding of miR-143-3p to *NOTCH1/2*'s 3' untranslated region (UTR). Depletion of *lncND* induced neuronal differentiation of neuroblastoma cells, an effect reminiscent of miR-143-3p overexpression (Figure 1D; Rani et al., 2016). Circular RNAs or transcripts of pseudogenes might also behave as ceRNAs (see next section for more details).

## lncRNAs in Myelination

lncRNAs also have roles in glial cell fate determination. Oligodendrocytes provide support and insulation to axons in the central nervous system of some vertebrates by creating the myelin sheath. Diseases that result in injury to the oligodendrocytes include demyelinating diseases such as multiple sclerosis and various leukodystrophies. Many lncRNAs are dynamically expressed during oligodendrocyte (OL) lineage specification, neuronal-glial fate switches, and OL lineage determination such as myelination (Mercer et al., 2010; Dong et al., 2015; He et al., 2017). In an integrative analysis using transcriptomic and epigenetic data, Dong et al. characterized lncRNAs that are differentially expressed in the process of oligodendrocyte precursor cell (OPC) differentiation from mouse neural stem cells (NSCs) and that are potential regulators of

oligodendrogenesis (Dong et al., 2015). Co-expression network analyses associates distinct oligodendrocyte-expressing lncRNA clusters with protein-coding genes and predict lncRNA functions in oligodendrocyte myelination. Genetic ablation of *lncOL1*, a chromatin-associate lncRNA, causes defects in CNS myelination and re-myelination following injury. *lncOL1* interacts with SUZ12, the core component of polycomb repressive complex 2 (PRC2), to promote oligodendrocyte maturation partly via transcriptional silencing of gene program that maintains the OL progenitor state (Figure 1I; He et al., 2017).

Notably, not all aforementioned lncRNAs have been exhaustively examined *in vivo*. Moreover, given lower sequence conservation of lncRNAs than protein-coding genes across species, functional validation using various model organisms, human neural cells and neural organoids are required to assign functions to lncRNAs (Table 1).

## lncRNAs' ROLES IN NEURITE OUTGROWTH, SYNAPTOGENESIS, AND SYNAPTIC PLASTICITY

After neural cells were generated in appropriate numbers, at right times, and in the correct places, they establish functional connections required for normal brain function. To form connections, neurons extend long processes, axons and dendrites, which allow synapse formation (synaptogenesis). Neurite outgrowth and synaptogenesis involve complex regulations on gene expression and signal transduction. Neurons can alter their synaptic connections and the relative strength of individual connections in response to increases or decreases in their activity. This so-called neural plasticity accounts for memory, learning, and cognition, as well as the brain's capability to recover from damage. Compared to studies on lncRNAs' roles in fate specifications of neural cells, little is known regarding lncRNAs' functions in modulating neurite growth, synaptogenesis and neural plasticity, which is partly due to their dynamic features, scarcity of research material and hurdles in functional validation (Puthanveetil et al., 2013).

Nonetheless, emerging evidence indicates both nuclear and synaptic lncRNAs are actively involved in these processes. Comparative sequence analysis of genomic regions covering 150 presynaptic genes discovered highly conserved elements in non-protein coding regions in eight vertebrate species. Many of these "non-exon-associated and non-protein-coding" elements can transcribe and were predicted to form a highly stable stem-loop RNA structure, whereas some conserved noncoding elements correlate with specific gene expression profiles (Hadley et al., 2006). This early work implied that non-coding transcripts are prevalent in genomic regions of presynaptic genes and may have regulatory roles in transcriptional regulation.

## Transcriptional Regulation of Neurotrophins and Synaptic Molecules

It has been shown neuronal activity may reshape the epigenetic landscape, thereby dynamically changing transcriptome and neuronal properties (Su et al., 2017). A genome-wide analysis

of hyperactive regions of human neocortex removed to treat intractable seizures identified numerous differentially expressed lncRNAs, with a fraction having expression profiles that matched activity-dependent coding genes. Among them, eight lncRNAs were overlapping with or adjacent to differentially expressed protein-coding genes, including reciprocal patterns between *BDNF* (Brain-derived neurotrophic factor) and *BDNF-AS* transcription (Lipovich et al., 2012). BDNF, a member of the neurotrophin family of growth factors, promotes differentiation of new neurons and growth of synapses during development and in adulthood. BDNF signaling is important for long-term memory and its dysfunction have implications in a number of neurodegenerative disorders including Alzheimer's disease (AD). LncRNA *BDNF-AS* negatively regulates BDNF expression by recruiting EZH2, a PRC2 core component, to the *BDNF* locus (**Figure 1J**; Lipovich et al., 2012). Knockdown of *BDNF-AS* induces BDNF expression in hippocampal neurosphere cultures, which leads to increased neuronal survival and neurite outgrowth (Modarresi et al., 2012). Moreover, *BDNF-AS* transcripts is markedly upregulated in Alzheimer's disease (Shi et al., 2017). In a recent study where differentiated SH-SY5Y neuroblastoma cells were treated with BDNF and subjected to microarray studies, several lncRNAs including *MALAT1/NEAT2* and *MIAT/GOMAFU* were found to differentially expressed. Interestingly, a few putative microRNA-lncRNA interactions were predicted and seven of the microRNAs are associated with psychiatric and neurodegenerative diseases (Aliperti and Donizetti, 2016).

*Malat1* (Metastasis-associated lung adenocarcinoma transcript 1) is a neuron-enriched nuclear-retained lncRNA associated with nuclear speckle, a structure implicated in pre-mRNA splicing and RNA transport (**Figure 1K**) (Bernard et al., 2010). Although *Malat1* knockout does not alter gross and histologic morphology of adult mouse brain (Eissmann et al., 2012), *Malat1* depletion in cultured hippocampal neurons resulted in a significant reduction in synaptic density. Conversely, *Malat1* overexpression increased presynaptic bouton density on dendrites. Accordingly, the levels for transcripts of Neurologin1 (NLGN1) and SynCAM1 post-synaptic proteins were significantly lower in *Malat1*-depleted hippocampal neurons (Bernard et al., 2010). Consistently, *Malat1* maintains survival and neurite outgrowth of Neuro-2a neuroblastoma cells probably via the ERK/MAPK signaling pathway (Chen et al., 2016). The discrepancy between these *in vivo* and *in vitro* results necessitates detail analyses of *Malat1* knockout animals, including structural and functional interrogation of neurites and synapses as well as behavioral studies.

Alterations in expression of genes encoding signaling proteins or ion-channel components can drastically change neuronal excitability. Therefore, regulations of these components can modulate neuronal plasticity. Scaffold protein CASK (Ca<sup>2+</sup>/calmodulin-dependent protein kinase) regulates synapse formation and plasticity during neural development (Martin and Ollo, 1996; Chen and Featherstone, 2011; Slawson et al., 2011). The expression of *Drosophila* CASK is positively regulated by its downstream overlapping lncRNA, *CRG* (CASK regulatory gene), which is neural-specific and is induced during embryonic

development. Loss of *CRG* leads to decreased locomotor activity and a defective climbing capability in fly—phenotypes reminiscent of *CASK* mutants and could be rescued by *CASK* overexpression (Li et al., 2012). *KCNA2* encodes a core potassium channel subunit and can be negatively regulated by its antisense RNA *Kcna2-as*. *Kcna2-as* transcript and *KCNA2* proteins are largely reciprocally expressed in DRG neurons. Upon peripheral nerve injuries, *Kcna2-as* expression was triggered by activation of the myeloid zinc finger protein 1 (MZF1) transcription factor. Upregulation of *Kcna2-as* decreased *Kcna2* mRNA, tempered total voltage-gated potassium current and elevated excitability in DRG neurons, ultimately leading to neuropathic pain symptoms. Moreover, blocking the induction of *Kcna2-as* attenuated neuropathic pain following peripheral nerve injury (Zhao et al., 2013).

## Post-transcriptional Regulation at Synapses

As dendrites and axons usually extend far away from the cell body, local protein translation appears particularly important for maintaining dendritic and axonal function (Job and Eberwine, 2001). Noncoding RNA *BC1/BC200* (*BC200* is the primate counterpart of rodent *BC1* RNA) is located in the dendrites and somata of a subset of neurons in the central and peripheral nervous system (Tiedge et al., 1991; Muslimov et al., 1997). The expression of *BC1* RNA in soma and dendrites of hippocampal neurons is induced during synapse formation, and is dependent upon neuronal activity (Muslimov et al., 1998). Accordingly, *BC1* controls protein translation in synaptodendritic microdomains. *BC1* RNA interacts directly with initiation factor eIF4A and poly(A)-binding protein (PABP), preventing association of the 48S pre-initiation complex with mRNA, thus inhibiting the formation of the 48S ribosomal translation initiation complex (Wang et al., 2002). Another report suggested *BC1* RNA binds to the fragile X syndrome protein FMRP to regulate the translation of specific FMRP target mRNAs at synapses (**Figure 1L**; Zalfa et al., 2003). The brains from *BC1*-null mice have no grossly morphological defects, as were the localization of CaMKII $\alpha$  and MAP2 dendritic mRNAs (Skryabin et al., 2003). But detailed behavioral studies found *BC1*-deficient mice have defects in exploratory behavior and higher levels of anxiety and increased neuronal excitability, probably due to hyperactive mGluRs (group I metabotropic glutamate receptor)-triggered translation in synapses (Lewejohann et al., 2004; Zhong et al., 2009). These studies implied *BC1/BC200* would contribute to the maintenance of synaptic plasticity. *BC200* levels were greatly reduced in aging brain cortices, but it was significantly up-regulated in AD brains. Relative *BC200* levels correlate with the progression of AD, and its mislocalization (clustered perikaryal localization but not somatodendritic) was observed in advanced AD brains (Mus et al., 2007).

Circular RNAs (circRNAs) have been identified in various species and categorized as a novel type of noncoding RNA (Salzman et al., 2012; Guo et al., 2014). Most circular RNAs arise from “back-splicing,” where a 5' splice donor joins an upstream 3' splice acceptor (Jeck and Sharpless, 2014; Xing et al.,

2016). Westholm et al. annotated more than 2,500 *Drosophila* circRNAs, many of which contain conserved canonical miRNA seed matches, indicating their impacts on posttranscriptional regulatory networks through acting as molecular sponges for miRNAs. Notably, circRNAs dominantly reside in the *Drosophila* nervous system and their levels increase with age (Westholm et al., 2014). Similarly, thousands of conserved circular RNAs (circRNAs) were found to be highly expressed in mammalian brain. The expression levels of many circRNAs are elevated in neurogenesis and they are more abundantly expressed in synaptic processes than their linear isoforms (Rybak-Wolf et al., 2015; You et al., 2015). Circular RNAs could behave as competing endogenous RNAs (ceRNAs): *ciRS-7* (circular RNA sponge for miR-7), a neuron-enriched circRNA, sequesters miR-7 and prevents miR-7's interactions with target mRNAs (Hansen et al., 2013). These findings point to prospective roles of circRNAs in the brain, especially in synaptogenesis and neural plasticity.

Current knowledge regarding lncRNAs' role in synaptogenesis and plasticity is relatively scarce. Studies found lncRNAs seem to regulate circadian or mating behavior in insects and worms (Soshnev et al., 2011; Gummalla et al., 2014), such evidence hasn't been found in mammals yet. Future genome-wide lncRNA knockout studies in mice would unveil the extent and mechanisms how lncRNAs are involved in circuitry dynamics.

## lncRNAs IN NEUROLOGICAL DISORDERS

Since lncRNAs regulate neural development and function, it's not surprising that mutation or dysregulation of lncRNAs has implications in pathogenesis of mental illness and neurodegenerative diseases such as autism spectrum disorder (ASD), depression, schizophrenia, amyotrophic lateral sclerosis (ALS), Alzheimer's disease and neuropathic pain. Some neural-specific lncRNAs have been emerged as potential therapeutic targets.

lncRNAs related to cognitive functions or synaptic plasticity or other psychiatry diseases, including *GOMAFU*, *BDNF-AS*, and *DISC2*, may potentially contribute to major depressive disorder (MDD) (Huang et al., 2017). In a microarray-based study, about two thousand lncRNAs were found to be differentially expressed in peripheral blood samples from major depression disorder (MDD) patients (Liu Z. et al., 2014), but their diagnostic and therapeutic implications remain to be elucidated. A recent genome-wide study characterized thousands of lncRNAs to be differentially expressed in ASD peripheral leukocytes. Gene ontology (GO) analysis of these lncRNA gene loci predicted neurological regulations of the synaptic vesicle cycling, long-term depression and potentiation to be mainly involved, including *SHANK2-AS* and *BDNF-AS* (Wang et al., 2015). Similarly, a large-scale study applied RNA sequencing (RNA-seq) of 251 post-mortem samples of frontal and temporal cortex and cerebellum from 48 individuals with ASD and 49 control human subjects, and identified 60 differentially expressed lncRNAs (Parikshak et al., 2016). Twenty of these lncRNAs were previously shown to

interact with microRNA (miRNA)-protein complexes, and 9 with the fragile X mental retardation protein (FMRP), whose mRNA targets are enriched in ASD risk genes (Parikshak et al., 2013; Iossifov et al., 2014). These data show that dysregulation of lncRNAs is an integral component of the transcriptomic signature of ASD (Parikshak et al., 2016). lncRNA *GOMAFU/MIAT* is downregulated in post-mortem cortical gray matter from schizophrenia (SZ) patients. *GOMAFU* associates with splicing factors SRSF1 (serine/arginine-rich splicing factor 1) and QKI and dysregulation of *GOMAFU* results in aberrant splicing of *DISC1* and *ERRB4*, two SZ-associated genes (Barry et al., 2014). Another study discovered that *GOMAFU* mediates mouse anxiety-like behavior probably via association with BMI1, a key member of the polycomb repressive complex 1 (PRC1), to repress the expression of *beta crystallin (Crybb1)*, one of the SZ-related genes (Spadaro et al., 2015).

Amyloid precursor protein (APP) is sequentially cleaved by beta-site APP-cleaving enzyme 1 (BACE1),  $\beta$ -secretase, and  $\gamma$ -secretase to generate the toxic A $\beta$ <sub>42</sub> peptide. Defective A $\beta$ <sub>42</sub> clearance and elevated BACE1 expression contribute to A $\beta$ <sub>42</sub> accumulation and AD progression. *BACE1-AS*, the antisense transcript of *BACE1*, can bind to and stabilize *BACE1* transcripts, thus increasing BACE1 protein levels. Interestingly, *BACE1-AS* is induced by A $\beta$ <sub>42</sub> peptide, leading to increased BACE1 mRNA stability and amyloid accumulation via a positive feedback loop. Consistently, expression of BACE1 and *BACE1-AS* is elevated in brains of AD, and knockdown of *BACE1-AS* reduced BACE1 levels *in vivo* (Faghihi et al., 2008; Liu T. et al., 2014). The neuromuscular disorder spinal muscular atrophy (SMA) is caused by insufficient expression of SMN (survival motor neuron) protein, and the primary goal of SMA therapeutics is to increase SMN levels (Lefebvre et al., 1997). lncRNA *SMN-AS1* is enriched in neurons and suppresses SMN expression by recruiting the polycomb repressive complex-2 (PRC2) to *SMN* promoter. Targeting *SMN-AS1* with antisense oligonucleotides (ASOs) increases SMN expression both in cultured neurons and in mice, indicating *SMN-AS1* has potential to be a novel therapeutic target for treating SMA (D'ydewalle et al., 2017).

Large-scale RNA dysregulations are essential molecular hallmarks in neurodegenerative diseases including ALS and FTL (Frontotemporal lobe dementia; Polymenidou et al., 2012). This is mostly due to the presence of aberrant protein states (proteinopathy) of two essential RNA/DNA binding proteins TDP-43 and FUS (Fused in sarcoma) in affected neurons, including cytosolic translocation, truncation, phosphorylation, ubiquitination, and aggregates formation (Lagier-Tourenne et al., 2010; Da Cruz and Cleveland, 2011). Although TDP-43 and FUS regulate the processing of an array of RNA molecules including non-coding RNAs, no specific RNA was yet identified as major causal factor of ALS and FTL (Tollervey et al., 2011). The association with TDP-43 or FUS/TLS could allow lncRNAs to carry out their cellular function. On the other hand, the dynamics of association/dissociation of RNAs with TDP-43 or FUS might contribute to TDP-43 and FUS proteinopathies (Yang et al., 2015).



The above findings implicate the correlation between dysregulation of lncRNAs and neurological diseases. However, many were *in vitro* studies with very few mechanistic hints. Thus, we are still far from understanding the extent and mechanisms that lncRNAs are involved in disease brains.

## ADVANCES AND CHALLENGES OF STUDYING lncRNAs *IN VIVO* AND *IN VITRO*

Although a number of lncRNAs have been found to be involved in most, if not all, aspects of neural circuitry assembly and plasticity, many essential questions remain to be answered. First, in contrast to the abundance of lncRNAs characterized, very few of them are essential for embryonic development, cell fate choices or circuitry functions. So, as many may ask, are lncRNAs largely transcriptional noise or non-functional? It's quite possible that most lncRNAs only play subtly regulatory roles, and that certain lncRNAs are not normally required but become essential upon neuronal activation or injury. Second, compared to proteins, most lncRNAs have low sequence conservation across species or among homologs, though evolutionary conservation of RNA secondary structures may exist across species. It's, therefore, hard to identify parallel or redundant pathways and related molecular mechanisms. Third, lncRNAs may exert functions *in cis* (transcripts dependent or independent), and/or *in trans* (chromatin remodeling, histone modification, DNA methylation, transcription and splicing regulation etc.). Moreover, the embedding DNA elements that transcribe lncRNAs may have cis-regulatory roles. These conditions greatly confound experiment design and data interpretation for functional studies of lncRNAs. Thus, it's not surprising loss-of-function studies *in vivo* or *in vitro* using different approaches (e.g., RNAi, antisense oligonucleotides, genomic deletion, polyadenylation insertion, promoter deletion/inversion and CRISPR-Cas9 mediated gene inactivation, etc.) may lead to distinct phenotypes (Bernard et al., 2010; Schorderet and Duboule, 2011; Eissmann et al., 2012; Li et al., 2013). Since each technique has advantages and limitations, researchers are required to exhaustively apply necessary approaches and develop new technologies to elucidate lncRNAs' roles and mechanisms.

Latest breakthroughs in genome engineering technology utilizing CRISPR (clustered regularly interspaced short palindromic repeats) and Cas9 system have dramatically accelerated biomedical researches (Doudna and Charpentier, 2014; Hsu et al., 2014). It has been widely used for generating genetic-modified cells, plants and animals (Cong et al., 2013; Niu et al., 2014; Peng et al., 2014); for disease modeling and genetic corrections (Platt et al., 2014; Cox et al., 2015); as well as for repressing (CRISPRi) or inducing (CRISPRa) gene expressions without altering genomic sequences (Gilbert et al., 2013, 2014; Konermann et al., 2014). In a few proof-of-principle studies, CRISPR-Cas9 has been successfully applied to lncRNA studies in cells and in animals (Ho et al., 2015; Ghosh et al., 2016; Zhu et al., 2016). A genome-scale deletion screening for functional lncRNAs were carried out using a lentiviral paired-guide RNA (gRNA) CRISPR-Cas9 library targeting hundreds of lncRNAs. This screen

identified fifty-one lncRNAs that can enhance or slow down human cancer cell growth. Next, nine lncRNA candidates were validated utilizing CRISPR-Cas9-mediated genomic deletion, CRISPRa or CRISPRi, functional rescue and transcriptome profiling. This study indicates high-throughput genome deletion method mediated by CRISPR-Cas9 enables rapid identification of functional non-coding elements (Zhu et al., 2016). Using the minimal CRISPRi (dCas9) system targeting the *roX* locus in the *Drosophila* cells leads to an efficient and specific knockdown of *roX1* and *roX2* lncRNAs. Moreover, this minimal CRISPRi system inhibits *roX* expression efficiently *in vivo* and leads to loss-of-function phenotype, thus validating the method in a multicellular model organism (Ghosh et al., 2016). To explore if certain RNA molecule can exert transactivation or adapter roles, Schechner et al. developed a targeted localization method called CRISPR-Display utilizing Cas9 to deploy RNA cargos to specific DNA loci. A distinct feature of CRISPR-Display is that it makes possible for multiplexing of different functions at multiple loci in the same cell (Schechner et al., 2015). The ever-growing innovation of CRISPR-Cas9 technique would also enable detection and editing of DNA and RNA with high specificity and sensitivity (Abudayyeh et al., 2016; Gootenberg et al., 2017; Qin et al., 2017). Nonetheless, caution must be taken when applying CRISPR-Cas9 techniques in lncRNA studies: First, deletion of lncRNA genes overlapping protein-coding genes' promoter/enhancer or intron should be avoided; Second, effects of CRISPRa and CRISPRi on promoter/enhancer elements that shared by both lncRNA and protein-coding genes should be taken into account when analyzing phenotypes. Finally, possible off-target effects can be addressed by applying multiple gRNAs and performing rescue experiments.

lncRNAs exert roles by associating with cellular macromolecules including chromatin, DNA, RNA or proteins. Current biochemical means using RNA-centric or protein-centric strategies can identify these molecules and been extensively reviewed elsewhere (Mchugh et al., 2014). Technology breakthroughs in physics, chemistry, molecular biology and neuroscience would allow researcher to carry on high-throughput investigations of lncRNAs at single-cell, circuitry and animal levels.

## CONCLUSIONS AND PERSPECTIVES

One of the primary aim of the BRAIN initiative is to “*identify and provide experimental access to the different brain cell types to determine their roles in health and disease*” (NIH, 2014). However, we are still far from identifying and characterizing the component cells comprising the neural circuits, especially for glial cells. This is partially due to lack of defined biomarkers and dynamic changes of cell properties upon stimulation or depression. Latest advances in cell labeling using genetic and viral means, high throughput purification, e.g., FACS (fluorescence-activated cell sorting) and microfluidic devices, enable researchers to isolate neural cells from embryonic and adult brain under a variety of conditions. Moreover, recent development in profiling transcriptomes and epigenomes from as few as single cells

markedly advanced molecular census of neural cell in embryonic and adult brains (Usoskin et al., 2015; Zeisel et al., 2015; Poulin et al., 2016; Tasic et al., 2016; Telley et al., 2016). Tasic et al. established a cellular classification of primary visual cortex in adult mice based on single-cell transcriptome analysis. A total of 49 transcriptomic cortical cell types, including 19 glutamatergic, 23 GABAergic, and 7 non-neuronal types were identified from around 1,600 cells labeled by *Cre* reporters. Interestingly, many transcriptomic cell types showed discrete anatomical and physiological characteristics, thus validating that the single-cell transcriptomic profiles can reflect specific properties of neural cells (Tasic et al., 2016).

Current annotations of brain lncRNAs are unfinished, partly because of the selection of polyadenylated (polyA) transcripts in most studies and RNA-seq libraries not preserving strand information (Miller et al., 2014; Darmanis et al., 2015). Single-cell RNA sequencing is even harder to detect lncRNAs because there's much less starting material and lncRNAs are generally less abundant than protein-coding transcripts. Nonetheless, latest single-cell transcriptome studies indeed correlate lncRNA profiles with developmental stages and cell identities. Using single cell RNA-seq to analyze roughly 100 individual cells from human embryonic stem cells (hESCs) and human preimplantation embryos, Yan *et al.* identified 2,733 previously uncharacterized lncRNAs, many of which are specifically expressed in developmental stages (Yan et al., 2013). In another single-cell study, more than 600 novel multi-exonic lncRNAs were discovered using micro-dissected adult subventricular zone (SVZ) tissues (Luo et al., 2015). Single-cell sequencing of hundreds of human cortical cells revealed that many lncRNAs are enriched in individual cells, and are cortical layer and cell type-specific (Liu et al., 2016), which coincides with previous studies showing lncRNAs provide more cell identity information during the development of mammalian cortex than protein-coding transcripts (Molyneaux et al., 2015). We speculate that lncRNA profiling at single-cell level, along with high-throughput single molecule fluorescent *in situ* hybridization (smFISH), would greatly advance the census of neurons and glial cell, especially astrocytes, in the context of neural development and plasticity (Femino et al., 1998; Raj et al., 2008). Moreover, transcriptome dissection of cells in cerebral organoid derived from human pluripotent cells or NSPCs would advance our understanding on lncRNAs in brain evolution, development and diseases (Lancaster et al., 2013; Fatehullah et al., 2016).

$N^6$ -Methyladenosine ( $m^6A$ ) is a widespread, reversible chemical modification of polyadenylated mRNAs and lncRNAs in eukaryotes, implicated in many aspects of RNA metabolism

including regulations of stability, transport and translation (Fu et al., 2014). Antibody-based  $N^6$ -methyladenosine ( $m^6A$ ) RNA immunoprecipitation followed by high-throughput sequencing (MeRIP-seq) has been developed to profile the transcriptome-wide distribution of  $m^6A$ , revealing  $m^6A$  is distributed in more than 7,000 mRNA and 250 lncRNA transcripts in human cells (Dominissini et al., 2012). In mouse brain,  $m^6A$  is present in mRNA at low levels throughout embryogenesis but increases dramatically by adulthood, suggesting that upregulation of  $m^6A$  levels accompanies neuronal maturation. Moreover, lncRNAs transcribed by RNA polymerase II are also subject to  $m^6A$  methylation, and long intergenic noncoding RNAs (lincRNAs) had significantly higher  $m^6A$  levels than mRNAs or pseudogenes, but its biological implication is largely unknown and awaits future explorations (Meyer et al., 2012; Molinie et al., 2016). Interesting, a long non-coding RNA antisense to *FOXMI* (*FOXMI-AS*) promotes the interaction of  $m^6A$  demethylase ALKBH5 with *FOXMI* nascent transcripts, which facilitates  $m^6A$  demethylation of *FOXMI* pre-mRNA at its 3'UTR. Demethylated *FOXMI* pre-mRNAs have higher affinity with HuR, a RNA binding protein, which stabilizes *FOXMI* to promote glioblastoma stem-like cells self-renewal and tumorigenesis (Zhang et al., 2017).

In summary, the knowledge of lncRNAs in neural circuitry assembly has been greatly expanded in recent years, whereas how lncRNAs exert roles in circuitry function in physiologic and pathologic conditions are much less known. Future studies would use modern genetic labeling, live-imaging, electrophysiology, behavioral and high-throughput means to explore how lncRNA-expressing neural cells are spatiotemporally participated in circuitry assembly and function, which can provide solid evidence that lncRNAs are essential fate and activity markers/determinants of neural cells.

## AUTHOR CONTRIBUTIONS

AW and JW collected references and wrote the review. YL and YZ wrote the review.

## ACKNOWLEDGMENTS

We thank all Zhou lab members for critical reading. This work was supported by grants from National Natural Science Foundation of China (No. 31671418 and No. 31471361), and Fundamental Research Funds for the Central Universities (2042017kf0205 and 2042017kf0242) to YZ.

## REFERENCES

- Abudayyeh, O. O., Gootenberg, J. S., Konermann, S., Joung, J., Slaymaker, I. M., Cox, D. B. T., et al. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* 353:aaf5573. doi: 10.1126/science.aaf5573
- Aliperti, V., and Donizetti, A. (2016). Long non-coding rna in neurons: new players in early response to BDNF stimulation. *Front. Mol. Neurosci.* 9:15. doi: 10.3389/fnmol.2016.00015
- Anderson, S. A., Kaznowski, C. E., Horn, C., Rubenstein, J. L. R., and McConnell, S. K. (2002). Distinct origins of neocortical projection neurons and interneurons *in vivo*. *Cerebral Cortex* 12, 702–709. doi: 10.1093/cercor/12.7.702

- Apra, J., and Calegari, F. (2015). Long non-coding RNAs in corticogenesis: deciphering the non-coding code of the brain. *EMBO J.* 34, 2865–2884. doi: 10.15252/embj.201592655
- Apra, J., Prenninger, S., Dori, M., Ghosh, T., Monasor, L. S., Wessendorf, E., et al. (2013). Transcriptome sequencing during mouse brain development identifies long non-coding RNAs functionally involved in neurogenic commitment. *EMBO J.* 32, 3145–3160. doi: 10.1038/emboj.2013.245
- Barry, G., Briggs, J. A., Vanichkina, D. P., Poth, E. M., Beveridge, N. J., Ratnu, V. S., et al. (2014). The long non-coding RNA Gomafu is acutely regulated in response to neuronal activation and involved in schizophrenia-associated alternative splicing. *Mol. Psychiatry* 19, 486–494. doi: 10.1038/mp.2013.45
- Belgard, T. G., Marques, A. C., Oliver, P. L., Abaan, H. O., Sirey, T. M., Hoerder-Suabedissen, A., et al. (2011). A transcriptomic atlas of mouse neocortical layers. *Neuron* 71, 605–616. doi: 10.1016/j.neuron.2011.06.039
- Berghoff, E. G., Clark, M. F., Chen, S., Cajigas, I., Leib, D. E., and Kohtz, J. D. (2013). *Evf2 (Dlx6as)* lncRNA regulates ultraconserved enhancer methylation and the differential transcriptional control of adjacent genes. *Development* 140, 4407–4416. doi: 10.1242/dev.099390
- Bernard, D., Prasanth, K. V., Tripathi, V., Colasse, S., Nakamura, T., Xuan, Z., et al. (2010). A long nuclear-retained non-coding RNA regulates synaptogenesis by modulating gene expression. *EMBO J.* 29, 3082–3093. doi: 10.1038/emboj.2010.199
- Birney, E., Stamatoyannopoulos, J. A., Dutta, A., Guigo, R., Gingeras, T. R., Margulies, E. H., et al. (2007). Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 447, 799–816. doi: 10.1038/nature05874
- Bond, A. M., Vangompel, M. J. W., Sametsky, E. A., Clark, M. F., Savage, J. C., Disterhoft, J. F., et al. (2009). Balanced gene regulation by an embryonic brain ncRNA is critical for adult hippocampal GABA circuitry. *Nat. Neurosci.* 12, 1020–1091. doi: 10.1038/nn.2371
- Briggs, J. A., Wolvetang, E. J., Mattick, J. S., Rinn, J. L., and Barry, G. (2015). Mechanisms of long non-coding RNAs in mammalian nervous system development, plasticity, disease, and evolution. *Neuron* 88, 861–877. doi: 10.1016/j.neuron.2015.09.045
- Carninci, P., Kasukawa, T., Katayama, S., Gough, J., Frith, M. C., Maeda, N., et al. (2005). The transcriptional landscape of the mammalian genome. *Science* 309, 1559–1563. doi: 10.1126/science.1112014
- Cesana, M., Cacchiarelli, D., Legnini, I., Santini, T., Sthandier, O., Chinappi, M., et al. (2011). A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell* 147, 358–369. doi: 10.1016/j.cell.2011.09.028
- Chalei, V., Sansom, S. N., Kong, L. S., Lee, S., Montiel, J., Vance, K. W., et al. (2014). The long non-coding RNA Dali is an epigenetic regulator of neural differentiation. *Elife* 3:e04530. doi: 10.7554/eLife.04530
- Chen, K., and Featherstone, D. E. (2011). Pre and postsynaptic roles for Drosophila CASK. *Mol. Cell Neurosci.* 48, 171–182. doi: 10.1016/j.mcn.2011.07.009
- Chen, L., Feng, P., Zhu, X., He, S., Duan, J., and Zhou, D. (2016). Long non-coding RNA Malat1 promotes neurite outgrowth through activation of ERK/MAPK signalling pathway in N2a cells. *J. Cell Mol. Med.* 20, 2102–2110. doi: 10.1111/jcmm.12904
- Cheng, J., Kapranov, P., Drenkow, J., Dike, S., Brubaker, S., Patel, S., et al. (2005). Transcriptional maps of 10 human chromosomes at 5-nucleotide resolution. *Science* 308, 1149–1154. doi: 10.1126/science.1108625
- Chodroff, R. A., Goodstadt, L., Sirey, T. M., Oliver, P. L., Davies, K. E., Green, E. D., et al. (2010). Long noncoding RNA genes: conservation of sequence and brain expression among diverse amniotes. *Genome Biol.* 11:R72. doi: 10.1186/gb-2010-11-7-r72
- Church, D. M., Goodstadt, L., Hillier, L. W., Zody, M. C., Goldstein, S., She, X. W., et al. (2009). Lineage-specific biology revealed by a finished genome assembly of the mouse. *PLoS Biol.* 7:e1000112. doi: 10.1371/journal.pbio.100112
- Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., et al. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819–823. doi: 10.1126/science.1231143
- Cox, D. B., Platt, R. J., and Zhang, F. (2015). Therapeutic genome editing: prospects and challenges. *Nat. Med.* 21, 121–131. doi: 10.1038/nm.3793
- Da Cruz, S., and Cleveland, D. W. (2011). Understanding the role of TDP-43 and FUS/TLS in ALS and beyond. *Curr. Opin. Neurobiol.* 21, 904–919. doi: 10.1016/j.conb.2011.05.029
- Darmanis, S., Sloan, S. A., Zhang, Y., Enge, M., Caneda, C., Shuer, L. M., et al. (2015). A survey of human brain transcriptome diversity at the single cell level. *Proc. Natl. Acad. Sci. U.S.A.* 112, 7285–7290. doi: 10.1073/pnas.1507125112
- Derrien, T., Johnson, R., Bussotti, G., Tanzer, A., Djebali, S., Tilgner, H., et al. (2012). The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res.* 22, 1775–1789. doi: 10.1101/gr.132159.111
- Desai, A. R., and McConnell, S. K. (2000). Progressive restriction in fate potential by neural progenitors during cerebral cortical development. *Development* 127, 2863–2872.
- Diederichs, S. (2014). The four dimensions of noncoding RNA conservation. *Trends Genet.* 30, 121–123. doi: 10.1016/j.tig.2014.01.004
- Dimitrova, N., Zamudio, J. R., Jong, R. M., Soukup, D., Resnick, R., Sarma, K., et al. (2014). *LincRNA-p21* activates *p21* In cis to promote polycomb target gene expression and to enforce the G1/S checkpoint. *Mol. Cell* 54, 777–790. doi: 10.1016/j.molcel.2014.04.025
- Djebali, S., Davis, C. A., Merkel, A., Dobin, A., Lassmann, T., Mortazavi, A., et al. (2012). Landscape of transcription in human cells. *Nature* 489, 101–108. doi: 10.1038/nature11233
- Dominissini, D., Moshitch-Moshkovitz, S., Schwartz, S., Salmon-Divon, M., Ungar, L., Osenberg, S., et al. (2012). Topology of the human and mouse m<sup>6</sup>A RNA methylomes revealed by m<sup>6</sup>A-seq. *Nature* 485, 201–206. doi: 10.1038/nature11112
- Dong, X., Chen, K., Cuevas-Diaz Duran, R., You, Y., Sloan, S. A., Zhang, Y., et al. (2015). Comprehensive identification of long non-coding RNAs in purified cell types from the brain reveals functional lncRNA in OPC fate determination. *PLoS Genet.* 11:e1005669. doi: 10.1371/journal.pgen.1005669
- Doudna, J. A., and Charpentier, E. (2014). The new frontier of genome engineering with CRISPR-Cas9. *Science* 346:1258096. doi: 10.1126/science.1258096
- D'ydewalle, C., Ramos, D. M., Pyles, N. J., Ng, S. Y., Gorz, M., Pilato, C. M., et al. (2017). The antisense transcript SMN-AS1 regulates SMN expression and is a novel therapeutic target for spinal muscular atrophy. *Neuron* 93, 66–79. doi: 10.1016/j.neuron.2016.11.033
- Eissmann, M., Gutschner, T., Hammerle, M., Gunther, S., Caudron-Herger, M., Gross, M., et al. (2012). Loss of the abundant nuclear non-coding RNA *MALAT1* is compatible with life and development. *RNA Biol.* 9, 1076–1087. doi: 10.4161/rna.21089
- Engreitz, J. M., Haines, J. E., Perez, E. M., Munson, G., Chen, J., Kane, M., et al. (2016). Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature* 539, 452–455. doi: 10.1038/nature20149
- Faghghi, M. A., Modarresi, F., Khalil, A. M., Wood, D. E., Sahagan, B. G., Morgan, T. E., et al. (2008). Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of  $\beta$ -secretase. *Nat. Med.* 14, 723–730. doi: 10.1038/nm1784
- Fatehullah, A., Tan, S. H., and Barker, N. (2016). Organoids as an *in vitro* model of human development and disease. *Nat. Cell Biol.* 18, 246–254. doi: 10.1038/ncb3312
- Femino, A. M., Fay, F. S., Fogarty, K., and Singer, R. H. (1998). Visualization of single RNA transcripts *in situ*. *Science* 280, 585–590. doi: 10.1126/science.280.5363.585
- Feng, J. C., Bi, C. M., Clark, B. S., Mady, R., Shah, P., and Kohtz, J. D. (2006). The *Evf-2* noncoding RNA is transcribed from the *Dlx-5/6* ultraconserved region and functions as a *Dlx-2* transcriptional coactivator. *Genes Dev.* 20, 1470–1484. doi: 10.1101/gad.1416106
- Franco, S. J., and Muller, U. (2013). Shaping our minds: stem and progenitor cell diversity in the mammalian neocortex. *Neuron* 77, 19–34. doi: 10.1016/j.neuron.2012.12.022
- Fu, Y., Dominissini, D., Rechavi, G., and He, C. (2014). Gene expression regulation mediated through reversible m<sup>6</sup>A RNA methylation. *Nat. Rev. Genet.* 15, 293–306. doi: 10.1038/nrg3724
- Gaspard, N., Bouschet, T., Hourez, R., Dimidschstein, J., Naeije, G., Van Den Ameel, J., et al. (2008). An intrinsic mechanism of corticogenesis from embryonic stem cells. *Nature* 455, 351–357. doi: 10.1038/nature07287

- Ghosh, S., Tibbit, C., and Liu, J. L. (2016). Effective knockdown of *Drosophila* long non-coding RNAs by CRISPR interference. *Nucleic Acids Res.* 44:e84. doi: 10.1093/nar/gkw063
- Gilbert, L. A., Larson, M. H., Morsut, L., Liu, Z., Brar, G. A., Torres, S. E., et al. (2013). CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 154, 442–451. doi: 10.1016/j.cell.2013.06.044
- Gilbert, L. A., Horlbeck, M. A., Adamson, B., Villalta, J. E., Chen, Y., Whitehead, E. H., et al. (2014). Genome-scale CRISPR-mediated control of gene repression and activation. *Cell* 159, 647–661. doi: 10.1016/j.cell.2014.09.029
- Gootenberg, J. S., Abudayyeh, O. O., Lee, J. W., Essletzbichler, P., Dy, A. J., Joung, J., et al. (2017). Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science* 56, 438–442. doi: 10.1126/science.aam9321
- Greig, L. C., Woodworth, M. B., Galazo, M. J., Padmanabhan, H., and Macklis, J. D. (2013). Molecular logic of neocortical projection neuron specification, development and diversity. *Nat. Rev. Neurosci.* 14, 755–769. doi: 10.1038/nrn3586
- Guil, S., and Esteller, M. (2012). Cis-acting noncoding RNAs: friends and foes. *Nat. Struct. Mol. Biol.* 19, 1068–1075. doi: 10.1038/nsmb.2428
- Gummalla, M., Galetti, S., Maeda, R. K., and Karch, F. (2014). Hox gene regulation in the central nervous system of *Drosophila*. *Front. Cell. Neurosci.* 8:96. doi: 10.3389/fncel.2014.00096
- Guo, J. U., Agarwal, V., Guo, H. L., and Bartel, D. P. (2014). Expanded identification and characterization of mammalian circular RNAs. *Genome Biol.* 15:409. doi: 10.1186/s13059-014-0409-z
- Guttman, M., Amit, I., Garber, M., French, C., Lin, M. F., Feldser, D., et al. (2009). Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* 458, 223–227. doi: 10.1038/nature07672
- Hadley, D., Murphy, T., Valladares, O., Hannehalli, S., Ungar, L., Kim, J., et al. (2006). Patterns of sequence conservation in presynaptic neural genes. *Genome Biol.* 7:R105. doi: 10.1186/gb-2006-7-11-r105
- Hansen, T. B., Jensen, T. I., Clausen, B. H., Bramsen, J. B., Finsen, B., Damgaard, C. K., et al. (2013). Natural RNA circles function as efficient microRNA sponges. *Nature* 495, 384–388. doi: 10.1038/nature11993
- He, D. Y., Wang, J. C., Lu, Y. L., Deng, Y. Q., Zhao, C. T., Xu, L. L., et al. (2017). lncRNA functional networks in Oligodendrocytes reveal stage-specific myelination control by an *IncOLL1/Suz12* complex in the CNS. *Neuron* 93, 362–378. doi: 10.1016/j.neuron.2016.11.044
- Ho, T. T., Zhou, N. J., Huang, J. G., Koirala, P., Xu, M., Fung, R. L., et al. (2015). Targeting non-coding RNAs with the CRISPR/Cas9 system in human cell lines. *Nucleic Acids Res.* 43:e17. doi: 10.1093/nar/gku198
- Hon, C. C., Ramilowski, J. A., Harshbarger, J., Bertin, N., Rackham, O. J. L., Gough, J., et al. (2017). An atlas of human long non-coding RNAs with accurate 5' ends. *Nature* 543, 199–204. doi: 10.1038/nature21374
- Hsu, P. D., Lander, E. S., and Zhang, F. (2014). Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157, 1262–1278. doi: 10.1016/j.cell.2014.05.010
- Huang, X., Luo, Y. L., Mao, Y. S., and Ji, J. L. (2017). The link between long noncoding RNAs and depression. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 73, 73–78. doi: 10.1016/j.pnpbp.2016.06.004
- Iossifov, I., O'roak, B. J., Sanders, S. J., Ronemus, M., Krumm, N., Levy, D., et al. (2014). The contribution of *de novo* coding mutations to autism spectrum disorder. *Nature* 515, 216–221. doi: 10.1038/nature13908
- Jeck, W. R., and Sharpless, N. E. (2014). Detecting and characterizing circular RNAs. *Nat. Biotechnol.* 32, 453–461. doi: 10.1038/nbt.2890
- Job, C., and Eberwine, J. (2001). Localization and translation of mRNA in dendrites and axons. *Nat. Rev. Neurosci.* 2, 889–898. doi: 10.1038/35104069
- Kapranov, P., Cheng, J., Dike, S., Nix, D. A., Dutttagupta, R., Willingham, A. T., et al. (2007a). RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 316, 1484–1488. doi: 10.1126/science.1138341
- Kapranov, P., Willingham, A. T., and Gingeras, T. R. (2007b). Genome-wide transcription and the implications for genomic organization. *Nat. Rev. Genet.* 8, 413–423. doi: 10.1038/nrg2083
- Katayama, S., Tomaru, Y., Kasukawa, T., Waki, K., Nakanishi, M., Nakamura, M., et al. (2005). Antisense transcription in the mammalian transcriptome. *Science* 309, 1564–1566. doi: 10.1126/science.1112009
- Keppetipola, N., Sharma, S., Li, Q., and Black, D. L. (2012). Neuronal regulation of pre-mRNA splicing by polypyrimidine tract binding proteins, PTB1 and PTB2. *Crit. Rev. Biochem. Mol. Biol.* 47, 360–378. doi: 10.3109/10409238.2012.691456
- Konermann, S., Brigham, M. D., Trevino, A. E., Joung, J., Abudayyeh, O. O., Barcena, C., et al. (2014). Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* 517, 583–588. doi: 10.1038/nature14136
- Lagier-Tourenne, C., Polymenidou, M., and Cleveland, D. W. (2010). TDP-43 and FUS/TLS: emerging roles in RNA processing and neurodegeneration. *Hum. Mol. Genet.* 19, R46–R64. doi: 10.1093/hmg/ddq137
- Lancaster, M. A., Renner, M., Martin, C.-A., Wenzel, D., Bicknell, L. S., Hurles, M. E., et al. (2013). Cerebral organoids model human brain development and microcephaly. *Nature* 501, 373–379. doi: 10.1038/nature12517
- Lefebvre, S., Bulet, P., Liu, Q., Bertrand, S., Clermont, O., Munnich, A., et al. (1997). Correlation between severity and SMN protein level in spinal muscular atrophy. *Nat. Genet.* 16, 265–269. doi: 10.1038/ng0797-265
- Lewejohann, L., Skryabin, B. V., Sachser, N., Prehn, C., Heiduschka, P., Thanos, S., et al. (2004). Role of a neuronal small non-messenger RNA: behavioural alterations in BC1 RNA-deleted mice. *Behav. Brain Res.* 154, 273–289. doi: 10.1016/j.bbr.2004.02.015
- Li, L., and Chang, H. Y. (2014). Physiological roles of long noncoding RNAs: insight from knockout mice. *Trends Cell Biol.* 24, 594–602. doi: 10.1016/j.tcb.2014.06.003
- Li, M., Wen, S., Guo, X., Bai, B., Gong, Z., Liu, X., et al. (2012). The novel long non-coding RNA CRG regulates *Drosophila* locomotor behavior. *Nucleic Acids Res.* 40, 11714–11727. doi: 10.1093/nar/gks943
- Li, L., Liu, B., Wapinski, O. L., Tsai, M. C., Qu, K., Zhang, J., et al. (2013). Targeted disruption of hotair leads to homeotic transformation and gene derepression. *Cell Rep.* 5, 3–12. doi: 10.1016/j.celrep.2013.09.003
- Lin, N. W., Chang, K. Y., Li, Z. H., Gates, K., Rana, Z. A., Dang, J. S., et al. (2014). An evolutionarily conserved long noncoding RNA TUNA controls pluripotency and neural lineage commitment. *Mol. Cell* 53, 1067–1067. doi: 10.1016/j.molcel.2014.03.013
- Lipovich, L., Dacht, F., Cai, J., Bagla, S., Balan, K., Jia, H., et al. (2012). Activity-dependent human brain coding/noncoding gene regulatory networks. *Genetics* 192, 1133–1148. doi: 10.1534/genetics.112.145128
- Liu, T., Huang, Y. Y., Chen, J. L., Chi, H. Y., Yu, Z. H., Wang, J., et al. (2014). Attenuated ability of BACE1 to cleave the amyloid precursor protein via silencing long noncoding RNA BACE1-AS expression. *Mol. Med. Rep.* 10, 1275–1281. doi: 10.3892/mmr.2014.2351
- Liu, Z., Li, X., Sun, N., Xu, Y., Meng, Y., Yang, C., et al. (2014). Microarray profiling and co-expression network analysis of circulating lncRNAs and mRNAs associated with major depressive disorder. *PLoS ONE* 9:e93388. doi: 10.1371/journal.pone.0093388
- Liu, S. J., Nowakowski, T. J., Pollen, A. A., Lui, J. H., Horlbeck, M. A., Attenello, F. J., et al. (2016). Single-cell analysis of long non-coding RNAs in the developing human neocortex. *Genome Biol.* 17:67. doi: 10.1186/s13059-016-0932-1
- Luo, Y. P., Coskun, V., Liang, A. B., Yu, J. H., Cheng, L. M., Ge, W. H., et al. (2015). Single-cell transcriptome analyses reveal signals to activate dormant neural stem cells. *Cell* 161, 1175–1186. doi: 10.1016/j.cell.2015.04.001
- Luo, S., Lu, Y. Y., Liu, L. C., Yin, Y. F., Chen, C. Y., Han, X., et al. (2016). Divergent lncRNAs regulate gene expression and lineage differentiation in pluripotent cells. *Cell Stem Cell* 18, 637–652. doi: 10.1016/j.stem.2016.01.024
- Martin, J. R., and Ollo, R. (1996). A new *Drosophila* Ca<sup>2+</sup> calmodulin-dependent protein kinase (Caki) is localized in the central nervous system and implicated in walking speed. *EMBO J.* 15, 1865–1876.
- Matsumoto, A., Pasut, A., Matsumoto, M., Yamashita, R., Fung, J., Monteleone, E., et al. (2017). mTORC1 and muscle regeneration are regulated by the LINC00961-encoded SPAR polypeptide. *Nature* 541, 228–232. doi: 10.1038/nature21034
- Mcconnell, S. K. (1995). Constructing the cerebral-cortex - neurogenesis and fate determination. *Neuron* 15:203 761–768. doi: 10.1016/0896-6273(95)90168-X
- McHugh, C. A., Russell, P., and Guttman, M. (2014). Methods for comprehensive experimental identification of RNA-protein interactions. *Genome Biol.* 15:203. doi: 10.1186/gb4152
- Mercer, T. R., Dinger, M. E., Sunken, S. M., Mehler, M. F., and Mattick, J. S. (2008). Specific expression of long noncoding RNAs in the mouse brain. *Proc. Natl. Acad. Sci. U.S.A.* 105, 716–721. doi: 10.1073/pnas.0706729105

- Mercer, T. R., Qureshi, I. A., Gokhan, S., Dinger, M. E., Li, G. Y., Mattick, J. S., et al. (2010). Long noncoding RNAs in neuronal-glia fate specification and oligodendrocyte lineage maturation. *BMC Neurosci.* 11:14. doi: 10.1186/1471-2202-11-14
- Meyer, Kate. D., Saletore, Y., Zumbo, P., Elemento, O., Mason Christopher, E, et al. (2012). Comprehensive analysis of mRNA Methylation reveals enrichment in 3' UTRs and near stop Codons. *Cell* 149, 1635–1646. doi: 10.1016/j.cell.2012.05.003
- Miller, J. A., Ding, S. L., Sunkin, S. M., Smith, K. A., Ng, L., Szafer, A., et al. (2014). Transcriptional landscape of the prenatal human brain. *Nature* 508, 199–206. doi: 10.1038/nature13185
- Modarresi, F., Faghihi, M. A., Lopez-Toledano, M. A., Fatemi, R. P., Magistri, M., Brothers, S. P., et al. (2012). Inhibition of natural antisense transcripts *in vivo* results in gene-specific transcriptional upregulation. *Nat. Biotechnol.* 30, 453–459. doi: 10.1038/nbt.2158
- Molinie, B., Wang, J., Lim, K. S., Hillebrand, R., Lu, Z. X., Van Wittenberghe, N., et al. (2016). m<sup>6</sup>A-LAIC-seq reveals the census and complexity of the m<sup>6</sup>A epitranscriptome. *Nat. Methods* 13, 692–698. doi: 10.1038/nmeth.3898
- Molyneaux, B. J., Goff, L. A., Brettler, A. C., Chen, H. H., Brown, J. R., Hrvatin, S., et al. (2015). DeCoN: genome-wide analysis of *in vivo* transcriptional dynamics during pyramidal neuron fate selection in neocortex. *Neuron* 85, 275–288. doi: 10.1016/j.neuron.2014.12.024
- Mus, E., Hof, P. R., and Tiedge, H. (2007). Dendritic BC200 RNA in aging and in Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* 104, 10679–10684. doi: 10.1073/pnas.0701532104
- Muslimov, I. A., Santi, E., Homel, P., Perini, S., Higgins, D., and Tiedge, H. (1997). RNA transport in dendrites: a *cis*-acting targeting element is contained within neuronal BC1 RNA. *J. Neurosci.* 17, 4722–4733.
- Muslimov, I. A., Banker, G., Brosius, J., and Tiedge, H. (1998). Activity-dependent regulation of dendritic BC1 RNA in hippocampal neurons in culture. *J. Cell Biol.* 141, 1601–1611. doi: 10.1083/jcb.141.7.1601
- Ng, S. Y., Johnson, R., and Stanton, L. W. (2012). Human long non-coding RNAs promote pluripotency and neuronal differentiation by association with chromatin modifiers and transcription factors. *EMBO J.* 31, 522–533. doi: 10.1038/emboj.2011.459
- Ng, S.-Y., Lin, L., Soh, B. S., and Stanton, L. W. (2013a). Long noncoding RNAs in development and disease of the central nervous system. *Trends Genet.* 29, 461–468. doi: 10.1016/j.tig.2013.03.002
- Ng, S. Y., Bogu, G. K., Soh, B. S., and Stanton, L. W. (2013b). The long noncoding RNA RMST interacts with SOX2 to regulate neurogenesis. *Mol. Cell* 51, 349–359. doi: 10.1016/j.molcel.2013.07.017
- NIH (2014). *BRAIN 2025: A Scientific Vision*. Available online at: <https://www.braininitiative.nih.gov/2025/>
- Niu, Y., Shen, B., Cui, Y., Chen, Y., Wang, J., Wang, L., et al. (2014). Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated gene targeting in one-cell embryos. *Cell* 156, 836–843. doi: 10.1016/j.cell.2014.01.027
- Onoguchi, M., Hirabayashi, Y., Koseki, H., and Gotoh, Y. (2012). A noncoding RNA regulates the neurogenin1 gene locus during mouse neocortical development. *Proc. Natl. Acad. Sci. U.S.A.* 109, 16939–16944. doi: 10.1073/pnas.1202956109
- Parikhshak, N. N., Luo, R., Zhang, A., Won, H., Lowe, J. K., Chandran, V., et al. (2013). Integrative functional genomic analyses implicate specific molecular pathways and circuits in autism. *Cell* 155, 1008–1021. doi: 10.1016/j.cell.2013.10.031
- Parikhshak, N. N., Swarup, V., Belgard, T. G., Irimia, M., Ramaswami, G., Gandal, M. J., et al. (2016). Genome-wide changes in lncRNA, splicing, and regional gene expression patterns in autism. *Nature* 540, 423–427. doi: 10.1038/nature20612
- Peng, Y., Clark, K. J., Campbell, J. M., Panetta, M. R., Guo, Y., and Ekker, S. C. (2014). Making designer mutants in model organisms. *Development* 141, 4042–4054. doi: 10.1242/dev.102186
- Platt, R. J., Chen, S., Zhou, Y., Yim, M. J., Swiech, L., Kempton, H. R., et al. (2014). CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* 159, 440–455. doi: 10.1016/j.cell.2014.09.014
- Polymenidou, M., Lagier-Tourenne, C., Hutt, K. R., Bennett, C. F., Cleveland, D. W., and Yeo, G. W. (2012). Misregulated RNA processing in amyotrophic lateral sclerosis. *Brain Res.* 1462, 3–15. doi: 10.1016/j.brainres.2012.02.059
- Poo, M. M., Du, J. L., Ip, N. Y., Xiong, Z. Q., Xu, B., and Tan, T. (2016). China brain project: basic neuroscience, brain diseases, and brain-inspired computing. *Neuron* 92, 591–596. doi: 10.1016/j.neuron.2016.10.050
- Poulin, J. F., Tasic, B., Hjerling-Lefler, J., Trimarchi, J. M., and Awatramani, R. (2016). Disentangling neural cell diversity using single-cell transcriptomics. *Nat. Neurosci.* 19, 1131–1141. doi: 10.1038/nn.4366
- Puthanveetil, S. V., Antonov, I., Kalachikov, S., Rajasethupathy, P., Choi, Y. B., Kohn, A. B., et al. (2013). A strategy to capture and characterize the synaptic transcriptome. *Proc. Natl. Acad. Sci. U.S.A.* 110, 7464–7469. doi: 10.1073/pnas.1304422110
- Qian, X. M., Shen, Q., Goderie, S. K., He, W. L., Capela, A., Davis, A. A., et al. (2000). Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. *Neuron* 28, 69–80. doi: 10.1016/S0896-6273(00)00086-6
- Qin, P., Parlak, M., Kuscu, C., Bandaria, J., Mir, M., Szlachta, K., et al. (2017). Live cell imaging of low- and non-repetitive chromosome loci using CRISPR-Cas9. *Nat. Commun.* 8:14725. doi: 10.1038/ncomms14725
- Quinn, J. J., and Chang, H. Y. (2016). Unique features of long non-coding RNA biogenesis and function. *Nat. Rev. Genet.* 17, 47–62. doi: 10.1038/nrg.2015.10
- Qureshi, I. A., Mattick, J. S., and Mehler, M. F. (2010). Long non-coding RNAs in nervous system function and disease. *Brain Res.* 1338, 20–35. doi: 10.1016/j.brainres.2010.03.110
- Raj, A., Van Den Bogaard, P., Rifkin, S. A., Van Oudenaarden, A., and Tyagi, S. (2008). Imaging individual mRNA molecules using multiple singly labeled probes. *Nat. Methods* 5, 877–879. doi: 10.1038/nmeth.1253
- Ramos, A. D., Diaz, A., Nellore, A., Delgado, R. N., Park, K. Y., Gonzales-Roybal, G., et al. (2013). Integration of genome-wide approaches identifies lncRNAs of adult neural stem cells and their progeny *in vivo*. *Cell Stem Cell* 12, 616–628. doi: 10.1016/j.stem.2013.03.003
- Ramos, A. D., Andersen, R. E., Liu, S. J., Nowakowski, T. J., Hong, S. J., Gertz, C. C., et al. (2015). The long noncoding RNA *Pnky* regulates neuronal differentiation of embryonic and postnatal neural stem cells. *Cell Stem Cell* 16, 439–447. doi: 10.1016/j.stem.2015.02.007
- Rani, N., Nowakowski, T. J., Zhou, H. J., Godshalk, S. E., Lisi, V., Kriegstein, A. R., et al. (2016). A primate lncRNA mediates notch signaling during neuronal development by sequestering miRNA. *Neuron* 90, 1174–1188. doi: 10.1016/j.neuron.2016.05.005
- Rinn, J. L., and Chang, H. Y. (2012). Genome regulation by long noncoding RNAs. *Annu. Rev. Biochem.* 81, 145–166. doi: 10.1146/annurev-biochem-051410-092902
- Rybak-Wolf, A., Stottmeister, C., Glazar, P., Jens, M., Pino, N., Giusti, S., et al. (2015). Circular RNAs in the mammalian brain are highly abundant, conserved, and dynamically expressed. *Mol. Cell* 58, 870–885. doi: 10.1016/j.molcel.2015.03.027
- Salmena, L., Poliseno, L., Tay, Y., Kats, L., and Pandolfi, P. P. (2011). A ceRNA hypothesis: the Rosetta stone of a hidden RNA language? *Cell* 146, 353–358. doi: 10.1016/j.cell.2011.07.014
- Salzman, J., Gawad, C., Wang, P. L., Lacayo, N., and Brown, P. O. (2012). Circular RNAs are the predominant transcript isoform from hundreds of human genes in diverse cell types. *PLoS ONE* 7:e30733. doi: 10.1371/journal.pone.0030733
- Sauvageau, M., Goff, L. A., Lodato, S., Bonev, B., Groff, A. F., Gerhardinger, C., et al. (2013). Multiple knockout mouse models reveal lincRNAs are required for life and brain development. *Elife* 2:e01749. doi: 10.7554/eLife.01749
- Schlackow, M., Nojima, T., Gomes, T., Dhir, A., Carmo-Fonseca, M., and Proudfoot, N. J. (2017). Distinctive patterns of transcription and RNA processing for human lincRNAs. *Mol. Cell* 65, 25–38. doi: 10.1016/j.molcel.2016.11.029
- Schorderet, P., and Duboule, D. (2011). Structural and functional differences in the long non-coding RNA *hotair* in mouse and human. *PLoS Genet.* 7:e1002071. doi: 10.1371/journal.pgen.1002071
- Shechner, D. M., Hacisuleyman, E., Younger, S. T., and Rinn, J. L. (2015). Multiplexable, locus-specific targeting of long RNAs with CRISPR-Display. *Nat. Methods* 12, 664–670. doi: 10.1038/nmeth.3433
- Shen, Q., Wang, Y., Dimos, J. T., Fasano, C. A., Phoenix, T. N., Lemischka, I. R., et al. (2006). The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells. *Nat. Neurosci.* 9, 743–751. doi: 10.1038/nn1694

- Shi, C., Zhang, L., and Qin, C. (2017). Long Non-coding RNAs in brain development, synaptic biology, and Alzheimer's disease. *Brain Res. Bull.* 32, 160–169. doi: 10.1016/j.brainresbull.2017.03.010
- Skryabin, B. V., Sukonina, V., Jordan, U., Lewejohann, L., Sachser, N., Muslimov, I., et al. (2003). Neuronal untranslated BC1 RNA: targeted gene elimination in mice. *Mol. Cell Biol.* 23, 6435–6441. doi: 10.1128/MCB.23.18.6435-6441.2003
- Slawson, J. B., Kuklin, E. A., Ejima, A., Mukherjee, K., Ostrovsky, L., and Griffith, L. C. (2011). Central regulation of locomotor behavior of *Drosophila melanogaster* depends on a CASK isoform containing CaMK-like and L27 Domains. *Genetics* 187, 171–184. doi: 10.1534/genetics.110.123406
- Soshnev, A. A., Ishimoto, H., Mcallister, B. F., Li, X. G., Wehling, M. D., Kitamoto, T., et al. (2011). A conserved long noncoding RNA affects sleep behavior in *Drosophila*. *Genetics* 189, U455–U497. doi: 10.1534/genetics.111.131706
- Spadaro, P. A., Flavell, C. R., Widagdo, J., Ratnu, V. S., Troup, M., Ragan, C., et al. (2015). Long noncoding rna-directed epigenetic regulation of gene expression is associated with anxiety-like behavior in Mice. *Biol. Psychiatry* 78, 848–859. doi: 10.1016/j.biopsych.2015.02.004
- Su, Y., Shin, J., Zhong, C., Wang, S., Roychowdhury, P., Lim, J., et al. (2017). Neuronal activity modifies the chromatin accessibility landscape in the adult brain. *Nat. Neurosci.* 20, 476–483. doi: 10.1038/nn.4494
- Tasic, B., Menon, V., Nguyen, T. N., Kim, T. K., Jarsky, T., Yao, Z., et al. (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat. Neurosci.* 19, 335–346. doi: 10.1038/nn.4216
- Tay, Y., Rinn, J., and Pandolfi, P. P. (2014). The multilayered complexity of ceRNA crosstalk and competition. *Nature* 505, 344–352. doi: 10.1038/nature12986
- Telley, L., Govindan, S., Prados, J., Stevant, I., Nef, S., Dermitzakis, E., et al. (2016). Sequential transcriptional waves direct the differentiation of newborn neurons in the mouse neocortex. *Science* 351, 1443–1446. doi: 10.1126/science.aad8361
- Tiedge, H., Freneau, R. T., Weinstock, P. H., Arancio, O., and Brosius, J. (1991). Dendritic location of neural Bc1 RNA. *Proc. Natl. Acad. Sci. U.S.A.* 88, 2093–2097. doi: 10.1073/pnas.88.6.2093
- Tollervey, J. R., Curk, T., Rogelj, B., Briese, M., Cereda, M., Kayikci, M., et al. (2011). Characterizing the RNA targets and position-dependent splicing regulation by TDP-43. *Nat. Neurosci.* 14, 452–458. doi: 10.1038/nn.2778
- Uhde, C. W., Vives, J., Jaeger, I., and Li, M. (2010). Rmst Is a novel marker for the mouse ventral mesencephalic floor plate and the anterior Dorsal midline cells. *PLoS ONE* 5:641. doi: 10.1371/journal.pone.0008641
- Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lonnerberg, P., Lou, D., et al. (2015). Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat. Neurosci.* 18, 145–153. doi: 10.1038/nn.3881
- Vance, K. W., Sansom, S. N., Lee, S., Chalei, V., Kong, L., Cooper, S. E., et al. (2014). The long non-coding RNA Paupar regulates the expression of both local and distal genes. *EMBO J.* 33, 296–311. doi: 10.1002/embj.201386225
- Wang, H., Iacoangeli, A., Popp, S., Muslimov, I. A., Imataka, H., Sonenberg, N., et al. (2002). Dendritic BC1 RNA: functional role in regulation of translation initiation. *J. Neurosci.* 22, 10232–10241.
- Wang, X. T., Arai, S., Song, X. Y., Reichart, D., Du, K., Pascual, G., et al. (2008). Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature* 454, 126–130. doi: 10.1038/nature06992
- Wang, K. C., Yang, Y. W., Liu, B., Sanyal, A., Corces-Zimmerman, R., Chen, Y., et al. (2011). A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature* 472, U120–U158. doi: 10.1038/nature09819
- Wang, Y., Zhao, X., Ju, W., Flory, M., Zhong, J., Jiang, S., et al. (2015). Genome-wide differential expression of synaptic long noncoding RNAs in autism spectrum disorder. *Transl. Psychiatry* 5:e660. doi: 10.1038/tp.2015.144
- Westholm, J. O., Miura, P., Olson, S., Shenker, S., Joseph, B., Sanfilippo, P., et al. (2014). Genome-wide analysis of drosophila circular rnas reveals their structural and sequence properties and age-dependent neural accumulation. *Cell Rep.* 9, 1966–1980. doi: 10.1016/j.celrep.2014.10.062
- Wonders, C. P., and Anderson, S. A. (2006). The origin and specification of cortical interneurons. *Nat. Rev. Neurosci.* 7, 687–696. doi: 10.1038/nrn1954
- Wu, T., Chen, C., Yang, L., Zhang, M., Zhang, X., Jia, J., et al. (2015). Distinct lncRNA expression profiles in the prefrontal cortex of SD rats after exposure to methylphenidate. *Biomed. Pharmacother.* 70, 239–247. doi: 10.1016/j.biopha.2015.01.023
- Xing, Y. H., Bai, Z. Q., Liu, C. X., Hu, S. B., Ruan, M. H., and Chen, L. L. (2016). Research progress of long noncoding RNA in China. *Iubmb Life* 68, 887–893. doi: 10.1002/iub.1564
- Xue, Y. C., Ouyang, K. F., Huang, J., Zhou, Y., Ouyang, H., Li, H. R., et al. (2013). Direct conversion of fibroblasts to neurons by reprogramming PTB-regulated microRNA circuits. *Cell* 152, 82–96. doi: 10.1016/j.cell.2012.11.045
- Yan, L. Y., Yang, M. Y., Guo, H. S., Yang, L., Wu, J., Li, R., et al. (2013). Single-cell RNA-Seq profiling of human preimplantation embryos and embryonic stem cells. *Nat. Struct. Mol. Biol.* 20, 1131–1139. doi: 10.1038/nsmb.2660
- Yang, Y. W., Flynn, R. A., Chen, Y., Qu, K., Wan, B. B., Wang, K. C., et al. (2014). Essential role of lncRNA binding for WDR5 maintenance of active chromatin and embryonic stem cell pluripotency. *Elife* 3:e02046. doi: 10.7554/eLife.02046
- Yang, L., Zhang, J., Kamelgarn, M., Niu, C., Gal, J., Gong, W., et al. (2015). Subcellular localization and RNAs determine FUS architecture in different cellular compartments. *Hum. Mol. Genet.* 24, 5174–5183. doi: 10.1093/hmg/ddv239
- You, X. T., Vlatkovic, I., Babic, A., Will, T., Epstein, I., Tushev, G., et al. (2015). Neural circular RNAs are derived from synaptic genes and regulated by development and plasticity. *Nat. Neurosci.* 18, 603–610. doi: 10.1038/nn.3975
- Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., et al. (2003). The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell* 112, 317–327. doi: 10.1016/S0092-8674(03)00079-5
- Zeisel, A., Muñoz-Manchado, A. B., Codeluppi, S., Lönnerberg, P., La Manno, G., Jureus, A., et al. (2015). Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* 347, 1138–1142. doi: 10.1126/science.aaa1934
- Zhang, S., Zhao, B. S., Zhou, A., Lin, K., Zheng, S., Lu, Z., et al. (2017). m6A Demethylase ALKBH5 maintains tumorigenicity of Glioblastoma stem-like cells by sustaining FOXM1 expression and cell proliferation program. *Cancer Cell* 31, 591.e596–606.e596. doi: 10.1016/j.ccell.2017.02.013
- Zhao, X. L., Tang, Z. X., Zhang, H. K., Atianjoh, F. E., Zhao, J. Y., Liang, L. L., et al. (2013). A long noncoding RNA contributes to neuropathic pain by silencing Kcna2 in primary afferent neurons. *Nat. Neurosci.* 16, U1024–U1076. doi: 10.1038/nn.3438
- Zhao, Y., Li, H., Fang, S. S., Kang, Y., Wu, W., Hao, Y. J., et al. (2016). NONCODE 2016: an informative and valuable data source of long non-coding RNAs. *Nucleic Acids Res.* 44, D203–D208. doi: 10.1093/nar/gkv1252
- Zhong, J., Chuang, S. C., Bianchi, R., Zhao, W., Lee, H., Fenton, A. A., et al. (2009). BC1 regulation of metabotropic glutamate receptor-mediated neuronal excitability. *J. Neurosci.* 29, 9977–9986. doi: 10.1523/JNEUROSCI.3893-08.2009
- Zhou, Y. (2012). Cortical development and asymmetric cell divisions. *Front. Biol.* 7:1235. doi: 10.1007/s11515-012-1235-x
- Zhu, S. Y., Li, W., Liu, J. Z., Chen, C. H., Liao, Q., Xu, P., et al. (2016). Genome-scale deletion screening of human long non-coding RNAs using a paired-guide RNA CRISPR-Cas9 library. *Nat. Biotechnol.* 34, 1279–1286. doi: 10.1038/nbt.3715

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Wang, Wang, Liu and Zhou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.