

Cognitive neuroscience of human counterfactual reasoning

Nicole Van Hoeck¹, Patrick D. Watson² and Aron K. Barbey^{2,3,4,5,6,7*}

¹ Psychology and Educational Sciences, Vrije Universiteit Brussel, Brussels, Belgium, ² Decision Neuroscience Laboratory, Beckman Institute for Advanced Science and Technology, University of Illinois, Urbana, IL, USA, ³ Department of Internal Medicine, University of Illinois, Champaign, IL, USA, ⁴ Department of Psychology, University of Illinois, Champaign, IL, USA, ⁵ Department of Speech and Hearing Science, University of Illinois, Champaign, IL, USA, ⁶ Neuroscience Program, University of Illinois, Champaign, IL, USA, ⁷ Carle R. Woese Institute for Genomic Biology, University of Illinois, Champaign, IL, USA

Counterfactual reasoning is a hallmark of human thought, enabling the capacity to shift from perceiving the immediate environment to an alternative, imagined perspective. Mental representations of counterfactual possibilities (e.g., imagined past events or future outcomes not yet at hand) provide the basis for learning from past experience, enable planning and prediction, support creativity and insight, and give rise to emotions and social attributions (e.g., regret and blame). Yet remarkably little is known about the psychological and neural foundations of counterfactual reasoning. In this review, we survey recent findings from psychology and neuroscience indicating that counterfactual thought depends on an integrative network of systems for affective processing, mental simulation, and cognitive control. We review evidence to elucidate how these mechanisms are systematically altered through psychiatric illness and neurological disease. We propose that counterfactual thinking depends on the coordination of multiple information processing systems that together enable adaptive behavior and goal-directed decision making and make recommendations for the study of counterfactual inference in health, aging, and disease.

Keywords: counterfactual thought, mental simulation, adaptive behavior, clinical disorders

Introduction

“Without considering alternatives to reality, we must accept the past as having been inevitable and must believe that the future will be no different from the past. The generation of counterfactuals gives us the flexibility in thinking about possible futures and prepares us better for those futures”
(Johnson and Sherman, 1999, p. 150).

Humans have the remarkable ability to infer how an event might have unfolded differently, without directly experiencing this alternative reality. We employ such counterfactual reasoning to make sense of the past, plan courses of action, make emotional and social judgments, and guide adaptive behavior. In this review article, we introduce an integrative cognitive neuroscience framework for understanding the psychological and neural foundations of counterfactual thought, drawing upon cognitive and neuroscience evidence to suggest that counterfactual thinking depends on an integrative network of systems for affective processing, mental simulation, and cognitive control.

We begin by surveying the psychological literature on counterfactual reasoning, followed by a review of the neuroscience literature on affective processing, mental simulation, and cognitive control networks supporting counterfactual thought. Finally, we review evidence to elucidate how these mechanisms are systematically altered in

OPEN ACCESS

Edited by:

Lutz Jäncke,
University of Zurich, Switzerland

Reviewed by:

Roberto Colom,
Universidad Autonoma
de Madrid, Spain
Marie Hennecke,
University of Zurich, Switzerland

*Correspondence:

Aron K. Barbey,
Decision Neuroscience Laboratory,
Beckman Institute for Advanced
Science and Technology, University
of Illinois, Urbana-Champaign,
405 North Mathews Avenue, Urbana,
IL 61801, USA
Barbey@illinois.edu;
www.DecisionNeuroscienceLab.org

Received: 02 January 2015

Accepted: 07 July 2015

Published: 23 July 2015

Citation:

Van Hoeck N, Watson PD and
Barbey AK (2015) Cognitive
neuroscience of human
counterfactual reasoning.
Front. Hum. Neurosci. 9:420.
doi: 10.3389/fnhum.2015.00420

psychiatric illness and neurological disease and draw conclusions about the importance of counterfactual thought for understanding adaptive behavior and goal-directed decision making.

Stages of Counterfactual Thought

We propose that counterfactual reasoning depends on three processing stages: *Activation*, *Inference*, and *Adaptation* (see **Figure 1**; e.g., Byrne, 2002; Epstude and Roese, 2008; Barbey et al., 2009). To illustrate these stages, consider the example of receiving a rejection letter for a new job. The letter would *activate* prior memories (e.g., the process of submitting an application and interviewing for the position) and elicit a mental simulation of the event (e.g., representing relevant knowledge and expectations; for example, the job requirements). As this activation spreads, it triggers mental simulations of similar situations (previous applications), allowing one to *infer* how this scenario would have played out under different circumstances (e.g., if one had spent more time preparing for the interview; i.e., the counterfactual event). These nearby counterfactual simulations influence one's interpretation of the factual experience (e.g., the committee's decision was caused by my failure to adequately prepare for the interview) and promote *adaptive behavior* that guides future planning and problem solving. The process of counterfactual reasoning therefore supports inferences about the factual event that guide understanding, modulate emotional responses, and provide the basis for future planning and decision-making.

Our proposal is that the coordination of systems for: (i) mental simulation; (ii) inference; and (iii) learning and adaptation together enable counterfactual reasoning. In the following sections, we review each of these processes and survey evidence from psychology and neuroscience to elucidate their contributions to counterfactual reasoning.

Activation of Mental Simulations

Counterfactual thoughts are automatically employed in response to real world experiences, in particular those situations where negative emotions are linked to violations of expectations and motivations (e.g., an unsuccessful job-application), in the form of implicit or explicit goal failures (e.g., answering the interview questions weakly), or close calls (physical, temporal or numerical proximity; e.g., being the runner-up) trigger counterfactual thoughts, representing a probable alternative state of affairs (Kahneman and Tversky, 1982; Davis et al., 1995; Sanna and Turley, 1996; Roese and Olson, 1997; McGraw et al., 2005; Epstude and Roese, 2008; Kühberger et al., 2011).

We propose that counterfactual thought depends on mental models of alternative possibilities represented in the form of mental simulations (Damasio, 1989; Barsalou et al., 2003a,b; Markman et al., 2009). Simulations provide the basis for constructing mental models of events and of imagining alternative realities “if only” different decisions were made or actions taken. For example, the counterfactual inference that “If I was to vacation in Europe rather than continue writing, then I wouldn't be this stressed” activates a mental simulation, which represents


relevant agents (e.g., the author), objects (e.g., museums), actions (e.g., sightseeing), mental states (e.g., freedom) and background settings (e.g., Florence). These simulations provide the basis for evaluating the consequences of the real and alternative courses of action, with the simulation of the author “vacationing in Europe” resulting in being more relaxed but not completing the article.

Our proposal shares much in common with previous frameworks of counterfactual reasoning, emphasizing the generative/constructive nature of counterfactual thought. For example, norm theory suggests that such reasoning is driven by simulations of previously encoded exemplars (Kahneman and Tversky, 1982; Kahneman and Miller, 1986). This theory emphasized the role of counterfactual thought in *reframing* such scenarios—generating alternative possibilities that change the norms (and expectations) used to interpret a state of affairs. The Model Theory of counterfactual thought (Byrne, 2002, 2007) emphasizes individuals' ability to entertain multiple parallel models corresponding to alternative possibilities, and suggests that counterfactual thought is engaged to search the space of possible alternatives (e.g., in simulating permissible or forbidden actions). Building upon these frameworks, the structured event complex theory proposes that counterfactual thought engages a network of regions within prefrontal cortex (PFC) that represent alternative goals, behavioral intentions, mindsets, motivations, and self-inferences that enable behavioral change and adaptation (Barbey et al., 2009).

Counterfactual Inference

In contrast to other types of conditional and hypothetical reasoning, an important feature of counterfactual inference is that it adheres to a “nearest possible world” constraint (Lewis, 1979, 1986; Hendrickson, 2010; Rafetseder et al., 2010, 2013; Van Hoeck et al., 2012). A counterfactual must closely model one's own experience of the real state of the world. It ties to specific situational features and prior knowledge of the situation (i.e., it requires the fewest independent changes to the actual circumstances and be maximal coherent with prior history). This constraint differentiates counterfactual thought from less constrained fantasy or imagination (Rao and Foo, 1989; Lebow, 2000; Revlin et al., 2001; Roese and Summerville, 2005; Byrne, 2007; Hendrickson, 2010; Dehghani et al., 2012). The alternative outcome suggested by the counterfactual reflects a small deviation from reality and is therefore probable (Over et al., 2007; Epstude and Roese, 2008; Petrocelli et al., 2011).

For example, counterfactual thought often focuses on the factor that: (1) played the strongest role in the (un-)desired outcome; (2) deviated most from expectancies; or (3) is most under the participants control (i.e., “mutable” factor; McCloy and Byrne, 2000; McGraw et al., 2005; Byrne, 2007; Over et al., 2007; Dehghani et al., 2012; Rips and Edwards, 2013). Counterfactual thought rarely suggests alterations from natural or social laws (Byrne, 1997, 2007; Roese, 1997; Nasco and Marsh, 1999; McCloy and Byrne, 2000; Walsh and Byrne, 2004; Tykocinski and Steinberg, 2005; Dehghani et al., 2012). Furthermore, counterfactual thought is tied to the meaning and relevance of specific events rather than to general situations or tendencies. Consider an example in which one failed an



exam by studying the wrong material. In this situation, the generalization that studying promotes academic achievement is unhelpful.

In addition to being tied to a specific event, counterfactual thought is often motivated by a specific individual perspective. An individual's implicit *belief of attainability* or self-efficacy will play an important role (Sanna, 1997; Wong et al., 2012; Dyczewski and Markman, 2012; Tyser et al., 2012; Zhang and Covey, 2014). For example, an individual who studied the wrong material for an exam would not believe they would otherwise pass, failing to endorse the counterfactual, "If I had studied the correct material, I would have passed the exam", when they held the implicit belief that they are not smart enough to master the correct material. To this individual, the semifactual, "Even if I had studied the correct material, I would have failed the exam", would be more realistic. Implicit belief can therefore explain why individuals sometimes alter other aspects than their own behavior (e.g., task-characteristics) when they are explicitly cued to generate counterfactuals (Pighin et al., 2011; Ferrante et al., 2013).

Learning and Adaptation

Once the counterfactual state of affairs has been inferred, this information will be incorporated in the representation of the factual state of affairs (re-evaluation) and update prior beliefs and action-values, resulting in behavioral and affective consequences. Counterfactual inference supports adaptive behavior and is known to:

- Enhance memory distortions, such as hindsight bias and source confusion (retrospective overestimation of the outcome's likelihood), contributing to suboptimal decision making (e.g., investment practices; Roese and Olson, 1996; Roese et al., 2004; Kruger et al., 2005; Nestler and von Collani, 2008; Petrocelli and Sherman, 2010; Petrocelli and Harris, 2011; Strahilevitz et al., 2011; Gerlach et al., 2013).
- Promote a relational/analytical or an expansive/creative processing style (Markman et al., 2007).
- Enable learning from past experience (Byrne, 1997; Epstude and Roese, 2008; Smallman and McCulloch, 2012) and the formation of behavioral intentions (c.f., Functional Theory of Counterfactual Thinking; Epstude and Roese, 2008).
- Support future planning and prediction (Roese, 1999; Barbey and Sloman, 2007; Markman et al., 2008; Smallman and Roese, 2009; Tobia et al., 2014).
- Provide the basis for creativity and insight (Sternberg and Gastel, 1989; Gomez Beldarrain et al., 2005; Markman et al., 2007; Kray et al., 2009; Roese and Morrison, 2009).
- Impart meaning to important life events by increasing perceptions of fate, benefit and growth (Davis et al., 1998; Koo et al., 2008; Kray et al., 2010; Teigen and Jensen, 2011).
- Generate emotions and social ascriptions (e.g., guilt, regret, blame and relief) that are central for managing and regulating social behavior (Davis et al., 1995; Roese and Olson, 1997, 2007; Pieters and Zeelenberg, 2005; Alicke et al., 2008; Coricelli and Rustichini, 2010; Brassens et al., 2012; Miller et al., 2013).

Several interacting factors influence the behavioral and affective consequences of counterfactual inference. In the first place the counterfactual outcome *value* will be important. Is the alternative outcome better or worse (valence; upward vs. downward counterfactual) and by how much (magnitude)? This valuation will impact how an individual perceives the factual, experienced outcome and its relative value. Secondly, does the individual perceive a *favorable opportunity* to improve performance? Will a similar situation take place in the future (repeatability, e.g., midterm vs. final exams) and does the individual believe it is feasible to change the outcome: do they perceive to be in control of the situation and that the counterfactual state is attainable? What is the probability that the alternative action(s) can bring about the desired outcome? The third factor is the perspective or *processing mode* in which counterfactual states are represented. Markman and McMullen (2003) suggest in their Reflective and Evaluative Model that a counterfactual representation can be further processed in two ways: the counterfactual representation can function as a standard to which to compare the real world (evaluative mode) or the counterfactual representation can be further processed in a more concrete-experiential way without using it as a standard of comparison (reflective mode). These different modes are very similar to the distinction between imagining an event from the first- vs. third-person perspective. From a first-person perspective (similar to reflective mode), the individual imagines an event as if it is unfolding before them (the individual is part of the experience). However, when the event is imagined from a third-person perspective (similar to evaluative mode), the individual visualizes the event from an observer's standpoint and examines the broader context (Libby and Eibach, 2011a,b; Valenti et al., 2011). This suggests that an evaluative focus takes current beliefs, traits, and goals more into account, whereas a reflective mode focuses more on sensory details of the counterfactual representation. We now turn to a discussion of

how the combination among the reviewed factors (i.e., value, opportunity, and processing mode) influences counterfactual inference.

Evaluations of how a counterfactual possibility could have been better (i.e., upward counterfactual thoughts), and how we might have personally effected such an outcome (perceived attainability) often elicit an emotional and social reaction. For example, regret will increase as we feel more responsible for the outcome and as the discrepancy between the experienced and counterfactual outcome value increases (Zeelenberg et al., 1998; van Dijk and Zeelenberg, 2002; Nicolle et al., 2011b). Like feelings of regret, the social ascription of blame arises from counterfactual reasoning about an action that caused or contributed to the negative experience and that is perceived to be under the person's control (see example about avoided car crash). However, blame demands an additional negative evaluation of the action itself (e.g., intentionally causing harm, breaking social norms, negligence; i.e., culpable control; c.f. Alicke et al., 2008).

Counterfactual thinking combined with an opportunity to improve future performance (repeatability of a similar event and belief in feasibility of different outcome), elicits *behavioral motivations* to pursue the counterfactual outcome, and modulates one's perception of control and preparedness, boosting persistence and performance (Roese and Olson, 1997; Sanna, 1997; Nasco and Marsh, 1999; Markman and Miller, 2006; Quelhas et al., 2008; Smallman and Roese, 2009; Dyczewski and Markman, 2012; Smallman, 2013; Zhang and Covey, 2014). It is assumed that the affect coupled with the experience will be *reappraised* in light of one's new feelings, and *inhibit* further counterfactual thoughts (Roese and Olson, 1997; Sanna, 1997; McMullen and Markman, 2002; Quelhas et al., 2008; Libby and Eibach, 2011b; Libby et al., 2011; Valenti et al., 2011). The reappraisal of the situation might also explain why the emotional intensity experienced during episodic counterfactual thinking is lower than during episodic past and future thinking (De Brigard and Giovanello, 2012).

However, in situations without opportunity for improvement (e.g., if the change is infeasible or the scenario cannot reoccur), upward counterfactual thinking will not be beneficial (Sanna, 1997; McMullen and Markman, 2002; Branscombe et al., 2003; Alicke et al., 2008; McCrea, 2008; Libby and Eibach, 2011a; Nicolle et al., 2011b; Dyczewski and Markman, 2012; Tyser et al., 2012; Wong et al., 2012; Zhang and Covey, 2014). Consider the case of breast cancer patients 3–4 months after diagnosis. In this group a high level of upward counterfactual thought is related to a high level of psychological distress (Gilbar and Hevroni, 2007). During this phase of the illness the shock of this life-threatening diagnosis is still being processed, for which upward counterfactuals would not be helpful. Similar results have been found with assault-victims (Branscombe et al., 2003; El Leithy et al., 2006). When one perceives no favorable opportunities to obtain a desired outcome or implement corrective behavior, goal disengagement is an adaptive coping mechanism (Wrosch et al., 2013). As such, downward counterfactual thoughts (i.e., thoughts about how the event could have turned out worse) instead might help regulate emotions in these circumstances and promote the inference that

an individual is a “lucky survivor” instead of an “unlucky victim” (Markman et al., 1993; Roese, 1994; Davis et al., 1998; Teigen and Jensen, 2011).

Markman and McMullen (2003) suggest that counterfactual information can also be processed in a more concrete, experiential manner (see also Libby and Eibach, 2011a,b). Effects of this processing match *affective assimilation* effects: the experienced affect is pulled into the direction of the counterfactual outcome. Processing an upward counterfactual in this manner will yield more positive affect (less intense negative emotions), but less behavioral benefits than evaluative upward counterfactuals. Vividly imagining a worse counterfactual outcome can instead lead to increase in negative affect, which can enhance persistence and performance to avoid this potentially worse outcome in the future (McMullen and Markman, 2000; Markman et al., 2008).


While most of the cases outlined above result in persistent behavioral change or re-evaluation of beliefs, not every instance of counterfactual reasoning will result in a persistent or positive behavioral change. Yet even in these cases, we suggest that counterfactual reasoning may influence the outcome. For example, if our counterfactual reasoning effort indicates a change in behavior would not result in a more favorable outcome, we still adapt our future behavior to this counterfactual information. Thus, we argue that counterfactual reasoning makes an obligatory contribution to learning, by providing alternative realities from which to draw conclusions. Individuals therefore adapt to counterfactuals as well as real feedback—and whether this is ultimately helpful depends upon future experiences.

These diverse effects derive from essential characteristics of counterfactual thought, resulting from the construction of a coherent and plausible representation of a nearby alternative that is maintained alongside an accurate representation of the actual event. The engagement of attentional and executive control processes enables comparison between these parallel representations. This comparison ultimately drives the evaluation of the final outcome and its contribution to learning.

Thus, counterfactual thinking is a constructive process. It draws for content upon general semantic knowledge, and specific memories of past experiences. It requires spatial and temporal integration of the elements of experience. It utilizes working memory in the storage and manipulation of items. It requires emotional processing, and integrates motivations and goals. These higher cognitive functions interact to construct an internal representation of the target scenario.

Neural Networks

We review neuroscience evidence to elucidate three neural networks that together support counterfactual thought: (1) the mental simulation network; (2) the cognitive control network; and (3) the reward network (see **Figure 2**). As such counterfactual reasoning is supported by “networks that underlie domain general functions that cut across different psychological domains” (Barrett and Satpute, 2013). It is the *interaction* between these networks that make the complex



psychological phenomena of counterfactual reasoning, including its effect on our behavior and emotions, possible. This view indicates that there is no distinctive “counterfactual reasoning” network nor that there is a one-on-one association between one of the processing stages of counterfactual reasoning we discussed above and one specific neural network. However, the engagement of these interactive networks, and regions within it (e.g., hippocampus or amygdala) depend on the type and amount of information that needs to be processed at a particular time.

Mental Simulation Network

Counterfactual thought engages a neural network that supports core processes for mentally undoing the present state of affairs and imagining alternative realities “if only” different decisions were made or actions taken (e.g., Nieuwland, 2012; Urrutia et al., 2012; De Brigard et al., 2013; Kulakova et al., 2013; Van Hoeck et al., 2013, 2014). This mental simulation network, also called the default mode network, engages regions in the medial frontal and temporal lobes, the posterior cingulate cortex, precuneus, and the lateral parietal and temporal lobes. This network is active when individuals are engaged in internally focused tasks, self-projection and scene construction, including

autobiographical memory retrieval, envisioning the future, and conceiving the perspectives of others (e.g., Buckner and Carroll, 2007; Hassabis and Maguire, 2009; Spreng et al., 2009; Spreng and Grady, 2010; Summerfield et al., 2010; Schacter et al., 2012, 2015; Spreng and Mar, 2012; Martinelli et al., 2013). Thus, this network is engaged when a situation is observed and interpreted, mentally altered and re-evaluated (i.e., all stages of counterfactual thought). Medial temporal lobe regions, such as the hippocampus, provide information from prior experiences in the form of memories and associations that are the core building blocks of mental simulation. Medial PFC regions facilitate the flexible use of this information during the construction of self-relevant mental simulations. These two subnetworks converge on important nodes of integration including the posterior cingulate cortex. Here we review emerging neuroscience evidence linking counterfactual thought to this network’s function.

Van Hoeck et al. (2010, 2013) directly compared past, future and counterfactual simulations of episodic events (i.e., specific personal experienced events) in this fMRI study. Healthy college students performed three tasks: (1) Recall a negative past event (e.g., “Imagine/recall the car accident from your holiday in France”); (2) Imagine how a similar future event turns out

positive instead (e.g., “Imagine a future car trip that goes well”); and (3) Imagine how the past event might have turned out better (a positive upward counterfactual) given a specific additive counterfactual action (e.g., “Imagine what would have happened if you had paid more attention to the road during that car trip in France”). Extensive neural overlap between these tasks was expected and observed. Episodic past, future and counterfactual simulation commonly activated the mental simulation network. But compared to past episodic thinking in particular, counterfactual simulation of episodic events engaged the mental simulation network to a larger extent (see Van Hoeck et al., 2013 for more information). Similar results were also documented by De Brigard et al. (2013), who likewise compared episodic past and counterfactual simulation. This stronger engagement might be understood as counterfactual thinking being a more complex or strenuous form of mental simulation, whereby both factual as counterfactual elements need to be represented and inferred.

A follow-up study conducted by Van Hoeck et al. (2014) demonstrated that counterfactual reasoning about hypothetical social scenarios engage this mental simulation network as well. Participants read a short scenario involving two agents, one of whom moved an object (e.g., moving a wallet from place to place) or changed the content or characteristics of an object (e.g., changing the contents of the wallet). Participants made counterfactual inferences (e.g., what if the object had not been moved?), false belief inferences (e.g., what would the absent agent expect?), and conditional inferences (e.g., what used to be in the wallet?). All three types of inference engaged the mental simulation network, confirming the general role of this network in the construction of mental representations.

During episodic counterfactual simulation the engagement of ventral medial prefrontal cortex (vmPFC) probably supports the construction of the “nearest possible world” by inferring relevant information from prior experience and background beliefs. vmPFC can also integrate the information from the constructed counterfactual event into existing representations, updating prior expectations and action-value associations, which serve to guide future behavior (see vmPFC/mOFC discussion in “Emotion and Value Processing Network” Section). This integration of counterfactual simulations with memory processes can also explain how counterfactual inference can contribute to false memories (e.g., De Brigard, 2013; Gerlach et al., 2013). Indeed, vmPFC involvement during counterfactual simulation may depend on the subject’s engagement in affective, self-reflective inference making (see also D’Argembeau et al., 2013).

We suggest that the hippocampus is more likely to be engaged during counterfactual thinking in situations requiring extensive *arbitrary relational binding* (Nieuwland, 2012; Urrutia et al., 2012; De Brigard et al., 2013; Kulakova et al., 2013; Van Hoeck et al., 2013, 2014). For example, when objects are randomly arranged on a table, the hippocampus will be more critically involved in remembering which object went in which location (location is not an inherent characteristic of an object, since objects can be moved and any could be bound to any location; therefore this arbitrary relationship needs to be encoded), than it would be involved in reconstructing the

general gestalt of the figure formed by the objects (this could be mapped to semantic knowledge for “squares” or “triangles” c.f. Watson et al., 2013). Previous work indeed indicates that the hippocampal memory network (e.g., parahippocampal gyrus, fusiform gyrus, perirhinal cortex, vmPFC) is more strongly activated when elements that were previously not related are bound together, for instance, during novel scenario simulation or the generation of new concepts (Hassabis et al., 2007; Addis et al., 2009; Hassabis and Maguire, 2009; Summerfield et al., 2009; Maguire and Hassabis, 2011; Zeithamova et al., 2012; Maguire and Mullally, 2013; Van Mulukom et al., 2013; Watson et al., 2013). During the construction of counterfactual scenarios the hippocampus will be engaged when information about the relations between elements of the scenario cannot be easily inferred from prior knowledge (Henke, 2010). The more changes we make to prior history (e.g., the remembered or experienced event) while creating the counterfactual representation the more the hippocampus is involved. The hippocampus’ role in scene-construction and relational binding also explains why constructing a future episode, and not always a counterfactual episode (which adheres to the “nearest possible world” constraint), engaged the hippocampal area more strongly than constructing a past episode (Addis and Schacter, 2012; Van Hoeck et al., 2013).

That counterfactual reasoning is strongly constrained by prior knowledge explains the increased engagement of the lateral temporal lobe (Nieuwland, 2012; Urrutia et al., 2012; Van Hoeck et al., 2013, 2014). This is often interpreted as an engagement of semantic processing, using general knowledge to construct the counterfactual scenario rather than information tied to a particular event or arbitrary configuration of elements (Addis et al., 2009; Viard et al., 2011; Irish et al., 2012; Schacter et al., 2012; De Brigard et al., 2013). Counterfactual reasoning that involves many layers of contextual abstraction (Gilead et al., 2012) may engage dorsal mPFC (dmPFC; Scott et al., 2003; Binder et al., 2009; Urrutia et al., 2012; Baetens et al., 2013; Van Hoeck et al., 2013, 2014). Goal-directed behavior, a common component of counterfactual thought, have been shown to engage inferior parietal lobe (IPL) and premotor cortex (Glover, 2004; Gazzola and Keysers, 2009; Van Overwalle and Baetens, 2009; Moody and Gennari, 2010; Stosic et al., 2014). It is worth emphasizing that because the content of counterfactual scenarios varies, so too will the engagement of different brain networks. For example, when there is a strong social component present in the situation we can expect the temporo-parietal junction to be more engaged. This in contrast to when the social component is absent and instead there is a stronger focus on a contextual and goal-directed action component (IPL). Depending upon the level of detail required, the number of relations between elements to configure, and the connection to specific or general events, different counterfactual paradigms will elicit slightly different configurations of the mental simulation network. The contribution of mental simulations to counterfactual reasoning then is in the construction and updating of an accurate representation of the scenario in question. Manipulation, reasoning, and decision making that

follow from that representation tend to engage other cognitive control regions, which we turn to now.

Cognitive Control Networks

Although it has never been the direct focus of a neuroscience study on counterfactual reasoning, we can assume that counterfactual thinking engages cognitive control and reasoning processes, and the outcomes of these processes contribute to overt and covert behavior regulation. We discuss two main control networks, the fronto-parietal network and the cingulo-opercular network, that have been engaged during counterfactual reasoning. We suggest that these networks: (1) enable switching between immediate experiences and counterfactual scenarios; (2) perform the mental transforms required for counterfactual inference and reasoning; and (3) monitor and update representations of the relative values of actions to steer subsequent behavior.

Fronto-Parietal Control Network

The fronto-parietal control network consists of the lateral PFC (lPFC), middle cingulate cortex, IPL and precuneus. This network enables the integration of multiple sources of information and supports the regulation and control of thought and behavior (e.g., Miller and Cohen, 2001; Seeley et al., 2007; Dosenbach et al., 2008; Vincent et al., 2008; Nelson et al., 2010; Spreng and Grady, 2010; Shackman et al., 2011; Barbey et al., 2012, 2013; Cabeza et al., 2012; Chein and Schneider, 2012; Schacter et al., 2012; Spreng, 2012; Whitman et al., 2013). The IPL and dorsal lateral PFC (dlPFC) play an important role in simulating and processing counterfactual information (e.g., Coricelli et al., 2005, 2007; Chua et al., 2009; Fujiwara et al., 2009; Van Hoeck et al., 2010, 2014; Nieuwland, 2012; Xue et al., 2012; De Brigard et al., 2013), but this representational role seems to be in service of executive, goal-directed functions. For example, the dlPFC (and posterior parietal cortex) was more strongly activated when processing information related to counterfactual *outcomes*, especially in situations with mixed appraisals (e.g., although you won, you could have won more; c.f. Henderson and Norris, 2013). In addition, the left dlPFC plays a central role in the manipulation of cognitive representations in the service of goal-directed behavior (Barbey et al., 2009, 2013; Ruh et al., 2012).

Knight and Grabowecy (1995; p. 1367) provide evidence from a patient with dlPFC damage who demonstrated a “complete absence of counterfactual expression”. Likewise, Gomez Beldarrain et al. (2005) observed significantly fewer spontaneous counterfactual thoughts in a group of dlPFC and orbital frontal cortex (OFC) lesion patients. However, these patients were able to construct hypothetical scenarios in response to specific cues, emphasizing the distinction between the mental simulation network and the more outcome-oriented, comparative PFC network.

Cingulo-Opercular Control Network

The cingulo-opercular network includes the dorsal anterior cingulate cortex (dACC), the posterior medial frontal cortex (pmFC), anterior insula (aI)/frontal operculum (posterior lateral OFC), and the anterior PFC (aPFC). This network is consistently

linked to maintaining task goals and monitoring actions, as well as contributing to slow behavioral adjustments over time (e.g., Seeley et al., 2007; Dosenbach et al., 2008; Power et al., 2011; Chein and Schneider, 2012). For example, Boorman et al. (2011) provide evidence that activity within this network (pmFC and aPFC) positively correlates with the reward-probability of a counterfactual choice option. Lateral aPFC activation also correlated positively with the reward-probability of the most favorable counterfactual option, and negatively with worst counterfactual option, suggesting that the lateral aPFC encodes and updates information associated with the best alternative and provides a signal for switching to an advantageous alternative (Boorman et al., 2009, 2011; Rushworth et al., 2011). These results fit well with previous findings of pmPFC involvement in adaptive behavior during tasks that entail performance monitoring, decision related uncertainty, and detection of response conflict and unfavorable outcomes (Ridderinkhof et al., 2004; Danielmeier et al., 2011). Similarly, in an episodic counterfactual simulation task, IOFC and aI activity is associated with increasing evidence of the counterfactual simulation's probability (De Brigard et al., 2013). This finding helps to explain why IOFC activation increased in a stimuli-identification task when subjects were aware that their upcoming performance would not be rewarded (there are better opportunities/trials out there) and *vice versa* (Ursu and Carter, 2005). Together these results suggest that the cingulo-opercular network critically contributes to behavioral regulation and feelings of regret by updating and maintaining counterfactual action-outcome information and signaling when it becomes profitable to change behavior (Liu et al., 2007; Nicolle et al., 2011b; Hampshire et al., 2012; Rudebeck et al., 2013).

Emotion and Value Processing Network

We have reviewed evidence to elucidate the neural mechanisms that support the construction of counterfactual scenarios and for evaluating and switching between these nearby possible realities to guide behavior toward rewarding alternatives. We now turn to the question of where these evaluative and rewarding signals originate: the networks underlying emotional and value processing traditionally associated with affective learning, valuation, reward processing, and autonomic/endocrine control (for an extensive overview see Roy et al., 2012). These networks include the vmPFC (including the medial orbitofrontal cortex, mOFC), amygdala, basal ganglia, IOFC and lPFC. We selectively review neuroscience studies that provide insight into how this network supports each of the central affect-related components of the counterfactual reasoning process (see especially the activation and adaptation-stage).

The amygdaloid complex is most frequently associated with emotional arousal and evaluative judgments (Berntson et al., 2011), especially in integrating sensory and associative information to process negative stimuli (e.g., LeDoux, 2000; Dolan, 2002; Ochsner and Gross, 2005; Phelps, 2006; Lewis et al., 2007; Kim et al., 2011; Bzdok et al., 2013; Denny et al., 2013). In addition, while the amygdala is important for reflexive or unconscious emotional processing, it is also involved in judgments of self awareness and personal responsibility (Sander

et al., 2003; Li et al., 2011; Nicolle et al., 2011b; Zalla and Sperduti, 2013). For example, Nicolle et al. (2011b) demonstrated that the amygdala responded to receiving the worse outcome from two gamble-options (counterfactual outcome was better) when these were associated with high, vs. low, personal responsibility.

Encoding the *relative value* of the obtained outcome of an event is supported by components of the basal ganglia and aI/pOFC. In a sequential decision-making study that asked subjects to invest money in a market based on its history (D'Ardenne et al., 2013), Blood Oxygen Level Dependent (BOLD) activity in two dopaminergic basal ganglia structures (the substantia nigra and the ventral tegmental area) was positively correlated with the relative values of the obtained outcome (the difference with prior expectations and with the counterfactual outcome). The BOLD response in these regions also tracked the influence of these relative values on subsequent behavior, indicating their contribution to experiential and counterfactual learning.

Further upstream in the dopaminergic pathway, striatum activity has been associated with *approach motivation*: encoding the degree of “wanting” of “liking” particular outcomes, displaying activation when relative outcome values favor an upcoming decision (i.e., no change of decision-making behavior is needed; current plan of action is favorable) and a stronger deactivation with increasing evidence against current action (i.e., change of decision making is needed; alternative/counterfactual actions need to be considered; Büchel et al., 2011; Nicolle et al., 2011a; Palminteri et al., 2012; D'Ardenne et al., 2013; Henderson and Norris, 2013; Tobia et al., 2014). aI/pOFC activity, on the other hand, is associated with sub-optimal evaluations (relative loss or threatening), contributing to feelings of regret and promoting corrective behavior (O'Doherty et al., 2003; Chua et al., 2009; Henderson and Norris, 2013). The aI/pOFC has also been implicated in encoding unanticipated avoidance of negative outcome (relief) and may thus in general contribute to assignment of *evaluative value* (Chandrasekhar et al., 2008; Berntson et al., 2011; Palminteri et al., 2012).

Patients with lesions in frontal regions demonstrate this interplay of action-value associations. Levens et al. (2014) examined pOFC and vmPFC/mOFC lesion patients' ability to generate reward expectations prior to making a decision and to assign a value to each outcome once the decision was made. Patients selected between two alternative “wheels of fortune” gambles. Each gamble had two possible outcomes (win or lose) with different reward-probabilities. This task doubly dissociated the behavior of patients with damage in the posterior IOFC from patients with vmPFC/mOFC damage. Patients with damage confined to pOFC successfully generated reward expectations and used them to calculate the expected value of each choice but were impaired in processing post-decision feedback in favor of the counterfactual choice option. By contrast, patients with vmPFC/mOFC damage selected financially worse gambles, demonstrating difficulty in computing expected value but their

post-decision affect reflected both the true and counterfactual outcomes.

These findings illustrate the distinction between assigning/learning evaluative value (undergirded by aI/pOFC) and the computation of expected values (vmPFC/mOFC). The literature broadly supports vmPFC/mOFC involvement in inferring the affective meaning of a situation by connecting conceptual information (based on personal prior experiences, beliefs, traits, and goals) with its subjective value (e.g., Coricelli et al., 2007; Van Overwalle, 2011; D'Argembeau and Salmon, 2012; Roy et al., 2012; D'Argembeau et al., 2013; Moore et al., 2014; Zaki et al., 2014). Through these value associations, vmPFC/mOFC can represent specific *expectations* about the outcome of an event and *update* these representations in light of new information (e.g., expectancy violations of counterfactual outcome values; O'Doherty et al., 2003; Alexander and Brown, 2011; Tobia et al., 2014). These findings indicate that vmPFC/mOFC abnormalities result in impairments in expectation-based regulation of emotions and behavior (Mellers et al., 1997; Sutton and Barto, 1998; Levens et al., 2014). This is generally consistent with the role of vmPFC/mOFC in the top-down modulation of emotional responses by ascribing affective meaning to the sensory information processed in the amygdaloid complex (Coricelli et al., 2007; Canessa et al., 2009; Kim et al., 2011; Zalla and Sperduti, 2013).

Taken together, these networks provide a basis for the diverse emotional and evaluative processing required during counterfactual thought.

Clinical Considerations

While these cognitive and neuropsychological studies have begun to treat counterfactual thought as an important part of human reasoning, little is known about how counterfactual thought is altered by psychiatric illness and neurological disease. We now turn to a review of the literature on counterfactual impairments and brain networks in clinical populations, including individuals with psychopathy, depression, schizophrenia and Parkinson's disease, and propose, in each case, that patients demonstrate systematic impairments in the cognitive and neural mechanisms for counterfactual inference.

Psychopathy

Psychopathic traits can be parsed into at least two interrelated components, the affective-interpersonal features (Factor 1; e.g., callousness, lack of empathy or emotional depth, and lack of genuine remorse) and the impulsive-antisocial traits (Factor 2; e.g., aggression). The former Factor one traits distinguish psychopathy from other antisocial syndromes and make people with such traits particularly pernicious (e.g., prone to repeated violence despite sanctions).

Adults with psychopathic traits display normal executive functions but decreased arousal for aversive stimuli and aberrant moral behavior (Koenigs et al., 2012; Young et al., 2012), possibly exacerbated by abnormalities in selective attention: ignoring elements outside their central focus of attention (early bottleneck; Baskin-Sommers and Newman, 2013). In addition,

Blair (1995) suggests that because individuals with psychopathic traits lack the emotional responses that lead ordinary people to imbue moral rules with genuine, authority-independent moral legitimacy, they fail to distinguish between moral and conventional rules and see all rules as mere rules. This indicates that, especially during goal-pursuit (focused attention), immoral behavior will not produce regular emotional arousal and/or will be evaluated differently. These affective and moral value related deficits have been connected with abnormal shape, size, activity and connectivity of regions of the emotion and value processing network and of the cognitive control networks (Blair, 2007; Glenn et al., 2009, 2012; Koenigs et al., 2011; Motzkin et al., 2011; Koenigs, 2012; Marazziti et al., 2013).

Within a framework of counterfactual thought, we argue that adults with psychopathic traits demonstrate deficits primarily attributable to abnormalities within the emotion and value processing network chiefly related to amygdala and vmPFC. These individuals are likely to exhibit decreased arousal for negative stimuli and abnormal selective attention (reducing the saliency of a rule violation), which will avert the activation of counterfactual inference. Individuals with psychopathic traits may assign a different affective meaning to immoral behavior, impacting the content of the counterfactuals they construe (e.g., focusing on the victim). As a result of a reduced spontaneous elicitation of counterfactual thought and different counterfactual content, they may report less remorse and self-blame and correct their immoral behavior less frequently. We predict that these deficits will exist in the presence of largely intact mental simulation and executive processes, except where those processes interact with the impaired moral or emotional processing.

Depression

Depression is a mood disorder characterized by feelings of sadness, apathy and ruminative thoughts (Nolen-Hoeksema, 2000). Rumination shares many features with counterfactual thought. Rumination can be described as a stable thought process that is repetitive, general, self-focused and oriented toward negative emotions whereby the trigger and content likely involve the discrepancy between the desired and actual state of affairs (Nolen-Hoeksema, 2000; Nolen-Hoeksema et al., 2008; Watkins, 2008; Smith and Alloy, 2009). However, it is still unclear how rumination and counterfactual reasoning are related to each other (many rumination measures include “regret”—a counterfactual emotion- as an item). There is some evidence speaking to the mood related changes occurring during counterfactual thought in individuals with depressive symptoms. A study conducted by Quelhas et al. (2008; Experiment 2) demonstrated that, immediately after receiving a negative grade on an academic test, college students with moderate symptoms of depression (measured by Beck Depression Inventory [BDI]; $M_{BDI-II} = 18.95$) generated a smaller number of *spontaneous* counterfactual thoughts relative to controls ($M_{BDI-II} = 3.08$). This difference was eliminated by an *explicit cue* to generate upward counterfactuals regarding a recent negative academic experience (Markman and Miller, 2006). In addition, upward counterfactuals generated by individuals with symptoms of depression and controls, had similar levels of focus on

controlling the situation by changing one’s own behavior. What differed was the feasibility of counterfactuals generated by the depression group: their alternative scenarios included less plausible actions and/or actions less likely to change the outcome. Depression *severity* also influenced number of counterfactuals focused on chronic or enduring aspects of the self (which are unlikely to change or be under immediate control). In addition, consistent with the earlier statement that processing upward counterfactuals with low feasibility only contributes to psychological distress but does not result in beneficial reappraisal effects, the post-counterfactual mood of depression groups was more negative than controls. Moreover, subjects with mild-to-moderate symptoms (BDI-II cutoff: 10–23) did not perceive an increase in their ability to exert control over the environment (see also Quelhas et al., 2008) and in the severe group (BDI-II cutoff: 24+) the perception of control decreased and re-evaluation of the event produced a more negative assessment than before the counterfactual thinking.

The altered spontaneous counterfactual inference pattern in depression might be related to other irregularities associated with depression, including deficits in removing negative stimuli from short-term memory (Joormann and Gotlib, 2010) and decreased attentional control when confronted with self-relevant negative information (De Raedt et al., 2010; Koster et al., 2011; Beckwé et al., 2013), which might prevent them from disengaging from their immediate emotional experience and ruminative thought and reallocating resources and attention to the (next) counterfactual inference process of how the event could have unfolded differently (Dolcos et al., 2008; Iordan et al., 2013). In addition, their rumination tendencies and overgeneral autobiographical memory may also contribute to generating counterfactuals that focus stronger on enduring traits than on specific behavior (Williams, 2006; Williams et al., 2007; Watkins, 2008; Sumner, 2012).

Depression may also impact processing of counterfactual information when the experimenter provides this information, rather than it being a product of an individual’s own counterfactual construction. Subjects with mild symptoms of depression ($M_{BDI} = 14.66$) reported more regret about a *hypothetical* hiring decision than controls when they were informed about another, better applicant (Monroe et al., 2005). This increase in regret was not driven by the absolute difference in applicant quality or by the individual’s initial confidence in the hiring decision. Howlett and Paulus (2013) attributed this increased experience of regret in subjects with mild-to-moderate symptoms to: (1) an increased vigilance for potential threats to self-esteem, indicating that negative stimuli are associated with a high self-relevancy and thus increase their affective meaning; (2) a stronger decrease of the relative value of the obtained outcome; and (3) increased damage to self-esteem when considering a better counterfactual alternative.

However, major depression disorder displays a different emotional profile, wherein sensitivity to relative losses (e.g., one could have won more) is muted and this response is related to self-reported apathy-scores (Chase et al., 2010). This might reflect changes in approach motivation and goal-processing (Shankman et al., 2007; Eddington et al., 2009), and suggests

that this disengagement might be related to coping mechanisms developed to protect their unstable self-esteem (Roese et al., 2009; Howlett and Paulus, 2013). This pattern of results might also be linked to the finding that individuals' with severe symptoms of depression, when cued to generate counterfactual scenarios, tend to construe less feasible, non-beneficial upward counterfactuals (focused on chronic or enduring aspects of the self instead of actions that would likely change the outcome), and experience afterwards a more negative mood than controls (Markman and Miller, 2006). In this case, disengaging from counterfactual reasoning may prevent increased psychological distress (Wrosch and Miller, 2009; Brassens et al., 2012; Wrosch et al., 2013), but may also prevent improvement from status-quo (see Tykocinski and Steinberg, 2005).

The literature on the neurobiology of depression is complex, with differences related to severity, time of onset, and chronicity. Changes are most consistently observed in emotion and value processing regions, such as amygdala, vmPFC, striatum and OFC (Johnstone et al., 2007; Stein et al., 2007; Eddington et al., 2009; Johnson et al., 2009; Townsend et al., 2010; Murray et al., 2011; Price and Drevets, 2012; Howlett and Paulus, 2013; Cole et al., 2014). Depression has also been associated with increased connectivity between and alterations within the aforementioned mental simulation network (e.g., Zhang et al., 2011). For example, severity of depression correlates positively with activity in ventromedial polar cortex and amygdala, possibly representing an increased vigilance for potential threats to self-esteem (Drevets, 2007; Murray et al., 2011; Price and Drevets, 2012). On the other hand, activity within the posterior OFC, a region we identified as playing a key role in processing counterfactual action-outcome information, decreases with severity. This finding is largely consistent with data from posterior OFC lesions patients indicating that damage to this region influences expectations about their post-decision affect but not for the counterfactual outcome (Levens et al., 2014).

In summary, depression symptoms seem to interfere with all three aspects of counterfactual thought: mental simulation, executive control and valuation. This interference is complex, and interacts with severity, chronicity, and onset of time of depression. Disruption is especially apparent for negative and self-relevant judgments, such as disengaging from self-relevant negative experiences. However, looking at depression symptoms through a lens of counterfactual thinking helps to identify disruptions that would not be immediately apparent from an evaluation of depression as a mood disorder alone. For example, subjects' generation of less feasible, likely, or detached counterfactuals is highly characteristic of depression but not closely linked to mood. Counterfactual thought can therefore provide a useful technique for understanding depressive pathology and symptoms.

Schizophrenia and Parkinson's Disease

Schizophrenia and Parkinson disease patients both exhibit primary deficits in executive functions and social cognition, stemming from disruptions within dopaminergic pathways in the brain. We discuss similarities in the impairments of counterfactual thought exhibited by these clinical groups.

Schizophrenia, a psychopathology with significant neurological dysfunction and symptoms including auditory hallucinations, paranoia, and other delusional thoughts, is also known to impair counterfactual thinking. Hooker et al. (2000) had 14 residential Schizophrenia patients (11 males and 3 females taking antipsychotic medication; with no history of substance abuse, head injury or co-morbid Axis one disorder) recall a negative personal event from the past year. They were explicitly asked if, as they recalled the past, they had any thoughts about how this event could have turned out differently. In comparison to healthy controls, schizophrenia patients mentioned fewer different counterfactual thoughts (see also Caño et al., 2011). Their performance could not be explained by a generalized cognitive deficit: in comparison to the control subjects they did not differ on Wechsler Adult Intelligence Scale (WAIS-R) measure of Vocabulary and Digit Span, or on the FAS Verbal Fluency test (verbal production). Key components for counterfactual simulations are broadly impaired by Schizophrenia. Research on episodic simulation indicate that these patients are impaired in the coherent recall of episodic events, inferring self-relevant meaning, and (re)construction of episodic future scenarios (D'Argembeau et al., 2008; Raffard et al., 2010a,b; Bennouna-Greene et al., 2012).

In addition, deficits in cognitive control may aggravate these problems (Cole et al., 2014). Savla et al. (2012) examined the performance of patients on measures of cognitive flexibility and abstraction (Silver and Bilker, 2013). Decreased cognitive flexibility was related to increase of positive symptoms associated with schizophrenia. This factor also includes failures of inhibition and control of attention (Clark et al., 2010; Goldberg et al., 2011; Waters et al., 2012). Furthermore, abstraction correlated negatively with the duration of illness and positively with everyday functioning. These results suggest broad impairments in counterfactual thought involving both failures to simulate and regulate counterfactual scenarios.

Consistent with this pattern of results, Schizophrenia patients display impairments the Counterfactual Inference Test (CIT; Hooker et al., 2000; Larquet et al., 2010). The CIT assesses social cognition across four vignettes, each of which describes a negative event in which two protagonists differ in their proximity to a better outcome or the normality of the action preceding the negative outcome (routine vs. uncommon action). To illustrate, consider the following example. "Janet is attacked by a mugger only 10 feet from her house. Susan is attacked only a mile from her house. Who is more upset by the mugging? (a) Janet (b) Susan (c) Same/Cant' tell" (Hooker et al., 2000, p. 330). Subjects were asked to judge whom of the two protagonists felt worse after experiencing this event, regrets their action the most, or thinks most (counterfactually) about the event. Schizophrenia patients were near chance in selecting appropriate CIT answers. Their performance could not be explained by a generalized cognitive deficit (WAIS-R: Vocabulary and Digit Span; FAS Verbal Fluency test). However, performance on CIT did mediate the group difference in social competence (measured by the Zigler scale). In addition, patients showed decreased activity in brain networks associated with reasoning about the cognitive states of others (i.e., Theory of Mind; temporo-parietal junction,

mPFC and lower portion of the precuneus; Amodio and Frith, 2006; Van Overwalle and Baetens, 2009; Mar, 2011; Schilbach et al., 2012) during completion of false belief tasks (which require inferring others' representations of reality, which does not correspond with their own perception reality; Lee et al., 2011). The observed pattern of findings suggest that Theory of Mind is not in general impaired in Schizophrenia, but that it instead reflects deficits in reasoning when complex non-factual elements are needed to understand the social environment (Kern et al., 2009; Sparks et al., 2010). Both counterfactual and false belief inference entail simulation of an alternative state of affairs different from the perceived reality. Van Hoeck et al. (2014) further confirmed that false belief and counterfactual inference about social hypothetical scenarios activate the mental simulation and frontoparietal networks (see also Canessa et al., 2009). These authors further suggest that additive counterfactual inference may reflect a more complex process that places additional demands on cognitive control (increased activation in dlPFC and pmPFC).

An abundance of neuroscience studies confirm abnormalities within the simulation and cognitive control networks, especially those regions supporting contextual coherence, relational processing and flexible cognitive control. For example, structural, functional, and neurochemical hippocampus abnormalities have been associated with Schizophrenia, therefore contributing to observed problems in scene-construction and relational processing during episodic simulation (Ongür et al., 2006; Boyer et al., 2007; Williams et al., 2010; Adriano et al., 2012; Herold et al., 2013). dmPFC, a region involved in contextual associations/abstractions and contributing to the coherency of the mental representation, shows altered interactions with vmPFC and posterior midline structures in Schizophrenia (Raffard et al., 2009; Shad et al., 2011; Alonso-Solis et al., 2012), and also decreased anticorrelation with the dlPFC, a frontoparietal control component (Whitfield-Gabrieli et al., 2009).

Similar disruptions are found in Parkinson patients. Parkinson is a progressive neurodegenerative disorder associated with substantial decrease in dopamine levels in the basal ganglia, frontal lobes and hippocampus. As in the case of Schizophrenia, Parkinson patients demonstrate Theory of Mind dysfunction (especially with respect to the cognitive component; Yu and Wu, 2013), impaired cognitive flexibility, selective attention and concept formation (Kudlicka et al., 2011). Parkinson patients, in comparison to age-matched controls, also generate a smaller number of upward episodic counterfactual thoughts and demonstrate decreased performance on the CIT (at chance level; McNamara et al., 2003). Their counterfactual performance could not be explained by a general cognition deficit, but did correlate with social function and cognitive

flexibility (as assessed by Stroop color inference task: inhibition, resistance to cognitive interference; Tower of London task: ill-structured problem space requiring planning and working memory).

Thus, evidence from psychology and neuroscience demonstrate impairments in the mental simulation and executive function components of counterfactual thought in Schizophrenia and Parkinson disease. Both patient groups demonstrate impairments in cognitive flexibility, contextual abstraction and relational binding. These deficits are especially apparent within complex, ill-structured environments, and in the context of goal-directed behavior and adaptive decision making. Since all of these functions are required in counterfactual thought, tests of counterfactual thinking may, once again, serve as a valuable diagnostic tool.

Conclusion

Counterfactual reasoning entails three processing stages: Activation, Inference and Adaptation. Counterfactual scenarios are automatically constructed in response to everyday events and are linked to violations of prior beliefs and motivations, adhering to a "nearest possible world" constraint. These mental simulations influence our emotional responses and future behavior, by integrating information about the counterfactual state of affairs into the representation of the current situation, prior beliefs, and expectations thereby allowing one to learn through counterfactual inference. This complex psychological phenomena of counterfactual reasoning is made possible by the coordinated interaction between three networks: (1) mental simulation in MTL/PFC; (2) cognitive control in the frontoparietal and cingulo-opercular network; and (3) motivation and valuation in limbic regions and vmPFC.

Because counterfactual reasoning depends upon so many inter-related brain systems and mental processes, it provides a broad window onto global brain function. Disruptions to counterfactual reasoning are a productive scientific and diagnostic tool. By identifying the specific sub-components of counterfactual reasoning that are affected clinicians can better understand the day-to-day challenges faced by different patient populations. In our everyday lives, counterfactual reasoning is a ubiquitous source of new insights into nearby possible worlds, and different circumstances, scientific understanding of this process is a critical component of insight into the complex alternate worlds that make up our rich, inner lives.

Acknowledgment

NVH was funded by the Vrije Universiteit Brussel, project SRP15.

References

- Addis, D. R., Pan, L., Vu, M.-A., Laiser, N., and Schacter, D. L. (2009). Constructive episodic simulation of the future and the past: distinct subsystems of a core brain network mediate imagining and remembering. *Neuropsychologia* 47, 2222–2238. doi: 10.1016/j.neuropsychologia.2008.10.026
- Addis, D. R., and Schacter, D. L. (2012). The hippocampus and imagining the future: where do we stand? *Front. Hum. Neurosci.* 5:173. doi: 10.3389/fnhum.2011.00173
- Adriano, F., Caltagirone, C., and Spalletta, G. (2012). Hippocampal volume reduction in first-episode and chronic schizophrenia: a review and meta-analysis. *Neuroscientist* 18, 180–200. doi: 10.1177/1073858410395147

- Alexander, W. H., and Brown, J. W. (2011). Medial prefrontal cortex as an action-outcome predictor. *Nature Neuroscience* 14, 1338–1344. doi: 10.1038/nn.2921
- Alicke, M. D., Buckingham, J., Zell, E., and Davis, T. (2008). Culpable control and counterfactual reasoning in the psychology of blame. *Pers. Soc. Psychol. Bull.* 34, 1371–1381. doi: 10.1177/0146167208321594
- Alonso-Solís, A., Corripio, I., de Castro-Manglano, P., Duran-Sindreu, S., Garcia-Garcia, M., Proal, E., et al. (2012). Altered default network resting state functional connectivity in patients with a first episode of psychosis. *Schizophr. Res.* 139, 13–18. doi: 10.1016/j.schres.2012.05.005
- Amodio, D. M., and Frith, C. D. (2006). Meeting of minds: the medial frontal cortex and social cognition. *Nat. Rev. Neurosci.* 7, 268–277. doi: 10.1038/nrn1884
- Baetens, K., Ma, N., Steen, J., and Van Overwalle, F. (2013). Involvement of the mentalizing network in social and non-social high construal. *Soc. Cogn. Affect. Neurosci.* 9, 817–824. doi: 10.1093/scan/nst048
- Barbey, A. K., Colom, R., and Grafman, J. (2013). Dorsolateral prefrontal contributions to human intelligence. *Neuropsychologia* 51, 1361–1369. doi: 10.1016/j.neuropsychologia.2012.05.017
- Barbey, A. K., Colom, R., Solomon, J., Krueger, F., Forbes, C., and Grafman, J. (2012). An integrative architecture for general intelligence and executive function revealed by lesion mapping. *Brain* 135, 1154–1164. doi: 10.1093/brain/aws021
- Barbey, A. K., Krueger, F., and Grafman, J. (2009). Structured event complexes in the medial prefrontal cortex support counterfactual representations for future planning. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 1291–1300. doi: 10.1098/rstb.2008.0315
- Barbey, A. K., and Sloman, S. A. (2007). Base-rate respect: from ecological rationality to dual processes. *Behav. Brain Sci.* 30, 241–254; discussion 255–297. doi: 10.1017/s0140525x07001653
- Barrett, L. F., and Satpute, A. B. (2013). Large-scale brain networks in affective and social neuroscience: towards an integrative functional architecture of the brain. *Curr. Opin. Neurobiol.* 23, 361–372. doi: 10.1016/j.conb.2012.12.012
- Barsalou, L. W., Kyle Simmons, W., Barbey, A. K., and Wilson, C. D. (2003a). Grounding conceptual knowledge in modality-specific systems. *Trends Cogn. Sci.* 7, 84–91. doi: 10.1016/s1364-6613(02)00029-3
- Barsalou, L. W., Niedenthal, P. M., Barbey, A. K., and Ruppert, J. (2003b). “Social embodiment,” in *The Psychology of Learning and Motivation*, ed. B. Ross (San Diego, CA: Academic Press), 43–92.
- Baskin-Sommers, A. R., and Newman, J. P. (2013). “Differentiating the cognition-emotion interactions that characterize psychopathy vs. externalizing disorders,” in *Handbook of Cognition and Emotion*, eds M. D. Robinson, E. R. Watkins, and E. Harmon-Jones (New York, NY: Guilford Press), 501–520.
- Beckwé, M., Deroost, N., Koster, E. H. W., De Lissnyder, E., and De Raedt, R. (2013). Worrying and rumination are both associated with reduced cognitive control. *Psychol. Res.* 78, 651–660. doi: 10.1007/s00426-013-0517-5
- Bennouna-Greene, M., Berna, F., Conway, M. A., Rathbone, C. J., Vidailhet, P., and Danion, J.-M. (2012). Self-images and related autobiographical memories in schizophrenia. *Conscious. Cogn.* 21, 247–257. doi: 10.1016/j.concog.2011.10.006
- Berntson, G. G., Norman, G. J., Bechara, A., Bruss, J., Tranel, D., and Cacioppo, J. T. (2011). The insula and evaluative processes. *Psychol. Sci.* 22, 80–86. doi: 10.1177/0956797610391097
- Binder, J. R., Desai, R. H., Graves, W. W., and Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cereb. Cortex* 19, 2767–2796. doi: 10.1093/cercor/bhp055
- Blair, R. (1995). A cognitive developmental approach to morality: investigating the psychopath. *Cognition* 57, 1–29. doi: 10.1016/0010-0277(95)00676-p
- Blair, R. J. R. (2007). The amygdala and ventromedial prefrontal cortex in morality and psychopathy. *Trends Cogn. Sci.* 11, 387–392. doi: 10.1016/j.tics.2007.07.003
- Boorman, E. D., Behrens, T. E. J., and Rushworth, M. F. S. (2011). Counterfactual choice and learning in a neural network centered on human lateral frontopolar cortex. *PLoS Biol.* 9:e1001093. doi: 10.1371/journal.pbio.1001093
- Boorman, E. D., Behrens, T. E. J., Woolrich, M. W., and Rushworth, M. F. S. (2009). How green is the grass on the other side? Frontopolar cortex and the evidence in favor of alternative courses of action. *Neuron* 62, 733–743. doi: 10.1016/j.neuron.2009.05.014
- Boyer, P., Phillips, J. L., Rousseau, F. L., and Ilivitsky, S. (2007). Hippocampal abnormalities and memory deficits: new evidence of a strong pathophysiological link in schizophrenia. *Brain Res. Rev.* 54, 92–112. doi: 10.1016/j.brainresrev.2006.12.008
- Branscombe, N. R., Wohl, M. J. A., Owen, S., Allison, J. A., and N’gbala, A. (2003). Counterfactual thinking, blame assignment and well-being in rape victims. *Basic Appl. Soc. Psych.* 25, 265–273. doi: 10.1207/s15324834basps2504_1
- Brassen, S., Gamer, M., Peters, J., Gluth, S., and Büchel, C. (2012). Don’t look back in anger! Responsiveness to missed chances in successful and unsuccessful aging. *Science* 336, 612–614. doi: 10.1126/science.1217516
- Büchel, C., Brassen, S., Yacubian, J., Kalisch, R., and Sommer, T. (2011). Ventral striatal signal changes represent missed opportunities and predict future choice. *Neuroimage* 57, 1124–1130. doi: 10.1016/j.neuroimage.2011.05.031
- Buckner, R. L., and Carroll, D. C. (2007). Self-projection and the brain. *Trends Cogn. Sci.* 11, 49–57. doi: 10.1016/j.tics.2006.11.004
- Byrne, R. M. J. (1997). Cognitive processes in counterfactual thinking about what might have been. *Psychol. Learn. Motiv.* 37, 105–154. doi: 10.1016/s0079-7421(08)60501-0
- Byrne, R. M. J. (2002). Mental models and counterfactual thought about what might have been. *Trends Cogn. Sci.* 6, 426–431. doi: 10.1016/s1364-6613(02)01974-5
- Byrne, R. M. J. (2007). Précis of the rational imagination: how people create alternatives to reality. *Behav. Brain Sci.* 30, 439–453; discussion 453–476. doi: 10.1017/S0140525X07002579
- Bzdok, D., Laird, A. R., Zilles, K., Fox, P. T., and Eickhoff, S. B. (2013). An investigation of the structural, connectional and functional subspecialization in the human amygdala. *Hum. Brain Mapp.* 34, 3247–3266. doi: 10.1002/hbm.22138
- Cabeza, R., Ciaramelli, E., and Moscovitch, M. (2012). Cognitive contributions of the ventral parietal cortex: an integrative theoretical account. *Trends Cogn. Sci.* 16, 338–352. doi: 10.1016/j.tics.2012.04.008
- Canessa, N., Motterlini, M., Di Dio, C., Perani, D., Scifo, P., Cappa, S. F., et al. (2009). Understanding others’ regret: a fMRI study. *PLoS One* 4:e7402. doi: 10.1371/journal.pone.0007402
- Caño, A., Contreras, F., and Menchón, J. M. (2011). P01–410—counterfactual thinking in schizophrenia patients. *Eur. Psychiatry* 26:413. doi: 10.1016/s0924-9338(11)72121-6
- Chandrasekhar, P. V. S., Capra, C. M., Moore, S., Noussair, C., and Berns, G. S. (2008). Neurobiological regret and rejoice functions for aversive outcomes. *Neuroimage* 39, 1472–1484. doi: 10.1016/j.neuroimage.2007.10.027
- Chase, H. W., Camille, N., Michael, A., Bullmore, E. T., Robbins, T. W., and Sahakian, B. J. (2010). Regret and the negative evaluation of decision outcomes in major depression. *Cogn. Affect. Behav. Neurosci.* 10, 406–413. doi: 10.3758/CABN.10.3.406
- Chein, J. M., and Schneider, W. (2012). The brain’s learning and control architecture. *Curr. Dir. Psychol. Sci.* 21, 78–84. doi: 10.1177/0963721411434977
- Chua, H. F., Gonzalez, R., Taylor, S. F., Welsh, R. C., and Liberzon, I. (2009). Decision-related loss: regret and disappointment. *Neuroimage* 47, 2031–2040. doi: 10.1016/j.neuroimage.2009.06.006
- Clark, L. K., Warman, D., and Lysaker, P. H. (2010). The relationships between schizophrenia symptom dimensions and executive functioning components. *Schizophr. Res.* 124, 169–175. doi: 10.1016/j.schres.2010.08.004
- Cole, M. W., Repovš, G., and Anticevic, A. (2014). The frontoparietal control system: a central role in mental health. *Neuroscientist* 20, 652–664. doi: 10.1177/1073858414525995
- Coricelli, G., Critchley, H. D., Joffily, M., O’Doherty, J. P., Sirigu, A., and Dolan, R. J. (2005). Regret and its avoidance: a neuroimaging study of choice behavior. *Nat. Neurosci.* 8, 1255–1262. doi: 10.1038/nn1514
- Coricelli, G., Dolan, R. J., and Sirigu, A. (2007). Brain, emotion and decision making: the paradigmatic example of regret. *Trends Cogn. Sci.* 11, 258–265. doi: 10.1016/j.tics.2007.04.003
- Coricelli, G., and Rustichini, A. (2010). Counterfactual thinking and emotions: regret and envy learning. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365, 241–247. doi: 10.1098/rstb.2009.0159
- Damasio, A. R. (1989). Time-locked multiregional retroactivation: a systems-level proposal for the neural substrates of recall and recognition. *Cognition* 33, 25–62. doi: 10.1016/0010-0277(89)90005-x

- Danielmeier, C., Eichele, T., Forstmann, B. U., Tittgemeyer, M., and Ullsperger, M. (2011). Posterior medial frontal cortex activity predicts post-error adaptations in task-related visual and motor areas. *J. Neurosci.* 31, 1780–1789. doi: 10.1523/JNEUROSCI.4299-10.2011
- D'Ardenne, K., Lohrenz, T., Bartley, K. A., and Montague, P. R. (2013). Computational heterogeneity in the human mesencephalic dopamine system. *Cogn. Affect. Behav. Neurosci.* 13, 747–756. doi: 10.3758/s13415-013-0191-5
- D'Argembeau, A., Cassol, H., Phillips, C., Balteau, E., Salmon, E., and Van der Linden, M. (2013). Brains creating stories of selves: the neural basis of autobiographical reasoning. *Soc. Cogn. Affect. Neurosci.* 9, 646–652. doi: 10.1093/scan/nst028
- D'Argembeau, A., Raffard, S., and Van der Linden, M. (2008). Remembering the past and imagining the future in schizophrenia. *J. Abnorm. Psychol.* 117, 247–251. doi: 10.1037/0021-843x.117.1.247
- D'Argembeau, A., and Salmon, E. (2012). The neural basis of semantic and episodic forms of self-knowledge: insights from functional neuroimaging. *Adv. Exp. Med. Biol.* 739, 276–290. doi: 10.1007/978-1-4614-1704-0_18
- Davis, C. G., Lehman, D. R., Wortman, C. B., Silver, R. C., and Thompson, S. C. (1995). The undoing of traumatic life events. *Pers. Soc. Psychol. Bull.* 21, 109–124. doi: 10.1177/0146167295212002
- Davis, C. G., Nolen-Hoeksema, S., and Larson, J. (1998). Making sense of loss and benefiting from the experience: two construals of meaning. *J. Pers. Soc. Psychol.* 75, 561–574. doi: 10.1037/0022-3514.75.2.561
- De Brigard, F. (2013). Is memory for remembering? Recollection as a form of episodic hypothetical thinking. *Synthese* 191, 155–185. doi: 10.1007/s11229-013-0247-7
- De Brigard, F., Addis, D. R., Ford, J. H., Schacter, D. L., and Giovanello, K. S. (2013). Remembering what could have happened: neural correlates of episodic counterfactual thinking. *Neuropsychologia* 51, 2401–2414. doi: 10.1016/j.neuropsychologia.2013.01.015
- De Brigard, F., and Giovanello, K. S. (2012). Influence of outcome valence in the subjective experience of episodic past, future and counterfactual thinking. *Conscious. Cogn.* 21, 1085–1096. doi: 10.1016/j.concog.2012.06.007
- Dehghani, M., Iliev, R., and Kaufmann, S. (2012). Causal explanation and fact mutability in counterfactual reasoning. *Mind Lang.* 27, 55–85. doi: 10.1111/j.1468-0017.2011.01435.x
- Denny, B. T., Ochsner, K. N., Weber, J., and Wager, T. D. (2013). Anticipatory brain activity predicts the success or failure of subsequent emotion regulation. *Soc. Cogn. Affect. Neurosci.* 9, 403–411. doi: 10.1093/scan/nss148
- De Raedt, R., Koster, E. H. W., and Joormann, J. (2010). Attentional control in depression: a translational affective neuroscience approach. *Cogn. Affect. Behav. Neurosci.* 10, 1–7. doi: 10.3758/CABN.10.1.1
- Dolan, R. J. (2002). Emotion, cognition and behavior. *Science* 298, 1191–1194. doi: 10.1126/science.1076358
- Dolcos, F., Diaz-Granados, P., Wang, L., and McCarthy, G. (2008). Opposing influences of emotional and non-emotional distracters upon sustained prefrontal cortex activity during a delayed-response working memory task. *Neuropsychologia* 46, 326–335. doi: 10.1016/j.neuropsychologia.2007.07.010
- Dosenbach, N. U. F., Fair, D. A., Cohen, A. L., Schlaggar, B. L., and Petersen, S. E. (2008). A dual-networks architecture of top-down control. *Trends Cogn. Sci.* 12, 99–105. doi: 10.1016/j.tics.2008.01.001
- Drevets, W. C. (2007). Orbitofrontal cortex function and structure in depression. *Ann. N. Y. Acad. Sci.* 1121, 499–527. doi: 10.1196/annals.1401.029
- Dyczewski, E. A., and Markman, K. D. (2012). General attainability beliefs moderate the motivational effects of counterfactual thinking. *J. Exp. Soc. Psychol.* 48, 1217–1220. doi: 10.1016/j.jesp.2012.04.016
- Eddington, K. M., Dolcos, F., McLean, A. N., Krishnan, K. R., Cabeza, R., and Strueman, T. J. (2009). Neural correlates of idiographic goal priming in depression: goal-specific dysfunctions in the orbitofrontal cortex. *Soc. Cogn. Affect. Neurosci.* 4, 238–246. doi: 10.1093/scan/nsp016
- El Leithy, S., Brown, G. P., and Robbins, I. (2006). Counterfactual thinking and posttraumatic stress reactions. *J. Abnorm. Psychol.* 115, 629–635. doi: 10.1037/0021-843x.115.3.629
- Epstude, K., and Roese, N. J. (2008). The functional theory of counterfactual thinking. *Pers. Soc. Psychol. Rev.* 12, 168–192. doi: 10.1177/1088868308316091
- Ferrante, D., Giroto, V., Stragà, M., and Walsh, C. (2013). Improving the past and the future: a temporal asymmetry in hypothetical thinking. *J. Exp. Psychol. Gen.* 142, 23–27. doi: 10.1037/a0027947
- Fujiwara, J., Tobler, P. N., Taira, M., Iijima, T., and Tsutsui, K.-I. (2009). A parametric relief signal in human ventrolateral prefrontal cortex. *NeuroImage* 44, 1163–1170. doi: 10.1016/j.neuroimage.2008.09.050
- Gazzola, V., and Keysers, C. (2009). The observation and execution of actions share motor and somatosensory voxels in all tested subjects: single-subject analyses of unsmoothed fMRI data. *Cereb. Cortex* 19, 1239–1255. doi: 10.1093/cercor/bhn181
- Gerlach, K. D., Dornblaser, D. W., and Schacter, D. L. (2013). Adaptive constructive processes and memory accuracy: consequences of counterfactual simulations in young and older adults. *Memory* 22, 145–162. doi: 10.1080/09658211.2013.779381
- Gilbar, O., and Hevroni, A. (2007). Counterfactuals, coping strategies and psychological distress among breast cancer patients. *Anxiety Stress Coping* 20, 383–392. doi: 10.1080/10615800701384439
- Gilead, M., Liberman, N., and Maril, A. (2012). Construing counterfactual worlds: the role of abstraction. *Eur. J. Soc. Psychol.* 42, 391–397. doi: 10.1002/ejsp.1862
- Glenn, A. L., Raine, A., Schug, R. A., Young, L., and Hauser, M. (2009). Increased DLPFC activity during moral decision-making in psychopathy. *Mol. Psychiatry* 14, 909–911. doi: 10.1038/mp.2009.76
- Glenn, A. L., Yang, Y., and Raine, A. (2012). “Neuroimaging in psychopathy and antisocial personality disorder: functional hypothesis,” in *Neuroimaging in Forensic Psychiatry: From the Clinic to the Courtroom*, ed. J. R. Simpson (Chichester, UK: John Wiley and Sons, Ltd.), 81–98.
- Glover, S. (2004). Separate visual representations in the planning and control of action. *Behav. Brain Sci.* 27, 3–78. doi: 10.1017/s0140525x04000020
- Goldberg, T. E., David, A., and Gold, J. M. (2011). “Neurocognitive impairments in schizophrenia: their character and role in symptom formation,” in *Schizophrenia*, ed. D. R. Weinberger and P. J. Harrison 3rd Edn. (Oxford, UK: Wiley-Blackwell), 142–162.
- Gomez Beldarrain, M., Garcia-Monco, J. C., Astigarraga, E., Gonzalez, A., and Grafman, J. (2005). Only spontaneous counterfactual thinking is impaired in patients with prefrontal cortex lesions. *Brain Res. Cogn. Brain Res.* 24, 723–726. doi: 10.1016/j.cogbrainres.2005.03.013
- Hampshire, A., Chudhry, A. M., Owen, A. M., and Roberts, A. C. (2012). Dissociable roles for lateral orbitofrontal cortex and lateral prefrontal cortex during preference driven reversal learning. *NeuroImage* 59, 4102–4112. doi: 10.1016/j.neuroimage.2011.10.072
- Hassabis, D., Kumaran, D., Vann, S. D., and Maguire, E. A. (2007). Patients with hippocampal amnesia cannot imagine new experiences. *Proc. Natl. Acad. Sci. U S A* 104, 1726–1731. doi: 10.1073/pnas.0610561104
- Hassabis, D., and Maguire, E. A. (2009). The construction system of the brain. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 1263–1271. doi: 10.1098/rstb.2008.0296
- Henderson, S. E., and Norris, C. J. (2013). Counterfactual thinking and reward processing: an fMRI study of responses to gamble outcomes. *NeuroImage* 64, 582–589. doi: 10.1016/j.neuroimage.2012.08.078
- Hendrickson, N. (2010). Counterfactual reasoning and the problem of selecting antecedent scenarios. *Synthese* 185, 365–386. doi: 10.1007/s11229-010-9824-1
- Henke, K. (2010). A model for memory systems based on processing modes rather than consciousness. *Nat. Rev. Neurosci.* 11, 523–532. doi: 10.1038/nrn2850
- Herold, C. J., Lässer, M. M., Schmid, L. A., Seidl, U., Kong, L., Fellhauer, I., et al. (2013). Hippocampal volume reduction and autobiographical memory deficits in chronic schizophrenia. *Psychiatry Res.* 211, 189–194. doi: 10.1016/j.psychres.2012.04.002
- Hooker, C., Roese, N. J., and Park, S. (2000). Impoverished counterfactual thinking is associated with schizophrenia. *Psychiatry* 63, 326–335.
- Howlett, J. R., and Paulus, M. P. (2013). Decision-making dysfunctions of counterfactuals in depression: who might I have been? *Front. Psychiatry* 4:143. doi: 10.3389/fpsy.2013.00143
- Iordan, A. D., Dolcos, S., and Dolcos, F. (2013). Neural signatures of the response to emotional distraction: a review of evidence from brain imaging investigations. *Front. Hum. Neurosci.* 7:200. doi: 10.3389/fnhum.2013.00200
- Irish, M., Addis, D. R., Hodges, J. R., and Piguet, O. (2012). Considering the role of semantic memory in episodic future thinking: evidence from semantic dementia. *Brain* 135, 2178–2191. doi: 10.1093/brain/aws119

- Johnson, M. K., Nolen-Hoeksema, S., Mitchell, K. J., and Levin, Y. (2009). Medial cortex activity, self-reflection and depression. *Soc. Cogn. Affect. Neurosci.* 4, 313–327. doi: 10.1093/scan/nsp022
- Johnson, M. K., and Sherman, S. J. (1999). “Constructing and reconstructing the past and the future in the present.” Vol. 2, *Handbook of Motivation and Cognition: Foundation of Social Behavior*, eds E. T. Higgins and R. M. Sorrentino (New York: The Guilford Press), 482–526.
- Johnstone, T., van Reekum, C. M., Urry, H. L., Kalin, N. H., and Davidson, R. J. (2007). Failure to regulate: counterproductive recruitment of top-down prefrontal-subcortical circuitry in major depression. *J. Neurosci.* 27, 8877–8884. doi: 10.1523/jneurosci.2063-07.2007
- Joormann, J., and Gotlib, I. H. (2010). Emotion regulation in depression: relation to cognitive inhibition. *Cogn. Emot.* 24, 281–298. doi: 10.1080/02699930903407948
- Kahneman, D., and Miller, D. T. (1986). Norm theory: comparing reality to its alternatives. *Psychol. Rev.* 93, 136–153. doi: 10.1037/0033-295x.93.2.136
- Kahneman, D., and Tversky, A. (1982). “The simulation heuristic,” in *Judgement Under Uncertainty: Heuristics and Biases*, eds D. Kahneman, E. Slovic, and A. Tversky (New York: Cambridge University Press), 201–208.
- Kern, R. S., Green, M. F., Fiske, A. P., Kee, K. S., Lee, J., Sergi, M. J., et al. (2009). Theory of mind deficits for processing counterfactual information in persons with chronic schizophrenia. *Psychol. Med.* 39, 645–654. doi: 10.1017/s0033291708003966
- Kim, M. J., Loucks, R. A., Palmer, A. L., Brown, A. C., Solomon, K. M., Marchante, A. N., et al. (2011). The structural and functional connectivity of the amygdala: from normal emotion to pathological anxiety. *Behav. Brain Res.* 223, 403–410. doi: 10.1016/j.bbr.2011.04.025
- Knight, R. T., and Grabowecky, M. (1995). “Escape from linear time: prefrontal cortex and conscious experience,” in *The Cognitive Neuroscience*, ed. M. S. Gazzaniga (Cambridge, MA: MIT Press), 1357–1371.
- Koenigs, M. (2012). The role of prefrontal cortex in psychopathy. *Rev. Neurosci.* 23, 253–262. doi: 10.1515/revneuro-2012-0036
- Koenigs, M., Baskin-Sommers, A., Zeier, J., and Newman, J. P. (2011). Investigating the neural correlates of psychopathy: a critical review. *Mol. Psychiatry* 16, 792–799. doi: 10.1038/mp.2010.124
- Koenigs, M., Kruepke, M., Zeier, J., and Newman, J. P. (2012). Utilitarian moral judgment in psychopathy. *Soc. Cogn. Affect. Neurosci.* 7, 708–714. doi: 10.1093/scan/nsr048
- Koo, M., Algoe, S. B., Wilson, T. D., and Gilbert, D. T. (2008). It’s a wonderful life: mentally subtracting positive events improves people’s affective states, contrary to their affective forecasts. *J. Pers. Soc. Psychol.* 95, 1217–1224. doi: 10.1037/a0013316
- Koster, E. H. W., De Lissnyder, E., Derakshan, N., and De Raedt, R. (2011). Understanding depressive rumination from a cognitive science perspective: the impaired disengagement hypothesis. *Clin. Psychol. Rev.* 31, 138–145. doi: 10.1016/j.cpr.2010.08.005
- Kray, L. J., Galinsky, A. D., and Markman, K. D. (2009). Counterfactual structure and learning from experience in negotiations. *J. Exp. Soc. Psychol.* 45, 2007–2010. doi: 10.1016/j.jesp.2009.03.008
- Kray, L. J., George, L. G., Liljenquist, K. A., Galinsky, A. D., Tetlock, P. E., and Roese, N. J. (2010). From what might have been to what must have been: counterfactual thinking creates meaning. *J. Pers. Soc. Psychol.* 98, 106–118. doi: 10.1037/a0017905
- Kruger, J., Wirtz, D., and Miller, D. T. (2005). Counterfactual thinking and the first instinct fallacy. *J. Pers. Soc. Psychol.* 88, 725–735. doi: 10.1037/0022-3514.88.5.725
- Kudlicka, A., Clare, L., and Hindle, J. V. (2011). Executive functions in Parkinson’s disease: systematic review and meta-analysis. *Mov. Disord.* 26, 2305–2315. doi: 10.1002/mds.23868
- Kühberger, A., Großbichler, C., and Wimmer, A. (2011). Counterfactual closeness and predicted affect. *Think. Reason.* 17, 137–155. doi: 10.1080/13546783.2011.562079
- Kulakova, E., Aichhorn, M., Schurz, M., Kronbichler, M., and Perner, J. (2013). Processing counterfactual and hypothetical conditionals: an fMRI investigation. *Neuroimage* 72, 265–271. doi: 10.1016/j.neuroimage.2013.01.060
- Larquet, M., Coricelli, G., Opolczynski, G., and Thibaut, F. (2010). Impaired decision making in schizophrenia and orbitofrontal cortex lesion patients. *Schizophr. Res.* 116, 266–273. doi: 10.1016/j.schres.2009.11.010
- Lebow, R. N. (2000). What’s so different about a counterfactual? *World Politics* 52, 550–585. doi: 10.1017/S0043887100020104
- LeDoux, J. E. (2000). Emotion circuits in the brain. *Annu. Rev. Neurosci.* 23, 155–184. doi: 10.1146/annurev.neuro.23.1.155
- Lee, J., Quintana, J., Nori, P., and Green, M. F. (2011). Theory of mind in schizophrenia: exploring neural mechanisms of belief attribution. *Soc. Neurosci.* 6, 569–581. doi: 10.1080/17470919.2011.620774
- Levens, S. M., Larsen, J. T., Bruss, J., Tranel, D., Bechara, A., and Mellers, B. A. (2014). What might have been? The role of the ventromedial prefrontal cortex and lateral orbitofrontal cortex in counterfactual emotions and choice. *Neuropsychologia* 54, 77–86. doi: 10.1016/j.neuropsychologia.2013.10.026
- Lewis, D. (1979). Counterfactual dependence and time’s arrow. *Noûs* 13, 455–476. doi: 10.2307/2215339
- Lewis, D. (1986). Probabilities of conditionals and conditional probabilities II. *Philos. Rev.* 4, 581–589.
- Lewis, P. A., Critchley, H. D., Rotshtein, P., and Dolan, R. J. (2007). Neural correlates of processing valence and arousal in affective words. *Cereb. Cortex* 17, 742–748. doi: 10.1093/cercor/bhk024
- Li, J., Schiller, D., Schoenbaum, G., Phelps, E. A., and Daw, N. D. (2011). Differential roles of human striatum and amygdala in associative learning. *Nat. Neurosci.* 14, 1250–1252. doi: 10.1038/nn.2904
- Libby, L. K., and Eibach, R. P. (2011a). Self-enhancement or self-coherence? Why people shift visual perspective in mental images of the personal past and future. *Pers. Soc. Psychol. Bull.* 37, 714–726. doi: 10.1177/0146167211400207
- Libby, L. K., and Eibach, R. P. (2011b). Visual perspective in mental imagery: a representational tool that functions in judgment. *Adv. Exp. Soc. Psychol.* 44, 185–245. doi: 10.1016/B978-0-12-385522-0.00004-4
- Libby, L. K., Valenti, G., Pfent, A., and Eibach, R. P. (2011). Seeing failure in your life: imagery perspective determines whether self-esteem shapes reactions to recalled and imagined failure. *J. Pers. Soc. Psychol.* 101, 1157–1173. doi: 10.1037/a0026105
- Liu, X., Powell, D. K., Wang, H., Gold, B. T., Corbly, C. R., and Joseph, J. E. (2007). Functional dissociation in frontal and striatal areas for processing of positive and negative reward information. *J. Neurosci.* 27, 4587–4597. doi: 10.1523/jneurosci.5227-06.2007
- Maguire, E. A., and Hassabis, D. (2011). Role of the hippocampus in imagination and future thinking. *Proc. Natl. Acad. Sci. U S A* 108:E39. doi: 10.1073/pnas.1018876108
- Maguire, E. A., and Mullally, S. L. (2013). The hippocampus: a manifesto for change. *J. Exp. Psychol. Gen.* 142, 1180–1189. doi: 10.1037/a0033650
- Mar, R. A. (2011). The neural bases of social cognition and story comprehension. *Annu. Rev. Psychol.* 62, 103–134. doi: 10.1146/annurev-psych-120709-145406
- Marazziti, D., Baroni, S., Landi, P., Ceresoli, D., and Dell’osso, L. (2013). The neurobiology of moral sense: facts or hypotheses? *Ann. Gen. Psychiatry* 12:6. doi: 10.1186/1744-859x-12-6
- Markman, K. D., Gavanski, I., Sherman, S. J., and McMullen, M. N. (1993). The mental simulation of better and worse possible worlds. *J. Exp. Soc. Psychol.* 29, 87–109. doi: 10.1006/jesp.1993.1005
- Markman, K. D., Klein, W. M. P., and Suhr, J. A. (Eds.). (2009). *Handbook of Imagination and Mental Simulation*. New York: Psychology Press.
- Markman, K. D., Lindberg, M. J., Kray, L. J., and Galinsky, A. D. (2007). Implications of counterfactual structure for creative generation and analytical problem solving. *Pers. Soc. Psychol. Bull.* 33, 312–324. doi: 10.1177/0146167206296106
- Markman, K. D., and McMullen, M. N. (2003). A reflection and evaluation model of comparative thinking. *Pers. Soc. Psychol. Rev.* 7, 244–267. doi: 10.1207/s15327957pspr0703_04
- Markman, K. D., McMullen, M. N., and Elizaga, R. A. (2008). Counterfactual thinking, persistence and performance: a test of the reflection and evaluation model. *J. Exp. Soc. Psychol.* 44, 421–428. doi: 10.1016/j.jesp.2007.01.001
- Markman, K. D., and Miller, A. K. (2006). Depression, control and counterfactual thinking: functional for whom? *J. Soc. Clin. Psychol.* 25, 210–227. doi: 10.1521/jscp.2006.25.2.210
- Martinelli, P., Sperduti, M., and Piolino, P. (2013). Neural substrates of the self-memory system: new insights from a meta-analysis. *Hum. Brain Mapp.* 34, 1515–1529. doi: 10.1002/hbm.22008

- McCloy, R., and Byrne, R. M. J. (2000). Counterfactual thinking about controllable events. *Mem. Cognit.* 28, 1071–1078. doi: 10.3758/bf03209355
- McCrea, S. M. (2008). Self-Handicapping, excuse making and counterfactual thinking: consequences for self-esteem and future motivation. *J. Pers. Soc. Psychol.* 95, 274–292. doi: 10.1037/0022-3514.95.2.274
- McGraw, A. P., Mellers, B. A., and Tetlock, P. E. (2005). Expectations and emotions of olympic athletes. *J. Exp. Soc. Psychol.* 41, 438–446. doi: 10.1016/j.jesp.2004.09.001
- McMullen, M. N., and Markman, K. D. (2000). Downward counterfactuals and motivation: the wake-up call and the pangloss effect. *Pers. Soc. Psychol. Bull.* 26, 575–584. doi: 10.1177/0146167200267005
- McMullen, M. N., and Markman, K. D. (2002). Affective impact of close counterfactuals: implications of possible futures for possible pasts. *J. Exp. Soc. Psychol.* 38, 64–70. doi: 10.1006/jesp.2001.1482
- McNamara, P., Durso, R., Brown, A., and Lynch, A. (2003). Counterfactual cognitive deficit in persons with Parkinson's disease. *J. Neurol. Neurosurg. Psychiatr.* 74, 1065–1070. doi: 10.1136/jnnp.74.8.1065
- Mellers, B. A., Schwartz, A., Ho, K., and Ritov, I. (1997). Decision affect theory: emotional reactions to the outcomes of risky options. *Psychol. Sci.* 8, 423–429. doi: 10.1111/j.1467-9280.1997.tb00455.x
- Miller, E. K., and Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci.* 24, 167–202. doi: 10.1146/annurev.neuro.24.1.167
- Miller, A. K., Markman, K. D., Wagner, M. M., and Hunt, A. N. (2013). Mental simulation and sexual prejudice reduction: the debiasing role of counterfactual thinking. *J. Appl. Soc. Psychol.* 43, 190–194. doi: 10.1111/j.1559-1816.2012.00992.x
- Monroe, M. R., Skowronski, J. J., Macdonald, W., and Wood, S. E. (2005). The mildly depressed experience more post-decisional regret than the non-depressed. *J. Soc. Clin. Psychol.* 24, 665–690. doi: 10.1521/jscp.2005.24.5.665
- Moody, C. L., and Gennari, S. P. (2010). Effects of implied physical effort in sensory-motor and pre-frontal cortex during language comprehension. *Neuroimage* 49, 782–793. doi: 10.1016/j.neuroimage.2009.07.065
- Moore, W. E., Merchant, J. S., Kahn, L. E., and Pfeifer, J. H. (2014). "Like me?": ventromedial prefrontal cortex is sensitive to both personal relevance and self-similarity during social comparisons. *Soc. Cogn. Affect. Neurosci.* 9, 421–426. doi: 10.1093/scan/nst007
- Motzkin, J. C., Newman, J. P., Kiehl, K. A., and Koenigs, M. (2011). Reduced prefrontal connectivity in psychopathy. *J. Neurosci.* 31, 17348–17357. doi: 10.1523/jneurosci.4215-11.2011
- Murray, E. A., Wise, S. P., and Drevets, W. C. (2011). Localization of dysfunction in major depressive disorder: prefrontal cortex and amygdala. *Biol. Psychiatry* 69, e43–e54. doi: 10.1016/j.biopsych.2010.09.041
- Nasco, S. A., and Marsh, K. L. (1999). Gaining control through counterfactual thinking. *Pers. Soc. Psychol. Bull.* 25, 557–569. doi: 10.1177/0146167299025005002
- Nelson, S. M., Cohen, A. L., Power, J. D., Wig, G. S., Miezin, F. M., Wheeler, M. E., et al. (2010). A parcellation scheme for human left lateral parietal cortex. *Neuron* 67, 156–170. doi: 10.1016/j.neuron.2010.05.025
- Nestler, S., and von Collani, G. (2008). Hindsight bias and the activation of counterfactual mind-sets. *Exp. Psychol.* 55, 342–349. doi: 10.1027/1618-3169.55.5.342
- Nicolle, A., Bach, D. R., Driver, J., and Dolan, R. J. (2011a). A role for the striatum in regret-related choice repetition. *J. Cogn. Neurosci.* 23, 845–856. doi: 10.1162/jocn.2010.21510
- Nicolle, A., Bach, D. R., Frith, C. D., and Dolan, R. J. (2011b). Amygdala involvement in self-blame regret. *Soc. Neurosci.* 6, 178–189. doi: 10.1080/17470919.2010.506128
- Nieuwland, M. S. (2012). Establishing propositional truth-value in counterfactual and real-world contexts during sentence comprehension: differential sensitivity of the left and right inferior frontal gyri. *Neuroimage* 59, 3433–3440. doi: 10.1016/j.neuroimage.2011.11.018
- Nolen-Hoeksema, S. (2000). The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *J. Abnorm. Psychol.* 109, 504–511. doi: 10.1037/0021-843x.109.3.504
- Nolen-Hoeksema, S., Wisco, B. E., and Lyubomirsky, S. (2008). Rethinking rumination. *Perspect. Psychol. Sci.* 3, 400–424. doi: 10.1111/j.1745-6924.2008.00088.x
- Ochsner, K. N., and Gross, J. J. (2005). The cognitive control of emotion. *Trends Cogn. Sci.* 9, 242–249. doi: 10.1016/j.tics.2005.03.010
- O'Doherty, J. P., Critchley, H. D., Deichmann, R., and Dolan, R. J. (2003). Dissociating valence of outcome from behavioral control in human orbital and ventral prefrontal cortices. *J. Neurosci.* 23, 7931–7939.
- Ongür, D., Cullen, T. J., Wolf, D. H., Rohan, M., Barreira, P., Zalesak, M., et al. (2006). The neural basis of relational memory deficits in schizophrenia. *Arch. Gen. Psychiatry* 63, 356–365. doi: 10.1001/archpsyc.63.4.356
- Over, D. E., Hadjichristidis, C., Evans, J. S. B. T., Handley, S. J., and Sloman, S. A. (2007). The probability of causal conditionals. *Cogn. Psychol.* 54, 62–97. doi: 10.1016/j.cogpsych.2006.05.002
- Palminteri, S., Justo, D., Jauffret, C., Pavlicek, B., Dauta, A., Delmaire, C., et al. (2012). Critical roles for anterior insula and dorsal striatum in punishment-based avoidance learning. *Neuron* 76, 998–1009. doi: 10.1016/j.neuron.2012.10.017
- Petrocelli, J. V., and Harris, A. K. (2011). Learning inhibition in the monty hall problem: the role of dysfunctional counterfactual prescriptions. *Pers. Soc. Psychol. Bull.* 37, 1297–1311. doi: 10.1177/0146167211410245
- Petrocelli, J. V., Percy, E. J., Sherman, S. J., and Tormala, Z. L. (2011). Counterfactual potency. *J. Pers. Soc. Psychol.* 100, 30–46. doi: 10.1037/a0021523
- Petrocelli, J. V., and Sherman, S. J. (2010). Event detail and confidence in gambling: the role of counterfactual thought reactions. *J. Exp. Soc. Psychol.* 46, 61–72. doi: 10.1016/j.jesp.2009.09.013
- Phelps, E. A. (2006). Emotion and cognition: insights from studies of the human amygdala. *Annu. Rev. Psychol.* 57, 27–53. doi: 10.1146/annurev.psych.56.091103.070234
- Pieters, R., and Zeelenberg, M. (2005). On bad decisions and deciding badly: when intention-behavior inconsistency is regrettable. *Organ. Behav. Hum. Decis. Process.* 97, 18–30. doi: 10.1016/j.obhdp.2005.01.003
- Pighin, S., Byrne, R. M. J., Ferrante, D., Gonzalez, M., and Giroto, V. (2011). Counterfactual thoughts about experienced, observed and narrated events. *Think. Reason.* 17, 197–211. doi: 10.1080/13546783.2011.561598
- Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., et al. (2011). Functional network organization of the human brain. *Neuron* 72, 665–678. doi: 10.1016/j.neuron.2011.09.006
- Price, J. L., and Drevets, W. C. (2012). Neural circuits underlying the pathophysiology of mood disorders. *Trends Cogn. Sci.* 16, 61–71. doi: 10.1016/j.tics.2011.12.011
- Quelhas, A. C., Power, M. J., Juhas, C., and Senos, J. (2008). Counterfactual thinking and functional differences in depression. *Clin. Psychol. Psychother.* 15, 352–365. doi: 10.1002/cpp.593
- Rafetseder, E., Cristi-Vargas, R., and Perner, J. (2010). Counterfactual reasoning: developing a sense of "nearest possible world". *Child Dev.* 81, 376–389. doi: 10.1111/j.1467-8624.2009.01401.x
- Rafetseder, E., Schitalla, M., and Perner, J. (2013). Counterfactual reasoning: from childhood to adulthood. *J. Exp. Child Psychol.* 114, 389–404. doi: 10.1016/j.jecp.2012.10.010
- Raffard, S., D'Argembeau, A., Bayard, S., Boulenger, J.-P., and Van der Linden, M. (2010a). Scene construction in schizophrenia. *Neuropsychology* 24, 608–615. doi: 10.1037/a0019113
- Raffard, S., D'Argembeau, A., Lardi, C., Bayard, S., Boulenger, J.-P., and Van der Linden, M. (2010b). Narrative identity in schizophrenia. *Conscious. Cogn.* 19, 328–340. doi: 10.1016/j.concog.2009.10.005
- Raffard, S., D'Argembeau, A., Lardi, C., Bayard, S., Boulenger, J.-P., and Van Der Linden, M. (2009). Exploring self-defining memories in schizophrenia. *Memory* 17, 26–38. doi: 10.1080/09658210802524232
- Rao, A. S., and Foo, N. Y. (1989). "Minimal change and maximal coherence: a basis for belief revision and reasoning about actions," in *Proceedings of the 11th International Joint Conference on Artificial Intelligence* (Detroit, MI), 966–971.
- Revlin, R., Cate, C. L., and Rouss, T. S. (2001). Reasoning counterfactually: combining and rending. *Mem. Cognit.* 29, 1196–1208. doi: 10.3758/bf03206389

- Ridderinkhof, K. R., Ullsperger, M., Crone, E. A., and Nieuwenhuis, S. (2004). The role of the medial frontal cortex in cognitive control. *Science* 306, 443–447. doi: 10.1126/science.1100301
- Rips, L. J., and Edwards, B. J. (2013). Inference and explanation in counterfactual reasoning. *Cogn. Sci.* 37, 1107–1135. doi: 10.1111/cogs.12024
- Roese, N. J. (1994). The functional basis of counterfactual thinking. *J. Pers. Soc. Psychol.* 66, 805–818. doi: 10.1037/0022-3514.66.5.805
- Roese, N. J. (1997). Counterfactual thinking. *Psychol. Bull.* 121, 133–148.
- Roese, N. J. (1999). Counterfactual thinking and decision making. *Psychon. Bull. Rev.* 6, 570–578. doi: 10.3758/bf03212965
- Roese, N. J., Epstude, K., Fessel, F., Morrison, M., Smallman, R., Summerville, A., et al. (2009). Repetitive regret, depression and anxiety: findings from a nationally representative survey. *J. Soc. Clin. Psychol.* 28, 671–688. doi: 10.1521/jscp.2009.28.6.671
- Roese, N. J., Koehler, D. J., and Harvey, N. (2004). “Twisted pair: counterfactual thinking and the hindsight bias,” in *Blackwell Handbook of Judgement and Decision Making*, eds D. J. Koehler and N. Harvey (Malden, MA: Blackwell Publishing Ltd.), 258–273.
- Roese, N. J., and Morrison, M. (2009). The psychology of counterfactual thinking. *Historical Social Research* 34, 16–26.
- Roese, N. J., and Olson, J. M. (1996). Counterfactuals, causal attributions and the hindsight bias: a conceptual integration. *J. Exp. Soc. Psychol.* 32, 197–227. doi: 10.1006/jesp.1996.0010
- Roese, N. J., and Olson, J. M. (1997). Counterfactual thinking: the intersection of affect and function. *Adv. Exp. Soc. Psychol.* 29, 1–59. doi: 10.1016/s0065-2601(08)60015-5
- Roese, N. J., and Olson, J. M. (2007). Better, stronger, faster: self-serving judgment, affect regulation and the optimal vigilance hypothesis. *Perspect. Psychol. Sci.* 2, 124–141. doi: 10.1111/j.1745-6916.2007.00033.x
- Roese, N. J., and Summerville, A. (2005). What we regret most... and why. *Pers. Soc. Psychol. Bull.* 31, 1273–1285. doi: 10.1177/0146167205274693
- Roy, M., Shohamy, D., and Wager, T. D. (2012). Ventromedial prefrontal-subcortical systems and the generation of affective meaning. *Trends Cogn. Sci.* 16, 147–156. doi: 10.1016/j.tics.2012.01.005
- Rudebeck, P. H., Saunders, R. C., Prescott, A. T., Chau, L. S., and Murray, E. A. (2013). Prefrontal mechanisms of behavioral flexibility, emotion regulation and value updating. *Nat. Neurosci.* 16, 1140–1145. doi: 10.1038/nn.3440
- Ruh, N., Rahm, B., Unterrainer, J. M., Weiller, C., and Kaller, C. P. (2012). Dissociable stages of problem solving (II): first evidence for process-contingent temporal order of activation in dorsolateral prefrontal cortex. *Brain Cogn.* 80, 170–176. doi: 10.1016/j.bandc.2012.02.012
- Rushworth, M. F. S., Noonan, M. P., Boorman, E. D., Walton, M. E., and Behrens, T. E. J. (2011). Frontal cortex and reward-guided learning and decision-making. *Neuron* 70, 1054–1069. doi: 10.1016/j.neuron.2011.05.014
- Sander, D., Grafman, J., and Zalla, T. (2003). The human amygdala: an evolved system for relevance detection. *Rev. Neurosci.* 14, 303–316. doi: 10.1515/revneuro.2003.14.4.303
- Sanna, L. J. (1997). Self-efficacy and counterfactual thinking: up a creek with and without a paddle. *Pers. Soc. Psychol. Bull.* 23, 654–666. doi: 10.1177/0146167297236008
- Sanna, L. J., and Turley, K. J. (1996). Antecedents to spontaneous counterfactual thinking: effects of expectancy violation and outcome valence. *Pers. Soc. Psychol. Bull.* 22, 906–919. doi: 10.1177/0146167296229005
- Savla, G. N., Twamley, E. W., Delis, D. C., Roesch, S. C., Jeste, D. V., and Palmer, B. W. (2012). Dimensions of executive functioning in schizophrenia and their relationship with processing speed. *Schizophr. Bull.* 38, 760–768. doi: 10.1093/schbul/sbq149
- Schacter, D. L., Addis, D. R., Hassabis, D., Martin, V. C., Spreng, R. N., and Szpunar, K. K. (2012). The future of memory: remembering, imagining and the brain. *Neuron* 76, 677–694. doi: 10.1016/j.neuron.2012.11.001
- Schacter, D. L., Benoit, R. G., De Brigard, F., and Szpunar, K. K. (2015). Episodic future thinking and episodic counterfactual thinking: intersections between memory and decisions. *Neurobiol. Learn. Mem.* 117, 14–21. doi: 10.1016/j.nlm.2013.12.008
- Schilbach, L., Bzdok, D., Timmermans, B., Fox, P. T., Laird, A. R., Vogeley, K., et al. (2012). Introspective minds: using ALE meta-analyses to study commonalities in the neural correlates of emotional processing, social & unconstrained cognition. *PLoS One* 7:e30920. doi: 10.1371/journal.pone.0030920
- Scott, S. K., Leff, A. P., and Wise, R. J. (2003). Going beyond the information given: a neural system supporting semantic interpretation. *Neuroimage* 19, 870–876. doi: 10.1016/s1053-8119(03)00083-1
- Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., et al. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *J. Neurosci.* 27, 2349–2356. doi: 10.1523/jneurosci.5587-06.2007
- Shackman, A. J., Salomons, T. V., Slagter, H. A., Fox, A. S., Winter, J. J., and Davidson, R. J. (2011). The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nat. Rev. Neurosci.* 12, 154–167. doi: 10.1038/nrn2994
- Shad, M. U., Brent, B. K., and Keshavan, M. S. (2011). Neurobiology of self-awareness deficits in schizophrenia: a hypothetical model. *Asian J. Psychiatr.* 4, 248–254. doi: 10.1016/j.ajp.2011.09.002
- Shankman, S. A., Klein, D. N., Tenke, C. E., and Bruder, G. E. (2007). Reward sensitivity in depression: a biobehavioral study. *J. Abnorm. Psychol.* 116, 95–104. doi: 10.1037/0021-843x.116.1.95
- Silver, H., and Bilker, W. B. (2013). Pathways to similar executive impairment: comparison of schizophrenia patients and healthy aging individuals. *Psychiatry Res.* 210, 694–701. doi: 10.1016/j.psychres.2013.07.031
- Smallman, R. (2013). It's what's inside that counts: the role of counterfactual content in intention formation. *J. Exp. Soc. Psychol.* 49, 842–851. doi: 10.1016/j.jesp.2013.05.004
- Smallman, R., and McCulloch, K. C. (2012). Learning from yesterday's mistakes to fix tomorrow's problems: when functional counterfactual thinking and psychological distance collide. *Eur. J. Soc. Psychol.* 42, 383–390. doi: 10.1002/ejsp.1858
- Smallman, R., and Roese, N. J. (2009). Counterfactual thinking facilitates behavioral intentions. *J. Exp. Soc. Psychol.* 45, 845–852. doi: 10.1016/j.jesp.2009.03.002
- Smith, J. M., and Alloy, L. B. (2009). A roadmap to rumination: a review of the definition, assessment and conceptualization of this multifaceted construct. *Clin. Psychol. Rev.* 29, 116–128. doi: 10.1016/j.cpr.2008.10.003
- Sparks, A., McDonald, S., Lino, B., O'Donnell, M., and Green, M. J. (2010). Social cognition, empathy and functional outcome in schizophrenia. *Schizophr. Res.* 122, 172–178. doi: 10.1016/j.schres.2010.06.011
- Spreng, R. N. (2012). The fallacy of a “task-negative” network. *Front. Psychol.* 3:145. doi: 10.3389/fpsyg.2012.00145
- Spreng, R. N., and Grady, C. L. (2010). Patterns of brain activity supporting autobiographical memory, prospection and theory of mind and their relationship to the default mode network. *J. Cogn. Neurosci.* 22, 1112–1123. doi: 10.1162/jocn.2009.21282
- Spreng, R. N., and Mar, R. A. (2012). I remember you: a role for memory in social cognition and the functional neuroanatomy of their interaction. *Brain Res.* 1428, 43–50. doi: 10.1016/j.brainres.2010.12.024
- Spreng, R. N., Mar, R. A., and Kim, A. S. N. (2009). The common neural basis of autobiographical memory, prospection, navigation, theory of mind and the default mode: a quantitative meta-analysis. *J. Cogn. Neurosci.* 21, 489–510. doi: 10.1162/jocn.2008.21029
- Stein, M. B., Simmons, A. N., Feinstein, J. S., and Paulus, M. P. (2007). Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *Am J. Psychiatry* 164, 318–327. doi: 10.1176/appi.ajp.164.2.318
- Sternberg, R. J., and Gastel, J. (1989). If dancers ate their shoes: inductive reasoning with factual and counterfactual premises. *Mem. Cognit.* 17, 1–10. doi: 10.3758/bf03199551
- Stosic, M., Brass, M., Van Hoeck, N., Ma, N., and Van Overwalle, F. (2014). Brain activation related to the perception of minimal agency cues: the role of the mirror system. *Neuroimage* 86, 364–369. doi: 10.1016/j.neuroimage.2013.10.007
- Strahilevitz, M. A., Odean, T., and Barber, B. M. (2011). Once burned, twice shy: how naive learning, counterfactuals and regret affect the repurchase of stocks previously sold. *J. Mark. Res.* 48, S102–S120. doi: 10.1509/jmkr.48.spl.s102
- Summerfield, J. J., Hassabis, D., and Maguire, E. A. (2009). Cortical midline involvement in autobiographical memory. *Neuroimage* 44, 1188–1200. doi: 10.1016/j.neuroimage.2008.09.033

- Summerfield, J. J., Hassabis, D., and Maguire, E. A. (2010). Differential engagement of brain regions within a “core” network during scene construction. *Neuropsychologia* 48, 1501–1509. doi: 10.1016/j.neuropsychologia.2010.01.022
- Sumner, J. A. (2012). The mechanisms underlying overgeneral autobiographical memory: an evaluative review of evidence for the CaR-FA-X model. *Clin. Psychol. Rev.* 32, 34–48. doi: 10.1016/j.cpr.2011.10.003
- Sutton, R. S., and Barto, A. G. (1998). Reinforcement learning: an introduction. *IEEE Trans. Neural Netw.* 9, 1054–1054.
- Teigen, K. H., and Jensen, T. K. (2011). Unlucky victims or lucky survivors? *European Psychologist* 16, 48–57. doi: 10.1027/1016-9040/a000033
- Tobia, M. J., Guo, R., Schwarze, U., Boehmer, W., Gläscher, J., Finckh, B., et al. (2014). Neural systems for choice and valuation with counterfactual learning signals. *Neuroimage* 89, 57–69. doi: 10.1016/j.neuroimage.2013.11.051
- Townsend, J. D., Eberhart, N. K., Bookheimer, S. Y., Eisenberger, N. I., Folland-Ross, L. C., Cook, I. A., et al. (2010). fMRI activation in the amygdala and the orbitofrontal cortex in unmedicated subjects with major depressive disorder. *Psychiatry Res.* 183, 209–217. doi: 10.1016/j.psychres.2010.06.001
- Tykocinski, O. E., and Steinberg, N. (2005). Coping with disappointing outcomes: retroactive pessimism and motivated inhibition of counterfactuals. *J. Exp. Soc. Psychol.* 41, 551–558. doi: 10.1016/j.jesp.2004.12.001
- Tyser, M. P., McCrea, S. M., and Knüpfel, K. (2012). Pursuing perfection or pursuing protection? Self-evaluation motives moderate the behavioral consequences of counterfactual thoughts. *Eur. J. Soc. Psychol.* 42, 372–382. doi: 10.1002/ejsp.1864
- Urrutia, M., Gennari, S. P., and de Vega, M. (2012). Counterfactuals in action: an fMRI study of counterfactual sentences describing physical effort. *Neuropsychologia* 50, 3663–3672. doi: 10.1016/j.neuropsychologia.2012.09.004
- Ursu, S., and Carter, C. S. (2005). Outcome representations, counterfactual comparisons and the human orbitofrontal cortex: implications for neuroimaging studies of decision-making. *Brain Res. Cogn. Brain Res.* 23, 51–60. doi: 10.1016/j.cogbrainres.2005.01.004
- Valenti, G., Libby, L. K., Eibach, R. P., and Dickens, C. (2011). Looking back with regret: visual perspective in memory images differentially affects regret for actions and inactions. *J. Exp. Soc. Psychol.* 47, 730–737. doi: 10.1016/j.jesp.2011.02.008
- van Dijk, W. W., and Zeelenberg, M. (2002). Investigating the appraisal patterns of regret and disappointment. *Motiv. Emot.* 26, 321–331. doi: 10.1023/A:1022823221146
- Van Hoeck, N., Begtas, E., Steen, J., Kestemont, J., Vandekerckhove, M., and Van Overwalle, F. (2014). False belief and counterfactual reasoning in a social environment. *Neuroimage* 90, 315–325. doi: 10.1016/j.neuroimage.2013.12.043
- Van Hoeck, N., Ma, N., Ampe, L., Baetens, K., Vandekerckhove, M. M. P., and Van Overwalle, F. (2013). Counterfactual thinking: an fMRI study on changing the past for a better future. *Soc. Cogn. Affect. Neurosci.* 8, 556–564. doi: 10.1093/scan/nss031
- Van Hoeck, N., Ma, N., Van Overwalle, F., and Vandekerckhove, M. M. P. (2010). Counterfactual thinking and the episodic system. *Behav. Neurol.* 23, 225–227. doi: 10.3233/BEN-2010-0301
- Van Hoeck, N., Revlin, R., Dieussaert, K., and Schaeken, W. (2012). The development of counterfactual reasoning in belief revision. *Psychol. Belg.* 52, 407–434. doi: 10.5334/pb-52-4-407
- Van Mulukom, V., Schacter, D. L., Corballis, M. C., and Addis, D. R. (2013). Re-imagining the future: repetition decreases hippocampal involvement in future simulation. *PLoS One* 8:e69596. doi: 10.1371/journal.pone.0069596
- Van Overwalle, F. (2011). A dissociation between social mentalizing and general reasoning. *Neuroimage* 54, 1589–1599. doi: 10.1016/j.neuroimage.2010.09.043
- Van Overwalle, F., and Baetens, K. (2009). Understanding others’ actions and goals by mirror and mentalizing systems: a meta-analysis. *Neuroimage* 48, 564–584. doi: 10.1016/j.neuroimage.2009.06.009
- Viard, A., Chételat, G., Lebreton, K., Desgranges, B., Landeau, B., de La Sayette, V., et al. (2011). Mental time travel into the past and the future in healthy aged adults: an fMRI study. *Brain Cogn.* 75, 1–9. doi: 10.1016/j.bandc.2010.10.009
- Vincent, J. L., Kahn, I., Snyder, A. Z., Raichle, M. E., and Buckner, R. L. (2008). Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *J. Neurophysiol.* 100, 3328–3342. doi: 10.1152/jn.90355.2008
- Walsh, C. R., and Byrne, R. M. J. (2004). Counterfactual thinking: the temporal order effect. *Mem. Cognit.* 32, 369–378. doi: 10.3758/bf03195831
- Waters, F., Allen, P., Aleman, A., Fernyhough, C., Woodward, T. S., Badcock, J. C., et al. (2012). Auditory hallucinations in schizophrenia and nonschizophrenia populations: a review and integrated model of cognitive mechanisms. *Schizophr. Bull.* 38, 683–693. doi: 10.1093/schbul/sbs045
- Watkins, E. R. (2008). Constructive and unconstructive repetitive thought. *Psychol. Bull.* 134, 163–206. doi: 10.1037/0033-2909.134.2.163
- Watson, P. D., Voss, J. L., Warren, D. E., Tranel, D., and Cohen, N. J. (2013). Spatial reconstruction by patients with hippocampal damage is dominated by relational memory errors. *Hippocampus* 23, 570–580. doi: 10.1002/hipo.22115
- Whitfield-Gabrieli, S., Thermenos, H. W., Milanovic, S., Tsuang, M. T., Faraone, S. V., McCarley, R. W., et al. (2009). Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia. *Proc. Natl. Acad. Sci. U S A* 106, 1279–1284. doi: 10.1073/pnas.0809141106
- Whitman, J. C., Metzack, P. D., Lavigne, K. M., and Woodward, T. S. (2013). Functional connectivity in a frontoparietal network involving the dorsal anterior cingulate cortex underlies decisions to accept a hypothesis. *Neuropsychologia* 51, 1132–1141. doi: 10.1016/j.neuropsychologia.2013.02.016
- Williams, J. M. G. (2006). Capture and rumination, functional avoidance and executive control (CaRFAX): three processes that underlie overgeneral memory. *Cogn. Emot.* 20, 548–568. doi: 10.1080/02699930500450465
- Williams, J. M. G., Barnhofer, T., Crane, C., Hermans, D., Raes, F., Watkins, E., et al. (2007). Autobiographical memory specificity and emotional disorder. *Psychol. Bull.* 133, 122–148. doi: 10.1037/0033-2909.133.1.122
- Williams, L. E., Must, A., Avery, S., Woolard, A., Woodward, N. D., Cohen, N. J., et al. (2010). Eye-movement behavior reveals relational memory impairment in schizophrenia. *Biol. Psychiatry* 68, 617–624. doi: 10.1016/j.biopsych.2010.05.035
- Wong, E. M., Haselhorn, M. P., and Kray, L. J. (2012). Improving the future by considering the past: the impact of upward counterfactual reflection and implicit beliefs on negotiation performance. *J. Exp. Soc. Psychol.* 48, 403–406. doi: 10.1016/j.jesp.2011.07.014
- Wrosch, C., and Miller, G. E. (2009). Depressive symptoms can be useful: self-regulatory and emotional benefits of dysphoric mood in adolescence. *J. Pers. Soc. Psychol.* 96, 1181–1190. doi: 10.1037/a0015172
- Wrosch, C., Scheier, M. F., and Miller, G. E. (2013). Goal adjustment capacities, subjective well-being and physical health. *Soc. Personal. Psychol. Compass* 7, 847–860. doi: 10.1111/spc3.12074
- Xue, G., Juan, C.-H., Chang, C.-F., Lu, Z.-L., and Dong, Q. (2012). Lateral prefrontal cortex contributes to maladaptive decisions. *Proc. Natl. Acad. Sci. U S A* 109, 4401–4406. doi: 10.1073/pnas.1111927109
- Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., and Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nat. Methods* 8, 665–670. doi: 10.1038/nmeth.1635
- Young, L., Koenigs, M., Kruepke, M., and Newman, J. P. (2012). Psychopathy increases perceived moral permissibility of accidents. *J. Abnorm. Psychol.* 121, 659–667. doi: 10.1037/a0027489
- Yu, R.-L., and Wu, R.-M. (2013). Social brain dysfunctions in patients with Parkinson’s disease: a review of theory of mind studies. *Transl. Neurodegener.* 2:7. doi: 10.1186/2047-9158-2-7
- Zaki, J., López, G., and Mitchell, J. P. (2014). Activity in ventromedial prefrontal cortex co-varies with revealed social preferences: evidence for person-invariant value. *Soc. Cogn. Affect. Neurosci.* 9, 464–469. doi: 10.1093/scan/nst005
- Zalla, T., and Sperduti, M. (2013). The amygdala and the relevance detection theory of autism: an evolutionary perspective. *Front. Hum. Neurosci.* 7:894. doi: 10.3389/fnhum.2013.00894
- Zeelenberg, M., van Dijk, W. W., van der Pligt, J., Manstead, A. S., van Empelen, P., and Reinderman, D. (1998). Emotional reactions to the outcomes of decisions: the role of counterfactual thought in the experience of regret and disappointment. *Organ. Behav. Hum. Decis. Process.* 75, 117–141. doi: 10.1006/obhd.1998.2784
- Zeithamova, D., Dominick, A. L., and Preston, A. R. (2012). Hippocampal and ventral medial prefrontal activation during retrieval-mediated learning supports novel inference. *Neuron* 75, 168–179. doi: 10.1016/j.neuron.2012.05.010
- Zhang, Q., and Covey, J. (2014). Past and future implications of near-misses and their emotional consequences. *Exp. Psychol.* 61, 118–126. doi: 10.1027/1618-3169/a000231

Zhang, J., Wang, J., Wu, Q., Kuang, W., Huang, X., He, Y., et al. (2011). Disrupted brain connectivity networks in drug-naive, first-episode major depressive disorder. *Biol. Psychiatry* 70, 334–342. doi: 10.1016/j.biopsych.2011.05.018

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Van Hoeck, Watson and Barbey. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.