



Neuropeptides Exert Neuroprotective Effects in Alzheimer's Disease

Xin-Yi Chen^{1,2†}, Yi-Feng Du² and Lei Chen^{1*}

¹ Department of Physiology and Pathophysiology, Qingdao University, Qingdao, China, ² Department of Neurology, Provincial Hospital Affiliated to Shandong University, Jinan, China

OPEN ACCESS

Edited by:

Christian Gonzalez-Billault,
Universidad de Chile, Chile

Reviewed by:

Luis Gerardo Aguayo,
Universidad de Concepción, Chile
Jing Chen,
University of Warwick,
United Kingdom

*Correspondence:

Lei Chen
chenleiqd@163.com

†Present Address:

Xin-Yi Chen,
Department of Medicine and
Therapeutics, The Chinese University
of Hong Kong, Hong Kong,
Hong Kong

Received: 05 May 2018

Accepted: 21 December 2018

Published: 11 January 2019

Citation:

Chen X-Y, Du Y-F and Chen L (2019)
Neuropeptides Exert Neuroprotective
Effects in Alzheimer's Disease.
Front. Mol. Neurosci. 11:493.
doi: 10.3389/fnmol.2018.00493

Alzheimer's disease (AD) is an age-related neurodegenerative disorder characterized by cognitive deficits and neuronal loss. Deposition of beta-amyloid peptide (A β) causes neurotoxicity through the formation of plaques in brains of Alzheimer's disease. Numerous studies have indicated that the neuropeptides including ghrelin, neurotensin, pituitary adenylate cyclase-activating polypeptide (PACAP), neuropeptide Y, substance P and orexin are closely related to the pathophysiology of Alzheimer's disease. The levels of neuropeptides and their receptors change in Alzheimer's disease. These neuropeptides exert neuroprotective roles mainly through preventing A β accumulation, increasing neuronal glucose transport, increasing the production of neurotrophins, inhibiting endoplasmic reticulum stress and autophagy, modulating potassium channel activity and hippocampal long-term potentiation. Therefore, the neuropeptides may function as potential drug targets in the prevention and cure of Alzheimer's disease.

Keywords: neuropeptide, Alzheimer's disease, ghrelin, neurotensin, pituitary adenylate cyclase-activating polypeptide, neuropeptide Y, substance P, orexin

INTRODUCTION

Alzheimer's disease (AD) is an age-related neurodegenerative disorder which is clinically characterized by cognitive deficits, memory impairment, disorientation, and behavioral issues (Burns and Iliffe, 2009). Commonly the Alzheimer's disease begins in people over the age of 65 years. The neuropathological features in Alzheimer's disease are amyloid plaques, neurofibrillary tangles, and neuronal loss (Tiraboschi et al., 2004). Beta-amyloid peptide (A β) oligomers have been reported to be the primary pathogenic forms of A β , which change the structure of synapses and eventually disrupt neuronal communication (Lacor et al., 2007). However, the pathogenesis remains unknown.

Neuropeptides are molecules which function as endogenous active substances within central nervous system and peripheral nervous system. Neuropeptides play important roles in a wide range of brain functions, including food intake, metabolism, reproduction, social behaviors, reward, learning and memory, sleep and wakefulness. Recent studies revealed that many neuropeptides including ghrelin, neurotensin, pituitary adenylate cyclase-activating polypeptide (PACAP), neuropeptide Y, substance P, and orexin may be associated with the pathophysiology and potential therapy of Alzheimer's disease. In this article, we review the recent advances about the involvement of neuropeptides in Alzheimer's disease.

GHRELIN AND ALZHEIMER'S DISEASE

Ghrelin, a 28-amino acid brain-gut peptide, activates the growth hormone secretagogue receptors which are expressed widely in the brain (Guan et al., 1997). Recent studies revealed that the receptor expression is developmental, with a stronger staining in the early stages and a weaker expression in the later stages of development (Lattuada et al., 2013). Ghrelin exerts critical roles in the regulation of energy homeostasis, neuroendocrine and neurodegenerative processes, especially in higher brain functions, such as learning and memory consolidation (Spitznagel et al., 2010; Rak-Mardyla, 2013; Murray et al., 2014; Panagopoulos and Ralevski, 2014; Jiao et al., 2017). Ghrelin is also involved in mitochondrial respiration and neuroprotection, which can be developed as biomarkers or drug targets for prevention and treatment of neurological disorders, including Parkinson's disease, stroke, epilepsy and Alzheimer's disease (dos Santos et al., 2013a; Shi et al., 2014, 2017; Stoyanova, 2014).

Ghrelin has been demonstrated to have a close relationship with Alzheimer's disease. One of the single nucleotide polymorphisms of the ghrelin gene, rs4684677 (Leu90Gln), has been proved to be associated with the onset age of Alzheimer's disease (Shibata et al., 2011). Early study revealed that the level of serum ghrelin is negatively related to several cognitive domains in older adults (Spitznagel et al., 2010). The level of the functional form of ghrelin, acylated ghrelin, is associated with Alzheimer's disease risk factors and mild cognitive impairment (Gahete et al., 2010; Cao et al., 2018). However, a significant reduction of ghrelin mRNA was observed in the temporal gyrus of Alzheimer's disease patients, suggesting the possible involvement of ghrelin in the cognitive deficit of Alzheimer's disease (Gahete et al., 2010).

Administration of ghrelin or ghrelin agonist decreases the level of A β and attenuates Alzheimer's disease-related cognitive impairment (Gahete et al., 2010, 2011; Dhurandhar et al., 2013; Kunath et al., 2015). Recent *in vitro* study demonstrated that pre-administration of ghrelin or ghrelin analog protects SH-SY5Y cellular models of Alzheimer's disease against methylglyoxal-induced toxicity and apoptosis, suggesting the potential treatment for neurodegenerative disorders (Cecarini et al., 2016; Popelová et al., 2018). Ghrelin exerts neuroprotective effects through different mechanisms. Central application of acylated ghrelin prevents A β -induced impairments of memory and energy and glucose metabolisms, probably through increase of AMPK and GSK phosphorylation and decrease of tau phosphorylation (Kang et al., 2015). Further study revealed that acylated ghrelin also blunts A β -induced depression of long-term potentiation (LTP) in hippocampus and therefore prevents impairments of recognition and spatial orientation (Santos et al., 2017). Ghrelin ameliorates neurogenesis impairment in hippocampus and improves memory deficits by prevention of synaptic degeneration including cholinergic fiber loss (Moon et al., 2011, 2014). Recent studies revealed that ghrelin exerts protective effects against A β -induced toxicity via preventing superoxide production, calcium elevation and mitochondrial membrane depolarization (Martins et al., 2013;





FIGURE 1 | A schematic diagram describing the possible pathways of ghrelin-induced neuroprotective effects in Alzheimer's disease. Ghrelin protects against A β -induced neurotoxicity through prevention of calcium elevation, superoxide production and mitochondrial membrane depolarization. Ghrelin also increases neuronal glucose uptake by activating Akt/GSK phosphorylation and improving tau hyperphosphorylation. Moreover, ghrelin prevents cholinergic synaptic degeneration and therefore protects against A β -induced memory deficits. Acylated ghrelin improves A β -induced deterioration of memory through increase of AMPK and GSK phosphorylation and decrease of tau phosphorylation. Analog of ghrelin protects against MGO-induced neurotoxicity and apoptosis in cellular models of Alzheimer's disease. GHS-R, growth hormone secretagogue receptors, also known as ghrelin receptors; $\Delta\psi_m$, mitochondrial membrane potential; MGO, methylglyoxal; ACh, acetylcholine; AMPK, adenosine 5'-monophosphate (AMP)-activated protein kinase; GSK, glycogen synthase kinase; AChR, cholinergic receptors; -P, phosphorylation. The internal and external circles represent the inner and outer leaflets of the cellular membrane. The dotted line in ACh synapse represents the degenerated synapse.

Gomes et al., 2014). In addition, ghrelin restores the proteasome functionality in Alzheimer's disease and thus contributes to the elimination of toxic aggregates (Cecarini et al., 2016). It is known that brain insulin resistance is closely related to the cognitive impairment and neurodegeneration, particularly within Alzheimer's disease. Recently, it is demonstrated that ghrelin increases neuronal glucose uptake and improves tau hyperphosphorylation by activating Akt and GSK-3 beta phosphorylation in cultured hippocampal neurons (Chen et al., 2010).

In summary, the levels of ghrelin change in Alzheimer's disease. The functional form of ghrelin, acylated ghrelin, increases in mild cognitive impairment. A single nucleotide polymorphism of ghrelin gene is associated with the onset age of Alzheimer's disease. Ghrelin protects against A β -induced neurotoxicity through multiple pathways. Ghrelin prevents calcium elevation, superoxide production and mitochondrial membrane depolarization. Moreover, ghrelin increases neuronal glucose uptake and improves A β -induced deterioration of memory through activating AMPK and GSK phosphorylation and decreasing tau phosphorylation. Furthermore, ghrelin prevents cholinergic synaptic degeneration. Therefore, ghrelin is considered as a potential drug in the treatment of Alzheimer's disease (**Figure 1**).



NEUROTENSIN AND ALZHEIMER'S DISEASE

Neurotensin is a tridecapeptide which binds with two neurotensin receptors, type-1 and type-2, in the brain. Neurotensin plays multiple effects in central nervous system and is involved in the pathophysiology of several central nervous system disorders, including schizophrenia (Garver et al., 1991; Kinkead and Nemeroff, 2004), Parkinson's disease (Bissette et al., 1985; Fernandez et al., 1994), as well as Alzheimer's disease (Constantinidis et al., 1983; Struble et al., 1987; Gahete et al., 2010; Xiao et al., 2014). The levels of neurotensin and neurotensin receptors change in several brain regions of Alzheimer's disease patients. Neurotensin influences the formation of senile plaques and therefore is associated with the pathogenesis of Alzheimer's disease through different pathways (Figure 2).

Previous studies revealed a decreased level of neurotensin in the septum (Ferrier et al., 1983), suprachiasmatic nucleus (Hu et al., 2013) and amygdala (Benzing et al., 1990), and a low level of neurotensin receptors in the entorhinal area (Jansen et al., 1990), dentate gyrus (Rowe et al., 2006) and temporal gyrus (Gahete et al., 2010) of Alzheimer's disease. In Alzheimer's disease patients, the amyloid plaque density in the occipital cortex is negatively associated with the neurotensin-positive neurons in the suprachiasmatic nucleus, i.e., the more plaques, the fewer neurotensin-positive neurons, which suggests the involvement of neurotensin in fractal activity disruption in Alzheimer's disease (Hu et al., 2013). The levels of amygdala

neurotensin in males are significantly higher than that in females (Biggins et al., 1983), which suggests that the intrinsic function of the amygdala may be associated with hormonal regulation of other sex-dependent mechanism. For example, Skup et al. (2011) reported that the males and the females show different patterns in amygdala volume decline over time. Previous studies revealed that men exhibit greater volumes and neuronal densities in amygdala (Giedd et al., 1996; Goldstein et al., 2001; Witte et al., 2010). Furthermore, the density of neurotensin and acetylcholine containing fibers is reduced dramatically in the regions of amygdala with greatest senile plaques (Benzing et al., 1992, 1993). However, in the case of high plaque non-demented, no significant reduction of neurotensinergic fibers and other neurotransmitter fibers was observed in the amygdala (Benzing et al., 1993).


It is well-known that the entorhinal cortex is a crucial brain region involved in the earliest pathological change of Alzheimer's disease (Mann, 1989; Beach et al., 1997). Recently, Xiao et al. (2014) found that neurotensin persistently increases the spontaneous firing rate of neurons in the entorhinal cortex. This facilitation is mediated by neurotensin type 1 receptor and TREK-2K⁺ channels. Further behavioral studies revealed that activation of neurotensin type 1 receptors enhances spatial learning and therefore improves memory status in APP/PS1 Alzheimer's disease mice model. Recent electrophysiological studies further demonstrated that neurotensin increases glutamate release and spontaneous firing rate of dentate gyrus through both presynaptic and post-synaptic neurotensin receptors, respectively (Zhang et al., 2015, 2016).

It has recently been shown that the precursor form of nerve growth factor (proNGF) forming complex with p75 and sortilin contributes to neuronal death in basal forebrain neurons (Al-Shawi et al., 2007, 2008). Sortilin plays an important role in proNGF-mediated neurotoxicity. As a sortilin ligand, neurotensin blocks the sortilin-induced effects of proNGF and further rescues the survival of old neurons (Al-Shawi et al., 2008).

In summary, the levels of neurotensin and neurotensin receptors decrease in many brain areas of Alzheimer's disease. The amyloid plaque density in the occipital cortex is negatively associated with the neurotensin-positive neurons in the suprachiasmatic nucleus. Neurotensin increases the excitability of entorhinal cortex neurons and therefore improves spatial learning and memory. Furthermore, neurotensin rescues the survival of aged neurons through blocking sortilin-induced proNGF neurotoxicity.

PACAP AND ALZHEIMER'S DISEASE


PACAP belongs to the superfamily of secretin/glucagon/vasoactive intestinal polypeptide (Miyata et al., 1989). High level of PACAP is expressed in the hypothalamus, hippocampus, cerebellum and several brainstem nuclei (Hannibal, 2002). By activating three types of receptors (PAC1, VPAC1, and VPAC2), PACAP functions as a neurohormone, neurotransmitter, or neurotrophic factor in central nervous system (Lee and Seo, 2014). There is growing evidence (Reglodi et al., 2011) suggests



that PACAP is closely associated with the pathology of Alzheimer's disease (Figure 3).

By using three different mouse models of Alzheimer's disease, early study showed downregulation of PACAP genes (Wu et al., 2006). Moreover, the PACAP levels are reduced in brain areas including entorhinal cortex, middle temporal gyrus, superior frontal gyrus, and primary visual cortex in Alzheimer's disease patients (Han et al., 2014a). Further study revealed that the lower PACAP levels are correlated with higher amyloid burden, tau protein, and the declined recognition memory with aging (Han et al., 2014b; An et al., 2017).

It is demonstrated that PACAP exerts neuroprotective effects through multiple mechanisms in Alzheimer's disease patients and mouse models (Vaudry et al., 2004; Rat et al., 2011). Onoue et al. (2002) first demonstrated that PACAP protects against A β -induced neuronal toxicity. Treatment with PACAP rescues 80% of A β -induced reduction of cell viability through increase of cAMP and deactivation of caspase-3 in PC12 cells. Intranasal application of PACAP increases the levels of brain-derived neurotrophic factor (BDNF) and antiapoptotic Bcl-2 protein. In addition, PACAP also increases the level of A β -degrading enzyme (Rat et al., 2011). Therefore, intranasal application of PACAP could be a useful therapeutic approach in treating Alzheimer's disease. It is known that, in the non-amyloidogenic pathway, the α -secretase cleaves the amyloid precursor protein and prevents amyloid plaque formation (Postina, 2012). Activation of PAC1 receptors increases APPs alpha secretion and enhances α -secretase cleavage of APP. Furthermore, both the MAPK and PI3K pathways are involved in PACAP-mediated α -secretase activation (Kojro et al., 2006).



Ginsenoside is a major component of the traditional herb ginseng which has been proved to exert neurotrophic and neuroprotective effects via preventing neuronal degeneration. Ginsenosides Rh2 increases the gene expression of PACAP in astrocytes of the brain and therefore ameliorates A β -induced growth inhibition of astrocytes (Shieh et al., 2008).

In summary, the level of PACAP decreases in brain areas of Alzheimer's disease patients and mouse models, which is correlated with higher amyloid burden, tau protein, and the declined recognition memory. PACAP protects against A β -induced neuronal toxicity through activation of cAMP, Bcl2, BDNF, A β -degrading enzyme, and deactivation of caspase-3. In addition, PACAP enhances α -secretase cleavage of APP via both MAPK and PI3K pathways.

NEUROPEPTIDE Y AND ALZHEIMER'S DISEASE

Neuropeptide Y is a 36-amino acid neuropeptide. Neuropeptide Y receptors are classified into five subtypes: Y1, Y2, Y4, Y5, and Y6 (Hsieh et al., 2013; Mittapalli and Roberts, 2014; Pérez-Fernández et al., 2014). Neuropeptide Y plays functions associated with modulation of food ingestion, mood, learning and memory (dos Santos et al., 2013b) and also plays an important role in neuroprotection against neurodegenerative diseases (Figure 4).

The expression of neuropeptide Y changes under neurodegenerative diseases including Alzheimer's disease (Duarte-Neves et al., 2016). An anomalous high expression

level of neuropeptide Y with aging was detected in the hippocampal circuits of mouse model of Alzheimer's disease (Diez et al., 2003; Krezymon et al., 2013). Recently, Mahar et al. (2017) reported that the number of neuropeptide Y immunoreactive hippocampal interneurons reduces in presymptomatic TgCRND8 Alzheimer's disease mouse model.

Neuropeptide Y has neuroprotective effects in Alzheimer's disease (Croce et al., 2011, 2012, 2013; Angelucci et al., 2014; Duarte-Neves et al., 2016; Spencer et al., 2016). Intracerebroventricular application of neuropeptide Y prevents A β _{1–40}-induced depressive-like symptoms and spatial memory impairments through Y2 receptors. Neuropeptide Y produces the neuroprotection through inhibition of A β -induced lipid peroxidation, indicating the involvement of prevention of intracellular oxidative stress (dos Santos et al., 2013b). Moreover, neuropeptide Y induces protective effects against endoplasmic reticulum stress-mediated cell loss through activation of PI3K-XBP1-induced Gip78/BiP pathway as well as inhibition of caspase-3 and caspase-4 activities (Lee et al., 2018). Neuropeptide Y-induced neuroprotective effects are also associated with the production of neurotrophin family (Croce et al., 2011, 2012, 2013; Angelucci et al., 2014). Pretreatment with neuropeptide Y protects neurons against A β neurotoxicity which is accompanied by an increased intracellular level of NGF (Croce et al., 2012) and BDNF (Croce et al., 2013). Moreover, decrease of miR-30-5p (a membrane of miR-30a family regulating BDNF tuning expression) levels is involved in neuropeptide Y-induced modulation of BDNF.




FIGURE 5 | A model describing the multiple effects of substance P in Alzheimer's disease. Substance P inhibits A β -induced overexpression of K⁺ channel and A β -induced increase of K⁺ current, and therefore attenuates cognitive deficit and apoptosis in Alzheimer's disease. Furthermore, substance P exerts neuroprotective effects through inhibition of caspase-3-induced PARP-1 cleavage and enhancement of α -secretase activity. Neurokinin B plays a role in copper homeostasis. However, substance P receptor antagonist attenuates aluminum-induced spatial memory deficit probably through blockade of substance P-mediated neuroinflammation. SP, substance P; NKR, neurokinin receptors; PARP-1, poly ADP-ribose polymerase-1; sAPP α , soluble amyloid precursor protein α ; NKB, neurokinin B; Cu, copper; Al, aluminum. The internal and external circles represent the inner and outer leaflets of the cellular membrane.

The neutral endopeptidase (NEP) neprilysin is an A β degrading enzyme which is therefore involved in the pathogenesis of Alzheimer's disease. However, NEP has also been proved to cleave neuropeptide Y into C-terminal fragments. The NEP-produced C-terminal fragments of neuropeptide Y attenuate the neurodegenerative process in both transgenic Alzheimer's disease mice and A β treated human neurons (Rose et al., 2009). It is well-known that A β accumulation, senile plaque formation and neuronal dysfunction contribute to the cognitive decline in Alzheimer's disease. However, weight loss is an early sign of Alzheimer's disease. It is reported that high level of A β potentially devastates hypothalamic arcuate neuropeptide Y neurons and downregulates the leptin state in the early disease process, which may lead to the weight loss (dos Santos et al., 2013b; Ishii et al., 2014).

In summary, the level of neuropeptide Y increases significantly in hippocampus of mouse Alzheimer's disease models. Neuropeptide Y exerts neuroprotective effects through multiple pathways. Neuropeptide Y mitigates endoplasmic reticulum stress-induced neuronal cell death through activation of PI3K-XBP1-induced Gip78/BiP pathway and inhibition of caspase-3 and caspase-4 activities. Furthermore, neuropeptide Y suppresses oxidative stress via inhibition of A β -induced lipid peroxidation. In addition, neuropeptide Y plays neuroprotection via increasing the levels of BDNF and NGF.




FIGURE 6 | A scheme describing the complicated effects of orexin in Alzheimer's disease. Firstly, in microglial cells, orexin suppresses autophagosome-lysosome fusion process, leading to impaired A β degradation. Furthermore, orexin suppresses A β uptake through downregulating phagocytosis regulating molecules, such as PI3K, Akt, and p38-MAPK. A β -plaque formation and tau hyper-phosphorylation decrease the expression of orexin receptors in Alzheimer's disease. Secondly, orexin receptors and GPR103 form functional heterodimers. Orexin augments NF- κ B, PI3K-Akt, Jak-STAT signaling and induces ERK_{1/2} phosphorylation, and therefore is involved in neuroprotective functions. Finally, one of the orexin receptor 2 gene is likely a risk factor for Alzheimer's disease. In APP/PS1 Alzheimer's disease mice, orexin gene knock out decreases the amount of A β . Orexin modulates the hippocampal oscillation and the expression of clock-controlled-genes, Bace1 and Bace2, which are associated with the production of A β . OXR, orexin receptors; AD, Alzheimer's disease. The internal and external circles represent the inner and outer leaflets of the cellular membrane.

SUBSTANCE P AND ALZHEIMER'S DISEASE

The tachykinin family includes substance P, neurokinin A and neurokinin B. By activating neurokinin receptors,

tachykinin plays important roles including pain, depression, nausea and emesis (Mantyh, 2002). In addition, tachykinin protects against the neurotoxic processes of Alzheimer's disease by multiple pathways (Severini et al., 2016).

TABLE 1 | Changes in the level of neuropeptides and receptors in Alzheimer's disease.

Neuropeptides	Receptors	Levels	Biological sample	State of disease		References
				Human	Animal models	
Ghrelin mRNA		↓	Temporal gyrus	AD patient		Gahete et al., 2010
Acylated ghrelin		↑	Serum	MCI patient		Cao et al., 2018
	GHS-R1a	↓	Temporal gyrus	AD patient		Gahete et al., 2010
Neurotensin	GHS-R1b	↑	Temporal gyrus	AD patient		Gahete et al., 2010
		↓	Amygdala	AD patient		Benzing et al., 1990, 1992, 1993
Neurotensin		↓	Septum	AD patient		Ferrier et al., 1983
Neurotensin mRNA		↓	Temporal gyrus	AD patient		Gahete et al., 2010
PACAP gene	NTSR1, NTSR2	↓	Temporal gyrus	AD patient		Gahete et al., 2010
	NTSR	↓	Entorhinal area	AD patient		Jansen et al., 1990
	NTSR	↓	Dentate gyrus, SNc, VTA, PVN _h		Aged (24–25 months) rats with or without memory impairment	Rowe et al., 2006
PACAP gene		↓	Cortex		APP/PS-1 (18 months) and Tg2576/PS-1 (12 months) mice with Aβ deposition	Wu et al., 2006
PACAP		↓	Temporal cortex	AD patients		
		↓	Entorhinal cortex, middle temporal gyrus, superior frontal gyrus, primary visual cortex	AD patient		Han et al., 2014a
PACAP		↓	Hippocampus, cortex around hippocampus		hAPP mice in each age group	Han et al., 2017
Neuropeptide Y		↑	Dentate gyrus		Tg2576 mice (18 months)	Krezymon et al., 2013
Neuropeptide Y		↑	Hippocampus, cortex		APP23 mice (27 months) with Aβ plaques	Diez et al., 2003
Neuropeptide Y		↓	Hippocampal interneurons		TgCRND8 mice (1 month before amyloid deposition)	Mahar et al., 2017
Substance P		↓	Dentate gyrus	AD patient		Quigley and Kowall, 1991
Substance P		↓	Cortex, hippocampus		Rats with Aβ infusion	Nag et al., 1999
Substance P		↓	Cortex, hippocampus	AD patient		Bouras et al., 1990
Substance P		↑	Pallidum, substantia nigra	AD patient		Bouras et al., 1990
Substance P		↑	Astrocytes in hippocampal formation and thalamus		TgAPP751 mice (12 months with Aβ plaques)	Willis et al., 2007
Substance P		↑	CSF	Late onset AD patient		Rösler et al., 2001; Johansson et al., 2015
Orexin		↓	Hypothalamus	AD patient		Fronczek et al., 2012
Orexin		↑	CSF	AD patient		Liguori et al., 2016; Osorio et al., 2016; Gabelle et al., 2017
Orexin precursor gene		↑	Hypothalamus		APP/PS1dE9 mice (12–15 months)	Ma et al., 2016
	OX1R, OX2R	↓	Hippocampus, Aβ42 treated SH-SY5Y cells	AD patient	Aβ plaques and tau phosphorylation	Davies et al., 2015

AD, Alzheimer's disease; CSF, cerebrospinal fluid; GHS-R, ghrelin G-protein coupled receptor; MCI, mild cognitive impairment; NTSR, neurotensin receptor; OXR, orexin receptor; PACAP, pituitary adenylate cyclase-activating polypeptide; PVN_h, paraventricular nucleus of the hypothalamus; SNc, substantia nigra zona compacta; VTA, ventral tegmental area.

TABLE 2 | Neuroprotective effects and possible mechanisms of neuropeptides in Alzheimer's disease.

Neuropeptides	Reagents or treatments	Concentrations and times used	Neuroprotective effects	Mechanisms	Cellular and animal models	References
Ghrelin	Ghrelin agonist: LY444711 Ghrelin	30 mg/kg/day (in a chocolate pill) for 4 months 0.1 and 1 μ M for 24 h	Improves cognition Growth-promoting effect on neuronal cells	Reduces A β and microglial inflammation in dentate gyrus; Impairs glucose tolerance immediately Induces GHS-R1 expression; Activate the proteasome; Deregulates autophagy	Tg APPSwdJ mice APP-transfected SH-SY5Y cells	Dhurandhar et al., 2013; Kunath et al., 2015 Cecairni et al., 2016
	Acyl-ghrelin or DES-acyl ghrelin	0.2 nmol/h for 3 weeks	Reverses impairments of cognition and energy and glucose metabolism	Suppresses A β deposition; Increases the phosphorylation of AMPK and GSK, decreases the phosphorylation of tau	Rats with i.c.v infusion of A β	Kang et al., 2015
	Acyl-ghrelin	0.3 mg/kg i.p. daily for 7 days	Prevents impairment of recognition and spatial orientation Rescues memory deficits	Blunts A β -induced depression of LTP in hippocampus	Mice with i.c.v infusion of A β	Santos et al., 2017
	Ghrelin	80 μ g/kg i.p. daily for 7 days	Rescues memory deficits	Decreases microgliosis in hippocampus; Attenuates hippocampal neuronal loss; Prevents synaptic degeneration including cholinergic fiber loss	Intrahippocampal injection of A β	Moon et al., 2011
	Ghrelin	80 μ g/kg i.p. every 2 days for 30 days	Ameliorates neurogenesis impairment in hippocampus Improves cell survival	N/A	5XFAD mice	Moon et al., 2014
	Ghrelin	0.1–0.5 μ M for 24 h	Augments neuronal glucose uptake Improves spatial learning and memory; Increases firing rate of AP in stellate neurons of EC Rescues the survival of aged neurons	Reduces superoxide production and mitochondrial membrane depolarization; Prevents GSK 3 β	A β 0 treated primary hippocampal neurons and hypothalamic N42 cell line	Martins et al., 2013; Gomes et al., 2014
Neurotensin	Neurotensin or NTS1 agonist: PD149163 Neurotensin	10 nM for 1 h 0.25 μ M microinjection or bath application 40 μ M for 2 h	Rescues impaired recognition; Stimulates the non-amyloidogenic processing of APP; Reduces A β ₄₀ and A β ₄₂	Increases Akt/GSK phosphorylation Inhibits TREK-2K channels via PLC/PKC pathway	Primary hippocampal neurons treated with glucose APP/PS1 mice	Chen et al., 2010 Xiao et al., 2014
PACAP	PACAP38	10 μ g daily for 3 months intranasal	Rescues impaired recognition; Stimulates the non-amyloidogenic processing of APP; Reduces A β ₄₀ and A β ₄₂	Blocks sortilin-mediated neurotoxic role of proNGF in old age; Increases proNGF expression in frontal cortex and hippocampus Enhances gene expression of α -secretases; Enhances A β -degrading enzyme neprilysin; Increases PACAP and PAC1 expression; Increases expression of BDNF; Increases expression of antiapoptotic Bcl-2	Aged mice BFN neurons; AD patients APPV717]-transgenic mice; SK-N-MC PAC1 cells (BDNF study)	Al-Shawi et al., 2007, 2008 Rat et al., 2011
	PACAP27	1 nM for 72 h	Rescues A β -induced cell death Enhances secretion of sAPP α	Increases cAMP formation; Decreases caspase-3 activity Stimulates MAPK pathway and PI3K	A β -treated PC12 cells SK-N-MC cells; HEK 293 cells	Onoue et al., 2002 Kojro et al., 2006

(Continued)

TABLE 2 | Continued

Neuropeptides	Reagents or treatments	Concentrations and times used	Neuroprotective effects	Mechanisms	Cellular and animal models	References
Neuropeptide Y	Neuropeptide Y	0.0234 $\mu\text{M}/\mu\text{L}$, i.c.v.	Prevents depressive-like behavior and spatial memory deficits Attenuates ER stress-induced cell death	Blunts A β -induced increase in lipid peroxidation in hippocampus and prefrontal cortex Decreases caspase-3 and -4 activities; Suppresses the activation of three major ER stress sensors; Activates PI3K-XBP1 pathway	Mice treated with i.c.v. A β ₁₋₄₀ SK-N-SH cells; Mouse cortical neurons	dos Santos et al., 2013b Lee et al., 2018
Neuropeptide Y	Neuropeptide Y	100 nM for 12h	Rescues A β -induced cell death	Increases NGF synthesis and restores NGF release; Increases the levels of BDNF via inhibiting miR-30-5p	A β ₂₅₋₃₅ -treated SH-SY5Y cells; A β ₂₅₋₃₅ -treated primary cortical neurons	Croce et al., 2012, 2013
Substance P	Amidated NPY CTFs (NPY21-36 and 31-36)	0.5, 1, and 2 μM for 24 h	Ameliorates neurodegenerative pathology; Protects human neuronal cultures	NPY CTFs generated during NEP-mediated proteolysis exerts neuroprotective effects <i>in vivo</i>	APP tg mice; A β ₁₋₄₂ -treated primary human cortical neurons	Rose et al., 2009
Substance P	Substance P	120 μM i.c.v. for 28 days <i>in vivo</i> ; 10 nM for 24 h <i>in vitro</i>	Prevents cognitive impairments	Reduces A β -induced overexpression of Kv1.4 in hippocampus and cerebral cortex	Rats with i.c.v. infusion of A β ₂₅₋₃₅	Campolongo et al., 2013
Orexin	Orexin-A Dual orexin receptor antagonist: almorexant	200 nM	Reverses cell death; Reverses A β -induced increase of I_{K_A} current	Inhibits caspase-3-induced PAPP-1 cleavage through Akt-dependent mechanism; Prevents A β -induced upregulation of Kv4.2 and Kv4.3 Enhances α -secretase activity; Increases sAPP α ; Reduces A β ₁₋₄₂	A β -treated rat cerebellar granule cells	Pieri et al., 2010
Orexin	Orexin-A Dual orexin receptor antagonist: almorexant	200 nM	Reverses K ⁺ -induced apoptotic cell death and amyloidogenic processing of APP		Rat cerebellar granule cells	Marolda et al., 2012
Orexin	Orexin-A Dual orexin receptor antagonist: almorexant	1.5 pM i.c.v. for 6 h 13.9 nM i.c.v. for 24 h i.p. for 8 weeks	Increases ISF A β levels; Suppresses ISF A β levels; Reduces A β plaque deposition in the cortex	N/A	Human APP transgenic (Tg2576) mice; APPsw/PS1dE9 mice	Kang et al., 2009
Orexin	Orexin-A; Orexin-B	1 μM for 24 h	Suppresses the uptake of A β ; Reduces the degradation of A β	Downregulates phagocytosis regulating molecules: PI3K, Akt and p38-MAPK; Interferes autophagosome-lysosome fusion, and then suppresses autophagic flux	A β -treated BV2 microglial cells	An et al., 2017
Orexin	Orexin-A; Orexin-B	100 nM for 24h 100 nM for 1h	Augments NF- κ B signaling (OXA); Up-regulates PI3K-Akt and Jak-STAT signaling (OXB); Induces ERK1/2 phosphorylation	NF- κ B regulates cell survival, neurogenesis, learning and memory; PI3K-Akt is involved in neuroprotective functions; ERK is a key molecule in prevention of neurodegeneration	SH-SY5Y cells treated with retinoic acid (microarray); HEK293 cells with OXRs and GPR103 form functional heterodimers	Davies et al., 2015

A β O, amyloid- β oligomers; AD, Alzheimer's disease; AMPK, adenosine 5'-monophosphate (AMP)-activated protein kinase; AP, action potential; APP, amyloid precursor protein; BDNF, brain-derived neurotrophic factor; BRN1, basal forebrain nuclei; CTFs, C-terminal fragments; EC, entorhinal cortex; ER, endoplasmic reticulum; GHS-R1, growth hormone secretagogue receptor type 1; GSK, glycogen synthase kinase; i.c.v., intracerebroventricular; I κ A current, A-type K⁺ current; ISF, interstitial fluid; LTP, long-term potentiation; MAPK, mitogen-activated protein kinase; NEP, neutral endopeptidase; NGF, nerve growth factor; NPY, neuropeptide Y; NTS1, neurotensin receptor 1; OXA, orexin-A; OXB, orexin-B; PACAP, pituitary adenylylate cyclase-activating polypeptide; PAPP-1, poly ADP-ribose polymerase-1; PI3K, phosphatidylinositol 3-kinase; sAPP α , soluble amyloid precursor protein α ; SK-N-MC PAC1 cells, human neuroblastoma SK-N-MC cells overexpressing PAC1 receptor.

Early studies revealed that the expression of substance P changes in different brain regions of Alzheimer's disease. The level of substance P decreases in the cortex, hippocampus and striatum in Alzheimer's disease patients and animal models (Bouras et al., 1990; Quigley and Kowall, 1991; Nag et al., 1999; Ahmed et al., 2004), but increases in the pallidum and substantia nigra (Bouras et al., 1990). However, Willis et al. (2007) showed that in old age of Alzheimer's disease transgenic mice, the substance P-immunoreactivity exists in astrocytes of hippocampus and thalamus. In late onset Alzheimer's disease patients, the level of substance P in cerebrospinal fluid increases significantly (Rösler et al., 2001). Furthermore, the increased level of substance P is positively associated with the level of A β _{1–42} in Alzheimer's disease patients (Johansson et al., 2015). In addition to the change of substance P, the activity of neuropeptidases to metabolize substance P also alters in Alzheimer's disease. Waters and David (1995) demonstrated that the activity of neuropeptidases is decreased in the temporal cortex of senile dementia of the Alzheimer's disease, which therefore increases the metabolic half-life of substance P.

Substance P exerts neuroprotective effects in a variety of *in vitro* and *in vivo* studies (Figure 5). Potassium channel dysfunction is a possible mechanism underlying the pathophysiology of Alzheimer's disease. Some kinds of voltage-gated potassium channels are involved in substance P-induced neuroprotective effects. Application of substance P prevents A β -induced impairment of cognitive processes through inhibition of A β -induced overexpression of potassium channel subunits (Campolongo et al., 2013), as well as A β -induced enhancement of A-type K⁺ currents (Pieri et al., 2010). Substance P also protects cerebellar granule cells against A β -induced apoptosis through inhibition of caspase-3-induced PARP-1 cleavage (Pieri et al., 2010). In addition, substance P plays non-amyloidogenic effect through decreasing A β _{1–42}, increasing sAPP α and α -secretase activity (Marolda et al., 2012). A β _{25–35} reduces the expression of substance P in hippocampus before the neuronal loss of Alzheimer's disease. Memantine, a non-competitive NMDA receptor antagonist, attenuates A β _{25–35}-induced decrease of substance P (Arif et al., 2009). Furthermore, in ibotenic acid-treated Alzheimer's disease model, memantine treatment recovers the decreased substance P expression (Ahmed et al., 2004). Recently, Fernandes et al. (2018) reported that a substance P receptor antagonist attenuates aluminum-induced spatial memory deficit probably through blockade of substance P-mediated neuroinflammation. It is known that the dysfunction of metal ions, such as copper is a feature of Alzheimer's disease. Neurokinin B protects against copper-induced calcium channel opening (Russino et al., 2013), as well as the synaptic homeostasis (Grosas et al., 2014).

In summary, the levels of substance P decrease in brain regions including cortex and hippocampus, but increase in cerebrospinal fluid of late onset Alzheimer's disease patients. Substance P exerts neuroprotective effects through inhibition of A β -induced over-expression and activity of K⁺ channel. Furthermore, substance P inhibits caspase-3-induced PARP-1 cleavage and increases α -secretase activity, and is therefore involved in neuroprotection in Alzheimer's disease.

OREXIN AND ALZHEIMER'S DISEASE

Orexin is a hypothalamic neuropeptide which plays an important role in maintaining wakefulness. It is well-known that sleep disturbances are common clinical symptom in neurodegenerative disorders. Recently people focus on the involvement of orexinergic system in the pathophysiology, especially sleep disturbances, of Alzheimer's disease (Ferini-Strambi, 2014; Malkki, 2014; Liguori, 2017; Liguori et al., 2017). Early morphological study revealed that the orexinergic neurons are reduced significantly in postmortem hypothalamus of Alzheimer's disease patients (Fronczek et al., 2012). However, it is demonstrated recently that the level of orexin-A in cerebrospinal fluid is higher in Alzheimer's disease patients which is positively associated with Alzheimer's disease biomarkers including A β and Tau (Liguori et al., 2016; Osorio et al., 2016; Gabelle et al., 2017). In both Alzheimer's disease patients and A β ₄₂ treated-SH-SY5Y cells, amyloid deposition and tau phosphorylation decrease the expression of orexin-1 and orexin-2 receptors (Davies et al., 2015). *In vivo* microdialysis revealed that infusion of orexin-A increases the amount of A β in brain interstitial fluid, while orexin receptor antagonist suppresses A β levels and reduces A β plaque deposition in the cortex (Kang et al., 2009). In A β -treated microglial cells, both orexin-A and orexin-B reduce the uptake of A β through downregulation of phagocytosis regulating molecules including PI3K, Akt and p38-MAPK. In addition, orexins suppress autophagosome-lysosome fusion and lead to impaired A β degradation (An et al., 2017). Furthermore, the orexin receptors 2 gene, rs2653349 polymorphism, is likely to be a risk factor of Alzheimer's disease (Gallone et al., 2014). In APP/PS1 transgenic Alzheimer's disease mice, orexin gene knock out markedly decreases the amount of A β pathology, while rescue of orexinergic neurons increases the amount of A β pathology in the brain (Roh et al., 2014). Recent study revealed that orexinergic system modulates both the hippocampal clock and clock-controlled-genes, Bace1 and Bace2. Both the two genes are correlated with the production of A β (Ma et al., 2016).

In addition to the inhibition of A β uptake and degradation in microglial cells (An et al., 2017), orexin may also exert neuroprotective effects. Orexinergic neuron degeneration impairs long-term social memory in mice (Yang et al., 2013). Intracerebroventricular microinjection of orexin-A improves memory in SAMP8 Alzheimer disease mice (Jaeger et al., 2002). Orexin receptors have been demonstrated to exert neuroprotective effects in Alzheimer's disease via heterodimerization with GPR103. ERK is a key molecule in the prevention of neurodegeneration. Treatment with orexins induces ERK_{1/2} phosphorylation suggesting the neuroprotection of the heterodimerization. Microarray showed that orexin-A augments NF- κ B signaling and orexin-B up-regulates PI3K-Akt and Jak-STAT signaling, respectively (Davies et al., 2015).

The hippocampus is a crucial brain region that plays important roles in learning and memory. Intra-hippocampal administration of orexin-A attenuates pain-induced impairment of learning and memory (Raouf et al., 2015), while blocking orexin receptors in hippocampal CA1 regions impairs spatial

memory retrieval (Akbari et al., 2006, 2008). It has been demonstrated that activation of cholinergic transmission increases the neuronal excitability of CA1 pyramidal neurons and therefore enhances hippocampus-dependent learning in Alzheimer disease animal model (Disterhoft and Oh, 2006). Moreover, decreasing the firing rate of CA1 pyramidal neurons by potassium channel activator impairs associative learning in rats (McKay et al., 2012). Therefore, our recent study that orexin-A increases the firing rate of hippocampal CA1 neurons may provide direct *in vivo* electrophysiological evidence for the possible involvement of orexin-A in Alzheimer disease (Chen et al., 2017).

In summary, A β deposition and tau phosphorylation decrease the expression of orexin and the receptors in hypothalamus, while the levels of orexins in cerebrospinal fluid increase in Alzheimer's disease patients. In microglial cells, orexins increase A β level through suppression of A β uptake and degradation. However, in cells forming OXRs and GPR103 heterodimers, orexins exert neuroprotective effects through upregulation of NF- κ B, PI3K-Akt, Jak-STAT signaling, as well as ERK $_{1/2}$ phosphorylation. **Figure 6** illustrated the complicated effects of orexin in Alzheimer's disease.

CONCLUSION

In this review article, we provide a description of recent advances of neuropeptides including ghrelin, neurotensin, PACAP, neuropeptide Y, substance P and orexin in Alzheimer's disease. Based on current studies, the levels of these neuropeptides and their receptors change in Alzheimer's disease (**Table 1**).

REFERENCES

- Ahmed, M. M., Hoshino, H., Chikuma, T., Yamada, M., and Kato, T. (2004). Effect of memantine on the levels of glial cells, neuropeptides, and peptide-degrading enzymes in rat brain regions of ibotenic acid-treated Alzheimer's disease model. *Neuroscience* 126, 639–649. doi: 10.1016/j.neuroscience.2004.04.024
- Akbari, E., Motamedi, F., Naghdi, N., and Noorbakhshnia, M. (2008). The effect of antagonization of orexin 1 receptors in CA1 and dentate gyrus regions on memory processing in passive avoidance task. *Behav. Brain Res.* 187, 172–177. doi: 10.1016/j.bbr.2007.09.019
- Akbari, E., Naghdi, N., and Motamedi, F. (2006). Functional inactivation of orexin 1 receptors in CA1 region impairs acquisition, consolidation and retrieval in Morris water maze task. *Behav. Brain Res.* 173, 47–52. doi: 10.1016/j.bbr.2006.05.028
- Al-Shawi, R., Hafner, A., Chun, S., Raza, S., Crutcher, K., Thrasivoulou, C., et al. (2007). ProNGF, sortilin, and age-related neurodegeneration. *Ann. N. Y. Acad. Sci.* 1119, 208–215. doi: 10.1196/annals.1404.024
- Al-Shawi, R., Hafner, A., Olsen, J., Chun, S., Raza, S., Thrasivoulou, C., et al. (2008). Neurotoxic and neurotrophic roles of proNGF and the receptor sortilin in the adult and ageing nervous system. *Eur. J. Neurosci.* 27, 2103–2114. doi: 10.1111/j.1460-9568.2008.06152.x
- An, H., Cho, M. H., Kim, D. H., Chung, S., and Yoon, S. Y. (2017). Orexin impairs the phagocytosis and degradation of amyloid- β fibrils by microglial cells. *J. Alzheimers Dis.* 58, 253–261. doi: 10.3233/JAD-170108
- Angelucci, F., Gelfo, F., Fiore, M., Croce, N., Mathé, A. A., Bernardini, S., et al. (2014). The effect of neuropeptide Y on cell survival and neurotrophin expression in *in-vitro* models of Alzheimer's disease. *Can. J. Physiol. Pharmacol.* 92, 621–630. doi: 10.1139/cjpp-2014-0099
- Neuropeptides exert significant neuroprotective effects against A β -induced neuronal toxicity (**Table 2**). Multiple intracellular mechanisms, including activation of α -secretases and A β -degrading enzyme, production of neurotrophins, increase of neuronal glucose uptake, upregulation of NF- κ B, PI3K-Akt, MAPK, Jak-STAT, ERK $_{1/2}$ phosphorylation and downregulation of caspase-3, inhibition of endoplasmic reticulum stress and autophagy, modulation of LTP and potassium channel activity, appear to be involved in neuropeptide-induced neuroprotection in Alzheimer's disease. As the levels of neuropeptides and the receptors may change before A β deposition and neuronal loss, as well as positively/negatively correlate with cognitive impairment, future studies would focus on detection of neuropeptides as biomarkers in early diagnosis and treatment evaluation of Alzheimer's disease. Furthermore, for the notable neuroprotective effects, further researches are needed to explore the potential use of neuropeptides, especially via convenient administration method like intranasal application, in the prevention and cure of Alzheimer's disease.

AUTHOR CONTRIBUTIONS

X-YC wrote the manuscript. Y-FD and LC revised the manuscript.

FUNDING

This work was supported by grants from the National Natural Science Foundation of China (31671076, 81772448, 81200872), and Taishan Scholars Construction Project.

- Arif, M., Chikuma, T., Ahmed, M. M., Nakazato, M., Smith, M. A., and Kato, T. (2009). Effects of memantine on soluble Alpha(25-35)-induced changes in peptidergic and glial cells in Alzheimer's disease model rat brain regions. *Neuroscience* 164, 1199–1209. doi: 10.1016/j.neuroscience.2009.08.063
- Beach, T. G., Honer, W. G., and Hughes, L. H. (1997). Cholinergic fibre loss associated with diffuse plaques in the non-demented elderly: the preclinical stage of Alzheimer's disease? *Acta Neuropathol.* 93, 146–153. doi: 10.1007/s004010050595
- Benzing, W. C., Mufson, E. J., and Armstrong, D. M. (1993). Immunocytochemical distribution of peptidergic and cholinergic fibers in the human amygdala: their depletion in Alzheimer's disease and morphologic alteration in non-demented elderly with numerous senile plaques. *Brain Res.* 625, 125–138. doi: 10.1016/0006-8993(93)90145-d
- Benzing, W. C., Mufson, E. J., Jennes, L., and Armstrong, D. M. (1990). Reduction of neurotensin immunoreactivity in the amygdala in Alzheimer's disease. *Brain Res.* 537, 298–302. doi: 10.1016/0006-8993(90)90372-1
- Benzing, W. C., Mufson, E. J., Jennes, L., Stopa, E. G., and Armstrong, D. M. (1992). Distribution of neurotensin immunoreactivity within the human amygdaloid complex: a comparison with acetylcholinesterase- and Nissl-stained tissue sections. *J. Comp. Neurol.* 317, 283–297. doi: 10.1002/cne.903170306
- Biggins, J. A., Perry, E. K., McDermott, J. R., Smith, A. I., Perry, R. H., and Edwardson, J. A. (1983). Post mortem levels of thyrotropin-releasing hormone and neurotensin in the amygdala in Alzheimer's disease, schizophrenia and depression. *J. Neurol. Sci.* 58, 117–122. doi: 10.1016/0022-510X(83)90114-4
- Bissette, G., Nemeroff, C. B., Decker, M. W., Kizer, J. S., Agid, Y., and Javoy-Agid, F. (1985). Alterations in regional brain concentrations of neurotensin and bombesin in Parkinson's disease. *Ann. Neurol.* 17, 324–328. doi: 10.1002/ana.410170403

- Bouras, C., Vallet, P. G., Hof, P. R., Charnay, Y., Golaz, J., and Constantinidis, J. (1990). Substance P immunoreactivity in Alzheimer disease: a study in cases presenting symmetric or asymmetric cortical atrophy. *Alzheimer Dis. Assoc. Disord.* 4, 24–34. doi: 10.1097/00002093-19904010000003
- Burns, A., and Liffle, S. (2009). Alzheimer's disease. *BMJ* 338:b158. doi: 10.1136/bmj.b158
- Campolongo, P., Ratano, P., Ciotti, M. T., Florenzano, F., Nori, S. L., Marolda, R., et al. (2013). Systemic administration of substance P recovers beta amyloid-induced cognitive deficits in rat: involvement of Kv potassium channels. *PLoS ONE* 8:e78036. doi: 10.1371/journal.pone.0078036
- Cao, X., Zhu, M., He, Y., Chu, W., Du, Y., and Du, H. (2018). Increased serum acylated ghrelin levels in patients with mild cognitive impairment. *J. Alzheimers Dis.* 61, 545–552. doi: 10.3233/JAD-170721
- Cecarini, V., Bonfili, L., Cuccioloni, M., Keller, J. N., Bruce-Keller, A. J., and Eleuteri, A. M. (2016). Effects of ghrelin on the proteolytic pathways of Alzheimer's disease neuronal cells. *Mol. Neurobiol.* 53, 3168–3178. doi: 10.1007/s12035-015-9227-x
- Chen, X. Y., Chen, L., and Du, Y. F. (2017). Orexin-A increases the firing activity of hippocampal CA1 neurons through orexin-1 receptors. *J. Neurosci. Res.* 95, 1415–1426. doi: 10.1002/jnr.23975
- Chen, Y., Cao, C. P., Li, C. R., Wang, W., Zhang, D., Han, L. L., et al. (2010). Ghrelin modulates insulin sensitivity and tau phosphorylation in high glucose-induced hippocampal neurons. *Biol. Pharm. Bull.* 33, 1165–1169. doi: 10.1248/bpb.33.1165
- Constantinidis, J., Bouras, C., and Richard, J. (1983). Putative peptide neurotransmitters in human neuropathology: a review of topography and clinical implications. *Clin. Neuropathol.* 2, 47–54.
- Croce, N., Ciotti, M. T., Gelfo, F., Cortelli, S., Federici, G., Caltagirone, C., et al. (2012). Neuropeptide Y protects rat cortical neurons against β -amyloid toxicity and re-establishes synthesis and release of nerve growth factor. *ACS Chem. Neurosci.* 3, 312–318. doi: 10.1021/cn200127e
- Croce, N., Dinallo, V., Ricci, V., Federici, G., Caltagirone, C., Bernardini, S., et al. (2011). Neuroprotective effect of neuropeptide Y against β -amyloid 25–35 toxicity in SH-SY5Y neuroblastoma cells is associated with increased neurotrophin production. *Neurodegener. Dis.* 8, 300–309. doi: 10.1159/000323468
- Croce, N., Gelfo, F., Ciotti, M. T., Federici, G., Caltagirone, C., Bernardini, S., et al. (2013). NPY modulates miR-30a-5p and BDNF in opposite direction in an *in vitro* model of Alzheimer disease: a possible role in neuroprotection? *Mol. Cell Biochem.* 376, 189–195. doi: 10.1007/s11010-013-1567-0
- Davies, J., Chen, J., Pink, R., Carter, D., Saunders, N., Sotiropoulos, G., et al. (2015). Orexin receptors exert a neuroprotective effect in Alzheimer's disease (AD) via heterodimerization with GPR103. *Sci. Rep.* 5:12584. doi: 10.1038/srep12584
- Dhurandhar, E. J., Allison, D. B., van Groen, T., and Kadish, I. (2013). Hunger in the absence of caloric restriction improves cognition and attenuates Alzheimer's disease pathology in a mouse model. *PLoS ONE* 8:e60437. doi: 10.1371/journal.pone.0060437
- Diez, M., Danner, S., Frey, P., Sommer, B., Staufienbiel, M., Wiederhold, K. H., et al. (2003). Neuropeptide alterations in the hippocampal formation and cortex of transgenic mice overexpressing beta-amyloid precursor protein (APP) with the Swedish double mutation (APP23). *Neurobiol. Dis.* 14, 579–594. doi: 10.1016/j.nbd.2003.08.003
- Disterhoft, J. F., and Oh, M. M. (2006). Pharmacological and molecular enhancement of learning in aging and Alzheimer's disease. *J. Physiol. Paris* 99, 180–192. doi: 10.1016/j.jphysparis.2005.12.079
- dos Santos, V. V., Rodrigues, A. L., De Lima, T. C., de Bariogio, S. R., Raisman-Vozari, R., and Prediger, R. D. (2013a). Ghrelin as a neuroprotective and palliative agent in Alzheimer's and Parkinson's disease. *Curr. Pharm. Des.* 19, 6773–6790. doi: 10.2174/13816128113199990411
- dos Santos, V. V., Santos, D. B., Lach, G., Rodrigues, A. L., Farina, M., De Lima, T. C., et al. (2013b). Neuropeptide Y (NPY) prevents depressive-like behavior, spatial memory deficits and oxidative stress following amyloid- β [A β (1–40)] administration in mice. *Behav. Brain Res.* 244, 107–115. doi: 10.1016/j.bbr.2013.01.039
- Duarte-Neves, J., Pereira de Almeida, L., and Cavadas, C. (2016). Neuropeptide Y (NPY) as a therapeutic target for neurodegenerative diseases. *Neurobiol. Dis.* 95, 210–224. doi: 10.1016/j.nbd.2016.07.022
- Ferini-Strambi, L. (2014). Possible role of orexin in the pathogenesis of Alzheimer disease. *JAMA Neurol.* 71, 1478–1480. doi: 10.1001/jamaneurol.2014.2819
- Fernandes, J., Mudgal, J., Rao, C. M., Arora, D., Basu Mallik, S., Pai, K. S. R., et al. (2018). N-acetyl-L-tryptophan, a substance-P receptor antagonist attenuates aluminum-induced spatial memory deficit in rats. *Toxicol. Mech. Methods* 28, 328–334. doi: 10.1080/15376516.2017.1411412
- Fernandez, A., de Ceballos, M. L., Jenner, P., and Marsden, C. D. (1994). Neurotensin, substance P, delta and mu opioid receptors are decreased in basal ganglia of Parkinson's disease patients. *Neuroscience* 61, 73–79. doi: 10.1016/0306-4522(94)90061-2
- Ferrier, I. N., Cross, A. J., Johnson, J. A., Roberts, G. W., Crow, T. J., Corsellis, J. A., et al. (1983). Neuropeptides in Alzheimer type dementia. *J. Neurol. Sci.* 62, 159–170. doi: 10.1016/0022-510X(83)90196-X
- Fronczek, R., van Geest, S., Frölich, M., Overeem, S., Roelandse, F. W., Lammers, G. J., et al. (2012). Hypocretin (orexin) loss in Alzheimer's disease. *Neurobiol. Aging* 33, 1642–1650. doi: 10.1016/j.neurobiolaging.2011.03.014
- Gabelle, A., Jaussent, I., Hirtz, C., Vialaret, J., Navucet, S., Grasselli, C., et al. (2017). Cerebrospinal fluid levels of orexin-A and histamine, and sleep profile within the Alzheimer process. *Neurobiol. Aging* 53, 59–66. doi: 10.1016/j.neurobiolaging.2017.01.011
- Gahete, M. D., Córdoba-Chacón, J., Kineman, R. D., Luque, R. M., and Castaño, J. P. (2011). Role of ghrelin system in neuroprotection and cognitive functions: implications in Alzheimer's disease. *Peptides* 32, 2225–2228. doi: 10.1016/j.peptides.2011.09.019
- Gahete, M. D., Rubio, A., Córdoba-Chacón, J., Gracia-Navarro, F., Kineman, R. D., Avila, J., et al. (2010). Expression of the ghrelin and neurotensin systems is altered in the temporal lobe of Alzheimer's disease patients. *J. Alzheimers Dis.* 22, 819–828. doi: 10.3233/JAD-2010-100873
- Gallone, S., Boschi, S., Rubino, E., De Martino, P., Scarpini, E., Galimberti, D., et al. (2014). Is HCRTR2 a genetic risk factor for Alzheimer's disease? *Dement. Geriatr. Cogn. Disord.* 38, 245–253. doi: 10.1159/000359964
- Garver, D. L., Bissette, G., Yao, J. K., and Nemeroff, C. B. (1991). Relation of CSF neurotensin concentrations to symptoms and drug response of psychotic patients. *Am. J. Psychiatry* 148, 484–488. doi: 10.1176/ajp.148.4.484
- Giedd, J. N., Snell, J. W., Lange, N., Rajapakse, J. C., Casey, B. J., Kozuch, P. L., et al. (1996). Quantitative magnetic resonance imaging of human brain development: ages 4–18. *Cereb. Cortex* 6, 551–560. doi: 10.1093/cercor/6.4.551
- Goldstein, J., Seidman, L., Horton, N., Makris, N., Kennedy, D., Caviness, V., et al. (2001). Normal sexual dimorphism of the adult human brain assessed by *in vivo* magnetic resonance imaging. *Cereb. Cortex* 11, 490–497. doi: 10.1093/cercor/11.6.490
- Gomes, S., Martins, I., Fonseca, A. C., Oliveira, C. R., Resende, R., and Pereira, C. M. (2014). Protective effect of leptin and ghrelin against toxicity induced by amyloid- β oligomers in a hypothalamic cell line. *J. Neuroendocrinol.* 26, 176–185. doi: 10.1111/jne.12138
- Grosas, A. B., Kalimuthu, P., Smith, A. C., Williams, P. A., Millar, T. J., Bernhardt, P. V., et al. (2014). The tachykinin peptide neurokinin B binds copper(I) and silver(I) and undergoes quasi-reversible electrochemistry: towards a new function for the peptide in the brain. *Neurochem. Int.* 70, 1–9. doi: 10.1016/j.neuint.2014.03.002
- Guan, X. M., Yu, H., Palyha, O. C., McKee, K. K., Feighner, S. D., Sirinathsinghji, D. J., et al. (1997). Distribution of mRNA encoding the growth hormone secretagogue receptor in brain and peripheral tissues. *Brain Res. Mol. Brain Res.* 48, 23–29. doi: 10.1016/S0169-328X(97)00071-5
- Han, P., Liang, W., Baxter, L. C., Yin, J., Tang, Z., Beach, T. G., et al. (2014a). Pituitary adenylate cyclase-activating polypeptide is reduced in Alzheimer disease. *Neurology* 82, 1724–1728. doi: 10.1212/WNL.0000000000000417
- Han, P., Nielsen, M., Song, M., Yin, J., Permenter, M. R., Vogt, J. A., et al. (2017). The impact of aging on brain pituitary adenylate cyclase activating polypeptide, pathology and cognition in mice and rhesus macaques. *Front. Aging Neurosci.* 9:180. doi: 10.3389/fnagi.2017.00180
- Han, P., Tang, Z., Yin, J., Maalouf, M., Beach, T. G., Reiman, E. M., et al. (2014b). Pituitary adenylate cyclase-activating polypeptide protects against β -amyloid toxicity. *Neurobiol. Aging* 35, 2064–2071. doi: 10.1016/j.neurobiolaging.2014.03.022
- Hannibal, J. (2002). Pituitary adenylate cyclase-activating peptide in the rat central nervous system: an immunohistochemical and *in situ* hybridization study. *J. Comp. Neurol.* 453, 389–417. doi: 10.1002/cne.10418

- Hsieh, Y. S., Chen, P. N., Yu, C. H., Liao, J. M., and Kuo, D. Y. (2013). The neuropeptide Y Y1 receptor knockdown modulates activator protein 1-involved feeding behavior in amphetamine-treated rats. *Mol. Brain* 6:46. doi: 10.1186/1756-6606-6-46
- Hu, K., Harper, D. G., Shea, S. A., Stopa, E. G., and Scheer, F. A. (2013). Noninvasive fractal biomarker of clock neurotransmitter disturbance in humans with dementia. *Sci. Rep.* 3:2229. doi: 10.1038/srep02229
- Ishii, M., Wang, G., Racchumi, G., Dyke, J. P., and Iadecola, C. (2014). Transgenic mice overexpressing amyloid precursor protein exhibit early metabolic deficits and a pathologically low leptin state associated with hypothalamic dysfunction in arcuate neuropeptide Y neurons. *J. Neurosci.* 34, 9096–9106. doi: 10.1523/jneurosci.0872-14.2014
- Jaeger, L. B., Farr, S. A., Banks, W. A., and Morley, J. E. (2002). Effects of orexin-A on memory processing. *Peptides* 23, 1683–1688. doi: 10.1016/s0196-9781(02)00110-9
- Jansen, K. L., Faull, R. L., Dragunow, M., and Synek, B. L. (1990). Alzheimer's disease: changes in hippocampal N-methyl-D-aspartate, quisqualate, neurotensin, adenosine, benzodiazepine, serotonin and opioid receptors—an autoradiographic study. *Neuroscience* 39, 613–627. doi: 10.1016/0306-4522(90)90246-Z
- Jiao, Q., Du, X., Li, Y., Gong, B., Shi, L., Tang, T., et al. (2017). The neurological effects of ghrelin in brain diseases: Beyond metabolic functions. *Neurosci. Biobehav. Rev.* 73, 98–111. doi: 10.1016/j.neubiorev.2016.12.010
- Johansson, P., Almqvist, E. G., Wallin, A., Johansson, J. O., Andreasson, U., Blennow, K., et al. (2015). Cerebrospinal fluid substance P concentrations are elevated in patients with Alzheimer's disease. *Neurosci. Lett.* 609, 58–62. doi: 10.1016/j.neulet.2015.10.006
- Kang, J. E., Lim, M. M., Bateman, R. J., Lee, J. J., Smyth, L. P., Cirrito, J. R., et al. (2009). Amyloid-beta dynamics are regulated by orexin and the sleep-wake cycle. *Science* 326, 1005–1007. doi: 10.1126/science.1180962
- Kang, S., Moon, N. R., Kim, D. S., Kim, S. H., and Park, S. (2015). Central acylated ghrelin improves memory function and hippocampal AMPK activation and partly reverses the impairment of energy and glucose metabolism in rats infused with β -amyloid. *Peptides* 71, 84–93. doi: 10.1016/j.peptides.2015.07.005
- Kincaid, B., and Nemeroff, C. B. (2004). Neurotensin, schizophrenia, and antipsychotic drug action. *Int. Rev. Neurobiol.* 59, 327–349. doi: 10.1016/S0074-7742(04)59013-X
- Kojro, E., Postina, R., Buro, C., Meiringer, C., Gehrig-Burger, K., and Fahrenholz, F. (2006). The neuropeptide PACAP promotes the alpha-secretase pathway for processing the Alzheimer amyloid precursor protein. *FASEB J.* 20, 512–514. doi: 10.1096/fj.05-4812je
- Krezymin, A., Richetin, K., Halley, H., Roybon, L., Lassalle, J. M., Francès, B., et al. (2013). Modifications of hippocampal circuits and early disruption of adult neurogenesis in the tg2576 mouse model of Alzheimer's disease. *PLoS ONE* 8:e76497. doi: 10.1371/journal.pone.0076497
- Kunath, N., van Groen, T., Allison, D. B., Kumar, A., Dozier-Sharp, M., and Kadish, I. (2015). Ghrelin agonist does not foster insulin resistance but improves cognition in an Alzheimer's disease mouse model. *Sci. Rep.* 5:11452. doi: 10.1038/srep11452
- Lacor, P. N., Buniel, M. C., Furlow, P. W., Clemente, A. S., Velasco, P. T., Wood, M., et al. (2007). Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J. Neurosci.* 27, 796–807. doi: 10.1523/jneurosci.3501-06.2007
- Lattuada, D., Crotta, K., Tonna, N., Casnici, C., Benfante, R., Fornasari, D., et al. (2013). The expression of GHS-R in primary neurons is dependent upon maturation stage and regional localization. *PLoS ONE* 8:e64183. doi: 10.1371/journal.pone.0064183
- Lee, D. Y., Hong, S. H., Kim, B., Lee, D. S., Yu, K., and Lee, K. S. (2018). Neuropeptide Y mitigates ER stress-induced neuronal cell death by activating the PI3K-XBP1 pathway. *Eur. J. Cell Biol.* 97, 339–348. doi: 10.1016/j.ejcb.2018.04.003
- Lee, E. H., and Seo, S. R. (2014). Neuroprotective roles of pituitary adenylate cyclase-activating polypeptide in neurodegenerative diseases. *BMB Rep.* 47, 369–375. doi: 10.5483/BMBRep.2014.47.7.086
- Liguori, C. (2017). Orexin and Alzheimer's disease. *Curr. Top. Behav. Neurosci.* 33, 305–322. doi: 10.1007/7854_2016_50
- Liguori, C., Chiaravalloti, A., Nuccetelli, M., Izzi, F., Sancesario, G., Cimini, A., et al. (2017). Hypothalamic dysfunction is related to sleep impairment and CSF biomarkers in Alzheimer's disease. *J. Neurol.* 264, 2215–2223. doi: 10.1007/s00415-017-8613-x
- Liguori, C., Nuccetelli, M., Izzi, F., Sancesario, G., Romigi, A., Martorana, A., et al. (2016). Rapid eye movement sleep disruption and sleep fragmentation are associated with increased orexin-A cerebrospinal-fluid levels in mild cognitive impairment due to Alzheimer's disease. *Neurobiol. Aging* 40, 120–126. doi: 10.1016/j.neurobiolaging.2016.01.007
- Ma, Z., Jiang, W., and Zhang, E. E. (2016). Orexin signaling regulates both the hippocampal clock and the circadian oscillation of Alzheimer's disease-risk genes. *Sci. Rep.* 6:36035. doi: 10.1038/srep36035
- Mahar, I., Albuquerque, M. S., Mondragon-Rodriguez, S., Cavanagh, C., Davoli, M. A., Chabot, J. G., et al. (2017). Phenotypic alterations in hippocampal NPY- and PV-expressing interneurons in a presymptomatic transgenic mouse model of Alzheimer's disease. *Front. Aging Neurosci.* 8:327. doi: 10.3389/fnagi.2016.00327
- Malkki, H. (2014). Alzheimer disease: increased orexin level correlates with sleep disruption and cognitive decline in Alzheimer disease. *Nat. Rev. Neurol.* 10:672. doi: 10.1038/nrneuro.2014.209
- Mann, D. M. (1989). The pathogenesis and progression of the pathological changes of Alzheimer's disease. *Ann. Med.* 21, 133–136.
- Mantyh, P. W. (2002). Neurobiology of substance P and the NK1 receptor. *J. Clin. Psychiatry* 63 (Suppl. 11), 6–10.
- Marolda, R., Ciotti, M. T., Matrone, C., Possenti, R., Calissano, P., Cavallaro, S., et al. (2012). Substance P activates ADAM9 mRNA expression and induces α -secretase-mediated amyloid precursor protein cleavage. *Neuropharmacology* 62, 1954–1963. doi: 10.1016/j.neuropharm.2011.12.025
- Martins, I., Gomes, S., Costa, R. O., Otvos, L., Oliveira, C. R., Resende, R., et al. (2013). Leptin and ghrelin prevent hippocampal dysfunction induced by A β oligomers. *Neuroscience* 241, 41–51. doi: 10.1016/j.neuroscience.2013.02.062
- McKay, B. M., Oh, M. M., Galvez, R., Burgdorf, J., Kroes, R. A., and Weiss, C. (2012). Increasing SK2 channel activity impairs associative learning. *J. Neurophysiol.* 108, 863–870. doi: 10.1152/jn.00025.2012
- Mittapalli, G. K., and Roberts, E. (2014). Ligands of the neuropeptide Y Y2 receptor. *Bioorg. Med. Chem. Lett.* 24, 430–441. doi: 10.1016/j.bmcl.2013.11.061
- Miyata, A., Arimura, A., Dahl, R. R., Minamino, N., Uehara, A., Jiang, L., et al. (1989). Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochem. Biophys. Res. Commun.* 164, 567–574. doi: 10.1016/0006-291X(89)91757-9
- Moon, M., Cha, M. Y., and Mook-Jung, I. (2014). Impaired hippocampal neurogenesis and its enhancement with ghrelin in 5XFAD mice. *J. Alzheimers Dis.* 41, 233–241. doi: 10.3233/jad-132417
- Moon, M., Choi, J. G., Nam, D. W., Hong, H. S., Choi, Y. J., Oh, M. S., et al. (2011). Ghrelin ameliorates cognitive dysfunction and neurodegeneration in intrahippocampal amyloid- β 1-42 oligomer-injected mice. *J. Alzheimers Dis.* 23, 147–159. doi: 10.3233/jad-2010-101263
- Murray, S., Tulloch, A., Gold, M. S., and Avena, N. M. (2014). Hormonal and neural mechanisms of food reward, eating behaviour and obesity. *Nat. Rev. Endocrinol.* 10, 540–552. doi: 10.1038/nrendo.2014.91
- Nag, S., Yee, B. K., and Tang, F. (1999). Reduction in somatostatin and substance P levels and choline acetyltransferase activity in the cortex and hippocampus of the rat after chronic intracerebroventricular infusion of beta-amyloid (1-40). *Brain Res. Bull.* 50, 251–262. doi: 10.1016/S0361-9230(99)00196-3
- Onoue, S., Endo, K., Ohshima, K., Yajima, T., and Kashimoto, K. (2002). The neuropeptide PACAP attenuates beta-amyloid (1-42)-induced toxicity in PC12 cells. *Peptides* 23, 1471–1478. doi: 10.1016/S0196-9781(02)00085-2
- Osorio, R. S., Ducca, E. L., Wohlleber, M. E., Tanzi, E. B., Gumb, T., Twumasi, A., et al. (2016). Orexin-A is associated with increases in cerebrospinal fluid phosphorylated-tau in cognitively normal elderly subjects. *Sleep* 39, 1253–1260. doi: 10.5665/sleep.5846
- Panagopoulos, V. N., and Ralevski, E. (2014). The role of ghrelin in addiction: a review. *Psychopharmacology (Berl.)* 231, 2725–2740. doi: 10.1007/s00213-014-3640-0
- Pérez-Fernández, J., Megias, M., and Pombal, M. A. (2014). Cloning, phylogeny, and regional expression of a Y5 receptor mRNA in the brain of the sea lamprey (*Petromyzon marinus*). *J. Comp. Neurol.* 522, 1132–1154. doi: 10.1002/cne.23481

- Pieri, M., Amadoro, G., Carunchio, I., Ciotti, M. T., Quaresima, S., Florenzano, F., et al. (2010). SP protects cerebellar granule cells against beta-amyloid-induced apoptosis by down-regulation and reduced activity of Kv4 potassium channels. *Neuropharmacology* 58, 268–276. doi: 10.1016/j.neuropharm.2009.06.029
- Popelová, A., Kákonová, A., Hrubá, L., Kuneš, J., Maletínská, L., and Železná, B. (2018). Potential neuroprotective and anti-apoptotic properties of a long-lasting stable analog of ghrelin: an *in vitro* study using SH-SY5Y cells. *Physiol. Res.* 67, 339–346.
- Postina, R. (2012). Activation of α -secretase cleavage. *J. Neurochem.* 120 Suppl. 1, 46–54. doi: 10.1111/j.1471-4159.2011.07459.x
- Quigley, B. J. Jr., and Kowall, N. W. (1991). Substance P-like immunoreactive neurons are depleted in Alzheimer's disease cerebral cortex. *Neuroscience* 41, 41–60. doi: 10.1016/0306-4522(91)90199-X
- Rak-Mardyla, A. (2013). Ghrelin role in hypothalamus-pituitary-ovarian axis. *J. Physiol. Pharmacol.* 64, 695–704.
- Raouf, R., Esmaili-Mahani, S., Abbasnejad, M., Raouf, M., Sheibani, V., Kooshki, R., et al. (2015). Changes in hippocampal orexin 1 receptor expression involved in tooth pain-induced learning and memory impairment in rats. *Neuropeptides* 50, 9–16. doi: 10.1016/j.npep.2015.03.002
- Rat, D., Schmitt, U., Tippmann, F., Dewachter, I., Theunis, C., Wiczerczak, E., et al. (2011). Neuropeptide pituitary adenylate cyclase-activating polypeptide (PACAP) slows down Alzheimer's disease-like pathology in amyloid precursor protein-transgenic mice. *FASEB J.* 25, 3208–3218. doi: 10.1096/fj.10-180133
- Reglodi, D., Kiss, P., Lubics, A., and Tamas, A. (2011). Review on the protective effects of PACAP in models of neurodegenerative diseases *in vitro* and *in vivo*. *Curr. Pharm. Des.* 17, 962–972. doi: 10.2174/138161211795589355
- Roh, J. H., Jiang, H., Finn, M. B., Stewart, F. R., Mahan, T. E., Cirrito, J. R., et al. (2014). Potential role of orexin and sleep modulation in the pathogenesis of Alzheimer's disease. *J. Exp. Med.* 211, 2487–2496. doi: 10.1084/jem.20141788
- Rose, J. B., Crews, L., Rockenstein, E., Adame, A., Mante, M., Hersh, L. B., et al. (2009). Neuropeptide Y fragments derived from neprilysin processing are neuroprotective in a transgenic model of Alzheimer's disease. *J. Neurosci.* 29, 1115–1125. doi: 10.1523/jneurosci.4220-08.2009
- Rösler, N., Wichart, N., and Jellinger, K. A. (2001). *Ex vivo* lumbar and post mortem ventricular cerebrospinal fluid substance P-immunoreactivity in Alzheimer's disease patients. *Neurosci. Lett.* 299, 117–120. doi: 10.1016/S0304-3940(01)01514-2
- Rowe, W. B., Kar, S., Meaney, M. J., and Quirion, R. (2006). Neurotensin receptor levels as a function of brain aging and cognitive performance in the Morris water maze task in the rat. *Peptides* 27, 2415–2423. doi: 10.1016/j.peptides.2006.03.036
- Russino, D., McDonald, E., Hejazi, L., Hanson, G. R., and Jones, C. E. (2013). The tachykinin peptide neurokinin B binds copper forming an unusual [CuII(NKB)2] complex and inhibits copper uptake into 1321N1 astrocytoma cells. *ACS Chem. Neurosci.* 4, 1371–1381. doi: 10.1021/cn4000988
- Santos, V. V., Stark, R., Rial, D., Silva, H. B., Bayliss, J. A., Lemus, M. B., et al. (2017). Acyl ghrelin improves cognition, synaptic plasticity deficits and neuroinflammation following amyloid β (A β 1-40) administration in mice. *J. Neuroendocrinol.* 29, 1–11. doi: 10.1111/jne.12476
- Severini, C., Petrella, C., and Calissano, P. (2016). Substance P and Alzheimer's disease: emerging novel roles. *Curr. Alzheimer Res.* 13, 964–972. doi: 10.2174/1567205013666160401114039
- Shi, L., Du, X., Jiang, H., and Xie, J. (2017). Ghrelin and neurodegenerative disorders—a review. *Mol. Neurobiol.* 54, 1144–1155. doi: 10.1007/s12035-016-9729-1
- Shi, L. M., Jiang, H., and Xie, J. X. (2014). Neuroprotective effects of ghrelin in Parkinson's disease and Alzheimer's disease. *Sheng Li Ke Xue Jin Zhan.* 45, 140–144.
- Shibata, N., Ohnuma, T., Kuerban, B., Komatsu, M., and Arai, H. (2011). Genetic association between ghrelin polymorphisms and Alzheimer's disease in a Japanese population. *Dement. Geriatr. Cogn. Disord.* 32, 178–181. doi: 10.1159/000333075
- Shieh, P. C., Tsao, C. W., Li, J. S., Wu, H. T., Wen, Y. J., Kou, D. H., et al. (2008). Role of pituitary adenylate cyclase-activating polypeptide (PACAP) in the action of ginsenoside Rh2 against beta-amyloid-induced inhibition of rat brain astrocytes. *Neurosci. Lett.* 434, 1–5. doi: 10.1016/j.neulet.2007.12.032
- Skup, M., Zhu, H., Wang, Y., Giovanello, K. S., Lin, J. A., Shen, D., et al. (2011). Sex differences in grey matter atrophy patterns among AD and aMCI patients: results from ADNI. *Neuroimage* 56, 890–906. doi: 10.1016/j.neuroimage.2011.02.060
- Spencer, B., Potkar, R., Metcalf, J., Thrin, I., Adame, A., Rockenstein, E., et al. (2016). Systemic central nervous system (CNS)-targeted delivery of neuropeptide Y (NPY) reduces neurodegeneration and increases neural precursor cell proliferation in a mouse model of Alzheimer disease. *J. Biol. Chem.* 291, 1905–1920. doi: 10.1074/jbc.M115.678185
- Spitznagel, M. B., Benitez, A., Updegraff, J., Potter, V., Alexander, T., Glickman, E., et al. (2010). Serum ghrelin is inversely associated with cognitive function in a sample of non-demented elderly. *Psychiatry Clin. Neurosci.* 64, 608–611. doi: 10.1111/j.1440-1819.2010.02145.x
- Stoyanova, I. I. (2014). Ghrelin: a link between ageing, metabolism and neurodegenerative disorders. *Neurobiol. Dis.* 72 (Pt A), 72–83. doi: 10.1016/j.nbd.2014.08.026
- Struble, R. G., Powers, R. E., Casanova, M. F., Kitt, C. A., Brown, E. C., and Price, D. L. (1987). Neuropeptidergic systems in plaques of Alzheimer's disease. *J. Neuropathol. Exp. Neurol.* 46, 567–584. doi: 10.1097/00005072-198709000-00006
- Tiraboschi, P., Hansen, L. A., Thal, L. J., and Corey-Bloom, J. (2004). The importance of neuritic plaques and tangles to the development and evolution of AD. *Neurology* 62, 1984–1989. doi: 10.1212/01.wnl.0000129697.01779
- Vaudry, D., Cottet-Rousselle, C., Basille, M., Falluel-Morel, A., Fournier, A., Vaudry, H., et al. (2004). Pituitary adenylate cyclase-activating polypeptide inhibits caspase-3 activity but does not protect cerebellar granule neurons against beta-amyloid (25-35)-induced apoptosis. *Regul. Pept.* 123, 43–49. doi: 10.1016/j.regpep.2004.05.025
- Waters, S. M., and David, T. P. (1995). Alterations of substance P metabolism and neuropeptidases in Alzheimer's disease. *J. Gerontol. A. Biol. Sci. Med. Sci.* 50, B315–B319. doi: 10.1093/gerona/50a.5.b315
- Willis, M., Hutter-Paier, B., Wietzorrek, G., Windisch, M., Humpel, C., Knaus, H. G., et al. (2007). Localization and expression of substance P in transgenic mice overexpressing human APP751 with the London (P171I) and Swedish (K670M/N671L) mutations. *Brain Res.* 1143, 199–207. doi: 10.1016/j.brainres.2007.01.080
- Witte, A. V., Savli, M., Holik, A., Kasper, S., and Lanzenberger, R. (2010). Regional sex differences in grey matter volume are associated with sex hormones in the young adult human brain. *Neuroimage* 49, 1205–1212. doi: 10.1016/j.neuroimage.2009.09.046
- Wu, Z. L., Ciallella, J. R., Flood, D. G., O'Kane, T. M., Bozyczko-Coyne, D., and Savage, M. J. (2006). Comparative analysis of cortical gene expression in mouse models of Alzheimer's disease. *Neurobiol. Aging* 27, 377–386. doi: 10.1016/j.neurobiolaging.2005.02.010
- Xiao, Z., Cilz, N. I., Kurada, L., Hu, B., Yang, C., Wada, E., et al. (2014). Activation of neurotensin receptor 1 facilitates neuronal excitability and spatial learning and memory in the entorhinal cortex: beneficial actions in an Alzheimer's disease model. *J. Neurosci.* 34, 7027–7042. doi: 10.1523/jneurosci.0408-14.2014
- Yang, L., Zou, B., Xiong, X., Pascual, C., Xie, J., Malik, A., et al. (2013). Hypocretin/orexin neurons contribute to hippocampus-dependent social memory and synaptic plasticity in mice. *J. Neurosci.* 33, 5275–5284. doi: 10.1523/jneurosci.3200-12.2013
- Zhang, H., Dong, H., Cilz, N. I., Kurada, L., Hu, B., Wada, E., et al. (2016). Neurotensinergic excitation of dentate gyrus granule cells via G α q-coupled inhibition of TASK-3 channels. *Cereb. Cortex* 26, 977–990. doi: 10.1093/cercor/bhu267
- Zhang, H., Dong, H., and Lei, S. (2015). Neurotensinergic augmentation of glutamate release at the perforant path-granule cell synapse in rat dentate gyrus: Roles of L-Type Ca $^{2+}$ channels, calmodulin and myosin light-chain kinase. *Neuropharmacology* 95, 252–260. doi: 10.1016/j.neuropharm.2015.03.028

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Chen, Du and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.