



OPEN ACCESS

EDITED BY
Zhaojun Wei,
Hefei University of Technology, China

REVIEWED BY
Ningyang Li,
Shandong Agricultural University,
China
Zhen-quan Yang,
Yangzhou University, China
Xiaoyu Li,
Yanshan University, China

*CORRESPONDENCE
Zhi-gui He
906260300@qq.com

SPECIALTY SECTION
This article was submitted to
Nutrition and Food Science
Technology,
a section of the journal
Frontiers in Nutrition

RECEIVED 05 June 2022
ACCEPTED 29 June 2022
PUBLISHED 04 August 2022

CITATION
He Z-g, Zhang Y, Yang M-d, Zhang Y-q,
Cui Y-y, Du M-y, Zhao D and Sun H
(2022) Effect of different sweeteners
on the quality, fatty acid and volatile
flavor compounds of braised pork.
Front. Nutr. 9:961998.
doi: 10.3389/fnut.2022.961998

COPYRIGHT
© 2022 He, Zhang, Yang, Zhang, Cui,
Du, Zhao and Sun. This is an
open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

Effect of different sweeteners on the quality, fatty acid and volatile flavor compounds of braised pork

Zhi-gui He^{1*}, Ying Zhang², Ming-duo Yang², Yu-qing Zhang¹, Ying-ying Cui¹, Mi-ying Du¹, Dong Zhao¹ and Hui Sun¹

¹School of Leisure and Health, Guilin Tourism University, Guilin, China, ²School of Tourism and Cuisine, Harbin University of Commerce, Harbin, China

This study aimed to assess how several sweeteners (white sugar, *Siraitia grosvenorii* fruit, mogrosides, and stevia glycoside) affected the flavor, fatty acid composition, and quality of braised pork. The findings indicated that braised meat prepared with sweeteners differed from typical braised pork. When simmered for 60 min, the typical braised pork with white granulated sugar exhibited a significant cooking loss (CL) and little water content. Significantly more than in the group containing *Siraitia grosvenorii*, mogroside, and stevia glycoside, the Thiobarbituric acid (TBARS) value increased by 14.39% ($P < 0.05$). The sample in the group that included mogroside had a low CL rate. After 40 min of stewing, the lean pork has the highest L* value, but the 60-min stew sample is nicely colored and stretchy. Mogroside can prevent protein, and lipid oxidation, is thermally stable and reduces CL during stewing. Additionally, *Siraitia grosvenorii* and stevia glycosides help prevent oxidation from intensifying during stewing. When *Siraitia grosvenorii* is added, lipid oxidation is significantly inhibited, and stevia glycosides are more beneficial for enhancing meat color. With an increase in heating time, the fatty acids in braised pork reduced; the unsaturated fatty acid (UFA) of the *Siraitia grosvenorii* fruit (SF) and mg group also fell somewhat, and the UFA: SFA ratio was higher than that of the white sugar (WS) group. The SFA content of the braised meat in the stevia glycoside group was higher than that of the WS group. In all, 75 volatile flavor elements in braised pork were discovered by Gas chromatography-ion mobility spectrometry (GC-IMS). The sweetener increased alcohols, esters, and acids in the braised pork. As stewing time increased, ketones decreased, but aldehydes and esters increased. The pork formed antioxidant peptides with great nutritional value after cooking. Braised pork with mogroside and stevia glycoside additions primarily have some protein color protection and antioxidant effects. This study may offer fresh perspectives on applying natural sweeteners and enhancing braised pork's flavor.

KEYWORDS

braised pork, fatty acid, volatile compound, *Siraitia grosvenorii*, mogroside, stevia glycoside

Introduction

Because of its vivid color, redness, distinctive flavor, alluring flavor, and texture, the braised pork is a traditional Chinese dish with a long history in Chinese cuisine (1). By selecting the proper stewing conditions and preparation techniques, such as pre-frying, it is possible to enhance the nutritional benefits of fat, particularly the color, taste, and flavor of braised pork (2). Traditional braised pork is made with pork belly, white sugar, and soy sauce; the Maillard reaction takes place during processing. This reaction will result in a distinctive flavor but will also unavoidably generate hazardous compounds, including acrylamide, heterocyclic amine, and furan (3).

Unlike starch, sugar has a higher metabolic capacity and is an excellent energy source (4). However, sufficient scientific data shows that eating a lot of hot, sugary meals can quickly result in health hazards, including tooth decay, obesity, and diabetes. People are searching for low-calorie sweeteners as sugar alternatives as they become more aware of the value of health (5). Sweeteners are compounds that can make meals sweeter. According to various sources, they can be separated into artificial and natural sweeteners (6). Consumers prefer natural sweeteners with low heat and are non-toxic compared to artificial sweeteners; this has increasingly been the research focus of the industry (7). Studies demonstrate that natural sweeteners can lower blood sugar and enhance lipid metabolism. Fruit powder, dried fruit, and fruit extract or concentrate are essential sweeteners that increase the sweetness of meals and aid in producing bioactive components (8).

Mogroside and stevia glycoside, both natural sweeteners, are typical biological resources in Guangxi. *Siraitia grosvenorii* fruit has traditionally utilized boiling water to prepare tea. In Guangxi, stewed pork or ribs prepared with *Siraitia grosvenorii* has a distinct flavor. Mogroside, the primary extract of *Siraitia grosvenorii*, has 300 times the sweetness of sucrose. It has a low temperature, high stability, and a scent of *Siraitia grosvenorii*. It has significant antioxidant action, regulates blood sugar, inhibits bacteria and inflammation, and inhibits malondialdehyde production (9). Stevia glycoside is derived from the *Stevia rebaudiana* plant, which possesses the high sweetness and low heat. It is safe for baked goods, sauces, and other items since it does not promote dental cavities. Activating glucose-induced insulin secretion β -Cell function and hypoglycemic impact can improve islet function (10). Sweeteners are frequently utilized in manufacturing industrial foods and pharmaceuticals as a replacement for sucrose, although variations in their sensory qualities have yet to be studied (11).

Previous research has shown that various plant herbs' phenolic acids, which have potent antioxidant properties and pharmacological effects (12), can reduce the damaging Maillard reaction products. Additionally, it has been demonstrated that natural antioxidants such as tea polyphenol, flavonoid compounds, and spice extracts specifically limit the formation

of hazardous chemicals during the thermal processing of food, such as baking, frying, and boiling (13). Although mogroside and stevioside may be used as natural sweeteners and are produced from medicinal and food-related plants, their antioxidant action is rarely utilized in food preparation, particularly when preparing meat products. The influence of raw material types (14), cooking techniques (15), heating temperature and time (16), condiments (17), and storage conditions (18) are now the primary topics of study on the sensory, color, flavor, and other qualities of meat products. However, no research has been conducted on how different sugar kinds affect the flavor and quality of meat products in dish preparation.

This study compares the effects of various sweeteners and stewing times on the fatty acids, volatile taste compounds, and quality of braised pork to examine the differences between them and conventional braised pork. It serves as a theoretical foundation for the deep processing technology study and application development of sugar substitute food and functional food of *Siraitia grosvenorii*, as well as a guide for the study of the use of natural sweeteners and antioxidants.

Materials and methods

Sample preparations

The pork belly in this study was obtained from Liyuan fresh food chain supermarket (Guilin, Guangxi, China). The pork belly was cut into 4 cm \times 3 cm \times 2 cm pieces after being washed, drained, and cooked in boiling water for 2 min. Then, the samples were stir-fried with soybean oil at 180°C for 5 min using an induction cooker (WT2202, Midea, Foshan, China). According to the basis of the weight of meat, 120% water, 4% light soy sauce, 4.2% thick soy sauce, 5.3% cooking wine, 1% salt, 2% scallion, 2% ginger, 0.4% spices, and 1.7% sweeteners (white sugar, *Siraitia grosvenorii* fruit, mogroside and stevia glycoside) were added. These mixtures were braised at 95°C, and the time was set at 40 and 60 min. The soup was collected at 150°C for 5 min until thickened. Relevant indexes such as texture, color, fatty acid and flavor compounds were detected by taking braised pork samples. The specific design of sweeteners and stewing time are shown in **Table 1**.

Cooking loss

The pork was put on clean filter paper, and the soup on the surface of the meat sample was drained and weighed after 5 min of natural cooling. The cooking loss (CL) was determined as follows:

$$\text{Cooking loss rate (\%)} = (M_1 - M_2)/M_1 \times 100\%$$

TABLE 1 Cooking conditions of braised pork.

Numbering	Processing methods
WS-1	Add white sugar, 40 min stewed pork
WS-2	Add white sugar, 60 min stewed pork
SF-1	Add <i>Siraitia grosvenorii</i> fruit, 40 min stewed pork
SF-2	Add <i>Siraitia grosvenorii</i> fruit, 60 min stewed pork
MG-1	Add mogroside, 40 min stewed pork
MG-2	Add mogroside, 60 min stewed pork
SG-1	Add stevia glycoside, 40 min stewed pork
SG-2	Add stevia glycoside, 60 min stewed pork

Where: M_1 is the mass of raw meat, g; M_2 is the mass of the sample after heat treatment, g.

Moisture content

Moisture content in each group of meat samples was determined by drying to a constant weight at 105°C according to AOAC procedure 950.46 using an electric blast drying oven (9140AL; Keelrein, Shanghai, China) from the laboratory.

Measurement of pH

One gram of braised pork was chopped, transferred to a beaker, homogenized with 10 mL of distilled water for 35 s, filtered, and tested the pH. Prior to measurement, the pH meter was calibrated using matched buffers with various values of 4.00, 6.86, and 9.18. After the pH meter reading stabilized, the pH electrode was placed into the filtrate for testing, and the result was recorded.

Color

Using a spectrophotometer (CM-5; KonicaMinolta, Tokyo, Japan), the lightness (L^*), redness (a^*), and yellowness value (b^*) of the fat layer and thin layer of braised pork were measured. Self-inspection, zero calibration and whiteboard calibration were conducted before use, and each group was conducted in triplicate.

Texture profile analysis

The texture profile in terms of hardness, springiness, cohesiveness, chewiness, and gumminess of rectangular samples (2 cm length, 1 cm width, 1 cm height) cut from braised pork were assessed for texture profile analysis (TPA) using the food texture analyzer (TMS-PILOT; Food Technology Corporation,

Virginia, United States). The parameters of TPA were as follows: the trigger force of 0.5 N, a test speed of 60 mm/min, a return distance of 35 mm, and deformation of 40%. The shear force is measured in single knife shear mode, and the working parameters were set: the trigger force of 2 N, a test speed of 120 mm/min and a return speed of 200 mm/min. Each group of samples is measured with three parallel samples, and the average value is taken for the calculation.

Malondialdehyde content

The Malondialdehyde (MDA) content was determined by following the Díaz et al. (19) method with appropriate modifications. The minced meat (5 g) was homogenized in 25 mL of trichloroacetic acid and EDTA mixture (Macklint Biochemical Technology Co., Ltd, Shanghai, China) at 10,000 rpm for 30 s. Then the sample was centrifuged at the speed of 3,000 r/min for 5 min, and the supernatant was filtered with filter paper. Five milliliters of supernatant were mixed with 5 mL of 0.02 mol/L thiobarbituric acid (TBARS) solution. The reaction mixture was heated at 95°C for 30 min in a thermostat water bath cauldron, and then the sample was placed in ice for 30 min and rapidly cooled to room temperature. The absorbance was measured at 532 nm, and the TBARS value was expressed as mg MDA/kg (measured by meat sample).

Extraction of myofibrillar protein

The extraction of myofibrillar protein (MP) was determined according to Park et al. (20), with some modifications. The minced pork meat was homogenized on ice for 60 s (divided into three times, the 20 s each time) with four volumes (w/v) of protein extract buffer (10 mmol/L Na_3PO_4 , 0.1 mol/L NaCl, 2 mmol/L MgCl_2 , 1 mmol/L EGTA), then centrifuged at 3,500 rpm, 4°C for 15 min. The precipitate was taken out, and the crude MP was obtained by repeating twice above the operation. Then, the precipitated sample was washed three times with 4 volumes (w/v) of 0.1 mol/L NaCl under the same centrifugation conditions. Finally, the suspension was filtered through four layers of gauze and centrifuged again at 3,500 rpm, 4°C for 15 min. The final precipitate was MP, and the next test was carried out.

Sulfhydryl content and carbonyl content

Protein oxidation, assessed on extracted MP, was evaluated using multiple indexes, including carbonyl and sulfhydryl content. The sulfhydryl content was determined according to Yang et al. (21) with minor modification. Eight milliliters of

Tris-glycine solution (10.4 g/L Tris, 0.9 g/L glycine, 10 mmol/L EDTA, 8 mol/L urea, pH 8.0) was added to 1 mL diluted MP solution (5 mg/mL), then centrifuged at 4°C and 4,500 rpm for 15 min after mixing to remove insoluble protein. Then, 0.5 mL of 10 mol/L 5, 5'-dithiobis-2-nitrobenzoic acid (DTNB) was added to 4.5 mL of supernatant. The mixed samples were incubated at room temperature and away from light for 30 min, and the light absorption value was measured at 412 nm. The sulfhydryl content was calculated using the molar extinction coefficient of 13,600 L/(mol·cm).

Carbonyl content was measured as determined by Ortuño et al. (22). One milliliter of 10 mmol/L 2,4-dinitrophenylhydrazine (DNPH) was added to 3 mL of MP diluent (5 mg/ml). After an incubation of 1 h (shaking every 15 min), 1 mL of 20% w:w TCA solution was added. The sample was vortexed and centrifuged for 5 min at 5,000 rpm, 4°C. The supernatant was discarded, and the precipitate was washed with 1 mL of ethanol/ethyl acetate (1:1) three times to remove the unreacted reagents. Then the supernatant was dissolved in 3 mL 6 mol/L guanidine hydrochloride, shaken, and centrifuged for 3 min at 5,000 rpm, 4°C, to avoid insoluble fragments. The absorbance value of the obtained solution was measured at 370 nm, and the carbonyl content was calculated using the molar absorbance coefficient of 22,000 L/(mol·cm).

Fatty acid analysis

The fatty acid composition of braised pork was determined using gas chromatography (GC). The lipid portion of braised pork was extracted by adding ether/petroleum (1:1, vol/vol), and the lipid extract was collected by a rotary evaporator. The extracted sample was placed in a 250 mL round-bottomed flask, and the methyl ester preparation was carried out by the boron trifluoride-methanol method according to Bazina et al. (23). Then, quantitative analysis was carried out using the standard internal technique, and qualitative analysis was carried out using the comparing retention times of mixed standards and samples of fatty acid methyl ester. The condition parameters of GC were set according to Chen et al. (24).

GC-IMS analysis

The FlavourSpec GC-IMS flavor analysis system (G.A.S., Dortmund, Germany) is an automated headspace injection system used to analyze the volatile flavor components. The volatile flavors of braised pork cooked with various sweeteners and stewing time were tracked. Five grams of sample were weighed and placed in a 20 mL headspace bottle for analysis, and three repetitions were made for each sample. The specific system conditions are shown in Tables 2, 3. The qualitative analysis of volatile flavor substances is carried out through the NIST

TABLE 2 Analysis conditions.

System		Condition
FlavourSpec unit	Analysis time	30 min
	Column type	MXT-5 15 m 1 μm 0.53 mm
	Column temperature	60°C
	Carrier gas flow	N ₂
	IMS temperature IMS	45°C
The automatic headspace sampling unit	Incubation temperature	90°C
	Incubation time	15 min
	Injection volume	500 μl
	Syringe temperature	85°C
	Incubation speed	500 rpm

TABLE 3 System conditions of GC.

Time	E1	E2	R
00:00,000	150 mL/min	2 mL/min	Rec
02:00,000	150 mL/min	2 mL/min	-
10:00,000	150 mL/min	10 mL/min	-
20:00,000	150 mL/min	100 mL/min	-
30:00,000	150 mL/min	100 mL/min	Stop

database and IMS database built in the software, and difference spectra and fingerprints of volatile organic compounds were constructed by Reporter and Gallery Plot programs.

LC-MS/MS

The identification of peptides was determined using Jiang et al. (25) with few modifications. The peptide samples digested by enzymatic (trypsin, v5280) were eluted using C18 analytical columns (Acclaim PepMap RSLC, 75 μm × 15 cm, 2 μm, 100 Å, nano Viper, Thermo Fisher Scientific, Palo Alto, CA, United States). The peptide solution was collected after centrifugation and dried in a vacuum.

Subsequently, the peptide is dissolved by 20 μL of solution (0.1% formic acid, 5% ANC), then centrifuged at 13,500 rpm and 4°C for 20 min after sufficient oscillation. The supernatant was transferred to the sample pipe and identified using Dionex Ultimate 3000 RSLCnano Liquid chromatography tandem mass spectrometry (LC-MS/MS) (Thermo Fisher Scientific, Palo Alto, CA, United States). The liquid phase setting parameters are as follows: mobile phase A is 0.1% formic acid, mobile phase B is 0.1% formic acid and 80% ACN. In a word, the polypeptide liquid was eluted with a linear gradient at a flow rate of 300 nL/min. The gradient elution was performed as follows: 0–5 min (5–10%B), 5–43 min (10–28%B), 43–51 min (28–38%B), 51–53 min (38–100%B), 53–60 min (100–100%B).

The separated peptides were detected online by Thermo Scientific Q Exactive PLUS (Thermo Fisher Scientific, Palo Alto, CA, United States), and the specific conditions of mass spectrometry were as follows: primary mass spectrometry parameters: resolution: 12000, AGC target: 4e5, maximum Injection time: 50 ms, scan range: 350–1550 m/z. Secondary mass spectrometry parameters: resolution: 30000, AGC target: 1e5, TopN: 20, NCE/stepped NCE: 32. In addition, carbamidomethyl (C) was specified as the fixed modification, and oxidation (M), acetyl (Protein N-term) and NEM (C) were set as variable modifications. Fragment mass error tolerance was 0.03 Da, and precursor mass error tolerance was set at 15 ppm. LC-MS/MS spectra of peptides were matched using the Swiss-Prot database against *Sus scrofa*¹.

Statistical analysis

The data were analyzed by one-way analysis of variance (ANOVA) and Duncan's multiple range test, using SPSS 21.0 (IBM., Armonk, NY, United States). Results of each group were expressed as means \pm standard deviations. The figures were performed by using the Origin2018 software (OriginLab, United States) and GraphPad Prism 8 (GraphPad Software Inc., United States).

Results and discussion

Basic indicators

The effects of the primary physical and chemical indexes of braised pork with different sweeteners in various stewing times are shown in **Table 4**. Concerning the types of sweeteners, the CL rate of braised pork with white sugar was significantly

1 <http://www.uniprot.org/>

TABLE 4 Basic physical and chemical indexes of braised pork with different sweeteners in various stewing time.

	CL	MC	pH
WS-1	23.10 \pm 0.16 ^b	38.85 \pm 0.27 ^c	6.09 \pm 0.02 ^a
WS-2	25.85 \pm 0.55 ^a	36.46 \pm 0.87 ^d	6.00 \pm 0.01 ^b
SF-1	23.15 \pm 1.67 ^b	38.83 \pm 0.39 ^c	5.78 \pm 0.02 ^{cd}
SF-2	24.69 \pm 1.38 ^{ab}	37.53 \pm 0.38 ^{cd}	5.81 \pm 0.01 ^c
MG-1	22.94 \pm 0.72 ^b	46.42 \pm 0.54 ^a	5.76 \pm 0.06 ^{cd}
MG-2	23.32 \pm 0.34 ^{ab}	44.90 \pm 1.02 ^b	5.47 \pm 0.03 ^e
SG-1	23.27 \pm 0.51 ^b	46.72 \pm 0.21 ^a	5.71 \pm 0.01 ^d
SG-2	24.11 \pm 0.82 ^{ab}	43.55 \pm 0.40 ^b	5.75 \pm 0.02 ^{cd}

All values are means \pm SD. Significant differences between each group were indicated by different letters ($p < 0.05$). The following table is the same.

higher than that of other sweeteners ($p < 0.05$). From the aspect of stewing time, the CL rate increased significantly due to the extension of stewing time, which confirmed the law of the quality change in stewing pork. Namely, the meat protein was thermally deformed during heating with decreased muscle fiber density, which caused water evaporation and fat loss, eventually increasing CL (26). Besides, the higher temperature resulted in more significant CL but less MC in cooked samples (27). The MC of braised pork with white sugar decreased significantly with the extension of stewing time, and the sample added with *Siraitia grosvenorii* fruit was significantly lower than that added with mogroside and stevia glycoside at 60 min of stewing time. Previous studies have indicated that mogroside is relatively stable and has heat and high-temperature resistance (28). Therefore, it is difficult to decompose during stewing and remains longer than other components in *Siraitia grosvenorii*. The MC of SG-2 was 43.55%, which was 6.79% lower than that of at 40 min for stewing time. The pH of WS-1 and WS-2 were 6.09 and 6.00, while the other braised pork with other sweeteners was lower than 6.00, and the pH of SF-2 was the lowest at 5.47. It indicated that the braised pork with *Siraitia grosvenorii*, mogroside, and stevia glycoside was generally acidic, which might be affected by the acidic substances such as flavonoids and saponins contained in *Siraitia grosvenorii*. Moreover, the pH of braised pork was affected by condiments such as soy sauce, cooking wine and salt during cooking (29).

Color analysis

Maillard's reaction during cooking was affected frequently by the added sugar. The degree of Maillard reaction becomes lower when the amount of sugar added is small, which results in lighter color (30). The change of color of braised pork is shown in **Table 5**. The a^* value of the lean layer and L^* value of the fat layer of the white sugar (WS) group were significantly higher than those in other groups ($p < 0.05$). In order to achieve the effect of red and bright color, the traditional braised pork is colored and flavored through the Maillard reaction of white sugar in hot processing. The Lightness value (L^*) of WS-1 and WS-2 decreased with the extension of stewing time. At the same time, the a^* value of the lean layer increased, and the a^* and b^* values of the fat layer decreased. The drop in a^* value is due to the reduction in myoglobin produced by the rise in pork's internal temperature (31). The L^* value of the fat layer is generally more significant than that of the lean layer. Dong et al. (32) manifested that the fat content significantly impacts the meat color of meat products, especially the L^* value, which will increase markedly with the increase of fat content. The b^* value of other groups was lower than that of stewed pork with *Siraitia grosvenorii* fruit ($p > 0.05$). *Siraitia grosvenorii* contains antioxidant polyphenol, flavonoids compound and water-soluble yellow pigment, which has thermal stability and

TABLE 5 Color of braised pork with different sweeteners in various stewing times.

	L*		a*		b*	
	Lean layer	Fat layer	Lean layer	Fat layer	Lean layer	Fat layer
WS-1	41.67 ± 0.63 ^{cd}	59.13 ± 0.57 ^a	11.46 ± 0.82 ^b	4.85 ± 0.28 ^{ab}	20.15 ± 0.29 ^b	24.79 ± 1.83 ^a
WS-2	39.54 ± 0.31 ^{de}	54.60 ± 0.87 ^{abc}	13.76 ± 0.17 ^a	4.40 ± 0.42 ^b	19.71 ± 0.82 ^b	23.32 ± 0.24 ^a
SF-1	44.58 ± 0.29 ^{ab}	56.09 ± 0.97 ^{ab}	11.05 ± 0.51 ^{bc}	5.86 ± 0.72 ^{ab}	23.58 ± 0.83 ^a	25.62 ± 1.99 ^a
SF-2	37.78 ± 1.00 ^c	54.74 ± 0.82 ^{abc}	12.11 ± 0.63 ^{ab}	6.09 ± 0.85 ^{ab}	19.94 ± 0.63 ^b	25.08 ± 1.07 ^a
MG-1	46.18 ± 0.49 ^a	54.75 ± 0.33 ^{abc}	9.43 ± 0.49 ^c	6.63 ± 0.49 ^a	20.42 ± 0.19 ^b	22.59 ± 0.28 ^a
MG-2	43.04 ± 0.32 ^{bc}	50.90 ± 1.27 ^c	10.61 ± 0.95 ^{bc}	5.41 ± 1.48 ^{ab}	20.86 ± 1.45 ^b	21.68 ± 1.73 ^a
SG-1	45.22 ± 1.70 ^{ab}	56.47 ± 0.87 ^{ab}	10.40 ± 0.26 ^{bc}	5.27 ± 0.31 ^{ab}	21.33 ± 1.31 ^{ab}	21.73 ± 1.69 ^a
SG-2	43.88 ± 1.22 ^{abc}	53.02 ± 0.79 ^{bc}	10.35 ± 0.96 ^{bc}	4.67 ± 0.17 ^{ab}	21.09 ± 0.12 ^b	22.71 ± 1.43 ^a

affects fat oxidation (33). However, the L* value of braised pork with *Siraitia grosvenorii* fruit decreased, and the a* value increased at stewing for 60 min. The L* of the lean layer in MG-1 was the largest and remained at 43.04 after stewing, making braised pork's overall color brighter. The L* value of the lean layer of SG is greater than that of WS, indicating that stevia glycoside can improve the color of braised pork. Karp et al. (34) showed that the brightness, redness and yellowness of muffin skin increased by making a muffin by adding stevia glycoside, which can improve the quality of baking products.

The analysis of texture

Table 6 displays the textural characteristics of braised pork using various sugars and stewing periods. The gumminess of the group added with white sugar and stevia glycoside was more excellent, while the group's hardness, chewiness, and shear force added with mogroside and stevia glycoside were much lower than those of WS. Due to the development of caramel by the impact of temperature and time during the process of hot processing, white sugar can boost the sweetness and consistency of meat products. This influences braised pork's hardness, chewiness, and other textural qualities (35). The collagen in meat was heated to dissolve during the stewing procession. The cross-linking between protein molecules is reduced, resulting in the loss of MP and the fragmentation of tissue structure, which will eventually decrease the shear force of samples (36). The shear force of the lean layer of SF-2 was the lowest. It could be connected to the fact that *Siraitia grosvenorii* protease, which has a high hydrolysis activity and a stable structure, was present in *Siraitia grosvenorii* (37). Studies have shown that plant protease can improve the tenderness of meat products owing to the ability of tenderization (38). However, the effect of stewing temperature and time on the tenderness of meat might be greater in the process of stewing. Braised pork with mogroside is more malleable during the 40-min stewing stage, and the fat layer is more springy than the lean layer. Due to its antioxidant capabilities, mogroside can reduce lipid buildup,

fat degeneration, and fat oxidation (39), which can significantly improve the elasticity of the fat layer in the early stage of stewing. The hardness and chewiness of the stewed pork with stevia glycoside were superior to those with white sugar, but its cohesion and springiness were marginally inferior to those with *Siraitia grosvenorii* fruit and mogroside. In other words, stevia glycoside also has the benefit of replacing sugar in enhancing the texture and quality of meat products.

Fat oxidation

During heat processing, muscle protein denatured, myofibril constricted, meat moisture and quality decreased, and the TBARS value increased (40). Figure 1 demonstrates that the stewing time has a discernible effect on the TBARS value of each sample group. Specifically, the longer the cooking time, the more fat is oxidized. There was a significant difference in the TBARS value of stewed pork ($p < 0.05$) with adding different sweeteners, and the TBARS value of the WS group at stewing time for 40 min was higher than that of other groups. When the stewing time was 60 min, the TBARS value of WS increased by 14.39%, which was much higher than that of *Siraitia grosvenorii* fruit (SF), MG, and SG, indicating that this group had a significant degree of fat oxidation and produced more secondary products.

In addition, lipid content and the ratio of fatty acids impact fat oxidation (41). The TBARS concentration of samples supplemented with mogroside and stevia glycoside rose gradually. The TBARS value of samples supplemented with mogroside and stevia glycoside grew slowly because mogroside and stevia glycoside are both natural sweeteners that lower blood sugar and lipid, and their antioxidant qualities can limit fat oxidation during cooking (42). When stewed for 60 min, the MG TBARS value was 0.44 mg MDA/kg, much lower than other groups ($p < 0.05$). It demonstrates that mogroside inhibits fat oxidation and remains stable after prolonged exposure to heat. The rise in TBARS was smaller in SF than in WS and SG, which may be because polyphenols provided by *Siraitia grosvenorii* can

TABLE 6 Texture quality of braised pork with different sweeteners in various stewing times.

		Hardness	Springiness	Chewiness	Cohesiveness	Gumminess	Shear force
WS-1	Lean layer	32.49 ± 3.52 ^a	3.22 ± 0.39 ^b	28.50 ± 0.70 ^{ab}	0.48 ± 0.02 ^{abc}	10.38 ± 0.08 ^{bc}	39.28 ± 2.68 ^a
	Fat layer	17.06 ± 1.02 ^a	3.63 ± 0.17 ^b	20.82 ± 1.67 ^a	0.45 ± 0.02 ^{cd}	10.39 ± 0.44 ^{ab}	16.40 ± 1.64 ^a
WS-2	Lean layer	19.44 ± 1.38 ^b	3.52 ± 0.11 ^{ab}	26.23 ± 0.49 ^b	0.41 ± 0.02 ^c	8.55 ± 0.46 ^{cd}	25.66 ± 0.45 ^{bcd}
	Fat layer	10.42 ± 0.58 ^{de}	3.02 ± 0.48 ^{bcd}	10.32 ± 0.49 ^c	0.34 ± 0.03 ^e	9.37 ± 0.43 ^{bc}	10.15 ± 1.73 ^b
SF-1	Lean layer	29.33 ± 0.69 ^a	3.39 ± 0.10 ^b	25.76 ± 0.15 ^b	0.47 ± 0.02 ^{bc}	8.18 ± 0.39 ^{cd}	31.63 ± 5.32 ^{abc}
	Fat layer	13.82 ± 1.44 ^{bc}	2.81 ± 0.06 ^d	16.99 ± 0.11 ^b	0.52 ± 0.03 ^{bc}	9.76 ± 0.73 ^{ab}	14.70 ± 1.08 ^a
SF-2	Lean layer	19.23 ± 1.67 ^b	4.12 ± 0.08 ^a	25.23 ± 0.32 ^{ab}	0.41 ± 0.01 ^c	7.45 ± 0.51 ^d	21.47 ± 0.42 ^d
	Fat layer	7.96 ± 0.14 ^e	2.89 ± 0.04 ^{cd}	10.27 ± 0.14 ^c	0.43 ± 0.02 ^d	8.76 ± 1.03 ^{bc}	9.70 ± 0.61 ^b
MG-1	Lean layer	28.95 ± 0.76 ^a	3.27 ± 0.04 ^b	26.68 ± 1.07 ^{ab}	0.52 ± 0.02 ^{ab}	10.40 ± 1.67 ^{bc}	37.47 ± 1.79 ^a
	Fat layer	12.47 ± 1.35 ^{cd}	4.46 ± 0.14 ^a	19.62 ± 1.80 ^{ab}	0.67 ± 0.01 ^a	7.83 ± 0.21 ^{bcd}	16.93 ± 0.03 ^a
MG-2	Lean layer	19.26 ± 0.32 ^b	4.10 ± 0.17 ^a	21.89 ± 0.85 ^c	0.52 ± 0.06 ^{ab}	10.34 ± 1.51 ^{bc}	27.04 ± 5.28 ^{bcd}
	Fat layer	8.43 ± 1.18 ^e	3.56 ± 0.34 ^{bc}	10.98 ± 0.58 ^c	0.55 ± 0.04 ^b	6.79 ± 0.51 ^d	10.88 ± 0.23 ^b
SG-1	Lean layer	28.44 ± 2.08 ^a	3.64 ± 0.07 ^{ab}	29.07 ± 2.21 ^a	0.55 ± 0.02 ^a	14.30 ± 0.53 ^a	33.53 ± 1.27 ^{ab}
	Fat layer	16.12 ± 0.78 ^{ab}	3.00 ± 0.11 ^{bcd}	18.87 ± 0.64 ^{ab}	0.45 ± 0.03 ^{cd}	10.86 ± 0.88 ^a	16.12 ± 0.81 ^a
SG-2	Lean layer	20.23 ± 1.86 ^b	3.11 ± 0.50 ^b	22.61 ± 0.07 ^c	0.44 ± 0.01 ^c	11.40 ± 0.56 ^b	25.02 ± 0.58 ^{cd}
	Fat layer	8.00 ± 1.60 ^e	3.27 ± 0.27 ^{bcd}	11.62 ± 1.30 ^c	0.40 ± 0.05 ^{de}	7.89 ± 0.87 ^{bcd}	11.75 ± 0.56 ^b




FIGURE 1


TBRAS value of stewed pork with different sweeteners in various stewing time. All values are means ± SD. Significant differences between each group were indicated by different letters ($p < 0.05$). The following figure is the same.

replace free radicals and interact with unsaturated fatty acids to limit fat oxidation in meat products (43).

Protein oxidation

During cooking and processing, oxidants and tissue degradation impact meat products. In addition, meat protein is

exposed to an environment including oxidative stress, resulting in the breakdown of its secondary and tertiary structures and conformational alterations (44). As the most active group in protein amino acid residues, a change in sulfhydryl concentration can represent a change in protein structure. Figure 2A depicts the effects of sulfhydryl concentration on braised pork with varying sweeteners and stewing periods. The sulfhydryl level in the 60 min stew stage was much lower than



that in the 40 min stew stage due to the escalation of protein oxidation degree in the high-temperature stew process and the oxidation of free sulfhydryl to form disulfide link after heating (45). The sulfhydryl concentration of pork braised with white sugar for 40 min was more remarkable, but it fell dramatically with prolonged stewing and heating ($p < 0.05$), indicating that white sugar did not influence the loss of the sulfhydryl group during protein oxidation. However, the loss of sulfhydryl groups during stewing might be prevented by adding antioxidant sweeteners, with mogroside and stevia glycoside being the most prominent examples. Additionally, the sulfhydryl level may be connected to the amount of the four different sweeteners used. A high quantity of phenols has been found to reduce meat's sulfhydryl content (46). The fruit of *Siraitia grosvenorii* fruit includes phenolic chemicals that affect the variation of sulfhydryl concentration.

Carbonylation of proteins is a post-translational alteration involving reactive oxygen species. Most protein oxidation happens on the side chain of amino acid residues. Oxidation can result in structural modifications to proteins, such as aromatic hydroxylation, sulfhydryl oxidation, and carbonyl formation (47). Frequently, alterations in the muscle fiber structure of meat products are the result of heat treatment. When the duration of heating is prolonged, protein denaturation increases. Generally, the greater the carbonyl concentration, the greater the protein oxidation (48). Due to the rise in protein oxidation degree caused by the lengthening of heat treatment duration, the carbonyl content rose as stewing time increased (Figure 2B).

In addition, the active carbonyl compounds initially formed by lipid oxidation and the Maillard process interact with specific amino acids on the protein side-chain via carbonyl crosslinking to make carbonyl (49). Carbonyl concentration varied significantly when stewed pork was combined with various sweeteners and cooking durations. Among them, the

carbonyl content of the MG group grew less during stewing, showing that the addition of mogroside had a particular inhibitory impact on the generation of carbonyl; in other words, it may limit protein oxidation. The WS group's carbonyl content was much greater than that of the other groups, reaching 6.69 nmol/mg at the 60-min stewing stage. More reactive oxygen species will be produced during cooking, which can quickly increase carbonyl production. However, antioxidants can diminish the carbonyl concentration of cooked foods (50), which is consistent with the shift in sulfhydryl content. Cheng et al. (51) previously found that *Momordica grosvenori* extract (MGE) had a protective impact on the oxidative loss of pork protein, could considerably delay carbonyl formation during the storage of dried minced pork slices and decreased the loss of sulfhydryl dose-dependently. Liu et al. (52) also demonstrated that MGE exhibited significant antioxidant capacity by scavenging peroxy free radicals and intense anti-diabetic activity by inhibiting glucose-mediated protein glycosylation and cross-linking, which effectively inhibits the increase of carbonyl caused by glycosylation.

Composition and content of fatty acid

Table 7 displays the composition and concentration of fatty acids in braised pork. Generally, the higher the unsaturated fatty acid (UFA) level of a product, the greater its nutritional value (53). The fatty acids in braised pork comprise SFA and UFA, with MUFA being the most abundant, followed by SFA, and UFA being substantially more abundant than SFA. The primary components of fatty acids in braised pork are palmitic acid (C16:0), stearic acid (C18:0), oleic acid (C18:1n9c), and linoleic acid (C18:2n6c). A similar result is reached by Da et al. (54). Cooking methods and stewing

TABLE 7 Fatty acid composition (g/100 g lipids) of braised pork with different sweeteners in various stewing times.

	WS1	WS2	SF1	SF2	MG1	MG2	SG1	SG2
C8:0	0.013 ± 0.00 ^a	0.012 ± 0.00 ^a	0.010 ± 0.00 ^a	0.006 ± 0.00 ^a	0.015 ± 0.01 ^a	0.006 ± 0.00 ^a	0.008 ± 0.00 ^a	0.010 ± 0.00 ^a
C10:0	0.048 ± 0.01 ^a	0.049 ± 0.01 ^a	0.050 ± 0.01 ^a	0.039 ± 0.03 ^a	0.046 ± 0.02 ^a	0.039 ± 0.00 ^a	0.043 ± 0.01 ^a	0.044 ± 0.01 ^a
C12:0	0.060 ± 0.01 ^{ab}	0.065 ± 0.02 ^{ab}	0.054 ± 0.02 ^b	0.052 ± 0.01 ^b	0.078 ± 0.01 ^a	0.053 ± 0.01 ^b	0.073 ± 0.00 ^a	0.059 ± 0.01 ^{ab}
C14:0	0.573 ± 0.04 ^a	0.578 ± 0.02 ^a	0.574 ± 0.03 ^a	0.525 ± 0.01 ^{ab}	0.562 ± 0.01 ^a	0.525 ± 0.02 ^{ab}	0.607 ± 0.02 ^a	0.556 ± 0.03 ^{ab}
C15:0	0.013 ± 0.01 ^b	0.015 ± 0.01 ^b	0.012 ± 0.00 ^b	0.011 ± 0.00 ^b	0.011 ± 0.00 ^b	0.017 ± 0.00 ^b	0.053 ± 0.01 ^a	0.015 ± 0.01 ^b
C16:0	9.140 ± 0.16 ^a	8.150 ± 0.12 ^b	9.150 ± 0.17 ^a	8.500 ± 0.10 ^b	9.200 ± 0.17 ^a	8.640 ± 0.30 ^b	9.660 ± 0.06 ^a	8.940 ± 0.09 ^a
C17:0	0.065 ± 0.00 ^{ab}	0.058 ± 0.02 ^b	0.065 ± 0.01 ^{ab}	0.065 ± 0.01 ^{ab}	0.084 ± 0.02 ^a	0.070 ± 0.03 ^{ab}	0.091 ± 0.02 ^a	0.083 ± 0.02 ^a
C18:0	4.520 ± 0.39 ^a	3.890 ± 0.22 ^c	4.210 ± 0.12 ^{bc}	4.620 ± 0.24 ^a	4.440 ± 0.20 ^{ab}	4.350 ± 0.09 ^{ab}	4.526 ± 0.17 ^a	4.480 ± 0.19 ^{ab}
C20:0	0.668 ± 0.04 ^b	0.774 ± 0.10 ^a	0.674 ± 0.04 ^b	0.664 ± 0.02 ^b	0.762 ± 0.07 ^a	0.737 ± 0.05 ^a	0.791 ± 0.01 ^a	0.690 ± 0.02 ^{ab}
C21:0	nd	nd	nd	nd	nd	nd	0.986 ± 0.05 ^a	0.862 ± 0.12 ^a
C23:0	0.089 ± 0.01 ^{ab}	0.091 ± 0.03 ^a	0.099 ± 0.00 ^a	0.110 ± 0.02 ^a	0.091 ± 0.01 ^a	0.118 ± 0.01 ^a	0.072 ± 0.02 ^{bc}	0.069 ± 0.01 ^c
C24:0	0.006 ± 0.01 ^a	nd	0.012 ± 0.00 ^a	nd	nd	nd	nd	nd
SFA	15.195 ± 0.38 ^{bcd}	13.683 ± 0.08 ^d	14.910 ± 0.18 ^{cd}	14.594 ± 0.13 ^{cd}	15.288 ± 0.14 ^{bc}	14.555 ± 0.35 ^c	16.911 ± 0.24 ^a	15.808 ± 0.44 ^b
C14:1	0.110 ± 0.02 ^a	0.092 ± 0.01 ^a	0.074 ± 0.02 ^{ab}	0.100 ± 0.01 ^a	0.089 ± 0.01 ^{ab}	0.096 ± 0.00 ^a	0.031 ± 0.00 ^c	0.020 ± 0.01 ^c
C16:1	1.210 ± 0.10 ^a	1.110 ± 0.18 ^{ab}	1.190 ± 0.10 ^a	1.160 ± 0.06 ^{ab}	1.250 ± 0.05 ^a	0.980 ± 0.01 ^b	1.220 ± 0.04 ^a	1.110 ± 0.02 ^{ab}
C17:1	0.072 ± 0.01 ^{ab}	0.060 ± 0.01 ^b	0.080 ± 0.02 ^a	0.076 ± 0.01 ^{ab}	0.081 ± 0.01 ^a	0.072 ± 0.00 ^{ab}	0.080 ± 0.01 ^a	0.070 ± 0.00 ^{ab}
C18:1n9t	4.180 ± 0.16 ^{ab}	4.230 ± 0.23 ^a	4.280 ± 0.07 ^a	3.980 ± 0.09 ^c	4.390 ± 0.06 ^a	4.190 ± 0.01 ^{ab}	4.320 ± 0.03 ^a	4.100 ± 0.01 ^{ab}
C18:1n9c	13.260 ± 0.92 ^a	11.260 ± 0.52 ^{ab}	13.220 ± 0.11 ^a	12.230 ± 0.20 ^{ab}	13.660 ± 0.05 ^a	12.106 ± 0.10 ^{ab}	13.066 ± 0.06 ^a	11.250 ± 0.14 ^{ab}
C22:1n9	0.108 ± 0.01 ^b	0.108 ± 0.01 ^b	0.880 ± 0.03 ^a	0.111 ± 0.02 ^b	0.104 ± 0.01 ^b	0.094 ± 0.00 ^b	nd	nd
MUFA	18.941 ± 1.20 ^{ab}	16.860 ± 0.94 ^c	19.724 ± 0.25 ^a	17.656 ± 0.22 ^{bc}	19.574 ± 0.14 ^a	17.538 ± 0.12 ^{bc}	18.717 ± 0.09 ^{ab}	16.549 ± 0.15 ^c
C18:2n6c	7.700 ± 0.02 ^b	5.960 ± 0.04 ^c	7.890 ± 0.03 ^b	7.840 ± 0.03 ^b	8.200 ± 0.04 ^{ab}	7.300 ± 0.03 ^b	9.900 ± 0.05 ^a	8.100 ± 0.04 ^{ab}
C18:3n6	0.083 ± 0.01 ^{ab}	0.086 ± 0.01 ^{ab}	0.086 ± 0.00 ^{ab}	0.085 ± 0.00 ^{ab}	0.089 ± 0.00 ^{ab}	0.087 ± 0.00 ^{ab}	0.100 ± 0.00 ^a	0.088 ± 0.00 ^{ab}
C18:3n3	0.066 ± 0.01 ^a	0.061 ± 0.00 ^{ab}	0.070 ± 0.00 ^a	0.066 ± 0.00 ^a	0.071 ± 0.00 ^a	0.068 ± 0.00 ^a	0.070 ± 0.00 ^a	0.069 ± 0.00 ^a
C20:2n6	0.331 ± 0.01 ^b	0.369 ± 0.02 ^{ab}	0.369 ± 0.01 ^{ab}	0.424 ± 0.01 ^a	0.365 ± 0.01 ^{ab}	0.349 ± 0.00 ^{ab}	0.394 ± 0.01 ^a	0.347 ± 0.00 ^{ab}
C20:3n6	0.067 ± 0.01 ^{ab}	0.059 ± 0.01 ^b	0.070 ± 0.00 ^a	0.059 ± 0.00 ^b	0.069 ± 0.01 ^{ab}	0.072 ± 0.00 ^a	0.082 ± 0.00 ^a	0.078 ± 0.00 ^a
C20:3n3	0.027 ± 0.00 ^a	nd	nd	nd	nd	nd	nd	nd
C20:4n6	0.026 ± 0.00 ^a	0.011 ± 0.00 ^b	0.029 ± 0.01 ^a	0.021 ± 0.00 ^a	0.018 ± 0.00 ^b	0.016 ± 0.00 ^b	0.026 ± 0.00 ^a	0.012 ± 0.00 ^b
C22:2n6	0.010 ± 0.00 ^a	nd	0.006 ± 0.00 ^a	nd	nd	nd	nd	nd
C20:5n3	0.063 ± 0.00 ^{ab}	0.047 ± 0.01 ^{bc}	0.098 ± 0.00 ^a	0.048 ± 0.00 ^{bc}	0.050 ± 0.00 ^b	0.018 ± 0.00 ^c	0.052 ± 0.00 ^{ab}	0.060 ± 0.00 ^{ab}
C22:6n3	0.013 ± 0.01 ^a	0.012 ± 0.00 ^a	0.014 ± 0.00 ^a	0.012 ± 0.00 ^a	0.013 ± 0.00 ^a	0.020 ± 0.00 ^a	0.012 ± 0.00 ^a	0.012 ± 0.00 ^a
PUFA	8.386 ± 0.04 ^d	6.605 ± 0.05 ^e	8.631 ± 0.04 ^c	8.555 ± 0.03 ^c	8.874 ± 0.05 ^a	7.930 ± 0.04 ^c	10.636 ± 0.07 ^a	8.765 ± 0.04 ^b
UFA	27.326 ± 1.22 ^{bc}	23.464 ± 0.98 ^e	28.355 ± 0.29 ^{ab}	26.211 ± 0.21 ^{cd}	28.448 ± 0.18 ^{ab}	25.468 ± 0.15 ^d	29.353 ± 0.16 ^a	25.314 ± 0.17 ^d
UFA:SFA	1.798 ± 0.11 ^{ab}	1.715 ± 0.07 ^{bc}	1.902 ± 0.02 ^a	1.803 ± 0.03 ^{ab}	1.861 ± 0.02 ^{ab}	1.750 ± 0.04 ^b	1.735 ± 0.02 ^{bc}	1.601 ± 0.03 ^c
n-3	0.169 ± 0.02 ^a	0.120 ± 0.01 ^{cd}	0.182 ± 0.00 ^a	0.126 ± 0.01 ^{bc}	0.133 ± 0.00 ^{bc}	0.106 ± 0.00 ^d	0.134 ± 0.00 ^{bc}	0.140 ± 0.00 ^b
n-6	0.516 ± 0.01 ^d	0.524 ± 0.03 ^d	0.559 ± 0.02 ^{bc}	0.589 ± 0.01 ^{ab}	0.541 ± 0.00 ^{cd}	0.524 ± 0.01 ^d	0.602 ± 0.01 ^a	0.525 ± 0.01 ^d
n-6/n-3	3.052 ± 0.26 ^e	4.376 ± 0.35 ^{bc}	3.066 ± 0.08 ^e	4.671 ± 0.19 ^{ab}	4.062 ± 0.10 ^{cd}	4.940 ± 0.09 ^a	4.499 ± 0.04 ^{ab}	3.755 ± 0.02 ^d

nd Indicates that no substance was detected. Different letters within a row are significantly different ($p < 0.05$).

duration considerably affect the fatty acids in stewed pork. The loss of fatty acids was due to oxidation and degradation during cooking, and the SFA concentration typically declines as cooking time increases (55). Table 7 demonstrates that the SFA level of pork braised for 60 min was lower than that of pork cooked for 40 min.

Meanwhile, the SFA level of pork cooked with various sweeteners was greater than that of pork braised with white sugar (56). In addition, fatty acids in meat may reflect its nutritional worth and influence the taste development of cooked meat products. During heat processing, the Maillard

reaction and fatty acid oxidation proceed rapidly, leading to the formation of several volatile chemicals and the possible reduction of fatty acids (57).

Unsaturated fatty acids can lower cholesterol and protect against atherosclerosis (58). The amount of C20:2n6 in WS and SF grew as the stewing duration increased. It has been observed that the Maillard reaction products possess the antioxidant properties of clear hydroxyl radicals and chelating metal ions, which can limit the self-oxidation of unsaturated fatty acids during processing (59). During the same stewing period, the fatty acid content of the other three groups was greater than

that of the WS group for the pork braised with white sugar. The UFA of SF and MG fell somewhat, and their ratio of UFA to SFA was more significant than that of the other two groups, with SF-1 having the highest ratio at 1,902. Shen et al. (60) investigated the antioxidant capability of adding Chinese pickled and dried mustard (PDM) to steaming pork. As additional PDM increased, the ratio of unsaturated to saturated fatty acids rose, and lipid and protein oxidation reduced. The SFA concentration of pork braised in SG was greater than that of pork cooked in WS. Kaur et al. (61) shown that stevia glycoside inhibits the production of free fatty acids.

Also critical was the balance between n-3 and n-6 PUFA in the diet. N-3 PUFA has the potential to lessen the risk of cardiovascular disease, and DHA (C22:6n3) and EPA (C20:5n3) have garnered a great deal of interest due to their high nutritional value (62). The n-6/n-3 ratio of braised pork stayed between 3 and 4, and the WS was significantly less than that of other groups. The ratio of n-6 to n-3 was more excellent in WS, SF, and MG than after 40 min of stewing. Highest EPA concentration in SF1 was 0.098 g/100 g, while maximum DHA concentration in MG2 was 0.020 g/100 g. However, area, variety, and processing conditions will also impact the composition and amount of fatty acids, which distinguishes this study from others (63).

Volatile flavor compounds

Utilizing the Gas chromatography-ion mobility spectrometry (GC-IMS) system's high separation and sensitivity, the effects of various sweeteners and cooking time on the volatile taste components of braised pork were determined. The volatile organic chemicals in the sample can be easily recognized without needing specific sample preparation. The system recognized 75 volatile taste chemicals, including 17 aldehydes, 14 alcohols, 14 esters, 6 ketones, 2 acids, 4 hydrocarbons, and 3 nitrogen-containing compounds, based on the retention time and drift time. In Table 8, the recognized component's name, CAS number, formula, and a few taste descriptors are included, whereas compounds with signals that could not be determined are not. Figure 3 demonstrates that the composition of volatile compounds in braised pork cooked with different sweeteners and varied stewing durations varies significantly. Using the spectrum of the braised pork sample stewed for 40 min with white sugar as a reference, a small number of red spots represent the higher compound content at the peak position, whereas blue spots represent the lower compound content at the peak position in the spectrum of the braised pork sample stewed for 60 min with white sugar. More red and blue spots emerged in the spectra of cooked pork samples sweetened with *Siraitia grosvenorii*, mogroside, or stevia glycoside. This illustrates that the stewing duration affects the volatile flavor components of the sample, but adding various sweeteners significantly

affects the volatile components of braised pork. In other words, the signal strength of some chemicals is greater in samples including other sweeteners, but the concentration of specific compounds is lower in braised pork with white sugar.

In order to examine thoroughly and intuitively the impacts of sweeteners and cooking time on the volatile substance composition of braised pork, the program may generate a fingerprint-based on the signal peak of volatile substances in each sample spectrum (Figure 4). Each row in the picture depicts the overall signal peak, whereas each column depicts the signal peak of the same volatile organic component across many samples. Each dot represents a volatile component, and the color indicates the concentration of volatile compounds. The more the content, the brighter the hue. Homologous chemicals contain monomer and dimer structures, and the concentration-dependent signals and spots are also distinct. Numbers on the fingerprint denote unidentified compounds. As shown in Figure 5, the relative concentration of volatile chemicals in each sample is also estimated based on the peak volume of volatile organic compounds. As seen in Figure 4, the signal peak intensity in regions A, B, and C is more robust in the selected region than in the other two regions.

In area A, there are esters such as butyl 2-methylbutanoate, ethyl 2-methylbutanoate, ethyl acetate, ethyl butanoate, hexyl propanoate, isoamyl acetate, isobutyl propionate, *n*-Propyl Acetate, pentyl acetate, alcohols such as 2-ethyl-1-hexanol, 2-Hexanol, 1-octanol, benzyl alcohol, ethanol, octan-2-ol, organic acids such as 2-Methylbutyric acid and 3-methylpentanoic acid, and 2,3-butanedione, methylal, butanal. Their content was lower in the braised pork with white sugar as sweetener, higher in the braised pork samples with stevia glycoside as sweetener, and more in the braised pork with *Siraitia grosvenorii* and mogroside as a sweetener and stewing time of 40 min. Aldehydes such as 3-methylbutanal, 4-heptenal (z), benzaldehyde, heptanal, octanal and phenylacetaldehyde, ketones such as 1-octen-3-one, 1-penten-3-one, 2-butanone and 2-heptanone, as well as 2-ethyl-5-methylpyrazine and 2,4,6-trimethyl-pyridine in area B are generally low in braised pork with stevia glycoside as a sweetener. The content is relatively low in the braised pork with *Siraitia grosvenorii* as a sweetener and stewed for 40 min. It can be seen from area C that alcohols such as 1-heptanol, 1-pentanol and 2-hexen-1-ol, aldehydes such as 2-heptenal (E), 2-pentanal (E), hexanal and pentanal, and acetoin, delta-3-carene, gamma hexalactone and 1,8-cineole are primarily in high content in braised pork with white sugar as the sweetener. Moreover, the content of 1-pentanol, *n*-nonanal, 2-heptanone, ethyl hexanoate, *p*-cymene and other substances in the braised pork added with the mogroside group was higher.

The primary volatile taste components in braised pork are aldehydes. They are generated during heat processing and derive primarily from fat breakdown. Due to their large concentration and low threshold, aldehydes have a considerable impact on the

TABLE 8 Qualitative information of characteristic flavor compounds.

Volatiles	No.	Compounds	CAS	Formula	RI	Rt	Dt	Flavor character (64)	
Alcohol	1	Ethanol	C64175	C ₂ H ₆ O	512.1	102.0	1.051	Sweaty	
	24	1-Pentanol M	C71410	C ₅ H ₁₂ O	780.9	245.9	1.250		
	25	1-Pentanol D	C71410	C ₅ H ₁₂ O	780.1	245.2	1.507		
	26	1-Pentanol T	C71410	C ₅ H ₁₂ O	780.3	245.4	1.815		
	27	2-Hexanol	C626937	C ₆ H ₁₄ O	780.3	245.4	1.282		
	28	Cis-2-penten-1-ol	C1576950	C ₅ H ₁₀ O	795.2	258.5	1.460		
	33	2-Hexen-1-ol M	C2305217	C ₆ H ₁₂ O	859.3	324.2	1.180		
	34	2-Hexen-1-ol D	C2305217	C ₆ H ₁₂ O	859.8	324.7	1.520		
	38	1-Hexanol	C111273	C ₆ H ₁₄ O	891.6	364.0	1.326		
	63	1-Heptanol	C111706	C ₇ H ₁₆ O	1001.8	543.1	1.389		
	65	Octan-2-ol	C123966	C ₈ H ₁₈ O	1002.8	545.1	1.435		
	69	Octen-3-ol	C3391864	C ₈ H ₁₆ O	1010.2	560.1	1.161	Mushrooms	
	76	Benzylalcohol	C100516	C ₇ H ₈ O	1039.4	623.4	1.325		
	78	1,8-Cineole M	C470826	C ₁₀ H ₁₈ O	1045.2	637.0	1.289	Green	
	79	1,8-Cineole D	C470826	C ₁₀ H ₁₈ O	1046.6	640.2	1.731	Green	
	86	2-Ethyl-1-hexanol	C104767	C ₈ H ₁₈ O	1058.0	667.8	1.420		
	91	1-Octanol	C111875	C ₈ H ₁₈ O	1070.3	698.9	1.458		
	97	1-Non-anol	C143088	C ₉ H ₂₀ O	1163.9	988.8	1.535		
	Ketones	5	2,3-Butanedione	C431038	C ₄ H ₆ O ₂	592.9	131.0	1.164	
		6	2-Butanone	C78933	C ₄ H ₈ O	600.9	134.4	1.252	Spicy
13		1-Penten-3-one	C1629589	C ₅ H ₈ O	656.3	161.0	1.316		
18		Acetoin M	C513860	C ₄ H ₈ O ₂	735.0	209.8	1.205		
19		Acetoin D	C513860	C ₄ H ₈ O ₂	734.6	209.5	1.330		
41		2-Heptanone M	C110430	C ₇ H ₁₄ O	899.3	374.1	1.259	Fruity	
42		2-Heptanone D	C110430	C ₇ H ₁₄ O	900.1	375.3	1.630	Fruity	
61		1-Octen-3-one M	C4312996	C ₈ H ₁₄ O	995.1	529.9	1.266		
62		1-Octen-3-one D	C4312996	C ₈ H ₁₄ O	995.4	530.5	1.684		
Aldehyde		7	Butanal	C123728	C ₄ H ₈ O	607.8	137.4	1.292	Green fruity
	11	3-Methylbutanal M	C590863	C ₅ H ₁₀ O	660.7	163.4	1.407	Malty	
	12	3-Methylbutanal D	C590863	C ₅ H ₁₀ O	656.3	161.0	1.316	Malty	
	15	Pentanal M	C110623	C ₅ H ₁₀ O	695.2	183.3	1.180	Spicy, fruity	
	16	Pentanal D	C110623	C ₅ H ₁₀ O	696.7	184.3	1.423	Spicy, fruity	
	22	2-Pentenal(E) M	C1576870	C ₅ H ₈ O	757.9	227.1	1.110		
	23	2-Pentenal(E) D	C1576870	C ₅ H ₈ O	756.2	225.7	1.361		
	30	Hexanal M	C66251	C ₆ H ₁₂ O	799.8	262.7	1.254	Grassy	
	31	Hexanal D	C66251	C ₆ H ₁₂ O	801.3	264.0	1.342	Grassy	
	32	Hexanal T	C66251	C ₆ H ₁₂ O	800.6	263.4	1.563		
	43	4-Heptenal(Z)	C6728310	C ₇ H ₁₂ O	904.4	381.1	1.150		
	44	Methional	C3268493	C ₄ H ₈ OS	909.0	387.4	1.403		
	45	Heptanal M	C111717	C ₇ H ₁₄ O	909.8	388.6	1.328	Fruity	
	46	Heptanal D	C111717	C ₇ H ₁₄ O	909.8	388.6	1.695	Fruity	
	52	2-Heptenal (E) M	C18829555	C ₇ H ₁₂ O	972.9	488.6	1.257		
	53	2-Heptenal(E) D	C18829555	C ₇ H ₁₂ O	972.9	488.6	1.671		
	57	Benzaldehyde M	C100527	C ₇ H ₆ O	990.1	520.2	1.150	Fruity, berry	
	58	Benzaldehyde D	C100527	C ₇ H ₆ O	989.7	519.4	1.468	Fruity, berry	
	71	Octanal M	C124130	C ₈ H ₁₆ O	1021.8	584.5	1.400	Fatty, soapy	
72	Octanal D	C124130	C ₈ H ₁₆ O	1022.1	585.1	1.825	Fatty, soapy		
75	2,4-Heptadienal (E,E)	C4313035	C ₇ H ₁₀ O	1029.8	601.9	1.188			

(Continued)

TABLE 8 (Continued)

Volatiles	No.	Compounds	CAS	Formula	RI	Rt	Dt	Flavor character (64)
	81	Phenylacetaldehyde	C122781	C ₈ H ₈ O	1048.9	645.9	1.257	
	88	2-Octenal(E) M	C2548870	C ₈ H ₁₄ O	1069.6	697.3	1.334	
	89	2-Octenal(E) D	C2548870	C ₈ H ₁₄ O	1069.3	696.3	1.820	
	92	<i>n</i> -Nonanal M	C124196	C ₉ H ₁₈ O	1097.6	773.4	1.469	Greasy, orange
	93	<i>n</i> -Nonanal D	C124196	C ₉ H ₁₈ O	1098.0	774.6	1.511	Greasy, orange
	94	<i>n</i> -Nonanal T	C124196	C ₉ H ₁₈ O	1097.6	773.4	1.942	
	95	[E]-2-nonenal	C18829566	C ₉ H ₁₆ O	1133.6	883.9	1.409	
	99	2,4-decadienal	C2363884	C ₁₀ H ₁₆ O	1249.4	1357.8	1.413	
Esters	8	Ethylacetate M	C141786	C ₄ H ₈ O ₂	623.0	144.4	1.099	Pineapple
	9	Ethylacetate D	C141786	C ₄ H ₈ O ₂	621.0	143.4	1.339	Pineapple
	17	<i>n</i> -Propylacetate	C109604	C ₅ H ₁₀ O ₂	734.5	209.5	1.160	Pear, strawberry
	20	Ethylbutanoate M	C105544	C ₆ H ₁₂ O ₂	756.9	226.3	1.219	Pineapple
	21	Ethylbutanoate D	C105544	C ₆ H ₁₂ O ₂	756.9	226.3	1.555	Pineapple
	29	Ethyl2-methylbutanoate	C7452791	C ₇ H ₁₄ O ₂	799.1	262.0	1.658	
	35	Isobutylpropionate M	C540421	C ₇ H ₁₄ O ₂	859.9	324.9	1.296	
	36	Isobutylpropionate D	C540421	C ₇ H ₁₄ O ₂	858.9	323.7	1.703	
	39	Isoamyl acetate	C123922	C ₇ H ₁₄ O ₂	896.4	370.2	1.753	Apple, pear and banana
	50	Pentyl acetate	C628637	C ₇ H ₁₄ O ₂	941.1	435.2	1.316	
	64	Ethyl hexanoate	C123660	C ₈ H ₁₆ O ₂	1001.9	543.1	1.794	Pineapple
	77	γ-Hexalactone	C695067	C ₆ H ₁₀ O ₂	1040.1	625.2	1.190	
	80	Isoamyl butyrate	C106274	C ₉ H ₁₈ O ₂	1045.9	638.7	1.398	
	82	Butyl2-methylbutanoate	C15706737	C ₉ H ₁₈ O ₂	1049.5	647.1	1.372	Pear, pineapple
	87	Diethylmalonate	C105533	C ₇ H ₁₂ O ₄	1068.7	694.7	1.249	
	90	Hexyl propanoate	C2445763	C ₉ H ₁₈ O ₂	1069.5	696.9	1.990	Fruity
	98	Hexyl butanoate	C2639636	C ₁₀ H ₂₀ O ₂	1205.7	1154.7	1.488	
Acids	49	2-Methylbutyric acid	C116530	C ₅ H ₁₀ O ₂	913.3	393.5	1.215	
	59	3-Methylpentanoic acid	C105431	C ₆ H ₁₂ O ₂	990.2	520.3	1.282	
Hydrocarbons and their derivatives	56	δ-3-Carene	C13466789	C ₁₀ H ₁₆	986.7	513.8	1.215	
	66	<i>p</i> -Cymene	C99876	C ₁₀ H ₁₄	1004.8	549.0	1.730	Mild, pleasant
	67	<i>n</i> -Butylcyclohexane	C1678939	C ₁₀ H ₂₀	1008.2	555.8	1.258	
	85	2-Methoxy-phenol	C90051	C ₇ H ₈ O ₂	1054.6	659.4	1.246	
Nitrogenous compounds	3	Ammonia M	C7664417	H ₃ N	561.4	118.6	0.846	
	4	Ammonia D	C7664417	H ₃ N	566.7	120.6	0.891	
	60	2,4,6-trimethyl-pyridine	C108758	C ₈ H ₁₁ N	994.5	528.7	1.585	
	70	2-Ethyl-5-methylpyrazine	C13360640	C ₇ H ₁₀ N ₂	1019.2	578.8	1.666	Whisky

flavor of meat and contribute considerably to the scent. Pentanal has the odor of grass and banana, hexanal has a delicate odor of new grass, heptanal has the odor of fruit, and *n*-nonanal has the odor of grease and frying. By-products of the oxidation of oleic, linoleic, and arachidonic acids are hexanal and heptanal. Oleic acid oxidizes nonanal primarily (65).


A consequence of the breakdown of phenylalanine or linoleic acid is benzaldehyde. It quickly manifests while cooking and has a bitter almond odor. Phenylacetaldehyde is mainly metabolized via the Strecker route by phenylalanine (66). With an increase in cooking time, the degree of lipid oxidation, the generation of volatile flavor compounds, and the transformation of certain aldehydes into other taste components all rise.

Figure 5 demonstrates that the amount of aldehydes in braised pork rises after 60 min of cooking; that is, both the type and amount of aldehydes increase as the cooking time increases.

Alcohols, which have a metal smell and a mushroom scent, are mainly produced through Strecker degradation and fat oxidation. In the stewing process, braised pork was rich in alcohols, with octen-3-ol, 1-pentanol, ethanol, 1-heptanol, and octan-2-ol playing a significant role in the volatile taste components. The relative alcohol concentration in samples with various sweeteners was reduced after 60 min of stewing. Contrarily, the change in the relative alcohol content of braised pork added with *Siraitia grosvenorii* was reasonably steady, while the relative alcohol content of braised pork added with

TABLE 9 Antioxidant site of peptides derived from proteins after trypsin hydrolyzed.

Protein	Annotated sequence	Position	BIOSEP sequences	Activity	Quality PEP
ATP-dependent 6-phosphofructokinase	ALVFQPVTELK	A0A286ZIJ9[747–757]	LK[10–11]	Antioxidation	0.00026007
ATP synthase subunit	VELVPPTPAEIPTAIQSLK	A0A5G2QK91[31–49]	EL[2–3], LK[18–19]	Antioxidation	7.3605E-05
Alpha-1,4 glucan phosphorylase	APNDFNLK	A0A286ZMZ9[247–254]	LK[7–8]	Antioxidation	0.014763
Peroxiredoxin-2	ASGNAHIGKPAPEFQA TAVVNGAFK	A0A287AJ76[2–26]	KP[9–10]	Antioxidation	6.4107E-13
Phosphoglycerate mutase	FCSWVDQK	A0A286ZQ31[22–29]	FC[1–2]	Antioxidation	0.041327
	FCGWFDAELSEK	B5KJG2[22–33]	FC [22–23]	Antioxidation	3.0511E-06
	MEFDICYTSVLK	B5KJG2[50–61]	SVL[58–60]	Antioxidation	9.1214E-05
Ig-like domain-containing protein	DNSQNTAYLQMNSLR	A0A287BAB3[73–87]	AY[7–8]	Antioxidation	5.1175E-13
	ADAKPSVFIFPPSK	F1STC5[107–120]	KP[38–39]	Antioxidation	0.0039453
Myosin light chain 1	KPAAAAAPAPAPAPAP APAPAPPKEEK	A0A287BJF1[9–35]	KP[1–2]	Antioxidation	3.3804E-41
	AGKPVICATQMLESNIK	A0A287AQJ5[926–942]	KP[3–4]	Antioxidation	2.165E-17
Multifunctional fusion protein	ADYNVLPASENPLLR	F1SHX0[354–368]	LLR[13–15]	Antioxidation	2.5815E-07
	VIYQGVNTPGQPVFLEGQQQL	A0A287B5J2[1213–1233]	IY[2–3]	Antioxidation	6.942E-05
Myosin binding protein C1	EWSVGEPAGEEQDKQNA NSQLSTLFLVVEKPSGGEVK	A0A5G2QM90[39–74]	KP[29–30]	Antioxidation	0.0065639
	AGAGSATLSMAYAGAR	P00346[242–257]	AY[11–12]	Antioxidation	2.125E-15
Voltage-dependent anion-selective channel protein 3	AADFQLHTHVNDGTEFGGSYQK	Q29380[175–197]	YQK[21–23]	Antioxidation	0.038426
Glyceraldehyde-3-phosphate dehydrogenase	VPTPDVSVVDLTCR	F1RM74[302–315]	LTC[11–13]	Antioxidation	4.5225E-16
Myosin-4	AEAHFSLIHAYGTVDYNTIGWLDKQ	9TV62[576–599]	AH[3–4], YNI[16–18]	Antioxidation	1.6559E-52
L-lactate dehydrogenase A chain	ATLKDQLIHNLK	P00339[2–14]	LK[3–4]	Antioxidation	1.1487E-07







FIGURE 4 Fingerprints of characteristic flavor compounds in braised pork with different sweeteners in various stewing times generated by gallery plot.

white sugar reduced the greatest. The use of cooking wine and other ingredients may cause the stewed pork's high ethanol concentration. Alcohols also contribute significantly to the creation of esters.

Unsaturated fatty acid breakdown, the Maillard process, and amino acid breakdown all produce ketones. They have a low threshold and can alter flavor. The primary ketones in braised pork are acetoin, 2,3-butanedione, 2, butanone,



2-butanone, and 2-heptanone. They smell like pork and cream, as is customary. They are said to improve the scent of meat items because of their distinctive qualities. 2-Butanone tastes like cream and butter, 2-Heptanone tastes like spicy blue cheese, and Acetoin have a nice flavor with a potent cream and fat aroma (67). The relative content of ketones is largely steady throughout hot processing, and there is little change when the stewing period is extended, as can be seen by the changes in fingerprint and relative content of volatile chemicals. Additionally, the varieties of braised pork with stevia glycoside and their ketones content are the lowest, and the levels of ketones in the samples from the SF2 and MG2 groups are nearly identical.



Esters are created when free fatty acids from lipid oxidation in muscle tissue mix with alcohols (68). They have a distinctive aromatic odor or a fruity scent, a high threshold, and no impact on flavor. The flavor of long-chain esters, such as hexyl butanoate, is somewhat greasy, but the flavor of short-chain fatty acid and alcohol esters, like ethyl acetate, is fruity (69). The aromas of pineapple, apple, pear, and banana may be detected in *N*-propyl acetate, ethyl butanoate, and isoamyl acetate, respectively. Esters make up a relatively tiny portion of the overall volatile flavor compounds. The braised pork and stevia glycoside had a higher ester content than the other groups, with ethyl acetate's most excellent ester level. After stewing for 60 min, the esters in the braised pork rose due to the ongoing interaction between alcohol and acid to create esters.

Braised pork with stevia glycoside mainly comprises an acid, while pork braised with white sugar and mogroside has a higher concentration of hydrocarbons and nitrogenous substances. A few hydrocarbons are precursors to heterocyclic chemicals. The threshold of hydrocarbons in braised pork

is high, and while individual hydrocarbons contribute little to flavor, the synergistic interaction of different alkanes and olefins enhances the flavor of the meat as a whole. Seasonings, including sugar, soy sauce, and cooking wine will be added during the braised pork preparation. Thermal deterioration, caramelization, and the Maillard reaction all occur during the stewing process and result in the production of pyridine, pyrazine, and sulfur compounds, which give the barbecue a mild and pleasant scent and give braised pork as it is distinctive flavor (70).

Peptide identifications

Following the enzymatic hydrolysis of braised pork, peptides were discovered using LC-MS/MS, and **Figures 6, 7** display the amount of distinct proteins and peptides in each group. Seven hundred twenty-four proteins, consisting of 85% distinct proteins and 5,617 peptides, were discovered. Most of these peptides come from sarcoplasmic proteins such as phosphoglycerate kinase, L-lactate dehydrogenase A chain, alpha-1,4 glucan phosphorylase, and MPs like myosin-4, actinin alpha 2, MYL2, and desmin. Peptides are biologically active protein fragments that may supply the body with nutrients. Actin, myosin-1, myosin-4, and several glycolytic enzymes have all been demonstrated in studies to be relatively abundant proteins in pork. The amount of peptides in braised pork is more significantly impacted by the variation in stewing time than by the kind of sweetener. The kind of protein reduces as cooking time increases. It could be caused by a rise in oxidation levels during stewing, a reduction in the digestibility of animal proteins, and protein breakdown, which results in a loss of peptide segments (71). Myosin binding protein and



muscle fiber structure are strongly connected. Myosin oxidation will compromise the muscle fiber structure's integrity, lowering the meat's quality. The elastic modulus of fibrin continues to climb, generating an elastic gel network (72). As the heating temperature rises, it will encourage the expansion of heavy myosin and the cross-linking of the myosin head and the denaturation of light myosin and the expansion of the myosin tail.

According to a Venn diagram, braised pork with white sugar has 2,586 peptides, braised meat with siraitin and stevia glycoside contains 2,485 peptides, and braised pork with mogroside and stevia glycoside contains 2,493 peptides (Figure 7). The amount of one peptide dropped by 180, 365, and 297 when the stewing time was increased to 60 min for the particular proteins of braised pork supplemented with white sugar, mogroside, and stevia glycoside. According to Figure 8, braised pork stewed with mogroside for 40 min had more unique peptides than other groups at the same time, while braised pork stewed with Stevia glycosides for 60 min had a higher LFQ intensity than other treatment groups and roughly the same coverage of proteome sequences. In general, increasing the number of unique peptides and proteome coverage can

improve the detection of myosin complexes. Sweeteners may affect the quantity of myosin peptides in the experiment.

Bioactive peptides, such as antioxidant and antibacterial peptides, which can control physiological activity and have a variety of uses in functional foods, are typically non-toxic and very stable (73). Cooking will impact the release of bioactive peptides and the digestibility of meat products following thermal processing. When mogroside or stevia glycoside was added to braised pork, several antioxidant peptides showed more prominent expression peaks, and stew for 60 min produced higher peak intensities than stewing for 40 min. Bioactive peptides can more accurately depict the changes that braised pork undergoes throughout stewing and can serve as a foundation for evaluating the meat's nutritional worth. The research also demonstrates a connection between glyceraldehyde-3-phosphate dehydrogenase and changes in meat color and myoglobin levels. By preventing lipid oxidation, peroxiredoxin can also lessen the impact on myoglobin oxidation (74). These proteins are primarily found in braised pork with mogroside and stevia glycoside, which may give braised meat with sugar replacements additional nutritional worth or practical benefits.




FIGURE 8

(A) The numbers of unique peptides counts per protein, (B) LFQ intensity of peptide, (C) sequence coverage, and (D) the number of peptides of samples with different sweeteners.

Correlation

The physical and chemical characteristics of braised pork with various sweeteners at various stewing durations are connected (Figure 9A). Positive correlations between SH and CH, negative correlations between CH and TBA, positive correlations between CH and CL, and negative correlations between MC and CL can be found. Braised pork suffers severe CL during a protracted stewing period. The degree of oxidation increases with increasing TBA value and carbonyl concentration while decreasing sulfhydryl content. The WS2 and SF2 groups are negatively linked with other samples, and there is a positive association between other groups, according to the correlation coefficient (Figure 9B). While braised pork does vary in quality during the first 40 min of stewing, the difference becomes more noticeable after the first 60 min.


Figure 9C, which depicts the results of a correlation analysis between volatile substances, fatty acid changes, and physical and chemical indexes, revealed that PUFA was positively correlated with octan-2-ol, *n*-nonanal, butanal, and 1-hexanol, negatively correlated with 2-butanone, 1,8-cineole, and acetin, MUFA was positively correlated with 2-octanal (E), hept Aldehydes, ketones, and other chemicals all depend on fatty acids as their precursors. UFA has a more significant effect on taste compounds than SFA does. Due to the oxidation and breakdown

of fat during the stewing process, aldehydes, alcohols, and ketones are more likely to be formed. Among them, aldehydes are the primary byproducts of the breakdown of unsaturated fatty acids. The relative abundances of butyric acid and nonanoic acid were positively influenced by MUFA and C18:1n9c, according to prior research.

However, there was a negative correlation between PUFA and C18:2n6c and the relative amounts of butyric acid, (E) – 2-octanoic acid, capric acid, and acetic acid (75). Some alcohols and acids continue to react to generate esters during a prolonged stewing process, which causes a drop in alcohols and a rise in esters. While TBA and carbonyl concentration were favorably connected with volatile flavoring agents like aldehydes and alcohols, SH showed a negative correlation with dimethyl malonate and isoamyl butyrate (Figure 9D). Fatty acids, TBA, sulfur concentration, and other indicators correlate strongly with volatile chemicals. They will react during the braised pork's stewing process and particularly affect its quality.

Discussion

Pork braised with white sugar has a high CL rate, low MC, and a pH significantly higher than other sweeteners. The L* value of this set of samples was reduced as the stewing duration



rose, the a^* value of the meat's lean layer increased, the a^* and b^* values of the fat layer declined, and the color of the meat was bright red with different spring and a chewiness. According to Mena et al. (76), the texture of cooked pork products is influenced by pH, cooking techniques, and food additives. While cooking causes some components in meat to be lost, it may also enhance flavor and taste. The sample that had mogroside added had a low CL rate and a pH under 6.00, and the lean meat layer's L^* value was at its maximum after 40 min of stewing. After 60 min of stewing, the sample's overall color was brilliant, the springiness was good, and the level of softness was moderate, suggesting that mogroside might be used in place of white sugar to make braised pork with better sensory attributes.

Due to the effects of heating temperature and duration during stewing, lipid and protein oxidation in pork happens. The amount of oxidation increased with longer heating times, increasing TBARS value, carbonyl content, and decreasing sulfhydryl content. The amount of fatty acids, particularly SFA, is also reduced with longer heating times. The carbonyl content was the lowest at 60 min, and the changes in TBARS value, SH, and CH of stewed pork with mogroside were noticeably less than those of other groups. Mogroside possesses strong thermal stability and can prevent oxidation during stewing, as seen from the graph. The UFA: SFA ratio was similarly higher than that of braised pork with other sweets, suggesting that natural antioxidants have a protective impact on fatty acids. The unsaturated fatty acids of SF and MG were significantly

reduced, and the UFA: SFA ratio was also higher. According to research, fruit extract contains natural antioxidant active ingredients that can block fat oxidation, help preserve meat, and maintain the fatty acid composition of meat products in a stable state (77).

Additionally, stevia glycoside was utilized as a natural sweetener and antioxidant. The samples produced by varied stewing periods are superior to those containing white granulated sugar in terms of hardness, chewiness, and other characteristics. However, their springiness and cohesiveness were somewhat lower than those of the SF and MG groups, demonstrating that stevia glycoside has the same texture and quality advantages as a sugar replacement. Historically, stevia was primarily employed in dessert creation and less frequently in meat preparation. During various phases of stewing, the L^* value of the lean layer in this group was more significant than that of the group using white granulated sugar, showing that stevia glycoside can improve meat sheen and color. The TBARS value and carbonyl content of stewed pork with stevia glycoside were much lower than those of white granulated sugar. The change of SH and the other change trend were likewise more diminutive than white granulated sugar, showing that stevia glycoside has antioxidant characteristics and may suppress the oxidation reaction during the cooking process.

As a fruit sweetener, *Siraitia grosvenorii* may enhance the flavor and quality of stewed pork, and the CL rate and water content of braised pork are more than in the white sugar group. When adding *Siraitia grosvenorii* to pork stew for 60 min, the sheer force attained its minimal quality and texture level. Due to the presence of water-soluble pigment in *Siraitia grosvenorii*, the b^* value is much greater than that of other groups. After 60 min of stewing, the pork stewed with *Siraitia grosvenorii* had a low TBARS value, and its sulfhydryl and carbonyl content was lower than that of the white granulated sugar group and more than that of the other two groups. *Siraitia grosvenorii* includes other components, such as polyphenols and flavonoid compounds (78), which can significantly decrease lipid oxidation and protein oxidation during stewing.

There are considerable changes in the taste ingredient composition of pork cooked with various sugars. Under the impact of stewing duration, the composition and concentration of flavoring ingredients alter proportionally. Aldehydes, alcohols, and ketones are the primary volatile taste components of braised pork after cooking. Increase the pork stew's alcohols, esters, and acids using sweeteners. Aldehydes and esters rose when the stewing time was prolonged, whereas ketones dropped. Previously, Song et al. (79) prepared braised meat with a sauce containing a Maillard reaction intermediate; the resulting flavor composition was equal to that of braised pork made with white sugar, providing a better notion

for the use of sugar alternatives in braised meat. Xu et al. (80) have demonstrated that the antioxidants tert butyl hydroquinone, rosemary, and L-ascorbyl palmitat may block the degradation of optimal taste components and limit the production of odor chemicals during frying. As natural antioxidants, mogroside and stevia glycoside offer benefits as sugar substitutes; however, the oxidation level in suppressing odor formation or other phases of meat products needs more research.

Related to the protein of braised pork was its peptide. The alteration of myofibril structure significantly affects the quality of stewed meat. After braising, pork was broken down into polypeptide molecules by trypsin, which includes antioxidant peptides and has a distinct nutritional benefit. The research on polypeptide modification in meat concentrates mainly on the effects of cooking temperature, duration, and storage conditions. The sweetness was similar to that of braised pork with white sugar, which showed a declining tendency and may be more impacted by stewing duration, based only on the change in polypeptide amount. To improve the ability to recognize the unique peptides of braised meat stewed with various sweeteners, it is necessary to further study the digestive function and digestibility in the gastrointestinal tract. It is also simpler to screen the meat protein components rich in specific peptides and more nutrient-dense to more effectively realize the transformation of homemade food into industrial food.

Conclusion

This study compared and examined the effects of various sweeteners on the quality of stewed pork at various stewing periods. The outcomes demonstrated that the quality, fatty acids, and taste of braised pork were influenced differently by sweeteners and stewing duration. The flexibility and chewability of braised pork may be diminished by prolonging the stewing period. Braised pork's fatty acid content fell at the same time the muscle fiber structure changed, its sulfhydryl content rose, and its carbonyl content rose. By maintaining sweetness and improving the braised pig's sensory quality, the braised pork with SF, MG, and SG can lessen oxidation and create secondary products. The mogroside-added sample has a lower level of protein oxidation, *Siraitia grosvenorii* fruit has a more potent inhibitory impact on lipid oxidation, and stevia glycoside has better benefits for enhancing meat color. The SF and MG group's UFA: SFA ratio was more excellent than the WS group's, and the SG group's braised pork included more SFA than the WS group. By identifying peptides, one may more accurately pinpoint the peptide sequence that gives cooked meat its antioxidant benefits and improves braised pork's nutritional value. This study can guide using sugar substitutes in meals and turning traditional cuisines into processed foods.

Data availability statement

The supplementary data presented in this study are deposited in the Uniprot-Susscrofa repository. All original data included in this study are available upon request by contact with the corresponding author.

Author contributions

Z-gH and YZ conducted the experiments, data analyzing, and writing—original draft preparation. Y-yC and Y-qZ contributed by the investigation and data curation. M-yD, DZ, and HS conducted the methodology, supervising data, and project administration. All authors contributed to the article and approved the submitted version.

Funding

This study was supported by Guangxi Zhuang Autonomous Region level financial science and technology

plan project Guangxi key research and development plan (2018AB49021).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Jiang S, Xue D, Zhang Z, Shan K, Ke W, Zhang M, et al. Effect of Sous-vide cooking on the quality and digestion characteristics of braised pork. *Food Chem.* (2022) 375:131683. doi: 10.1016/j.foodchem.2021.131683
- Yingqiu L, Chunbao L, Fan Z, Xisha L, Yun B, Guanghong Z. The effects of long—duration stewing combined with different cooking and heating methods on the quality of pork belly. *J Food Process Preserv.* (2016) 40:94–102.
- Li H, Tang X, Wu C, Yu S. Maillard reaction in Chinese household-prepared stewed pork balls with brown sauce: potentially risky and volatile products. *Food Sci Hum Wellness.* (2021) 10:221–30.
- Castro-Muñoz R, Correa-Delgado M, Córdova-Almeida R, Lara-Nava D, Chávez-Muñoz M, Velásquez-Chávez VF, et al. Natural sweeteners: sources, extraction and current uses in foods and food industries. *Food Chem.* (2022) 370:130991. doi: 10.1016/j.foodchem.2021.130991
- Antonik N, Janda K, Jakubczyk K. Characteristics of sweeteners used in foods and their effects on human health. *Pomeranian J Life Sci.* (2020) 66:57–65.
- Yin KJ, Xie DY, Zhao L, Fan G, Ren JN, Zhang LL, et al. Effects of different sweeteners on behavior and neurotransmitters release in mice. *J Food Sci Technol.* (2020) 57:113–21.
- Mejia E, Pearlman M. Natural alternative sweeteners and diabetes management. *Curr Diabetes Rep.* (2019) 19:1–10.
- Chandini SK, Amanat A, Annamalai M. Health benefits of substituting added sugars with fruits in developing value-added food products: a review. *Int J Nutr Pharmacol Neurol Dis.* (2020) 10:75–90.
- Shivani, Thakur BK, Mallikarjun CP, Mahajan M, Kapoor P, Malhotra J, et al. Introduction, adaptation and characterization of monk fruit (*Siraitia grosvenorii*): a non-caloric new natural sweetener. *Sci Rep.* (2021) 11:1–13. doi: 10.1038/s41598-021-85689-2
- Li Y, Zhu W, Cai J, Liu W, Akihisa T, Li W, et al. The role of metabolites of steviol glycosides and their glucosylated derivatives against diabetes-related metabolic disorders. *Food Funct.* (2021) 12:8248–59. doi: 10.1039/d1fo01370j
- Mora MR, Dando R. The sensory properties and metabolic impact of natural and synthetic sweeteners. *Comprehensive Rev Food Sci Food Saf.* (2021) 20:1554–83. doi: 10.1111/1541-4337.12703
- Majid K, Huilin L, Jing W, Baoguo S. Inhibitory effect of phenolic compounds and plant extracts on the formation of advance glycation end products: a comprehensive review. *Food Res Int.* (2020) 130:108933. doi: 10.1016/j.foodres.2019.108933
- Liu L, Jiang Y, Shen M, Xie M, Nie S. Inhibitory effect of natural antioxidants on the formation of thermal processing hazards in red braised pork. *Food Sci.* (2021) 42:50–7.
- Li J, Zhang J, Yang Y, Zhu J, He W, Zhao Q, et al. Comparative characterization of lipids and volatile compounds of Beijing Heilium and Liaiwu Chinese black pork as markers. *Food Res Int.* (2021) 146:110433. doi: 10.1016/j.foodres.2021.110433
- Honegger LT, Bryan EE, Price HE, Ruth TK, Boler DD, Dilger AC. The effect of cooking method and cooked color on consumer acceptability of boneless pork chops. *Foods.* (2021) 11:106. doi: 10.3390/foods11010106
- Ruirui W, Feng H, Liang Z, Qiannan L, Chunjiang Z, Hong Z, et al. Changes in the texture, microstructures, colour and volatile compounds of pork meat loins during superheated steam cooking. *Int J Food Sci Technol.* (2019) 54:2821–30.
- Emilio AP, Gilberto MM, Laura ADLR, José Alberto LD, Abraham WM, Gustavo AGA. Antioxidant activity and prevention of pork meat lipid oxidation using traditional Mexican condiments (pasilla dry pepper, achiote, and mole sauce). *Food Sci Technol.* (2014) 34:371–8.
- Monica RM, Alonzo AG. Changes in the physicochemical and microbiological properties of pork and chicken meats at ambient storage condition. *Sci J Meat Technol.* (2020) 61:44–53.
- Díaz P, Linares MB, Egea M, Auqui SM, Garrido MD. TBARS distillation method: revision to minimize the interference from yellow pigments in meat products. *Meat Sci.* (2014) 98:569–73. doi: 10.1016/j.meatsci.2014.06.012
- Park D, Xiong YL. Oxidative modification of amino acids in porcine myofibrillar protein isolates exposed to three oxidizing systems. *Food Chem.* (2007) 103:607–16.
- Hua Y, Peipei M, Youling LX, Lizhen M, Changlu W, Yingchun Z. Oxidation in HiOx-packaged pork Longissimus muscle predisposes myofibrillar and sarcoplasmic proteins to N-nitrosamine formation in nitrite-curing solution. *Meat Sci.* (2013) 95:465–71. doi: 10.1016/j.meatsci.2013.05.038

22. Jordi O, Rafael S, María JJ, Sancho B. Shelf life of meat from lambs given essential oil-free rosemary extract containing carnosic acid plus carnosol at 200 or 400 mg kg⁻¹. *Meat Sci.* (2014) 96:1452–9.
23. Bazina N, He J. Analysis of fatty acid profiles of free fatty acids generated in deep-frying process. *J Food Sci Technol.* (2018) 55:3085–92.
24. Chen YJ, Zhou GH, Zhu XD, Xu XL, Tang XY, Gao F. Effect of low dose gamma irradiation on beef quality and fatty acid composition of beef intramuscular lipid. *Meat Sci.* (2007) 75:423–31. doi: 10.1016/j.meatsci.2006.08.014
25. Jiang S, Zhao D, Nian Y, Wu J, Zhang M, Li Q, et al. Ultrasonic treatment increased functional properties and in vitro digestion of actomyosin complex during meat storage. *Food Chem.* (2021) 352:129398. doi: 10.1016/j.foodchem.2021.129398
26. José ML, Aurora C, Paulo EM, Rubén D. Physicochemical properties of foal meat as affected by cooking methods. *Meat Sci.* (2015) 108:50–4. doi: 10.1016/j.meatsci.2015.05.021
27. Song Y, Huang F, Li X, Han D, Zhao L, Liang H, et al. Water status evolution of pork blocks at different cooking procedures: a two-dimensional LF-NMR T1-T2 relaxation study. *Food Res Int.* (2021) 148:110614. doi: 10.1016/j.foodres.2021.110614
28. Pawar RS, Krynskiy AJ, Rader JJ. Sweeteners from plants—with emphasis on *Stevia rebaudiana* (Bertoni) and *Siraitia grosvenorii* (Swingle). *Anal Bioanal Chem.* (2013) 405:4397–407. doi: 10.1007/s00216-012-6693-0
29. Talukder S. Effect of dietary fiber on properties and acceptance of meat products: a review. *Crit Rev Food Sci Nutr.* (2015) 55:1005–11.
30. Vilma Q, Juanita V, Valeska U, Nicol G, María JA, Carolina A, et al. The role of non-caloric sweeteners in sensory characteristics of pastry products. *Foods.* (2019) 8:329. doi: 10.3390/foods8080329
31. Suman SP, Nair MN, Joseph P, Hunt MC. Factors influencing internal color of cooked meats. *Meat Sci.* (2016) 120:133–44.
32. Dong J, Kou X, Liu L, Hou L, Li R, Wang S. Effect of water, fat, and salt contents on heating uniformity and color of ground beef subjected to radio frequency thawing process. *Innovat Food Sci Emerg Technol.* (2021) 68:102604.
33. Xu-Bing C, Jing-Jing Z, Jun-Hua L, Min L, Lei M, Jing C, et al. Potential AMPK activators of cucurbitane triterpenoids from *Siraitia grosvenorii* Swingle. *Bioorganic Med Chem.* (2011) 19:5776–81. doi: 10.1016/j.bmc.2011.08.030
34. Karp S, Wyrwiz J, Kurek M, Wierzbicka A. Physical properties of muffins sweetened with steviol glycosides as the sucrose replacement. *Food Sci Biotechnol.* (2016) 25:1591–6. doi: 10.1007/s10068-016-0245-x
35. Alam SS, Selamat J, Maimunah S, Alfi K. Chemical profiling of different types of soy sauce and the relationship with its sensory attributes. *J Food Qual.* (2016) 39:714–25.
36. Kim JH, Lee HJ, Shin DM, Kim TK, Kim YB, Choi YS. The dry-aging and heating effects on protein characteristics of beef *Longissimus dorsi*. *Korean J Food Sci Anim Resour.* (2018) 38:1101–8. doi: 10.5851/kosfa.2018.e43
37. Zhou X, Liang C, Xu Q, Liu G, Su X. Study on the separation, purification and some properties of the Protease from the Fruits of *Siraitia grosvenorii*. *Food Res Dev.* (2012) 33:36–40.
38. Gadekar YP, Shinde AK, Sonawane GG, Saxena VK, Bahire SV. Use of natural plant proteases in tenderization of mutton. *Ind J Small Ruminants.* (2019) 25:89–94.
39. Harada N, Ishihara M, Horiuchi H, Ito Y, Tabata H, Suzuki YA, et al. Mogrol derived from *Siraitia grosvenorii* mogrosides suppresses 3T3-L1 adipocyte differentiation by reducing cAMP-response element-binding protein phosphorylation and increasing AMP-activated protein kinase phosphorylation. *PLoS One.* (2016) 11:e0162252. doi: 10.1371/journal.pone.0162252
40. Ewa R, Jaroslawa R, Ewa CS, Krzysztof T. Effects of cooking methods on changes in fatty acids contents, lipid oxidation and volatile compounds of rabbit meat. *LWT.* (2019) 110:64–70.
41. Berardo A, De Maere H, Stavropoulou DA, Rysman T, Leroy F, De Smet S. Effect of sodium ascorbate and sodium nitrite on protein and lipid oxidation in dry fermented sausages. *Meat Sci.* (2016) 121:359–64. doi: 10.1016/j.meatsci.2016.07.003
42. Seki H, Tamura K, Muranaka T. Plant-derived isoprenoid sweeteners: recent progress in biosynthetic gene discovery and perspectives on microbial production. *Biosci Biotechnol Biochem.* (2018) 82:927–34. doi: 10.1080/09168451.2017.1387514
43. Ophélie F, Karim EK, Sandrine M. The natural antioxidant rosmarinic acid spontaneously penetrates membranes to inhibit lipid peroxidation in situ. *Biochim Biophys Acta (BBA) Biomembranes.* (2011) 1808:2973–80. doi: 10.1016/j.bbame.2011.08.011
44. Lin W, Liu F, Li Y, Ran L, Hou R, Zhong Y, et al. Inhibitory mechanism of plant polyphenols on protein oxidation and their application for shelf-life extension of meat products: a literature review. *Food Sci.* (2019) 40:266–72.
45. Longteng Z, Qian L, Shiliang J, Zhan H, Yongkang L. Effect of different stunning methods on antioxidant status, in vivo myofibrillar protein oxidation, and the susceptibility to oxidation of silver carp (*Hypophthalmichthys molitrix*) filets during 72 h postmortem. *Food Chem.* (2018) 246:121–8. doi: 10.1016/j.foodchem.2017.10.140
46. Jongberg S, Gislason NE, Lund MN, Skibsted LH, Waterhouse AL. Thiol-quinone adduct formation in myofibrillar proteins detected by LC-MS. *J Agric Food Chem.* (2011) 59:6900–5. doi: 10.1021/jf200965s
47. Zhang X, Yang X, Li Y, Wang Z, He X, Sun J. Effect of peroxy radical-induced oxidation on functional and structural characteristics of walnut protein isolates revealed by high-resolution mass spectrometry. *Foods.* (2022) 11:385. doi: 10.3390/foods11030385
48. Murphy RY, Marks BP. Effect of meat temperature on proteins, texture, and cook loss for ground chicken breast patties. *Poultry Sci.* (2000) 79:99–104. doi: 10.1093/ps/79.1.99
49. Zhang W, Xiao S, Ahn DU. Protein oxidation: basic principles and implications for meat quality. *Crit Rev Food Sci Nutr.* (2013) 53:1191–201. doi: 10.1080/10408398.2011.577540
50. Xiaojing T, Lele S, Qianqian Y, Wladimir SV, Xingmin L, Ruitong D. Comparative analysis of quality uniformity of ohmic and water bath heating treated pork batter with different fat content. *J Food Process Preserv.* (2020) 44:e14377.
51. Jing-Rong C, Xue-Ming L, You-Sheng Z, Ye-Hui Z, Zhi-Yi C, Dao-Bang T, et al. Protective effects of *Momordica grosvenori* extract against lipid and protein oxidation-induced damage in dried minced pork slices. *Meat Sci.* (2017) 133:26–35. doi: 10.1016/j.meatsci.2017.04.238
52. Liu H, Wang C, Qi X, Zou J, Sun Z. Antiglycation and antioxidant activities of mogroside extract from *Siraitia grosvenorii* (Swingle) fruits. *J Food Sci Technol.* (2018) 55:1880–8. doi: 10.1007/s13197-018-3105-2
53. Simoes JS, Mársico ET, Marshall MR, Simonne A, Yagiz Y, Xu C, et al. Applications of extracts from skin and seed muscadine grape (*Vitis rotundifolia* Michx.) waste on bacterial growth, autoxidation, and color in atlantic salmon (*Salmo salar* L.). *J Food Process Preserv.* (2019) 43:e13976.
54. Da D, Nian Y, Zou B, Zhao D, Zhang Z, Li C. Influence of induction cooking on the flavor of fat cover of braised pork belly. *J Food Sci.* (2021) 86:1997–2010. doi: 10.1111/1750-3841.15710
55. Yingqiu L, Chumbao L, He L, Xisha L, Shaolin D, Guanghong Z. Physicochemical and fatty acid characteristics of stewed pork as affected by cooking method and time. *Int J Food Sci Technol.* (2016) 51:359–69.
56. Hoa V, Seol K, Seo HW, Seong PN, Kang SM, Kim YS, et al. Meat quality characteristics of pork bellies in relation to fat level. *Anim Biosci.* (2021) 34:1663. doi: 10.5713/ab.20.0612
57. Han D, Zhang CH, Fauconnier ML, Jia W, Wang JF, Hu FF, et al. Characterization and comparison of flavor compounds in stewed pork with different processing methods. *LWT.* (2021) 144:11229.
58. Neff MR, Bhavsar SP, Braekvelt E, Arts MT. Effects of different cooking methods on fatty acid profiles in four freshwater fishes from the Laurentian Great Lakes region. *Food Chem.* (2014) 164:544–50. doi: 10.1016/j.foodchem.2014.04.104
59. In Guk H, Hyun YK, Koan SW, Junsoo L, Heon SJ. Biological activities of Maillard reaction products (MRPs) in a sugar-amino acid model system. *Food Chem.* (2011) 126:221–7.
60. Shen Q, Wang M, Tian J, Hu L, Ren S, Chen J, et al. Effects of Chinese pickled and dried mustard on nutritional quality, sensory quality, and shelf life of steamed pork belly. *Food Sci Nutr.* (2018) 6:747–56. doi: 10.1002/fsn.3612
61. Gurveer K, Tridib KG. Effect of stevia level on chemical, microbiological, and sensory properties of dairy dessert (rasgulla) at different storage periods and temperatures. *J Food Process Preserv.* (2020) 44:e14293.
62. Connor WE. Importance of n-3 fatty acids in health and disease. *Am J Clin Nutr.* (2000) 71:171S–5S.
63. Vehovskí K, Zadinová K, Stupka R, ěitek J, Lebedová N, Okrouhlá M, et al. Fatty acid composition in pork fat: de-novo synthesis, fatty acid sources and influencing factors—a review. *Agron Res.* (2018) 16:2211–28.
64. Mingquan H, Jiaying H, Jihong W, Mouming Z, Jinyun S, Fuping Z, et al. Structural characterization of a tetrapeptide from Sesame flavor-type Baijiu and its interactions with aroma compounds. *Food Res Int.* (2019) 119:733–40. doi: 10.1016/j.foodres.2018.10.055
65. Chen M, Chen T, Qi X, Lu D, Chen B. Analyzing changes of volatile components in dried pork slice by gas chromatography-ion mobility spectroscopy. *CyTA J Food.* (2020) 18:328–35.

66. Delgado RM, Hidalgo FJ, Zamora R. Antagonism between lipid-derived reactive carbonyls and phenolic compounds in the Strecker degradation of amino acids. *Food Chem.* (2016) 194:1143–8. doi: 10.1016/j.foodchem.2015.07.126
67. Yanan S, Xiang L, Aixiang H. Multivariate analysis approach for assessing coated dry-cured ham flavor quality during long-term storage. *J Food Sci Technol.* (2021) 58:651–9. doi: 10.1007/s13197-020-04579-z
68. Miaomiao H, Dongmei L, Xinglian X, Guanghong Z, Chunbao L. Effect of postmortem aging time on flavor profile of stewed pork rib broth. *Int J Food Properties.* (2018) 21:1449–62.
69. Carrapiso AI, Noseda B, García C, Reina R, del Pulgar JS, Devlieghere F. SIFT-MS analysis of Iberian hams from pigs reared under different conditions. *Meat Sci.* (2015) 104:8–13. doi: 10.1016/j.meatsci.2015.01.012
70. Han D, Zhang C, Fauconnier ML. Effect of seasoning addition on volatile composition and sensory properties of stewed pork. *Foods.* (2021) 10:83–113. doi: 10.3390/foods10010083
71. Zou X, He J, Zhao D, Zhang M, Xie Y, Dai C, et al. Structural changes and evolution of peptides during chill storage of pork. *Front Nutr.* (2020) 7:151. doi: 10.3389/fnut.2020.00151
72. Du J, Zhou C, Xia Q. The effect of fibrin on rheological behavior, gelling properties and microstructure of myofibrillar proteins. *LWT.* (2022) 153:112457.
73. Yang F, Chen X, Huang M, Yang Q, Cai X, Chen X, et al. Molecular characteristics and structure-activity relationships of food-derived bioactive peptides. *J Integr Agric.* (2021) 20:2313–32.
74. Tao Y, Ma L, Li D, Tian Y, Liu J, Liu D. Proteomics analysis to investigate the effect of oxidized protein on meat color and water holding capacity in Tan mutton under low temperature storage. *LWT.* (2021) 146:111429.
75. Da D, Nian Y, Shi J, Li Y, Zhao D, Zhang G, et al. Characterization of specific volatile components in braised pork with different tastes by SPME-GC/MS and electronic nose. *J Food Process Preserv.* (2021) 45:e15492.
76. Mena B, Fang Z, Ashman H, Hutchings S, Ha M, Shand PJ, et al. Influence of cooking method, fat content and food additives on physicochemical and nutritional properties of beef meatballs fortified with sugarcane fibre. *Int J Food Sci Technol.* (2020) 55:2381–90.
77. Pranav C, Arun KD, Annada D, Dipanwita B, Nanda PK. Effect of Black Cumin and Arjuna fruit extract on lipid oxidation in pork nuggets during refrigerated storage. *J Meat Sci.* (2018) 13:73–80.
78. Li Y, Sen Z, Zhen-Hong L, Ying-Ming P. Chemical components of the leaves of *Siraitia grosvenorii*. *Chem Nat Comp.* (2016) 52:891–2.
79. Shiqing S, Li F, Xiaodong X, Rui X, Qian J, Tao F. Aroma Patterns Characterization of braised pork obtained from a novel ingredient by sensory-guided analysis and gas-chromatography-olfactometry. *Foods.* (2019) 8:87. doi: 10.3390/foods8030087
80. Xu L, Mei X, Wu G, Karrar E, Jin Q, Wang X. Inhibitory effect of antioxidants on key off-odors in French fries and oils and prolong the optimum frying stage. *LWT.* (2022) 162:113417.