



OPEN ACCESS

EDITED BY

Alfonso Benítez-Páez,
Principe Felipe Research Center (CIPF),
Spain

REVIEWED BY

Silvia Arbolea,
Institute of Dairy Products of Asturias
(CSIC), Spain
Francois-Pierre Martin,
H&H Group, Switzerland

*CORRESPONDENCE

Isabelle Le Huërou-Luron
isabelle.luron@inrae.fr

SPECIALTY SECTION

This article was submitted to
Nutrition and Microbes,
a section of the journal
Frontiers in Nutrition

RECEIVED 23 June 2022

ACCEPTED 30 August 2022

PUBLISHED 21 September 2022

CITATION

Charton E, Bourgeois A, Bellanger A,
Le-Gouar Y, Dahirel P, Romé V,
Randuineau G, Cahu A, Moughan PJ,
Montoya CA, Blat S, Dupont D,
Deglaire A and Le Huërou-Luron I
(2022) Infant nutrition affects
the microbiota-gut-brain axis:
Comparison of human milk vs. infant
formula feeding in the piglet model.
Front. Nutr. 9:976042.
doi: 10.3389/fnut.2022.976042

COPYRIGHT

© 2022 Charton, Bourgeois, Bellanger,
Le-Gouar, Dahirel, Romé, Randuineau,
Cahu, Moughan, Montoya, Blat,
Dupont, Deglaire and Le
Huërou-Luron. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Infant nutrition affects the microbiota-gut-brain axis: Comparison of human milk vs. infant formula feeding in the piglet model

Elise Charton^{1,2}, Alexandre Bourgeois², Amandine Bellanger³, Yann Le-Gouar¹, Patrice Dahirel², Véronique Romé², Gwenaëlle Randuineau², Armelle Cahu², Paul J. Moughan⁴, Carlos A. Montoya^{4,5}, Sophie Blat², Didier Dupont¹, Amélie Deglaire¹ and Isabelle Le Huërou-Luron^{2*}

¹STLO, INRAE, Institut Agro, Rennes, France, ²Institut NuMeCan, INRAE, INSERM, Univ Rennes, Saint-Gilles, France, ³CHU Rennes, CIC-Inserm 1414, Rennes, France, ⁴Riddet Institute, Massey University, Palmerston North, New Zealand, ⁵Smart Foods and Bioproducts Innovation Centre of Excellence, AgResearch Limited, Palmerston North, New Zealand

Early nutrition plays a dominant role in infant development and health. It is now understood that the infant diet impacts the gut microbiota and its relationship with gut function and brain development. However, its impact on the microbiota-gut-brain axis has not been studied in an integrative way. The objective here was to evaluate the effects of human milk (HM) or cow's milk based infant formula (IF) on the relationships between gut microbiota and the collective host intestinal-brain axis. Eighteen 10-day-old Yucatan mini-piglets were fed with HM or IF. Intestinal and fecal microbiota composition, intestinal phenotypic parameters, and the expression of genes involved in several gut and brain functions were determined. Unidimensional analyses were performed, followed by multifactorial analyses to evaluate the relationships among all the variables across the microbiota-gut-brain axis. Compared to IF, HM decreased the α -diversity of colonic and fecal microbiota and modified their composition. Piglets fed HM had a significantly higher ileal and colonic paracellular permeability assessed by *ex vivo* analysis, a lower expression of genes encoding tight junction proteins, and a higher expression of genes encoding pro-inflammatory and anti-inflammatory immune activity. In addition, the expression of genes involved in endocrine function, tryptophan metabolism and nutrient transport was modified mostly in the colon. These diet-induced intestinal modifications were associated with changes in the brain tissue expression of genes encoding the blood-brain barrier, endocrine function and short chain fatty acid receptors, mostly in hypothalamic and striatal areas. The integrative approach underlined specific groups of bacteria

(Veillonellaceae, Enterobacteriaceae, Lachnospiraceae, Rikenellaceae, and Prevotellaceae) associated with changes in the gut-brain axis. There is a clear influence of the infant diet, even over a short dietary intervention period, on establishment of the microbiota-gut-brain axis.

KEYWORDS

human milk, infant nutrition, intestinal immune system, intestinal physiology, microbiota, brain, hypothalamus

Introduction

Human milk (HM) is assumed to meet the nutritional needs of infants, and to promote their optimal growth and development, including cognition, and favors beneficial bacteria related to health (1). Despite WHO recommendations (2), the exclusive breastfeeding rate is still low and world-wide reaches only 44% of infants aged 0 to 5-months (3). Infant formulas (IFs) have a close nutritional composition to HM. They are not identical, however, differing compositionally and structurally. Consequently, IF may not provide all of the physiological benefits associated with HM. For instance, breastfeeding reduces the risk of developing diarrhea or otitis within the first years of life (4, 5) and modulates the development of the intestinal immune system (6–9). Its effect on intestinal barrier function is still debated (10–13). Breastfeeding also benefits on brain development, resulting in improved language and motor function and learning abilities (14–16).

HM contains a myriad of bioactive substances including proteins and lipids, and acts as a prebiotic and probiotic due to its oligosaccharides and microbial composition that are not yet mimicked in IFs (1). IFs are mostly formulated with bovine proteins, which differ from HM proteins. Consequently, to cover the infant amino acid requirements, IF must contain a higher protein content than HM (on average: 1.3 vs. 0.8–1.2 g/100 mL, respectively) (17). This is partly due to the limiting content of tryptophan in bovine whey proteins. Tryptophan has been particularly studied over the past decades in relation to both intestinal and brain functions (18–22). While HM and IF lipid concentrations are usually similar (3.4 g/100 mL), lipid composition and structure differ due to their origin, with plant lipids mostly used in IFs (23), and due to IF homogenization that transforms the large Milk Fat Globule into smaller lipid droplets (24).

These differences between HM and IF could explain reported differences in intestinal bacterial ecosystems and health outcomes. The fecal microbiota of breastfed infants has a low α -diversity and is characterized by a low relative abundance of Firmicutes at 3–6 months of age (25, 26). Proteobacteria, scarcely abundant in the first weeks of life, persist as a low abundant phylum throughout the first 2.5 years of life (25–27). *Bifidobacterium* is highly abundant in the early postnatal

days and *Bacteroides* increases over the first month of life of breastfed infants (28). In contrast, a higher α -diversity is commonly reported in IF-fed compared to HM-fed infants up to 6 months of age (26, 27, 29). The most abundant genera also differ with a higher abundance of *Clostridium*, *Enterococcus*, and *Klebsiella* in 1- and 4-week-old IF-fed infants, unlike in HM-fed infants who have a higher abundance of *Bifidobacterium*, *Bacteroides*, and *Staphylococcus* (6). The major role of the microbiota on gut immune system and barrier maturation (30–33) as well as on brain development has been widely studied (34–37). Many studies have shown that the gut microbiota influences neuro-development during the first years of life notably by reducing the risk of developing some neuropsychiatric or neurodevelopmental diseases (35, 38–41). In 3 year-old term-children, negative correlations between dominance of Clostridiales and communication, personal and social skills, and between high abundance of *Bacteroides*, low abundance of *Escherichia/Shigella* and *Bifidobacterium* with fine motor skills have been observed (42).

During the past few years, many studies have analyzed the impact of the infant diet on the microbiota and its relationship with gut function or brain development. To date, however, no integrative study has evaluated collectively the modulation of the microbiota-gut-brain axis by the infant diet. This is the aim of the present paper, comparing HM vs. IF, using Yucatan piglets as a human infant model. The suckled piglet is well established as a suitable animal model for human infants (43–45) and the Yucatan miniature pig has the advantage of needing less amount of milk. Intestinal microbiota composition, intestinal phenotypic parameters and intestinal and brain gene expression patterns were determined before proceeding with multifactorial and correlation analyses to evaluate the relationships among all the variables across the microbiota-gut-brain axis.

Materials and methods

Human milk sample collection

The protocol for HM collection for the present pre-clinical study was approved by the Institutional Review Board of South Mediterranean V (no19.12.12.65653). Two types of HM pools

were used. Regarding the first pool, HM samples were obtained frozen from the donor milk bank of the Rennes University Hospital Centre. HM samples were heat-treated following the Holder pasteurization (62.5°C, 30 min). Milk from 22 donating mothers (range of women's lactation period: 0.30 – 5.61 months post-partum) were analyzed for crude protein, tryptophan and fat content and the heat-treated HM was pooled. This pooled HM had a similar chemical composition to that of the IF (Table 1). The heat-treated HM pool was stored at –20°C and thawed overnight before being given to the piglets as detailed below. For the second HM pool, fresh HM samples were collected from 50 healthy mothers (range of women's lactation period: 1.76 – 1.96 months post-delivery), pooled and stored at 4°C until use the day after collection.

Animal study and diets

The study was designed and conducted in compliance with the current ethical standards of the European and French guidelines. The ethics committees of CREEA (Rennes Committee of Ethics in Animal Experimentation) and of France's Ministry of Higher Education and Research approved the protocol (authorization #2020020610329770). Animals were observed daily throughout the experimental protocol to ensure their welfare and they did not receive medication or antibiotic treatment.

Eighteen 10-day-old healthy Yucatan piglets (10 female and 8 male) were housed individually in stainless steel metabolic cages. Room temperature was maintained at 26 ± 2°C with a 15:9 h light/dark cycle. Piglets received 345 mL of diet/kg

BW/day partitioned in 10 meals distributed from 7:30 to 22:00. The diet was given via a drinking trough. During the study, piglets were weighed every 3 days to adjust their food ration. Piglets were fed with an adaptation diet for 8 ± 2 days (adaptation period), based on a full fat bovine milk powder (Euroserum, Sodiaal, France, Supplementary Table 1) enriched with vitamins and minerals (0.5% powder, Piglet premix Step 1, Mg2Mix, France) and rehydrated with ultrapure water (14.5 g/100 mL diet). Then, piglets were randomly assigned to one of the two dietary groups and received their specific diets for 6 days (experimental period). Allocation to each diet was balanced between groups according to BW at 10 days of age, litter origin and sex. The IF piglet group was fed with a standard cow's milk based IF, produced within our laboratory such as previously described (46). IF powder was rehydrated with ultrapure water daily (11.5 g/100 mL diet) and stored at 4°C until feeding. Due to the limited access to fresh HM, the HM piglet group was fed with the pasteurized HM pool during the first 5 days and with the fresh HM pool on the last day of the experiment. Ytterbium and Cobalt-EDTA were added to IF and HM diets at a level of 0.3 g/100 g dry matter to allow determination of digestibility values (data not shown). All diets were supplemented with liquid vanilla (0.3 g/100 mL diet) to encourage intake during both adaptation and experimental (IF or HM) periods. Dietary intake was recorded at each meal. The experiment was undertaken over 3 independent blocks.

Sample collection

During the adaptation period, feces were collected over the last 2 days and stored at –80°C for microbiota analysis. On the last day of the experiment, animals received 6 meals distributed every hour and were euthanized 30 min after the last meal by electrical stunning immediately followed by exsanguination. Blood was collected in tubes containing K²-EDTA plus an anti-dipeptidyl-peptidase-IV (anti-DPP-IV, 10 µL/mL of blood) for GLP1 analysis (Millipore, Billerica, MA, United States). After centrifugation (10 min, 2500 g, 4°C), plasma samples were stored at –80°C. Ileal digesta and tissues were collected over 60 cm anterior to the ileocecal junction and colonic digesta and tissues were collected from the first third of the colon. Ileal and colonic digesta and feces were immediately frozen in liquid nitrogen and stored at –80°C for microbiota analysis. The ileum and proximal colon were dissected and rinsed with cold phosphate buffered saline (PBS). Ten-cm segments were kept in ice-cold Dulbecco's Minimum Essential Medium (Gibco, Thermo Fisher) for immediate Ussing chamber analysis. About 100 mg of ileal and proximal colonic tissues were sampled and kept in an RNA later solution for 24 h at 4°C and stored at –20°C for RNA extraction and gene expression analysis. Adjacent segments (10-cm) were fixed in 4% paraformaldehyde for 48 h until further dehydration in ethanol and embedding

TABLE 1 Protein, tryptophan, lipid, and HM oligosaccharide content of pasteurized human milk[§] (PHM) and the infant formula (IF) (mean ± SEM).

g/100 mL	Pasteurized human milk (PHM) [§]	Infant formula (IF)	P-value
Crude protein [□]	1.67 ± 0.05	1.44 ± 0.02	0.014
True protein [†]	1.28 ± 0.03	1.37 ± 0.02	0.085
Tryptophan	0.028 ± 0.001	0.026 ± 0.001	0.339
Lipids	3.21 ± 0.15	3.15 ± 0.06	0.714
HM oligosaccharides [‡]	0.87 ± 0.01	–	–

[§]Pool of 22 anonymous donations of human milk heat treated by Holder pasteurization range of women's lactation period: 0.3 – 5.61 months, gestational age range: 23–38 weeks.

[□]Crude protein = total nitrogen * 6.38.

[†]True protein = (total nitrogen – non-protein nitrogen) * 6.38.

[‡]Mean concentration (g/100 mL) of the top 20 oligosaccharides found in the highest concentrations: 2'FL: 0.163 ± 0.010; 3'FL: 0.090 ± 0.004; 6'SL/LNT: 0.086 ± 0.002; LNDFH1: 0.051 ± 0.006; MFLNH 3: 0.039 ± 0.001; LNFP2: 0.037 ± 0.003; LNFP1: 0.035 ± 0.007; 2HexNAc4Hex3Fuc: 0.029 ± 0.001; LDFT/DFL: 0.027 ± 0.001; DFLNH: 0.025 ± 0.002; Neu5Ac4Hex2HexNAcFuc: 0.021 ± 0.001; LNFP3: 0.020 ± 0.001; DFLNH2: 0.016 ± 0.001; 3'SL: 0.015 ± 0.0004; 6'SLNT: 0.014 ± 0.01; LnNT: 0.014 ± 0.001; DFLNH a: 0.013 ± 0.001; LNH/LnNH: 0.010 ± 0.0003; DFLNH3: 0.010 ± 0.01.

in paraffin, for morphometry analysis and GLP1, chromogranin and goblet cell counting. Finally, ileal and colonic tissues (1 cm) were immediately frozen in liquid nitrogen and stored at -80°C until GLP1 extraction and assay. Two pieces of liver (100 mg each) were collected, immediately frozen in liquid nitrogen and stored at -80°C for RNA extraction. Immediately after euthanasia, the brain was extracted and four regions of interest (hypothalamus, striatum, prefrontal cortex, and hippocampus from the right hemisphere) were sampled. Brain samples were immediately frozen in liquid nitrogen and stored at -80°C for RNA extraction and gene expression analysis.

Diet biochemical analysis

Crude and true protein content

The total nitrogen content was measured in duplicate using the Dumas method (ISO 14891:2008) on a LECO FP 828 analyzer after calibration using EDTA. A protein factor conversion of 6.38 was used to determine the crude protein content. Total N was corrected for non-protein nitrogen and then multiplied by the protein factor conversion of 6.38 to determine the true protein content.

Tryptophan content

The tryptophan content was determined based on the method of the European Commission (47) and the ISO 13904:2016 (48). HM and rehydrated IF were heated up to 35°C . Each diet sample, containing at least 20 mg of crude protein, was inserted in a 13 mL screw cap Teflon tube and diluted with 2 mL of ultrapure water containing 1.05 g of octa-hydrated barium hydroxide before a 16 h hydrolysis in autoclave at 110°C . After hydrolysis, 5-methyl-tryptophan was added as an internal standard at a level of 1.0 – 3.5 mg/L before mixing and cooling down on ice for 15 min. After centrifugation at 4,000 g for 2 min at 8°C , supernatants were diluted in acid acetic 10%, filtered through a 0.2 μm pore-size membrane (Chromafil Xtra Filter, 13 mm, PTFE) prior to HPLC analysis. Samples were analyzed by RP-HPLC using a Symmetry C18 (5 μm) column (2.1 mm \times 150 mm, WATERS) with an isocratic elution (95% Sodium acetate 0.4% pH = 4.5; 5% Pure Acetonitrile) at 0.25 mL/min. An L-tryptophan standard with a range of concentration from 0 to 10 mg/L, corrected by the internal 5-methyl-tryptophan standard (3.57 mg/L), was used for tryptophan content determination. The detection was made by fluorimetry, using excitation and emission wavelengths of 280 and 346 nm, respectively. Tryptophan content was corrected for losses occurring during basic hydrolysis, estimated after performing multiple hydrolysis of the samples.

Total lipid content

Diet samples (500 μL) were precisely weighed into a 15 mL screw cap Teflon tube before addition of 10 mL of Folch reagent (chloroform/methanol, 2/1, v/v). After 1 h of rotative agitation,

2 mL of KCL 0.8% was added. After centrifugation (5 min, 450 g, 20°C), the solvent phases were rinsed twice with a chloroform/methanol/KCl 0.8% solution (3/48/47, v/v/v) and filtered (Whatman filter paper, 1PS). The filtrate solvent was evaporated under a nitrogen flow in a 40°C water bath and the remaining total lipids were precisely weighed.

Microbiota analysis

Extraction of total bacterial DNA from feces, colonic and ileal digesta was performed as described in the instruction guide of the Quick-DNA Fecal/Soil Microbe Miniprep Kit (ZYMO Research). After extraction, bacterial DNA extracts were sequenced for 16s rRNA using Illumina Miseq protocol (49) (INRAE GenoToul platform, Toulouse).

Morphometry and immunohistochemical analyses

Morphometric analysis was performed after alcian blue and periodic acid Schiff staining on 7 μm sections of formalin-fixed, paraffin-embedded ileal and proximal colonic tissues. Sections were examined under a light microscope (Nikon Eclipse E400, Nikon Instruments, France) using image analysis software (NIS-Elements AR 3.0, Nikon Instruments) as described by Le Bourgot et al. (50). Villus and crypt length, width and surface area were measured in at least 15–20 crypt-villus units per piglet. Goblet cells were also counted per villus and per crypt using the same staining condition. Immunohistochemical analysis of ileum and colon sections was processed as previously described (51) to determine the number of enteroendocrine (chromogranin A-labeled) cells and GLP1-containing cells per area of mucosa.

Plasma and tissue GLP1 concentration

GLP1 content was extracted from ileal and proximal colonic tissue by homogenization of 1 g of tissue in 5 mL of ethanol acid solution (1% HCl 12 M, 74% absolute ethanol, 25% H_2O) (Polytron 3100, Kinematica, 24,000 rpm, 2 s \times 20 s). After 24 h at 4°C , samples were centrifuged (20 min, 2,000 g, 4°C) and supernatants diluted (1:1000 and 1:250 for ileum and colon, respectively). Intestinal and plasma GLP1 concentration was measured using a GLP1 active ELISA kit (Millipore), according to the manufacturer instructions.


Ex vivo permeability measurement

Ileal and colonic permeability measurements were performed using Ussing chambers (Physiological Instruments,

San Diego, CA, United States). Permeability was determined using tracer molecules Na-FITC for paracellular permeability and horseradish peroxidase (HRP) for transcellular permeability. The tracer molecules were added into the apical compartment and those transferred through the epithelium were analyzed in the serosal compartment. Concentration of Na-FITC in the samples collected at 30-min intervals for 120 min from the serosal buffer was measured by fluorimetry (fluorimeter LB940 Mithras, Berthold Technologies, Thoiry, France), while concentration of HRP was determined using spectrophotometry (Multiskan spectrum, Thermo Labsystem, Midland, Canada) after enzymatic reaction using *o*-dianisidine as substrate (Sigma-Aldrich). Mucosal-to-serosal fluxes were then calculated and expressed as ng/cm²/h (52).

Gene expression analysis

Total RNA extraction from intestinal tissues and liver was performed using the “NucleoSpin® RNA” kit (Macherey Nagel) or using the RNeasy Plus Universal Mini kit (Qiagen) for brain tissues. Extracted RNA were quantified using a Denovix spectrophotometer. RNA quality and integrity controls were performed with Agilent RNA 6000 Nano kit utilizing an Agilent 2100 Bioanalyzer (Agilent Technologies France, Massy, France) and by calculation of the RIN (RNA Integrity Number). All RIN had good quality. Reverse transcription was then performed on 2 µg of extracted RNAs with the High Capacity Complementary DNA Reverse Transcription Kit, as previously described (51).



The gene expression analysis was carried out in brain and intestine using the Smartchip Real time PCR through syber green technology using the Wafergen Smartchip cycler and Smartchip Multisample Nanodispenser (Biogenouest Genomics and the EcogenO core facility of Rennes, France). Two dedicated porcine smartchips were designed in-house to specifically investigate targeted gene expression in the intestine ('Porcine Gut Smartchip') and in the brain ('Porcine Brain Smartchip'). For the 'Porcine Gut Smartchip,' the expression of 106 genes targeted on specific intestinal functions and of 12 housekeeping genes was analyzed in ileum and colon (**Supplementary Table 2**). The porcine Gut Smartchip was focused on genes related to immune system, barrier function, endocrine function, digestion/nutrient carriers, and tryptophan pathways. For the 'Porcine Brain Smartchip,' the expression of 63 targeted genes and of 10 housekeeping genes was analyzed in the 4 areas of interest (**Supplementary Table 3**). The porcine Brain Smartchip was focused on genes related to barrier function, immune system, endocrine function, neurosynaptogenesis function, neurotransmitters, nutrient carriers and tryptophan pathways. The steadiest housekeeping genes, selected with *Genorm software*, were PPIA, RPL4, HPTR1, and POLR2G for the intestine (ileum and colon), and ACTB, ALDOA, B2M, HPTR1, PGK1, RPL4, and YWHAZ for the brain areas. In liver, the expression of 5 genes of interest (IDO, KMO, KYAT, TDO, and TPH1) were also analyzed using RT-qPCR technology of the PCR Step One Plus (Applied Biosystems). Relative expressions of the target genes were determined using the $2^{-\Delta\Delta C_t}$ method to compare the IF group to the HM group. Full dataset is available on an online dataverse <https://doi.org/10.57745/5FHAYQ>.

Statistical analysis

Data are presented as mean \pm standard error of the mean (SEM).

Microbiota data analysis

Raw sequences that were obtained from microbiota analysis were analyzed using the bioinformatic pipeline FROGS (Find Rapidly OTU with Galaxy Solution) software (53).

The descriptive analysis of the structure (α and β diversity) of microbiota was conducted with the Phyloseq function (EdgeR package, Bioconductor). The α -diversity indices used were Observed species and Chao1 representing the bacterial richness, and Shannon and InvSimpson representing the bacterial equitability. Significant differences between dietary groups were assessed using ANOVA (aov function). Phylogenetic β diversity was studied using the Unifrac distance and group differences was evaluated with principal co-ordinate analysis (PCoA) and permutational multivariate analysis (PERMANOVA) of

variance using distance matrices. Differences in phyla, families and genera were assessed with pairwise comparisons, after aggregation at the desired taxonomic rank (phyloseq' tax_glom function) by using EdgeR package (Bioconductor). Multiple testing corrections (False Discovery Rate) were used to avoid false positives (significance threshold = 0.05).

Unidimensional analysis

All statistical analyses were performed using the R software, version 3.6.2 (54).

A linear model was used to test the statistical significance of the dietary treatment on milk lipid, protein and tryptophan dietary contents. A linear mixed model was used to test the impact of diet, time, block, sex and their 2-by-2 interactions on piglet growth and food intake. Piglet was considered as a random effect, and other parameters were fixed effects. A linear model was used to test the impact of diet, block and sex on gene expression, permeability measurement and morphometry data where parameters were fixed effects. For both models, when the sex or block effects were non-significant ($p > 0.1$), these factors were removed from the linear model. The normality distribution and the homoscedasticity of the residuals of each linear model were tested using Shapiro–Wilk and Levene's tests, respectively (55). Models were considered as acceptable for $p > 0.05$. When the raw data did not fulfill these model assumptions, a natural logarithmic transformation of the data was performed prior to running the linear models. If the assumptions were still not satisfied, data were tested with a non-parametric Wilcoxon test. Differences were considered as statistically significant for $p < 0.05$ and a trend for difference at $0.05 < p < 0.1$.

Data correlation

Pearson correlation coefficients were determined between gut microbiota variables (phylum, family and genus abundances in ileum and colon), intestinal variables (morphometry, permeability, gene expression, goblet cell counting, GLP1

TABLE 2.1 Dietary impact on relative abundance of bacterial phyla in colon and feces of HM- and IF-fed piglets.

Comparison	Phylum	<i>p</i> -adjusted	Log ₂ (Fold Change)
HM vs. IF (colon)	Bacteroidetes	0.006	−2.18
	Deferribacteres	0.043	−6.58
	Fusobacteria	0.006	−3.18
HM vs. IF (feces)	Actinobacteria	0.013	2.30
	Bacteroidetes	0.001	−2.20
	Deferribacteres	0.001	−6.60
	Fusobacteria	0.004	−3.20

Log₂FoldChange, log₂(abundance HM/abundance IF). The Foldchange indicates the differential abundance of phyla in HM group compared to IF group. For example: *Bacteroidetes* abundance is lower in HM colon compared to IF one.

TABLE 2.2 Dietary impact on relative abundance of bacterial families in ileum, colon and feces of HM- and IF-fed piglets.

Comparison	Phylum	Family	<i>p</i> -adjusted	Log ₂ (FoldChange)
HM vs. IF (ileum)	Firmicutes	Erysipelotrichaceae	<0.001	-8.63
		Proteobacteria		
		Aeromonadaceae	<0.001	13.84
		Pseudomonadaceae	<0.001	8.45
HM vs. IF (colon)	Actinobacteria	Xanthomonadaceae	0.005	4.63
		Coriobacteriaceae	<0.001	4.29
	Bacteroidetes	Eggerthellaceae	0.030	-3.32
		Marinifilaceae	<0.001	-5.34
		Muribaculaceae	<0.001	-4.94
		Prevotellaceae	<0.001	-5.66
		Rikenellaceae	<0.001	-3.83
	Epsilonbacteraeota	Campylobacteraceae	0.006	7.16
	Firmicutes	Erysipelotrichaceae	0.001	-3.31
		Peptococcaceae	<0.001	-20.57
	Proteobacteria	Veillonellaceae	<0.001	6.05
		Aeromonadaceae	0.026	5.91
		Desulfovibrionaceae	0.006	2.27
		Enterobacteriaceae	<0.001	3.53
Pasteurellaceae		0.012	-1.96	
Unclassified_Bradymonadales order		0.005	-9.94	
HM vs. IF (feces)	Actinobacteria	Eggerthellaceae	0.009	-4.01
		Bacteroidetes		
		Marinifilaceae	<0.001	-6.9
		Muribaculaceae	<0.001	-6.25
		Prevotellaceae	0.013	-3.3
		Rikenellaceae	<0.001	-4.63
	Deferribacteres	Deferribacteraceae	0.012	-4.95
	Firmicutes	Family XIII	<0.001	-5.92
		Peptococcaceae	0.002	-7.74
	Proteobacteria	Veillonellaceae	<0.001	4.45
		Aeromonadaceae	0.007	3.98
		Enterobacteriaceae	<0.001	3.37
		Nitrosomonadaceae	<0.001	-5.39
Unclassified_Bradymonadales order		<0.001	-27.1	

HM, human milk; IF, infant formula. The Foldchange indicates the differential abundance of families in HM group compared to IF group. For example: Erysipelotrichaceae abundance is lower in HM ileum compared to IF.

content, and cell numbers), and brain variables (gene expression over the four brain areas). Correlations were considered significant when $p < 0.05$ and $|r| > 0.7$. Full dataset is available on an online dataverse <https://doi.org/10.57745/5FHAYQ>.

Multidimensional analysis

The Multi-factor analysis [FACTOMINE R package (56, 57)] was performed on all the data including microbiota, intestinal (ileum and colon) and brain (four areas) variables to integrate the global dietary effect on the microbiota-gut-brain axis. Variables were divided into three groups: (1) microbiota (ileum and colon) variables; (2) intestinal (ileum and colon) variables; (3) brain variables.

Results

Diet composition

The dietary crude protein content was significantly higher in HM than in IF (Table 1) while there was no statistically

significant difference for the content in true protein, tryptophan or lipid between diets. HM contained 0.87 ± 0.01 g/100 mL oligosaccharides while no oligosaccharides were added to the IF (Table 1).

Piglet growth and food intake

There was no significant diet effect on piglet growth, with an average daily weight gain of 54 ± 5 g/day. The dietary intake was significantly higher for HM on the 2nd day of the experimental period, while no further differences in intake between the HM and IF groups was observed. HM and IF daily intakes were on average 270 ± 10 and 240 ± 10 g/kg BW/day, respectively.

Intestinal microbiota

Before the start of the experimental diet period, the fecal α -diversity was similar between piglets (Figure 1A), while feeding HM vs. IF induced changes in fecal α -diversity

(Figure 1A). The fecal Shannon index of HM-fed piglets was significantly reduced after a 6-day dietary intervention, illustrating a significantly lower richness and equitability of HM-induced bacterial ecosystem compared to the IF-induced one. Significantly lower Chao1 (data not shown) and Shannon α -diversity indices were also observed in the colonic digesta for HM-fed piglets, but no difference was observed in ileal digesta (Figure 1A).

The fecal β -diversity, represented by the Unifrac index, confirmed that there was no difference before the start of the experimental period, whereas feeding piglets with HM or IF induced bacterial ecosystem changes in feces (Figure 1B) and colonic digesta (Figure 1C). In addition, the β -diversity analysis showed differences between ileal and colonic microbiota (Figure 1C), with a less apparent impact of the diet on the ileal microbiota.

TABLE 3.1 Dietary impact on relative abundance of bacterial genera in the ileum and colon of HM- and IF-fed piglets.

Comparison	Phylum	Family	Genus	<i>p</i> -adjusted	Log ₂ (FoldChange)
HM vs. IF (ileum)	Bacteroidetes	Muribaculaceae	<i>Muribaculum</i>	<0.001	17.24
		Prevotellaceae	<i>Prevotella</i>	0.047	5.28
			<i>Prevotellaceae NK3B31 group</i>	0.026	5.40
	Firmicutes	Erysipelotrichaceae	<i>Turcibacter</i>	<0.001	-8.04
		Lachnospiraceae	<i>Blautia</i>	0.047	-3.47
		Lactobacillaceae	<i>Lactobacillus</i>	<0.001	2.27
	Proteobacteria	Aeromonadaceae	<i>Aeromonas</i>	<0.001	14.29
			<i>Sutterella</i>	0.011	4.11
			<i>Kluyvera</i>	0.002	8.00
		Enterobacteriaceae	<i>Enterobacter</i>	<0.001	10.92
			<i>Raoultella</i>	<0.001	42.77
			<i>Salmonella</i>	<0.001	12.27
		Moraxellaceae	<i>Acinetobacter</i>	<0.001	6.18
		Pseudomonadaceae	<i>Pseudomonas</i>	<0.001	10.94
		Xanthomonadaceae	<i>Stenotrophomonas</i>	<0.001	9.69
HM vs. IF (colon)	Bacteroidetes	Marinifilaceae	<i>Butyricimonas</i>	<0.001	-6.87
		Muribaculaceae	<i>Muribaculum</i>	<0.001	14.63
		Prevotellaceae	<i>Alloprevotella</i>	<0.001	-9.49
		Rikenellaceae	<i>dgA-11 gut group</i>	0.015	-6.71
			<i>Rikenellaceae RC9 gut group</i>	<0.001	-5.77
	Deferribacteres	Deferribacteraceae	<i>Mucispirillum</i>	0.041	-4.40
	Firmicutes	Erysipelotrichaceae	<i>Catenisphaera</i>	<0.001	-5.79
			<i>Sharpea</i>	0.005	8.35
			<i>Turcibacter</i>	0.009	-8.53
		Family XIII	<i>Family XIII AD3011 group</i>	0.013	-6.33
		Lachnospiraceae	<i>Blautia</i>	0.001	-4.68
		Ruminococcaceae	<i>[Eubacterium] coprostanoligenes group</i>	<0.001	-6.81
			<i>Ruminiclostridium 9</i>	0.031	-4.77
			<i>Ruminococcaceae UCG-004</i>	0.002	-4.41
			<i>Subdoligranulum</i>	<0.001	-5.68
Veillonellaceae		<i>Megasphaera</i>	<0.001	7.12	
	<i>Mitsuokella</i>	0.002	11.72		
	<i>Veillonella</i>	<0.001	8.32		
Proteobacteria	Aeromonadaceae	<i>Aeromonas</i>	<0.001	17.31	
	Desulfovibrionaceae	<i>Bilophila</i>	<0.001	-3.77	
		<i>Unknown genus</i>	0.007	-4.79	
		<i>Escherichia-Shigella</i>	<0.001	3.23	
	Enterobacteriaceae	<i>Enterobacter</i>	0.002	4.13	
		<i>Salmonella</i>	<0.001	11.85	
Nitrosomonadaceae	<i>GOUTA6</i>	0.007	-5.17		

HM, human milk; IF, infant formula.

The microbial composition was diet-dependent in the colon (Figure 1D and Table 2.1) and in the feces (Table 2.1), but not in the ileum (Figure 1D). At the phylum level, the relative abundances of Bacteroidetes, Deferribacteres and Fusobacteria were lower in colon and feces (Table 2.1), but not different in the ileum, of HM-fed piglets compared to IF-fed piglets (Figure 1D). At the family level, 4, 15, and 13 taxa were differentially abundant in ileum, colon and feces of HM- vs. IF-fed piglets, respectively (Table 2.2). They corresponded to 15, 24, and 28 differential genus abundance in ileum, colon and feces, respectively (Tables 3.1, 3.2). In the ileum, all differential genera of Proteobacteria and Bacteroidetes were more abundant in HM- than in IF-fed piglets, while the Firmicutes genera were less abundant, except for *Lactobacillus*. In the colon, genera of Firmicutes and Proteobacteria represented the majority (50 and 29%, respectively) of the differentially abundant taxa. Similarly, the differential abundant genera in feces corresponded to genera of Firmicutes and Proteobacteria (39 and 32%, respectively). Fecal *Bifidobacterium* (Actinobacteria) was more abundant in HM-fed piglets than in IF-fed piglets.

Despite these different microbial profiles, fecal and colonic short chain fatty acid (SCFA) concentrations were not significantly different between HM- and IF-fed piglets with the highest concentrations being found for acetate and propionate (Supplementary Table 4). Total SCFA content was significantly higher in colon than in feces.

Morphometry, goblet cell number, and epithelial barrier permeability

The dietary impact on ileal and colonic villi and crypts was moderate. The ileal crypt length was reduced by 19% in HM- compared to IF-fed piglets (Table 4). There was no diet effect on the goblet cell number, with mean values of 22 ± 2 goblet cells per villus and 16 ± 0.5 goblet cells per crypt in ileum and 38 ± 2 goblet cells per crypt in colon, in agreement with unchanged expression of genes involved in mucosal defense (BD2, Lyz, MUC1, and MUC2) and proliferation (PCNA) (data not shown).

TABLE 3.2 Dietary impact on differential relative abundance of bacterial genera in feces of HM- and IF-fed piglets.

Comparison	Phylum	Family	Genus	p-adjusted	Log ₂ (FoldChange)	
HM vs. IF (feces)	Actinobacteria	Bifidobacteriaceae	<i>Bifidobacterium</i>	0.034	3.56	
		Coriobacteriaceae	<i>Collinsella</i>	<0.001	4.38	
	Bacteroidetes	Bacteroidaceae	<i>Bacteroides</i>	0.034	2.09	
		Marinifilaceae	<i>Butyricimonas</i>	0.002	-5.55	
		Prevotellaceae	<i>Prevotella 2</i>	<0.001	-5.94	
			<i>Alloprevotella</i>	0.033	-8.49	
		Rikenellaceae	<i>Rikenellaceae RC9 gut group</i>	<0.001	-5.31	
	Epsilonbacteraeota	Campylobacteraceae	<i>Campylobacter</i>	0.005	8.02	
	Firmicutes	Erysipelotrichaceae	<i>Turcibacter</i>	<0.001	-6.4	
			<i>Sharpea</i>	0.011	6.89	
		Lachnospiraceae	<i>Blautia</i>	<0.001	-4.91	
			<i>Lachnoclostridium</i>	0.010	2.82	
		Lactobacillaceae	<i>Lactobacillus</i>	0.026	1.55	
		Ruminococcaceae	<i>Subdoligranulum</i>	0.037	-4.81	
			<i>[Eubacterium] coprostanoligenes group</i>	<0.001	-4.49	
			<i>UBA1819</i>	0.008	6.32	
			Veillonellaceae	<i>Anaerovibrio</i>	0.007	6.35
				<i>Megasphaera</i>	<0.001	6.3
				<i>Veillonella</i>	<0.001	7.76
	Proteobacteria	Aeromonadaceae	<i>Aeromonas</i>	0.018	7.18	
			Desulfovibrionaceae	<i>Bilophila</i>	0.008	-3.16
		<i>Desulfovibrio</i>		0.008	2.6	
		<i>Mailhella</i>		0.043	-3.73	
Enterobacteriaceae		<i>Enterobacter</i>	<0.001	5.24		
		<i>Kluyvera</i>	0.014	6.22		
		<i>Raoultella</i>	<0.001	36.94		
		<i>Salmonella</i>	<0.001	14.79		
Moraxellaceae		<i>Acinetobacter</i>	0.022	3.41		

HM, human milk; IF, infant formula.

TABLE 4 Ileal and colonic tissue characteristics.

			Human milk	Infant formula	P-value
Morphometry	Ileum	Villous area (μm^2)	19442.9 \pm 1649.1	18468.9 \pm 2185.9	0.52
		Villous length (μm)	221.4 \pm 18.7	217.2 \pm 19.8	0.83
		Villous width (μm)	96.7 \pm 2.5	94.0 \pm 4.8	0.23
		Crypt area (μm^2)	4294.8 \pm 474.7	5175.0 \pm 481.3	0.21
		Crypt length (μm)	111.4 \pm 8.4	137.6 \pm 8.9	0.05
		Crypt width (μm)	46.2 \pm 2.3	43.3 \pm 2.4	0.39
	Colon	Crypt area (μm^2)	16416.9 \pm 1215.6	16964.5 \pm 1891.3	0.74
		Crypt length (μm)	270.6 \pm 13.3	292.9 \pm 29.5	0.61
		Crypt width (μm)	67.4 \pm 3.4	65.5 \pm 2.1	0.61
Permeability [†]	Ileum	Na-FITC permeability (ng/cm ² /h)	42.8 \pm 9.8	25.0 \pm 3.0	0.03
		HRP permeability (ng/cm ² /h)	61.3 \pm 13.3	33.3 \pm 6.8	0.07
	Colon	Na-FITC permeability (ng/cm ² /h)	57.7 \pm 5.3	42.3 \pm 3.8	0.03
		HRP permeability (ng/cm ² /h)	72.0 \pm 22.0	60.3 \pm 12.7	0.73
Goblet cells	Ileum	Mucin cell number per villous	23.3 \pm 2.4	20.7 \pm 2.7	0.76
		Mucin cell number per crypt	15.3 \pm 0.7	16.3 \pm 0.8	0.98
	Colon	Mucin cell number per crypt	37.4 \pm 1.0	39.0 \pm 3.9	0.70
Endocrine function	Ileum	GLP1 content (pM/g tissue)	127.9 \pm 20.0	256.5 \pm 24.1	<0.01
		GLP1 cell number per mm ²	71.0 \pm 3.9	65.4 \pm 3.6	0.40
		Chrg A cell number per mm ²	144.4 \pm 9.1	147.3 \pm 8.8	0.63
		Ratio GLP1/Chrg A cells per mm ² (%)	49.8 \pm 2.1	45.0 \pm 2.1	0.07
	Colon	GLP1 content (pM/g tissue)	105.2 \pm 15.2	122.8 \pm 14.2	0.26
		GLP1 cell number per mm ²	18.8 \pm 2.1	18.6 \pm 1.6	0.59
		Chrg A cell number per mm ²	49.2 \pm 6.3	52.8 \pm 4.4	0.66
		Ratio GLP1/Chrg A cells per mm ² (%)	39.0 \pm 1.3	35.7 \pm 2.0	0.08
	Plasma	GLP1 content (pM/g tissue)	26.8 \pm 2.7	24.8 \pm 3.8	0.68

Mean \pm SEM, $n = 9$ per group, except for permeability[†] $n_{HM} = 6$ piglets and $n_{IF} = 7$ piglets; HM, human milk; IF, infant formula, Chrg, chromogranin.

The ileal and colonic paracellular (Na-FITC passage) permeability was significantly higher in HM-fed than in IF-fed piglets (Table 4). Concomitantly, the HM diet significantly reduced the expression of genes encoding tight junction proteins of the epithelial barrier (CHD1, CLDN2, CLDN3, and MLCK) in colon (Table 5). The ileal transcellular (HRP passage) permeability tended to be higher in HM-fed piglets, while no statistically significant difference was observed between HM-fed and IF-fed piglets for the colon (Table 4).

Enteroendocrine function

Ileal tissue GLP1 concentration was two times lower in HM-fed than in IF-fed piglets (Table 4). In contrast, no statistically significant diet effect was observed either for plasma or colonic tissue GLP1 concentration, nor on GLP1-secreting cell densities. The percentage of GLP1-secreting cells compared to all enteroendocrine cells tended to be higher in the ileal and colonic tissue of HM-fed piglets (Table 4). Concomitantly, a diet effect was found for the

relative expression of genes involved in endocrine function (Table 5). Compared to IF-fed piglets, HM-fed piglets had upregulated ileal gene expression of GLP1R, GHSR, and NTS, and colonic gene expression of CCKBR, IRS2, and SOCS3, but downregulated colonic gene expression of CHGA, NPY2, and PCSK.

Immune gene expression

Ileal IL10 and DPPIV gene expression was significantly higher and BAFF, CCL2, ICAM1, TGF β 2R, TLR2, and TOLLIP gene expression tended to be higher for HM-fed piglets (Table 5). Similarly, half of the analyzed genes involved in immune function were significantly upregulated in the colon of HM-fed piglets compared to the colon of IF-fed piglets. For instance, genes encoding anti-inflammatory immune response proteins such as IL10 and its receptor (IL10Ra), TGF β , and BAFF, as well as genes encoding pro-inflammatory immune response proteins such as IL8, IL4, and TNF α were upregulated in the colon of HM-fed piglets. Likewise, TLR2, TLR4, and MYD88 genes were upregulated in the colon of HM-fed piglets.

TABLE 5 Relative expression of differentially expressed genes ($p < 0.1$) in the ileum and the colon of HM- and IF-fed piglets (mean \pm SEM).

Site	Function	Gene	Human milk	Infant formula	P-value
Ileum	Endocrine	DPPIV	1.04 \pm 0.27	0.44 \pm 0.11	0.047
		GHSR	1.08 \pm 0.17	0.70 \pm 0.20	0.043
		GLP1R	1.25 \pm 0.14	0.41 \pm 0.11	0.005
		NTS	1.05 \pm 0.13	0.49 \pm 0.12	0.028
	Immune system	IL10	1.02 \pm 0.09	0.66 \pm 0.06	0.003
		Tryptophan metabolism	IDO	1.15 \pm 0.61	0.55 \pm 0.49
	KYNU		1.07 \pm 0.16	0.49 \pm 0.18	0.005
	SERT		1.09 \pm 0.22	1.65 \pm 0.07	0.046
	Nutrient transporter		GLUT1	1.02 \pm 0.09	0.78 \pm 0.06
		PLA2G4	1.01 \pm 0.32	0.72 \pm 0.17	0.006
Colon	Barrier	Cdh1	1.04 \pm 0.09	1.30 \pm 0.08	0.046
		CLDN2	1.04 \pm 0.18	1.44 \pm 0.14	0.047
		CLDN3	1.04 \pm 0.09	1.41 \pm 0.11	0.007
		MLCK	1.04 \pm 0.10	1.42 \pm 0.12	0.029
	Endocrine	CCKBR	0.80 \pm 0.29	0.25 \pm 0.09	0.018
		CHGA	1.02 \pm 0.12	1.74 \pm 0.25	0.047
		IRS2	1.04 \pm 0.13	0.66 \pm 0.06	0.017
		NPY2R	0.77 \pm 0.08	1.06 \pm 0.21	0.028
		PCSK1	1.08 \pm 0.17	1.75 \pm 0.23	0.034
		Immune	BAFF	0.94 \pm 0.24	0.27 \pm 0.04
	CCL2		1.23 \pm 0.33	0.50 \pm 0.06	0.002
	CX3CL1		1.02 \pm 0.09	0.72 \pm 0.04	0.012
	ICAM1		1.06 \pm 0.27	0.44 \pm 0.06	0.002
	IL10		1.06 \pm 0.16	0.29 \pm 0.05	<0.001
	IL10Ra		1.11 \pm 0.24	0.48 \pm 0.06	0.003
	IL8		1.05 \pm 0.29	0.18 \pm 0.02	<0.001
	MYD88		1.04 \pm 0.10	0.73 \pm 0.02	0.005
	SOCS3		1.20 \pm 0.27	0.44 \pm 0.07	0.001
	TGF β		1.15 \pm 0.21	0.66 \pm 0.08	0.023
	Tryptophan metabolism	TLR2	1.02 \pm 0.11	0.57 \pm 0.04	0.002
		TLR4	1.09 \pm 0.22	0.57 \pm 0.07	0.029
		TNF α	1.09 \pm 0.22	0.39 \pm 0.12	0.007
		TNF α R1	1.00 \pm 0.08	0.72 \pm 0.06	0.010
		AAAD	1.05 \pm 0.15	1.31 \pm 0.10	0.043
		AANAT	1.00 \pm 0.28	0.49 \pm 0.14	0.049
		IDO	0.46 \pm 0.17	0.18 \pm 0.07	0.038
		KYNU	1.02 \pm 0.18	0.62 \pm 0.03	0.016
		MAO	1.01 \pm 0.08	1.84 \pm 0.16	<0.001
		Tph1	1.07 \pm 0.14	1.91 \pm 0.29	0.018
	Nutrient transporter	FFAR2	1.22 \pm 0.34	0.60 \pm 0.27	0.031
FFAR3		1.12 \pm 0.19	0.50 \pm 0.07	0.007	
GPR120		1.03 \pm 0.10	0.76 \pm 0.08	0.042	
MCT1		1.00 \pm 0.12	1.62 \pm 0.18	0.010	
MCT2		1.02 \pm 0.08	1.29 \pm 0.06	0.017	
MCT4		1.02 \pm 0.09	0.68 \pm 0.08	0.009	
NIACR1		0.59 \pm 0.19	0.22 \pm 0.06	0.019	
SLC38A5		1.02 \pm 0.08	1.28 \pm 0.07	0.019	
SLC6A19		1.17 \pm 0.07	1.01 \pm 0.20	<0.001	
Liver		Tryptophan metabolism	TDO	1.04 \pm 0.15	0.50 \pm 0.05

Tryptophan pathways and nutrient transporter gene expression

Among the selected genes involved in tryptophan metabolism, the expression of nine genes (three ileal and six colonic) were modulated by the dietary intervention, with upregulation of genes encoding proteins involved in the kynurenine pathway (KYNU, IDO, and AANAT) and

downregulation of genes encoding proteins in the serotonin pathway (SERT, MAO, TPH1, and AAAD) in HM-fed piglets compared to IF-fed piglets. In the liver, only the gene expression of TDO was significantly influenced by the diet, with upregulation in HM-fed piglets (Table 5).

Regarding nutrient transporters and digestion function, only GLUT1 and PLA2G4 gene expression was increased in the ileum of HM-fed piglets. In the colon, FFAR2 and FFAR3

TABLE 6 Relative expression of differentially expressed genes ($p < 0.1$) in four brain sites of HM- and IF-fed piglets (Mean \pm SEM, significant p -values are in bold).

Site	Function	Gene	Human milk	Infant formula	P -value	
Hypothalamus	Blood–brain barrier	CDH2	1.61 \pm 0.63	2.48 \pm 0.50	0.039	
		CLDN12	1.40 \pm 0.55	2.30 \pm 0.78	0.006	
		CTNNA1	1.51 \pm 0.50	1.91 \pm 0.35	0.038	
		Marveld2	0.99 \pm 0.16	3.07 \pm 1.03	0.005	
		OCLN	1.26 \pm 0.42	1.81 \pm 0.36	0.004	
	Endocrine	CCKBR	2.12 \pm 0.66	5.34 \pm 0.71	0.005	
		GLP1R	2.05 \pm 1.29	2.96 \pm 0.91	0.010	
		IRS1	1.22 \pm 0.33	2.27 \pm 0.60	0.001	
		MME	1.45 \pm 0.43	5.46 \pm 1.47	0.004	
		CX3CL1	1.04 \pm 0.22	1.67 \pm 0.21	0.009	
	Immune	TGF β	1.13 \pm 0.33	2.00 \pm 0.53	0.019	
		TLR4	1.56 \pm 0.94	2.80 \pm 0.94	0.047	
		Neurosynaptogenesis	BDNF	1.55 \pm 0.61	4.38 \pm 1.00	0.015
			CSF1R	1.44 \pm 0.52	2.68 \pm 0.76	0.004
			CYFIP2	1.22 \pm 0.28	2.40 \pm 0.41	0.002
	DGL4		1.75 \pm 0.79	3.07 \pm 1.38	0.011	
	FTO		1.11 \pm 0.37	2.00 \pm 0.46	0.010	
	Neurotransmitters	RANBP9	1.20 \pm 0.35	2.04 \pm 0.67	0.005	
		GABBR1	1.49 \pm 0.52	2.55 \pm 0.78	0.008	
		GRIN2B	0.68 \pm 0.26	1.03 \pm 0.17	0.007	
		NPY	1.68 \pm 0.69	3.71 \pm 0.88	0.009	
5HT2RB		1.52 \pm 0.49	2.34 \pm 0.54	0.053		
Tryptophan pathways	FFAR2	1.74 \pm 0.66	3.13 \pm 0.82	0.008		
	LRP1	1.22 \pm 0.42	2.00 \pm 0.53	0.005		
Nutrient transporter	MCT1	1.21 \pm 0.26	1.90 \pm 0.40	0.003		
	CDH2	0.99 \pm 0.23	2.47 \pm 0.53	0.015		
Striatum	Blood–brain barrier	CLDN12	1.54 \pm 0.30	2.73 \pm 0.34	0.003	
		LSR	0.98 \pm 0.21	2.73 \pm 0.74	0.031	
		CCKBR	1.28 \pm 0.30	3.35 \pm 0.80	0.017	
	Endocrine	IL1bR	1.30 \pm 0.27	2.96 \pm 0.97	0.024	
		HTR1F	0.98 \pm 0.34	2.01 \pm 0.32	0.040	
	Immune	TPH2	1.37 \pm 0.34	4.27 \pm 1.04	0.009	
		FFAR2	0.61 \pm 0.13	2.92 \pm 0.92	0.007	
	Tryptophan pathways	FFAR3	0.83 \pm 0.28	3.27 \pm 0.99	0.022	
		MCT2	0.99 \pm 0.45	2.82 \pm 0.53	0.007	
		MCT4	0.96 \pm 0.23	4.35 \pm 1.52	0.025	
		CDN5	1.04 \pm 0.13	1.70 \pm 0.15	0.004	
	Pre-frontal cortex	Blood–brain barrier	LSR	1.08 \pm 0.15	1.44 \pm 0.13	0.049
			INSR	1.07 \pm 0.16	1.61 \pm 0.25	0.036
Endocrine		CSF1R	1.02 \pm 0.12	1.55 \pm 0.17	0.012	
		MOG	1.08 \pm 0.16	2.00 \pm 0.40	0.019	
Nutrient transporter		SLC27A4	1.04 \pm 0.12	1.44 \pm 0.11	0.025	
Hippocampus	Endocrine	LEPR	0.83 \pm 0.38	1.56 \pm 0.29	0.012	
		5HTR2B	0.73 \pm 0.14	1.56 \pm 0.51	0.040	
	Nutrient transporter	LRP1	0.28 \pm 0.01	1.04 \pm 0.37	0.015	

A MFA Microbiote-Gut-Brain Axis




FIGURE 2
(Continued)

B

Distribution of variables significantly contributing to Dim1
Sorted by site and function





FIGURE 2
(Continued)



genes were upregulated and MCT1 and MCT2 genes were downregulated in HM-fed piglets (Table 5).

Brain gene expression

The proportion of differentially expressed genes between HM- and IF-fed piglets were high in the hypothalamus (38%) and striatum (18%) and low in the pre-frontal cortex (9%) and hippocampus (5%) (Table 6). All differentially expressed genes were downregulated in HM-fed piglets. For instance, in the hypothalamus, genes encoding tight junction proteins such as CDH2, CLDN12, OCLN, CTNBN1, or MARVELD2 were downregulated in HM- compared to IF-fed piglets. CDH2 and CLDN12 genes were also downregulated in the striatum of HM-fed piglets. In addition, the LSR gene was downregulated in HM-fed piglet striatum and pre-frontal cortex. The expression of genes involved in the endocrine function such as GLP1R, CCKBR, IRS1, and MME, was lower in HM hypothalamus. The pre-frontal cortex expression of INSR gene, the hippocampus expression of LEPR gene, and the striatum expression of CCKBR gene were lower in HM- vs. IF-fed piglets. Some genes encoding proteins of the immune function were significantly less expressed in hypothalamus (CX3CL1, TGF β , and TLR4) and striatum (IL1bR) of HM-fed piglets. Genes implicated in the neurosynaptogenesis were significantly downregulated in the hypothalamus (BDNF, CSF1R, CYFIP2, FTO, and RANBP9) and the cortex (CSF1R and MOG) of HM-fed piglets. The expression of genes encoding neurotransmitters in the hypothalamus, GABBR1, GRIN2B, and NPY, was significantly impacted by diet, with a lower expression in the HM-fed piglets. Genes encoding SCFA receptors (FFAR2 and MCT1 in hypothalamus; FFAR2, FFAR3, MCT2, and MCT4 in striatum) were less expressed in HM-fed piglets. Finally, the expression of the 5HT2B gene in hypothalamus and hippocampus, and of that HTR1F and TPH2 genes in the striatum, was downregulated in HM-fed piglets.


Overall impact of the diet on the microbiota-gut-brain axis

The multifactorial analysis aimed to analyze the relationships between diet and groups of variables ($n = 19$), gathered by site (brain, intestine, and microbiota) and function (seven functions) or phylum ($n = 7$). HM and IF variables were discriminated on both dimensions 1 and 3 (25% of variance recovered, Figure 2) unlike that on dimension 1 and 2 representation (data not shown). The groups of variables contributing the most in the discrimination between HM- and IF-fed piglets on the first dimension implicated brain functions (neurosynaptogenesis, nutrient carrier, blood-brain barrier, endocrine function, neurotransmitter, and immune

function). Those contributing the most in the separation of HM- and IF-fed piglets on the third dimension implicated microbiota (Firmicutes, Bacteroidetes, and Proteobacteria), brain functions (endocrine function, neurotransmitter, blood-brain barrier, and nutrient carrier), but also included variables representative of the intestinal functions (tryptophan metabolism, and endocrine function) (Figure 2A). Among all the variables included in this multifactorial analysis, 37% of them significantly contributed (% contribution > 0.14) to define the dimension 1, while 30% of them significantly contributed (% contribution > 0.14) to dimension 3. On a functional basis (Figures 2B,C), 15% (dimension 1) and 12% (dimension 3) of these significantly contributing variables were genes involved in brain and intestinal barrier function, respectively, 18% (dimension 1) and 11% (dimension 3) of them were genes involved in intestinal and brain immune function, 14% (dimension 1) and 14% (dimension 3) of them were genes involved in brain and intestinal endocrine functions, 14% (dimension 1) and 13% (dimension 3) of them were genes encoding nutrient carrier proteins in brain and intestine, and 9% (dimension 1) and 8% (dimension 3) of them were genes involved in intestinal and brain tryptophan pathways. An additional 6% (dimension 1) and 8% (dimension 3) of these variables were genes encoding for neurotransmitters in brain, and 10% (dimension 1) and 10% (dimension 3) were genes encoding for neurosynaptogenesis in brain. Finally, 15% (dimension 1) and up to 24% (dimension 3) of the variables significantly contributing to the definition of these dimensions were variables from the intestinal microbiota. Overall, this indicated that the diet affected the microbiota-gut-brain axis and their associated functions.

To provide a more thorough insight into the individual associations between variables, correlation analyses were performed between bacterial taxa (phyla, family, and genus levels) and ileal, colonic and brain tissue variables. Relative abundances of bacterial taxa were positively or negatively correlated ($p < 0.05$ and $|r| > 0.7$) with 10% of ileal genera correlated with 0.4% ileal variables analyzed, 33% of colonic genera with 1.0% colonic variables, and 48% of the gut genera with 0.6% cerebral variables.

At the ileal level, relative abundance of Microbacteriaceae and *Microbacterium* was positively correlated with genes related to immune function and nutrient carriers, and *Kluyvera* abundance was positively correlated to genes involved in endocrine function and tryptophan pathways (Figure 3). At the colonic level, *Anaerovibrio*, *Mitsuokella*, *Raoultella*, *Salmonella*, and *Veillonella* abundances were positively correlated and *Rikenellaceae* abundance negatively correlated with genes encoding for interleukins and their receptor, and toll-like receptors (TLR2 and TLR4) (Figure 4). Concerning the kynurenine pathway, *Anaerovibrio*, *Enterobacter*, *Mitsuokella*, *Raoultella*, *Salmonella*, and *Veillonella* were positively



correlated with KMO and KYNU genes, *Campylobacter*, *Epsilonproteobacteria*, *Enterobacter*, and *Raoultella* were positively correlated with IDO and AANAT genes, whereas *Enterobacter*, *Raoultella*, *Salmonella*, and *Veillonella* were negatively correlated with AAAD (Figure 4). It is noticeable that NPY gene expression was positively correlated with several families and genera, specifically with *Prevotella* and *Subdoligranulum* genera.

All matrix correlations between bacterial taxa (ileum and colon) and brain genes revealed positive correlations with brain site-dependent patterns. In the hypothalamus, correlations were found mainly with ileal *Acinetobacter* and *Anaerovibrio*, and colonic *Alistipes*, *Deferribacteraceae*, *Desulfovibrio*, *GOUTA6*, *Flavonifractor*, *Lachnoclostridium*, and *Mucispirillum* (Figure 5). All these taxa were correlated with the expression of 36 genes involved in functions distributed equally among all the cerebral functions studied.

The expression of hypothalamic HTR1a gene involved in tryptophan pathways and CDH5 involved in blood–brain barrier were highly correlated with ileal taxa, while the expression of MME (hormonal regulation), and BDNF and CYFIP2 (neurosynaptogenesis) were highly correlated with colonic family and genera. The expression of 30 genes in the striatum and of 36 genes in the hippocampus, related to all functions studied, was correlated with both ileal and colonic microbiota (Figures 6, 7). It is notable that the gene encoding for the dopamine receptor (DRD2b) was correlated with several ileal and colonic taxa in the striatum and the hippocampus (Figures 6, 7). For the prefrontal cortex, the expression of only 10 genes was correlated to a few ileal and colonic taxa (Figure 8).

Moreover, several functions expressed in both intestinal and brain tissues were correlated one to each other, such as genes involved in immune, barrier and endocrine functions, tryptophan pathways, nutrient receptors and transporters or synaptogenesis (data not shown). In addition, it is remarkable that nine pairs of gene expressions highly correlated in the ileum and thirteen pairs of gene expression highly correlated in the colon were also significantly correlated in brain areas (Table 7).

Discussion

For the first time, the impact of HM and IF diets on the collective microbiota-gut-brain axis in the piglet model has been demonstrated. Compared to IF, HM induced a different colonic and fecal microbiota profile, modulated intestinal gene expression, in particular those genes involved in the immune response, epithelial barrier, endocrine function, nutrients transporters and tryptophan metabolism. These results were in line with observed colonic physiological parameters, such as para-cellular permeability and the proportion of GLP-1 secreting cells. These diet-induced modifications were associated with modifications observed in the brain tissue expression of genes encoding the blood–brain barrier, endocrine function and SCFA receptors, mostly in the hypothalamic and striatal areas.

The microbiota

The present study highlighted an impact of diet on the colonic and fecal microbiota composition, such as previously reported in infants over the first months of life (30, 58). Consistent with results reported for 0 to 6 month-old breastfed infants and HM-fed piglets (26, 27, 59), the present study also showed that the fecal and colonic α -diversity decreased in HM-fed piglets compared to IF counterparts after a 5-day dietary intervention. More specifically, fecal and colonic Bacteroidetes and Fusobacteria phyla were significantly here





FIGURE 4
Correlations matrix between colonic relative gene expression and colonic microbiota ($|r| \geq 0.7, P < 0.05$). CO, colon; F, family; P, phylum; G, genera; Trp, tryptophan pathways; Horm, endocrine function; Barr, barrier function; Trsp, transport-nutrient carrier; Immu, immune function.

decreased in the HM-fed piglets, such as was observed in infants (26) and piglets (59). Among differentially abundant bacterial families, the lower abundance of Campylobacteraceae and the higher abundance of Prevotellaceae were previously observed in the colon of HM-fed piglets (60). Several dietary factors may explain the difference observed in the present study. Particularly, HM oligosaccharides likely contributed to the higher abundance of *Bifidobacterium*, *Lactobacillus*, and


Bacteroides in HM feces (61–63) as well as the high urea content in HM, which can be used as a substrate by *Bifidobacterium* (64). In addition, the quality of the dietary lipids (plant in IF vs. animal in HM) may have also contributed to the modulation of the intestinal microbiota composition, such as demonstrated previously (1). In the present study, the statistically significant correlations found between SCFA transporters or receptors and Bacteroidetes, Firmicutes and Proteobacteria phyla in the ileum



and colon are consistent with an effect of the origin (plant based vs. milk fat) of infant formula lipids (1, 51, 65, 66). In contrast, no difference in ileal microbial diversity was observed between HM- and IF-fed piglets, indicating that the most decipherable impact of the dietary treatments on the microbiota occurred in the large intestine where non-digestible molecules, especially HM oligosaccharides present in HM and urea are fermented by commensal bacteria (64, 67, 68).

Despite the pasteurization applied to the HM pool, the present study conducted to similar results

as those found in the literature for piglets and infants fed with fresh HM in comparison with an IF. The present comparison highlights the prebiotic and postbiotic role of HM in infant nutrition rather than its probiotic role. Moreover, the present observations are in compliance with previous work studying the effect of pasteurization on HM health outcomes (69). Therefore, changes in microbiota and gut-brain axis parameters are assumed to mainly result from the pasteurized milk feeding.




Intestinal physiology

The diet-induced changes in intestinal gene expression were greater in the colon (38% of total targeted genes) than in the ileum (8%), such as observed for the microbiota.

Immune and barrier functions

A remarkable observation in the present study was the higher colonic expression of genes encoding pro- but also anti-inflammatory cytokines and their receptors in HM-compared to IF-fed piglets. Such an observed HM-induced boost of the mucosal immune system agrees with data reported in breastfed infants (70–72). For instance, several studies have reported a higher fecal calprotectin content in HM-fed infants over the first weeks of life compared to IF-fed ones. Calprotectin has been considered as a valuable marker of intestinal mucosa inflammatory infiltration by neutrophils in response to early bacterial colonization (6–9). Moreover,


the importance of the Firmicutes-induced pre-weaning peak of intestinal inflammatory markers was demonstrated in rodents as an essential phase for both immune ontogeny and regulation of susceptibilities to immunopathologies later in life (32). Positive correlations between fecal calprotectin excretion and colonization by *Staphylococcus* and *Clostridium* (Firmicutes phylum) sustained the role of bacteria in the maturation of the intestinal immune system (73). Accordingly, our data support the relationship between microbiota and the mucosal immune system maturation. Veillonellaceae (Firmicutes) were more abundant in the colon and feces of HM-fed piglets. Several significant positive correlations were observed between *Anaerovibrio*, *Mitsuokella*, and *Veillonella* genera belonging to this family and genes encoding anti- and pro-inflammatory cytokines (IL10, IL10Ra, SOCS3, CCL2, IL1bR, IL8, and TNF α) and cellular signaling (ICAM1 and MYD88). Other immunomodulatory factors, such as lactoferrin or other minor proteins present in HM (74)



but not in IF, are likely to have contributed to the immune system boost.

The positive correlations found here between Lachnospiraceae (Firmicutes) and Moraxellaceae (Proteobacteria) and genes encoding barrier function (CLDN4 and MLCK) illustrate the relationship between the microbiota and intestinal barrier function. Colonic genes encoding tight junction proteins that sustain epithelial barrier integrity (75) were less expressed in HM- than in IF-fed piglets, in line with the observed increased colonic epithelial paracellular permeability. It is noteworthy that the measured permeability values of paracellular and transcellular permeabilities were in the physiological range of values reported in sow milk-fed piglets (52). The results are in agreement with the study of Lee et al. (10), who showed a higher permeability in breastfed infants, but do not corroborate other studies that reported no change or a reduction of the total gut permeability in breastfed infants compared to IF-fed infants (6, 11, 76, 77) or a reduced expression of genes encoding tight junction

proteins in HM-fed piglets (78) when determined at a specific time point. When paracellular permeability was measured in the first month of age in suckling piglets, a three- to ten-fold increase was observed in the ileum with a smaller increase in the colon (52, 76). Therefore, feeding with HM may promote an age-induced increase of intestinal permeability that coincides with the evolution of mucosal immune cells present in the intestine of young pigs. A high intestinal permeability may allow for an increased passage of molecules through the epithelium, thereby influencing the maturation of the immune system and promoting the acquisition of tolerance against indigenous bacterial and dietary antigens (76). It is well known that pig neonates are skewed toward a Th2 profile and the balance between Th1 and Th2 responses develops progressively during the first 3–4 weeks of age. An enhancement of the intestinal immune system development observed in suckling piglets whose mothers were fed a prebiotic-supplemented diet (50), was shown to have further beneficial consequences by strengthening gut defenses and vaccine



immune response post-weaning (79). Therefore, the higher ileal and colonic permeability, associated with an acceleration of the intestinal immune system maturation and changes in microbiota composition reported here in HM-fed piglets may constitute a key component of the lifelong breastfeeding health benefits.

Endocrine function

No dietary effect was observed on the concentrations of colonic and fecal SCFAs and GLP1, and GLP1R genes were not significantly correlated with bacterial taxa, suggesting that microbiota and bacterial metabolites did not play a key role in the modulation of GLP1 release in our study. Surprisingly, the ileal GLP1 content was two-fold lower in HM-fed piglets. Therefore, the reduced GLP1 content may result in a lower insulin secretion (80), corroborating the few data in the literature pointing to a lower insulinemic response in breastfed infants compared to IF fed infants (81). A similar effect of “pancreatic savings” has been previously observed in piglets supplemented with prebiotics or probiotics (50, 82). In addition to its effects on insulin secretion, GLP1 also takes part in central regulation of food intake, emotional eating and mood (83).

Further investigations are needed to characterize a diet effect on central regulation of emotional eating and mood.

Tryptophan metabolism

The diet contributed significantly to the modulation of the tryptophan metabolism pathways. The differential ileal and colonic expression of genes involved in tryptophan metabolism was in favor of the kynurenine pathway in HM-fed piglets and in favor of the serotonin pathway in IF-fed piglets in agreement with Brink et al. (59), who reported a significant increase of kynurenic acid in HM-fed piglets. In the present study, positive correlations were found between the expression of intestinal (ileal or colonic) IDO, KMO, and KYNU genes and pro-inflammatory genes (BAFF, CCL2, IL1B, IL6R, IL23A, IL8, and TNF α) in agreement with the inflammation-induced kynurenine pathway (33, 84).

There were also positive correlations in the colon between KMO and KYNU gene expression and *Anaerovibrio*, *Mitsuokella*, and *Veillonella* genera (Veillonellaceae family) and in the ileum between 5-HTR1A, AAANT, TPH1, and Enterobacteriaceae and Corynebacteriaceae families and *Propionivibrio*, which is in agreement with the known

TABLE 7 List of pairs of genes for which their expression was highly correlated ($r > 0.7$; $P < 0.05$) within the intestine and the brain.

Gene 1		Gene 2		Intestine		Brain				
Name	Function	Name	Function	r COLON	r ILEUM	r STR	r HYP	r HIP	r PFC	
CCKBR	Endocrine	FFAR3	Nutrient carrier	0.75		0.79				
CTNNB1	Barrier	EATP3	Nutrient carrier	0.70					0.70	
CTNNB1	Barrier	SLC27A4	Nutrient carrier	0.76			0.85	0.85		0.87
FFAR2	Nutrient carrier	IL1BR	Immunity	0.71			0.90	0.75		
FFAR2	Nutrient carrier	IRS2	Endocrine	0.74				0.91		
ICAM1	Immunity	IL1BR	Immunity	0.82				0.87	0.71	
ICAM1	Immunity	IRS2	Endocrine	0.82			0.94			
ICAM1	Immunity	SOCS3	Immunity	0.98				0.82		
TGF β	Immunity	IRS2	Endocrine	0.71				0.96		
TGF β	Immunity	MCT4	Nutrient carrier	0.72				0.82	0.94	
TGF β	Immunity	TLR4	Immunity	0.79				0.81		
TLR4	Immunity	IRS2	Endocrine	0.76				0.72		
TNFAR1	Immunity	MCT4	Nutrient carrier	0.75				0.71		
CTNNB1	Barrier	GLUT1	Nutrient carrier			0.74				0.84
CTNNB1	Barrier	IRS1	Endocrine		0.79				0.92	
CTNNB1	Barrier	OCLN	Barrier		0.84		0.79		0.73	
CTNNB1	Barrier	PDGFR	Barrier		0.86	0.73		0.81		
F11R	Barrier	OCLN	Barrier		0.91				0.72	
F11R	Barrier	PDGFR	Barrier		0.73		0.83			
F11R	Barrier	SLC27A4	Nutrient carrier		0.89	0.71				
GLUT1	Nutrient carrier	IRS1	Endocrine		0.76	0.87		0.81	0.73	
GLUT1	Nutrient carrier	PDGFR	Barrier		0.82				0.73	

r , coefficient of correlation; STR, striatum; HYP, hypothalamus; HIP, hippocampus; PFC, prefrontal cortex.

influence of gut bacteria in serotonin and kynurenine pathway induction (36, 85). Positive correlations between TPH1 gene expression and Erysipelotrichaceae and Lachnospiraceae families corroborate that observed in 6-month-old breastfed infants (86). Moreover, *Lactobacillus* and *Bacteroides* were more abundant in HM-fed piglets, which could influence indole metabolism, such as previously reported (33). Indole-3-propionic acid has been previously shown to be less abundant in HM-fed piglets (60). Further investigation of the specific HM components shaping the bacterial metabolomic profile is warranted. Despite a similar tryptophan content in HM and IF, the protein containing tryptophan differed, particularly regarding α -lactalbumin, present in HM and in much smaller quantity in IF. In addition, the protein structure differed in the diet, being altered in IF due to

the heat treatment applied for IF production. Whether the tryptophan release differs during HM and IF digestion remains unknown.

Brain gene expression

Interestingly, HM and IF diets induced differential expression profiles of genes encoding blood-brain barrier, endocrine and immune functions and neurosynaptogenesis in the four studied brain areas, and more particularly in the hypothalamus and striatum.

It is acknowledged that microbiota can modulate brain function (18). Accordingly, at a family level, Lachnospiraceae and Enterobacteriaceae were positively correlated with genes

encoding SCFAs and monocarboxylate transporters in the brain and blood-brain barrier function (87, 88). It is noteworthy that the genus *Alistipes* (Rikenellaceae) was correlated with numerous genes encoding blood-brain barrier, immune, endocrine and neurosynaptogenesis functions, particularly in the hypothalamus. This genus, generally considered as a commensal of human gut microbiota, ferments undigested proteins that escape digestion in the small intestine (89, 90), suggesting that a different amount of undigested or partially digested protein between HF and IF may reach the colon. Although there is contrasting evidence for on the critical role *Alistipes* plays in inflammation, gastrointestinal and behavior disorders (37, 91–93), the present results suggest that this genus may be a key actor in the microbiota-brain axis.

Microbiota-gut-brain axis: Global overview

Overall, the multifactorial analysis indicated that the brain variables, representative of most of the functions studied, the intestinal variables (mainly representative of the endocrine function and tryptophan metabolism), and the microbiota variables contributed to the differences observed for the HM-and IF-fed piglets. It illustrates the complexity of the relationships between the intestine and brain areas in interaction with the dietary treatment. Correlation analyses highlighted specific families (Veillonellaceae, Enterobacteriaceae, Lachnospiraceae, Rikenellaceae, and Prevotellaceae) whose relative abundance was correlated with several ileal, colonic and brain variables involved in different functions such as tryptophan metabolism, endocrine and barrier functions. It is remarkable that all statistically significant ileal and colonic taxa correlations with brain parameters were positive correlations. Moreover, specific ileal bacteria were correlated to the expression of similar genes in both ileum and brain content. These correlations concerned *Corynebacterium* and 5HTR1a, *Microbacterium* and LEPR, and *Actinotignum* and FFAR3. These data suggest that microbiota may act on similar functions (tryptophan metabolism, endocrine, and nutrient transport) shared on the gut-brain axis and may be key components of specific pathways. However, due to the limited data available in the literature, it is not possible to conclude on the reason of these correlations.

Finally, it should be born in mind that pasteurized HM had to be used for the piglet feeding as it was not ethically and technically possible to use such a large volume of fresh HM. It has been demonstrated that Holder pasteurization (30 min, 62.5°C) induces some protein denaturation, particularly for lactoferrin or for other bioactive and heat-sensitive proteins such as immunoglobulins or bile salt-stimulated lipase (69). This can in turn modulate their digestion (94, 95). However,

beneficial health outcomes of pasteurized HM are still acknowledged (69). Using fresh HM during the entire experimentation may have enhanced the observed differences.

Conclusion

The microbiota-gut-brain axis was modulated by the diet, with the microbiota probably playing an interface role between the diet and the host, especially in the colon. Particularly, the HM-associated microbiota profile likely improved the maturation of the intestinal epithelial barrier, immune system and endocrine functions, and modulated intestinal and cerebral tryptophan metabolism as well as several other cerebral functions in the early period of life. Undigestible nutrients such as HM oligosaccharides and urea may have contributed to the different microbiota profile. Other bioactive components of HM may also have likely contributed to the observed effects, either directly or through the microbiota. Further investigations focused on the dietary components of the IF would be useful.

Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://entrepot.recherche.data.gouv.fr/dataset.xhtml?persistentId=doi:10.57745/5FHAYQ> and <https://doi.org/10.57745/5FHAYQ>.

Ethics statement

The studies involving human participants were reviewed and approved by the Institutional Review Board of South Mediterranean V (no. 19.12.12.65653). The patients/participants provided their written informed consent to participate in this study. This animal study was reviewed and approved by Ethics Committees of CREEA (Rennes Committee of Ethics in Animal Experimentation) France's Ministry of Higher Education and Research approved the protocol (authorization #2020020610329770).

Author contributions

DD, AD, and IL contributed to the conception and design of the study. EC, AIB, AmB, YL-G, PD, VR, GR, AC, AD, and IL collected the data. EC and AIB organized the database. EC, AIB, IL, and AD performed statistical analysis. EC wrote the first draft of the manuscript. AD, IL, and EC wrote section of

the manuscript. PM, CM, DD, AD, and IL contributed to the funding of the study. All authors contributed to manuscript revision, read, and approved the submitted version.

Funding

This work was supported by the Brittany Region, France (Grant ARED, n°1318) and L'Institut Agro Rennes-Angers. Financial support was also received from the Riddet Institute, a New Zealand Centre of Research Excellence (CoRE) and from the Catalyst Fund of the New Zealand Ministry of Business, Innovation and Employment (MBIE).

Acknowledgments

We warmly thank the nurses of the milk bank and of the maternity department of Rennes University Hospital for their involvement in the project and all volunteer mothers for their breastmilk samples. We are most grateful to Biogenouest Genomics and the EcogenO core facility of Rennes (OSUR) for its technical support. We thank Sophie Le Gall and Loric Thoulouze from the BIBS facility for the HM oligosaccharides analysis (http://www.bibs.inrae.fr/bibs_eng/, UR1268 BIA, IBiSA, Phenome-Emphasis-FR ANR-11-INBS-

0012, PROBE/CaliS infrastructure, Biogenouest) and Marie-Bernadette Maillard from STLO (Rennes) for the SCFA analysis.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fnut.2022.976042/full#supplementary-material>

References

- Boudry G, Charton E, Le Huerou-Luron I, Ferret-Bernard S, Le Gall S, Even S, et al. The relationship between breast milk components and the infant gut microbiota. *Front Nutr.* (2021) 8:629740. doi: 10.3389/fnut.2021.629740
- World Health Organization. *WHO | Exclusive breastfeeding for six months best for babies everywhere*. Geneva: World Health Organization (2011).
- UNICEF. *UNICEF global database on infant and young child feeding*. New York, NY: UNICEF (2021).
- Mosca F, Gianni ML. Human milk: Composition and health benefits. *Pediatr Med Chir.* (2017) 39:155. doi: 10.4081/pmc.2017.155
- Horta BL, Victora CG, World Health Organization. Short-term effects of breastfeeding: A systematic review on the benefits of breastfeeding on diarrhoea and pneumonia mortality. *World Health Organ.* (2013):49.
- Castanet M, Costalos C, Haiden N, Hascoet J-M, Berger B, Sprenger N, et al. Early effect of supplemented infant formulae on intestinal biomarkers and microbiota: A randomized clinical trial. *Nutrients.* (2020) 12:1481. doi: 10.3390/nu12051481
- Asgarshirazi M, Shariat M, Nayeri F, Dalili H, Abdollahi A. Comparison of fecal calprotectin in exclusively breastfed and formula or mixed fed infants in the first six months of life. *Acta Med Iran.* (2017) 55:53–8.
- Li F, Ma J, Geng S, Wang J, Ren F, Sheng X. Comparison of the different kinds of feeding on the level of fecal calprotectin. *Early Hum Dev.* (2014) 90:471–5. doi: 10.1016/j.earlhumdev.2014.06.005
- Savino F, Castagno E, Calabrese R, Viola S, Oggero R, Miniero R. High faecal calprotectin levels in healthy, exclusively breast-fed infants. *Neonatology.* (2010) 97:299–304. doi: 10.1159/000255161
- Lee GO, McCormick BJJ, Seidman JC, Kosek MN, Haque R, Olortegui MP, et al. Infant nutritional status, feeding practices, enteropathogen exposure, socioeconomic status, and illness are associated with gut barrier function as assessed by the lactulose mannitol test in the MAL-ED birth cohort. *Am J Trop Med Hyg.* (2017) 97:281–90. doi: 10.4269/ajtmh.16-0830
- Catassi C, Bonucci A, Coppa GV, Carlucci A, Giorgi PL. Intestinal permeability changes during the first month: Effect of natural versus artificial feeding. *J Pediatr Gastroenterol Nutr.* (1995) 21:383–6. doi: 10.1097/00005176-199511000-00003
- Colomé G, Sierra C, Blasco J, García MV, Valverde E, Sánchez E. Intestinal permeability in different feedings in infancy. *Acta Paediatr.* (2007) 96:69–72. doi: 10.1111/j.1651-2227.2007.00030.x
- Weaver LT, Laker MF, Nelson R, Lucas A. Milk feeding and changes in intestinal permeability and morphology in the newborn. *J Pediatr Gastroenterol Nutr.* (1987) 6:351–8. doi: 10.1097/00005176-198705000-00008
- Deoni S, Dean D, Joelson S, O'Regan J, Schneider N. Early nutrition influences developmental myelination and cognition in infants and young children. *NeuroImage.* (2018) 178:649–59. doi: 10.1016/j.neuroimage.2017.12.056
- Deoni SCL, Dean DC, Piryatinsky I, O'Muircheartaigh J, Waskiewicz N, Lehman K, et al. Breastfeeding and early white matter development: A cross-sectional study. *NeuroImage.* (2013) 82:77–86. doi: 10.1016/j.neuroimage.2013.05.090
- Timby N, Domellöf M, Lönnerdal B, Hernell O. Supplementation of infant formula with bovine milk fat globule membranes. *Adv Nutr Bethesda Md.* (2017) 8:351–5. doi: 10.3945/an.116.014142
- European Commission. *Commission delegated regulation (EU) 2016/127 of 25 September 2015 supplementing regulation (EU) No 609/2013 of the European parliament and of the council as regards the specific compositional and information requirements for infant formula and follow-on formula and as regards requirements*

on information relating to infant and young child feeding (Text with EEA relevance). Luxembourg: European Commission (2016).

18. Cryan JF, O'Riordan KJ, Cowan CSM, Sandhu KV, Bastiaanssen TFS, Boehme M, et al. The microbiota-gut-brain axis. *Physiol Rev.* (2019) 99:1877–2013. doi: 10.1152/physrev.00018.2018
19. Jaglin M, Rhimi M, Philippe C, Pons N, Bruneau A, Goustard B, et al. Indole, a signaling molecule produced by the gut microbiota, negatively impacts emotional behaviors in rats. *Front Neurosci.* (2018) 12:216. doi: 10.3389/fnins.2018.00216
20. O'Rourke L, Clarke G, Nolan A, Watkins C, Dinan T, Stanton C, et al. Tryptophan metabolic profile in term and preterm breast milk: Implications for health. *J Nutr Sci.* (2018) 7:e13. doi: 10.1017/jns.2017.69
21. Kennedy PJ, Cryan JF, Dinan TG, Clarke G. Kynurenine pathway metabolism and the microbiota-gut-brain axis. *Neuropharmacology.* (2017) 112:399–412. doi: 10.1016/j.neuropharm.2016.07.002
22. Cubero J, Valero V, Sánchez J, Rivero M, Parvez H, Rodríguez AB, et al. The circadian rhythm of tryptophan in breast milk affects the rhythms of 6-sulfatoxymelatonin and sleep in newborn. *Neuro Endocrinol Lett.* (2005) 26: 657–61.
23. Le Huërou-Luron I, Lemaire M, Blat S. Quels bénéfices santé de la matière grasse laitière et des membranes des globules gras du lait (MFGM) dans les préparations pour nourrissons ? *Cah Nutr Diététique.* (2019) 54:52–60. doi: 10.1016/j.cnd.2018.10.004
24. Boulleu C, Deglaire A, De Oliveira S, Ménard O, Le Gouar Y, Carrière F, et al. Towards infant formula biomimetic of human milk structure and digestive behaviour. *OCL.* (2017) 24:D206. doi: 10.1051/ocl/2017010
25. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, et al. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci U.S.A.* (2011) 108:4578–85. doi: 10.1073/pnas.1000081107
26. Ho NT, Li F, Lee-Sarwar KA, Tun HM, Brown BP, Pannaraj PS, et al. Meta-analysis of effects of exclusive breastfeeding on infant gut microbiota across populations. *Nat Commun.* (2018) 9:4169. doi: 10.1038/s41467-018-06473-x
27. Ma J, Li Z, Zhang W, Zhang C, Zhang Y, Mei H, et al. Comparison of gut microbiota in exclusively breast-fed and formula-fed babies: A study of 91 term infants. *Sci Rep.* (2020) 10:15792. doi: 10.1038/s41598-020-72635-x
28. Jost T, Lacroix C, Braegger CP, Chassard C. New insights in gut microbiota establishment in healthy breast fed neonates. *PLoS One.* (2012) 7:e44595. doi: 10.1371/journal.pone.0044595
29. Donovan SM, Wang M, Li M, Friedberg I, Schwartz SL, Chapkin RS. Host-microbe interactions in the neonatal intestine: Role of human milk oligosaccharides. *Adv Nutr Bethesda Md.* (2012) 3:450S–5S. doi: 10.3945/an.112.001859
30. Wang M, Monaco MH, Donovan SM. Impact of early gut microbiota on immune and metabolic development and function. *Semin Fetal Neonatal Med.* (2016) 21:380–7. doi: 10.1016/j.siny.2016.04.004
31. Ximenez C, Torres J. Development of microbiota in infants and its role in maturation of gut mucosa and immune system. *Arch Med Res.* (2017) 48:666–80. doi: 10.1016/j.arcmed.2017.11.007
32. Al Nabhani Z, Dulauroy S, Marques R, Cousu C, Al Bounny S, Déjardin F, et al. Weaning reaction to microbiota is required for resistance to immunopathologies in the adult. *Immunity.* (2019) 50:1276.e–88.e. doi: 10.1016/j.immuni.2019.02.014
33. Gao J, Xu K, Liu H, Liu G, Bai M, Peng C, et al. Impact of the gut microbiota on intestinal immunity mediated by tryptophan metabolism. *Front Cell Infect Microbiol.* (2018) 8:13. doi: 10.3389/fcimb.2018.00013
34. Braniste V, Al-Asmakh M, Kowal C, Anuar F, Abbaspour A, Tóth M, et al. The gut microbiota influences blood-brain barrier permeability in mice. *Sci Transl Med.* (2014) 6:263ra158. doi: 10.1126/scitranslmed.3009759
35. Borre YE, O'Keefe GW, Clarke G, Stanton C, Dinan TG, Cryan JF. Microbiota and neurodevelopmental windows: Implications for brain disorders. *Trends Mol Med.* (2014) 20:509–18. doi: 10.1016/j.molmed.2014.05.002
36. Clarke G, Grenham S, Scully P, Fitzgerald P, Moloney RD, Shanahan F, et al. The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiatry.* (2013) 18:666–73. doi: 10.1038/mp.2012.77
37. Jiang H, Ling Z, Zhang Y, Mao H, Ma Z, Yin Y, et al. Altered fecal microbiota composition in patients with major depressive disorder. *Brain Behav Immun.* (2015) 48:186–94. doi: 10.1016/j.bbi.2015.03.016
38. Tomova A, Husarova V, Lakatosova S, Bakos J, Vlkova B, Babinska K, et al. Gastrointestinal microbiota in children with autism in Slovakia. *Physiol Behav.* (2015) 138:179–87. doi: 10.1016/j.physbeh.2014.10.033
39. Shaaban SY, El Gendy YG, Mehanna NS, El-Senousy WM, El-Feki HSA, Saad K, et al. The role of probiotics in children with autism spectrum disorder: A prospective, open-label study. *Nutr Neurosci.* (2018) 21:676–81. doi: 10.1080/1028415X.2017.1347746
40. Cerdó T, García-Valdés L, Altmäe S, Ruiz A, Suárez A, Campoy C. Role of microbiota function during early life on child's neurodevelopment. *Trends Food Sci Technol.* (2016) 57:273–88. doi: 10.1016/j.tifs.2016.08.007
41. Pärtty A, Kalliomäki M, Wacklin P, Salminen S, Isolauri E. A possible link between early probiotic intervention and the risk of neuropsychiatric disorders later in childhood: A randomized trial. *Pediatr Res.* (2015) 77:823–8. doi: 10.1038/pr.2015.51
42. Sordillo JE, Korrick S, Laranjo N, Carey V, Weinstock GM, Gold DR, et al. Association of the infant gut microbiome with early childhood neurodevelopmental outcomes: An ancillary study to the VDAART randomized clinical trial. *JAMA Netw Open.* (2019) 2:e190905. doi: 10.1001/jamanetworkopen.2019.0905
43. Moughan PJ, Birtles MJ, Cranwell PD, Smith WC, Pedraza M. The piglet as a model animal for studying aspects of digestion and absorption in milk-fed human infants. In: Simopoulos AP editor. *World review of nutrition and dietetics.* Berlin: Karger (1991). p. 40–113. doi: 10.1159/000419461
44. Darragh AJ, Moughan PJ. The three-week-old piglet as a model animal for studying protein digestion in human infants. *J Pediatr Gastroenterol Nutr.* (1995) 21:387–93. doi: 10.1097/00005176-199511000-00004
45. Roura E, Koopmans S-J, Lallès J-P, Le Huërou-Luron I, de Jager N, Schuurman T, et al. Critical review evaluating the pig as a model for human nutritional physiology. *Nutr Res Rev.* (2016) 29:60–90. doi: 10.1017/S0954422416000020
46. Yu X, Leconte N, Méjean S, Garric G, Even S, Henry G, et al. Semi-industrial production of a minimally processed infant formula powder using membrane filtration. *J Dairy Sci.* (2021) 104:5265–78. doi: 10.3168/jds.2020-19529
47. European Commission. *Commission directive 2000/45/EC of 6 July 2000 establishing community methods of analysis for the determination of vitamin A, vitamin E and tryptophan in feedingstuffs (Text with EEA relevance).* Luxembourg: European Commission (2000).
48. Agence Française de Normalisation. *Animal Feeding Stuffs – Determination Of Tryptophan Content.* ISO 13904:2016. La Plaine Saint-Denis: Agence Française de normalisation (2016) p. 14.
49. Illumina. *MiSeq system guide.* Illumina: San Diego, CA (2018).
50. Le Bourgot C, Ferret-Bernard S, Le Normand L, Savary G, Menendez-Aparicio E, Blat S, et al. Maternal short-chain fructooligosaccharide supplementation influences intestinal immune system maturation in piglets. *PLoS One.* (2014) 9:e107508. doi: 10.1371/journal.pone.0107508
51. Lemaire M, Dou S, Cahu A, Formal M, Le Normand L, Romé V, et al. Addition of dairy lipids and probiotic *Lactobacillus fermentum* in infant formula programs gut microbiota and entero-insular axis in adult minipigs. *Sci Rep.* (2018) 8:11656. doi: 10.1038/s41598-018-29971-w
52. Arnaud AP, Rome V, Richard M, Formal M, David-Le Gall S, Boudry G. Post-natal co-development of the microbiota and gut barrier function follows different paths in the small and large intestine in piglets. *FASEB J Off Publ Fed Am Soc Exp Biol.* (2020) 34:1430–46. doi: 10.1096/fj.201902514R
53. Escudié F, Auer L, Bernard M, Mariadassou M, Cauquil L, Vidal K, et al. FROGS: Find, rapidly, OTUs with galaxy solution. *Bioinformatics.* (2018) 34:1287–94. doi: 10.1093/bioinformatics/btx791
54. R Core Team. *R: A Language And Environment For Statistical Computing.* Vienna: R Foundation for Statistical Computing (2020).
55. Fox J, Weisberg S. *An R companion to applied regression.* 3rd ed. Thousand Oaks, CA: SAGE Publications (2019).
56. Lê S, Josse J, Husson F. FactoMineR: A package for multivariate analysis. *J Stat Softw.* (2008) 25:1–18. doi: 10.18637/jss.v025.i01
57. Escofier B, Pagès J. Aspects théoriques et techniques de l'analyse factorielle multiple. *Analyses Factorielles Simples Et Multiples.* Paris: Dunod (1998). p. 171–204.
58. Pannaraj PS, Li F, Cerini C, Bender JM, Yang S, Rollie A, et al. Association between breast milk bacterial communities and establishment and development of the infant gut microbiome. *JAMA Pediatr.* (2017) 171:647–54. doi: 10.1001/jamapediatrics.2017.0378
59. Brink LR, Matalaz K, Piccolo BD, Bowlin AK, Chintapalli SV, Shankar K, et al. Neonatal diet impacts bioregional microbiota composition in piglets fed human breast milk or infant formula. *J Nutr.* (2019) 149:2236–46. doi: 10.1093/jn/nx z170

60. Rosa F, Matazel KS, Bowlin AK, Williams KD, Elolimy AA, Adams SH, et al. Neonatal diet impacts the large intestine luminal metabolome at weaning and post-weaning in piglets fed formula or human milk. *Front Immunol.* (2020) 11:3167. doi: 10.3389/fimmu.2020.607609
61. Yang I, Corwin EJ, Brennan PA, Jordan S, Murphy JR, Dunlop A. The infant microbiome: Implications for infant health and neurocognitive development. *Nurs Res.* (2016) 65:76–88. doi: 10.1097/NNR.0000000000000133
62. Neville MC, Anderson SM, McManaman JL, Badger TM, Bunik M, Contractor N, et al. Lactation and neonatal nutrition: Defining and refining the critical questions. *J Mammary Gland Biol Neoplasia.* (2012) 17:167–88. doi: 10.1007/s10911-012-9261-5
63. Marques TM, Wall R, Ross RP, Fitzgerald GF, Ryan CA, Stanton C. Programming infant gut microbiota: Influence of dietary and environmental factors. *Curr Opin Biotechnol.* (2010) 21:149–56. doi: 10.1016/j.copbio.2010.03.020
64. Schimmel P, Kleinjans L, Bongers RS, Knol J, Belzer C. Breast milk urea as a nitrogen source for urease positive *Bifidobacterium infantis*. *FEMS Microbiol Ecol.* (2021) 97:fiab019. doi: 10.1093/femsec/fiab019
65. Babakobi MD, Reshef L, Gihaz S, Belgorodsky B, Fishman A, Bujanover Y, et al. Effect of maternal diet and milk lipid composition on the infant gut and maternal milk microbiomes. *Nutrients.* (2020) 12:2539. doi: 10.3390/nu12092539
66. Bourlieu C, Bouzeur K, Ferret–Bernard S, Bourgot CL, Chever S, Ménard O, et al. Infant formula interface and fat source impact on neonatal digestion and gut microbiota. *Eur J Lipid Sci Technol.* (2015) 117:1500–12. doi: 10.1002/ejlt.201500025
67. Donovan SM, Comstock SS. Human milk oligosaccharides influence neonatal mucosal and systemic immunity. *Ann Nutr Metab.* (2016) 69:42–51. doi: 10.1159/000452818
68. German JB, Freeman SL, Lebrilla CB, Mills DA. Human milk oligosaccharides: Evolution, structures and bioselectivity as substrates for intestinal bacteria. *Nestle Nutr Workshop Ser Paediatr Programme.* (2008) 62:205–18. doi: 10.1159/000146322
69. Peila C, Moro GE, Bertino E, Cavallarini L, Giribaldi M, Giuliani F, et al. The effect of holder pasteurization on nutrients and biologically-active components in donor human milk: A review. *Nutrients.* (2016) 8:477. doi: 10.3390/nu8080477
70. Jeon M, Kim K-S, Kim ES, Lee S, Kim J, Hoe H-S, et al. Emerging pathogenic role of peripheral blood factors following BBB disruption in neurodegenerative disease. *Ageing Res Rev.* (2021) 68:101333. doi: 10.1016/j.arr.2021.101333
71. Laubereau B, Brockow I, Zirngibl A, Koletzko S, Gruebl A, von Berg A, et al. Effect of breast-feeding on the development of atopic dermatitis during the first 3 years of life—results from the GINI-birth cohort study. *J Pediatr.* (2004) 144:602–7. doi: 10.1016/j.jpeds.2003.12.029
72. Calder PC, Krauss-Etschmann S, de Jong EC, Dupont C, Frick J-S, Frokiaer H, et al. Early nutrition and immunity – progress and perspectives. *Br J Nutr.* (2006) 96:774–90. doi: 10.1079/BJN20061917
73. Rougé C, Butel M, Piloquet H, Ferraris L, Legrand A, Vodovar M, et al. Fecal calprotectin excretion in preterm infants during the neonatal period. *PLoS One.* (2010) 5:e11083. doi: 10.1371/journal.pone.0011083
74. Chatterton DEW, Rasmussen JT, Heegaard CW, Sørensen ES, Petersen TE. In vitro digestion of novel milk protein ingredients for use in infant formulas: Research on biological functions. *Trends Food Sci Technol.* (2004) 15:373–83. doi: 10.1016/j.tifs.2003.12.004
75. Bazzoni G. The JAM family of junctional adhesion molecules. *Curr Opin Cell Biol.* (2003) 15:525–30. doi: 10.1016/s0955-0674(03)00104-2
76. Le Huërou-Luron I, Blat S, Boudry G. Breast- v. formula-feeding: Impacts on the digestive tract and immediate and long-term health effects. *Nutr Res Rev.* (2010) 23:23–36. doi: 10.1017/S0954422410000065
77. Weaver LT. The impact of milk and weaning diet on gastrointestinal permeability in English and Gambian infants. *Trans R Soc Trop Med Hyg.* (1988) 82:784–9. doi: 10.1016/0035-9203(88)90236-2
78. Elolimy AA, Washam C, Byrum S, Chen C, Dawson H, Bowlin AK, et al. Formula diet alters the ileal metagenome and transcriptome at weaning and during the postweaning period in a porcine model. *mSystems.* (2020) 5:e457–420. doi: 10.1128/mSystems.00457-20
79. Le Bourgot C, le normand L, Formal M, Respondek F, Blat S, Apper E, et al. Maternal short-chain fructo-oligosaccharide supplementation increases intestinal cytokine secretion, goblet cell number, butyrate concentration and *Lawsonia intracellularis* humoral vaccine response in weaned pigs. *Br J Nutr.* (2017) 117:1–10. doi: 10.1017/S0007114516004268
80. Ranganath LR. The entero-insular axis: Implications for human metabolism. *Clin Chem Lab Med.* (2008) 46:43–56. doi: 10.1515/CCLM.2008.008
81. Lucas A, Boyes S, Bloom SR, Aynsley-Green A. Metabolic and endocrine responses to a milk feed in six-day-old term infants: Differences between breast and cow's milk formula feeding. *Acta Paediatr.* (1981) 70:195–200. doi: 10.1111/j.1651-2227.1981.tb05541.x
82. Lemaire M, Ménard O, Cahu A, Nogret I, Briard-Bion V, Cudennec B, et al. Addition of dairy lipids and probiotic *Lactobacillus fermentum* in infant formulas modulates proteolysis and lipolysis with moderate consequences on gut physiology and metabolism in yucatan piglets. *Front Nutr.* (2021) 8:20. doi: 10.3389/fnut.2021.615248
83. Guerrero-Hreins E, Goldstone AP, Brown RM, Sumithran P. The therapeutic potential of GLP-1 analogues for stress-related eating and role of GLP-1 in stress, emotion and mood: A review. *Prog Neuropsychopharmacol Biol Psychiatry.* (2021) 110:110303. doi: 10.1016/j.pnpbp.2021.110303
84. Wirleitner B, Neurauder G, Schrocksnadel K, Frick B, Fuchs D. Interferon gamma-induced conversion of tryptophan: Immunologic and neuropsychiatric aspects. *Curr Med Chem.* (2003) 10:1581–91. doi: 10.2174/0929867033457179
85. De Vadder F, Grasset E, Mannerås Holm L, Karsenty G, Macpherson AJ, Olofsson LE, et al. Gut microbiota regulates maturation of the adult enteric nervous system via enteric serotonin networks. *Proc Natl Acad Sci U.S.A.* (2018) 115:6458–63. doi: 10.1073/pnas.1720017115
86. Brink LR, Mercer KE, Piccolo BD, Chintapalli SV, Elolimy A, Bowlin AK, et al. Neonatal diet alters fecal microbiota and metabolome profiles at different ages in infants fed breast milk or formula. *Am J Clin Nutr.* (2020) 111:1190–202. doi: 10.1093/ajcn/nqaa076
87. Flint HJ, Duncan SH, Scott KP, Louis P. Links between diet, gut microbiota composition and gut metabolism. *Proc Nutr Soc.* (2015) 74:13–22. doi: 10.1017/S0029665114001463
88. Zhang J, Song L, Wang Y, Liu C, Zhang L, Zhu S, et al. Beneficial effect of butyrate-producing Lachnospiraceae on stress-induced visceral hypersensitivity in rats. *J Gastroenterol Hepatol.* (2019) 34:1368–76. doi: 10.1111/jgh.14536
89. Brestoff JR, Artis D. Commensal bacteria at the interface of host metabolism and the immune system. *Nat Immunol.* (2013) 14:676–84. doi: 10.1038/ni.2640
90. Whitt DD, Demoss RD. Effect of microflora on the free amino acid distribution in various regions of the mouse gastrointestinal tract. *Appl Microbiol.* (1975) 30:609–15. doi: 10.1128/am.30.4.609-615.1975
91. Bangsgaard Bendtsen KM, Krych L, Sørensen DB, Pang W, Nielsen DS, Josefsen K, et al. Gut microbiota composition is correlated to grid floor induced stress and behavior in the BALB/c mouse. *PLoS One.* (2012) 7:e46231. doi: 10.1371/journal.pone.0046231
92. Naseribafrouei A, Hestad K, Avershina E, Sekelja M, Linløkken A, Wilson R, et al. Correlation between the human fecal microbiota and depression. *Neurogastroenterol Motil.* (2014) 26:1155–62. doi: 10.1111/nmo.12378
93. Parker BJ, Wearsch PA, Veloo ACM, Rodriguez-Palacios A. The genus *Alistipes*: Gut bacteria with emerging implications to inflammation, cancer, and mental health. *Front Immunol.* (2020) 11:906. doi: 10.3389/fimmu.2020.00906
94. De Oliveira SC, Bourlieu C, Ménard O, Bellanger A, Henry G, Rousseau F, et al. Impact of pasteurization of human milk on preterm newborn in vitro digestion: Gastrointestinal disintegration, lipolysis and proteolysis. *Food Chem.* (2016) 211:171–9. doi: 10.1016/j.foodchem.2016.05.028
95. De Oliveira SC, Bellanger A, Ménard O, Plady P, Le Gouar Y, Henry G, et al. Impact of homogenization of pasteurized human milk on gastric digestion in the preterm infant: A randomized controlled trial. *Clin Nutr ESPEN.* (2017) 20:1–11. doi: 10.1016/j.clnesp.2017.05.001