



Bubbles Quantified *In vivo* by Ultrasound Relates to Amount of Gas Detected *Post-mortem* in Rabbits Decompressed from High Pressure

Yara Bernaldo de Quirós^{1*}, Andreas Møllerlækken², Marianne B. Havnes², Alf O. Brubakk², Oscar González-Díaz³ and Antonio Fernández¹

¹ Veterinary Histology and Pathology, Department of Morphology, Veterinary School, Institute of Animal Health, University of Las Palmas de Gran Canaria, Las Palmas, Spain, ² Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway, ³ Physical and Chemical Instrumental Center for the Development of Applied Research Technology and Scientific Estate, University of Las Palmas de Gran Canaria, Las Palmas, Spain

OPEN ACCESS

Edited by:

Jean-Pierre Montani,
University of Fribourg, Switzerland

Reviewed by:

Jacek Kot,
Medical University of Gdansk, Poland
Neal William Pollock,
Divers Alert Network, USA
Costantino Balestra,
Haute Ecole Paul Henri Spaak,
Belgium

*Correspondence:

Yara Bernaldo de Quirós
yberaldo@becarios.ulpgc.es

Specialty section:

This article was submitted to
Integrative Physiology,
a section of the journal
Frontiers in Physiology

Received: 29 April 2016

Accepted: 07 July 2016

Published: 21 July 2016

Citation:

Bernaldo de Quirós Y, Møllerlækken A, Havnes MB, Brubakk AO, González-Díaz O and Fernández A (2016) Bubbles Quantified *In vivo* by Ultrasound Relates to Amount of Gas Detected *Post-mortem* in Rabbits Decompressed from High Pressure. *Front. Physiol.* 7:310. doi: 10.3389/fphys.2016.00310

The pathophysiological mechanism of decompression sickness is not fully understood but there is evidence that it can be caused by intravascular and autochthonous bubbles. Doppler ultrasound at a given circulatory location is used to detect and quantify the presence of intravascular gas bubbles as an indicator of decompression stress. In this manuscript we studied the relationship between presence and quantity of gas bubbles by echosonography of the pulmonary artery of anesthetized, air-breathing New Zealand White rabbits that were compressed and decompressed. Mortality rate, presence, quantity, and distribution of gas bubbles elsewhere in the body was examined postmortem. We found a strong positive relationship between high ultrasound bubble grades in the pulmonary artery, sudden death, and high amount of intra and extra vascular gas bubbles widespread throughout the entire organism. In contrast, animals with lower bubble grades survived for 1 h after decompression until sacrificed, and showed no gas bubbles during dissection.

Keywords: decompression sickness, blockage of circulation, bubble grade, gas bubbles, gas emboli

INTRODUCTION

Decompression illness is a term used to describe diseases caused by intravascular or extravascular bubbles that are formed as a result of reduction in environmental pressure (decompression). The term covers both arterial gas embolism and decompression sickness (DCS; Vann et al., 2011). Bubbles can have mechanical, embolic, and biochemical effects with manifestations ranging from trivial to fatal (Vann et al., 2011).

Bert noted that the presence of gas bubbles in the blood was not a necessary cause of death or even visible symptoms (Bert, 1878). However, he suggested that gas in the vascular system or in the tissues must cause pain and decompression symptoms. He further suggested that the risks of paralysis or death depends upon the size of the gas bubble and hence, the capability to block the circulation.

The presence and amount of intravascular circulating bubbles can be determined by ultrasound technology. This technology is used in scuba divers as a non-invasive system for detection of

intravascular bubbles after diving schedules, as an indicator of decompression stress (Doolette and Mitchell, 2001; Pollock, 2007; Germonpre et al., 2014; Møllerlækken et al., 2016). The absence of detectable bubbles has been found to highly correlate with the absence of DCS symptoms (Sawatzky, 1991), and large quantities of intravascular bubbles predispose for the development of DCS (Evans et al., 1972; Neuman et al., 1976; Spencer, 1976; Gardette, 1979; Sawatzky, 1991). However, the relationship between intravascular bubbles detected by ultrasound and DCS is largely probabilistic, with no absolute threshold for number or size of bubbles below which there is no risk (Weathersby et al., 1984). DCS has occurred in divers with few bubbles detected by Doppler (Bayne et al., 1985). In contrast, no symptoms of DCS were recorded in individuals with large quantities of intravascular bubbles (Nishi et al., 2003). A recent study has shown that venous gas emboli detected by two-dimensional echocardiography are an imperfect surrogate endpoint for DCS (Doolette, 2016). There are no studies on detectable bubbles and lethal decompression in human controlled studies for obvious ethical reasons; hence the relationship between detectable bubbles by ultrasound and fatal decompression remains unknown.

The relationship between intravascular gas bubbles in a given localization determined by ultrasound and the amount of gas bubbles and the distribution of these intravascularly or extravascularly elsewhere in the organism is unknown. This relationship might contribute to understand better the pathophysiology of DCS. In the present study, we compared the presence and amount of intravascular circulatory gas bubbles at the pulmonary artery of New Zealand White (NZW) rabbits after decompression using ultrasound, to the presence and amount of gas bubbles in different vascular and extravascular body locations found grossly *post mortem* (PM) during necropsy.

MATERIALS AND METHODS

Animals

Male NZW rabbits (4 from Animal Supply Center of the Negrin Hospital, Spain, and 14 from the Unit of Comparative Medicine, St Olav University Hospital NTNU, Norway) of 2.5–3.8 kg weigh were used in this experiment. All experiments were conducted in accordance with the European Union regulations for laboratory animals. Experimental protocols for the control and putrefaction studies were performed in Spain and approved by the Ethical Committee for Animal Experiments of the University of Las Palmas de Gran Canaria (Spain). The Norwegian Committee for Animal Experiments approved the protocol for the compression/decompression study where it was carried out.

Experimental Protocol

Animals were assigned into the following two experimental groups (a) control or gas putrefaction studies ($n = 4$), and (b) compression/decompression treatment ($n = 14$). All experiments were conducted under surgical anesthesia (Medetomidine ($0.5 \text{ mg}\cdot\text{kg}^{-1}$) and Ketamine ($25 \text{ mg}\cdot\text{kg}^{-1}$) subcutaneously).

(a) Control Group

After anesthesia animals were euthanized with an intraperitoneal injection of pentobarbital ($200 \text{ mg}\cdot\text{kg}^{-1}$). Dead animals were kept in hermetically sealed plastic boxes for biological material at room temperature ($24.0 \pm 0.8^\circ\text{C}$) for 1, 3, 6, and 12 h PM ($n = 1$ for each time) before necropsies were performed.

(b) Compression/Decompression Model

Anesthetized NZW rabbits were compressed in pairs, in a dry hyperbaric chamber (Animal Chamber System, NUT, Haugesund, Norway) to 8 atmospheres absolute (ATA) with 45 min bottom time followed by a fast decompression (0.33 m/s). During the dive, animals breathed compressed air (Medical air, Yara Praxair, Oslo, Norway). The dive profile was planned to be aggressive. Previous diving experiments in rabbits showed that a dive to 6ATA during 45 min showed a low mortality rate (Shim et al., 1967; Tanoue et al., 1987; Su et al., 2004). After decompression, the pulmonary artery and the aorta of the rabbits were monitored by ultrasound for *in vivo* bubble detection by a 10 MHz transducer connected to an ultrasound scanner (GE Vivid Five, Vingmed Ultrasound AS, Norway). Bubbles were detected as bright spots. The abundance of gas bubbles was evaluated using the Eftedal and Brubakk (EB) grading scale from 0 to 5 (Table 1; Eftedal and Brubakk, 1997). Ultrasound monitoring was repeated every 15 min for 1 h following decompression, or until death of the animal. Animals that survived for 1 h after decompression were euthanized with an intraperitoneal injection of pentobarbital ($200 \text{ mg}\cdot\text{kg}^{-1}$). The bodies were placed in hermetically sealed plastic boxes for biological material at room temperature ($23.3 \pm 1.3^\circ\text{C}$) for 0, 20, and 40 min, and 1, 3, 6, and 12 h ($n = 2$ for each time) PM.

Post-mortem Procedures

Animals were carefully dissected and the presence of gas was evaluated using a standardized gas score index (Bernaldo de Quirós et al., 2016). For this purpose a gas score from 0 to VI was given to different vascular locations (subcutaneous, mesenteric, femoral, and coronary veins as well as the right atrium) and a gas score from 0 to III was used to evaluate the presence of gas beneath the capsule of different tissues (subcapsular emphysema), and within adipose tissues (interstitial emphysema; Tables 2, 3). These scores were used to obtain a new gas score index to represent the global or total gas score for each animal. This new index was obtained by the summation of the gas scores

TABLE 1 | Grading code for ultrasonic images following Eftedal and Brubakk (1997).

Grade	Definition
0	No observable bubbles
1	Occasional bubble
2	At least 1 bubble every 4 heart cycles
3	At least 1 bubbles every heart cycle
4	At least 1 bubble per cm^2 in every image
5	"White-out": single bubbles cannot be discriminated

for each tissue, thus total gas score in each animal ranges from 0 to 42 (Table 3). The production of putrefaction gases with PM time was studied in a previous manuscript of this same experiment including animals with longer PM hours (Bernaldo de Quirós et al., 2013, 2016). Results from gas composition analyses and gas score analyses indicated that putrefaction gases were not significant in the animals with 27 h PM or less. Putrefaction gases were not detected in animals with 12 h PM or less. For this study we have included only those animals that were considered fresh and free of putrefaction gases following gas composition and gas scoring analyses (Bernaldo de Quirós et al., 2013, 2016).

Data Analysis

SPSS Statistics 17.0 software was used for statistical analysis. The correlation between EB bubble grade and PM gas scoring was analyzed by Spearman's rank correlation coefficient. PM gas scoring data was partitioned in two groups using K-means clustering. The statistical difference between these two groups was assessed by Mann-Whitney *U*-test. Statistical significance was set at $p < 0.05$.

RESULTS

Post-diving Observations

A large individual variability in EB bubble grade was found: 2/14 NZW rabbits (14.3%) presented with an EB bubble grade of 0; 2/14 NZW rabbits (14.3%) presented with an EB bubble grade of 1; 1/14 NZW rabbits (7.1%) presented with an EB bubble grade of 2; 1/14 NZW rabbits (7.1%) presented with an EB bubble grade of

3; 4/14 NZW rabbits (28.6%) presented with an EB bubble grade of 4; and 4/14 NZW rabbits (28.6%) presented with an EB bubble grade of 5. About half of the animals (42.9%) presented with an EB bubble grade of 0–3 while the other 57.1% of the animals presented with an EB bubble grade of 4 or 5. Animals with bubble grade 0–3 survived for 1 h and were euthanized according to the experimental protocol. In contrast, all the animals with bubble grade 4 or 5 showed severe respiratory distress signs and died within 5 to 35 min after decompression.

PM Examinations

Control Group

Grossly, there was an absence of intravascular and extravascular gas bubbles. Occasionally single or very few scattered gas bubbles were observed. Minimum and maximum gas score values for this group were 0 and 4, respectively. Mode gas score value was 0.

Diving Group: EB Bubble Grade 0–3

All these animals ($n = 6$) survived for 1 h after decompression and were euthanized following the protocol. There was an absence of intravascular and extravascular gas bubbles. Gas score was 0 for all these animals.

Diving Group: EB Bubble Grades 4 and 5

These animals ($n = 8$) died within 5–35 min after decompression. Gas was abundant and disseminated in all the venous circulatory system and in the tissues, including peripheral veins and the abdominal adipose tissue. There were veins completely filled with gas showing evident vascular obstruction. In many cases, lesions such as hemorrhages were observed in association with the bubbles (Figure 1). Minimum and maximum gas score values for this group were 29 and 39, respectively. Mode gas score value was 34.

Relationship between EB Bubble Grade and PM Gas Score

The K-means cluster test differentiated two clusters attending to PM gas scoring. Cluster 1 was composed by animals ($n = 6$) with gas score of 0, and center at 0 gas score. Cluster 2 was composed by animals ($n = 8$) with gas score of 29 or higher, and center at 33 gas score. Distances between final cluster centers was 33.3. Cluster 1 and 2 were statistically significant different ($P < 0.001$; Figure 2).

Cluster data was crossed with EB bubble grade. Animals from cluster 1 presented with EB bubble grade of 3 or lower. In contrast, animals from cluster 2 all presented with EB bubble grade of 4 or higher (Figure 2). EB bubble grade and PM gas scoring were found to be strongly correlated ($\rho = 0.883$; $P \leq 0.0001$).

TABLE 2 | Definition of gas score index for post-mortem examinations following Bernaldo de Quirós et al. (2016).

Gas score	Definition
INTRAVASCULAR	
0	Absence of bubbles
I	Occasional bubble
II	Few bubbles and/or discontinuities of blood
III	Few bubbles and large discontinuities of blood
IV	Moderate presence of bubbles
V	Abundant presence of bubbles
VI	Complete sections of vessels filled with gas
EXTRA-VASCULAR	
0	Absence of gas
I	Scarce presence: affecting only 1 organ
II	Moderate presence of gas: in more than 1 organ
III	Abundant presence of gas: systemic

TABLE 3 | Calculation of total gas score for each animal following Bernaldo de Quirós et al. (2016).

Identification	Subcutaneous v.	Mesenteric v.	Femoral v.	Vena cava	Right atrium	Coronary v.	Interstitial emphysema in fatty tissue	Subcapsular emphysema	Total
Animal n	0–VI	0–VI	0–VI	0–VI	0–VI	0–VI	0–III	0–III	0–42





FIGURE 1 | Pictures from PM examinations of the rabbits with bubble grades 4 and 5 showing gas in the abdominal adipose tissue (A) and extensive gas embolism including right atrium and coronary veins (B), mesenteric veins (C), gastric veins and filling of the spleen (D), subcutaneous veins (E), and urinary bladder veins (F) among other veins.



DISCUSSION

To the authors knowledge this is the first study describing the relationship between bubble grades detected by ultrasound in the pulmonary artery *in vivo*, and the presence of macroscopic bubbles elsewhere in the organism.

The present study showed a high inter-individual variability in bubble grades detected by ultrasound, despite having similar subjects (species, gender, weight) exposed to the same treatment. This result supports previous work by Eckenhoff et al. (1990) who reported a considerable inter and intra-individual variability in bubble formation at any level of exposure (Eckenhoff et al., 1990).

The main finding from this study was that EB bubble grade measured *in vivo* by ultrasound correlated to mortality and to gas score determined PM: animals with EB bubble grades 1–3 survived for 1 h after decompression until sacrificed according to protocol and presented with low gas score when examined PM, while animals with EB bubble grades 4–5 died within 5–35 min after decompression and presented with high gas score of gas bubbles widely dispersed in different veins, including peripheral veins, and in the adipose abdominal tissue when examined PM.

The mortality result was unexpected since large quantities of intravascular bubbles have been reported in asymptomatic apparently healthy divers (Nishi et al., 2003). PM examination showed only two treatment responses: large amount of gas bubbles within veins and tissues in animals dying shortly after decompression vs. absence or very few gas bubbles in animals which survived and were later sacrificed. Similar observations on survival and amount of bubbles detected PM have been previously reported in sparrows, mice, rats, pigs, cats, dogs, and rabbits (including NZW), that have been rapidly decompressed from high pressures (Bert, 1878; Eggleton et al., 1945; Lever et al., 1966; Shim et al., 1967). Most of these studies support our

findings regardless species and exposure pressure levels. Bert's results were more varied but they also showed that animals dying rapidly after sudden decompression always presented with large amounts of gas bubbles in the venous side of the circulatory system (Bert, 1878).

It is interesting to note that regardless the different sensitivity to decompression (demonstrated by higher prevalence of DCS symptoms and/or morbidity) of different species (Bert, 1878; Eggleton et al., 1945), necropsy findings were similar to our results showing a dual response in dead vs. surviving animals (Eggleton et al., 1945; Shim et al., 1967): all animals dying shortly after decompression showed large amounts of gas bubbles in the veins regardless of animal species, or hyperbaric exposure.

Our animals that died shortly after decompression (EB bubble grades 4 and 5) also presented with gas bubbles within the abdominal adipose tissue. This finding has been described before and used as a symptom to diagnose DCS in rats (Hyldegaard and Madsen, 1989).

The dual response (presence and high amounts of gas bubbles vs. absence of gas bubbles) of the gas score was very different from the high individual variability in EB bubbles grades measured by ultrasound. Differences found between EB bubble grade measured by ultrasound and PM gas score examination could be due to different resolution limits of the two methods. Monitoring with high-resolution ultrasound renders the possibility of detecting smaller bubbles than is possible by visual examination. Additionally, small bubbles are more susceptible to be trapped and excreted in the pulmonary capillaries (Francis and Simon, 2003) or simply get diluted. According to LaPlace equation, smaller bubbles have larger inner pressure, thus dissolve more quickly than large bubbles (Hrncir, 1996). Large bubbles are more stable and easier to see macroscopically. Additional limitation of this method is the low number of cases studied. More studies using fresh animals should be carried out to reassure the results presented in this manuscript.

Future pathological studies comparing animals with EB bubble grades 0–3 and EB bubble grades 4–5 might help us to understand better the unexpected mortality rate described in this study, as well as the pathophysiological mechanism of DCS.

This study has focused in lethal decompression, which is rare among human divers except for the most severe accidents, thus the application of these results in human clinical medicine might be limited. On the other hand, studying diseases in their most extreme expression might be helpful to detect better pathophysiological mechanisms of the disease that might not be detected otherwise. The results from the present experiment should be considered carefully due to the low number of individuals studied, but the results suggest a strong relationship between fatal decompression and high loads of gas emboli in NZW rabbits. This relationship remains unclear in less severe decompression cases in human divers (Evans et al., 1972; Neuman et al., 1976; Spencer, 1976; Gardette, 1979; Bayne et al., 1985; Sawatzky, 1991; Nishi et al., 2003).

Another limitation to our data is that there are to our knowledge no studies available today that describes where decompression induced vascular bubbles do originate from,

and whether there are specific tissues or organs that are more “bubble-producers” than others. Recent studies are attempting to detect small stationary bubbles in tissues using Dual-frequency ultrasound (Swan et al., 2011, 2014). In this regard, the total gas score estimated for each animal might underweight the pathophysiological importance of high bubble loads in a single tissue. The gas score for each tissue might be more informative for that respect.

In summary, in the present study we have reported a dual response of presence of gas bubbles in postmortem examinations (high amount of gas bubbles vs. absence of gas bubbles) in animals of the same species, gender, and similar weight, exposed to the same diving profile. *In vivo* ultrasound measurements at the pulmonary artery related to quantity of gas widely distributed in the venous system in decompression cases. High bubble grades were related to high gas score causing death within 35 min after decompression in NZW rabbits.

AUTHOR CONTRIBUTIONS

AM and MH contributed to the experiments, interpretation of the results and provide their expertise. AB, OG, and AF,

supervised the work, contributed to the discussion of the results and provide their respective expertise. YB carried out the experiments, data analyses and interpreted the results. All authors listed contributed to the writing of the manuscript.

ACKNOWLEDGMENTS

The authors would like to thank all colleagues from the University of Las Palmas de Gran Canaria (Spain) that contribute to this work and to the Barophysiology Group at the Norwegian University of Science and Technology (Norway) for its scientific contribution.

This work was supported by the Spanish Ministry of Science and Innovation with two research projects: (AGL 2005-07947) and (CGL 2009/12663), as well as the Government of Canary Islands (DG Medio Natural). The Spanish Ministry of Education contributed with personal financial support (the University Professor Formation fellowship). The work was also supported by the Liaison Committee between the Central Norway Regional Health Authority (RHA) and the Norwegian University of Science and Technology, NTNU, grant number 460 28600.

REFERENCES

- Bayne, C. G., Hunt, W. S., Johanson, D. C., Flynn, E. T., and Weathersby, P. K. (1985). Doppler bubble detection and decompression sickness - a prospective clinical trial. *Undersea Biomed. Res.* 12, 327–332.
- Bernaldo de Quirós, Y., González-Díaz, O., Møllerløkken, A., Brubakk, A. O., Hjelde, A., Saavedra, P., et al. (2013). Differentiation at autopsy between *in vivo* gas embolism and putrefaction using gas composition analysis. *Int. J. Legal Med.* 127, 437–445. doi: 10.1007/s00414-012-0783-6
- Bernaldo de Quirós, Y., Saavedra, P., Møllerløkken, A., Brubakk, A. O., Jørgensen, A., González-Díaz, O., et al. (2016). Differentiation at necropsy between *in vivo* gas embolism and putrefaction using a gas score. *Res. Vet. Sci.* 106, 48–55. doi: 10.1016/j.rvsc.2016.03.007
- Bert, P. (1878). *La Pression Barometrique: Recherches de Physiologie Expérimentale*. Paris: Masson.
- Doolette, D. J. (2016). Venous gas emboli detected by two-dimensional echocardiography are an imperfect surrogate endpoint for decompression sickness. *Diving Hyperb. Med.* 46, 4–10.
- Doolette, D. J., and Mitchell, S. J. (2001). The physiological kinetics of nitrogen and the prevention of decompression sickness. *Clin. Pharmacokinet.* 40, 1–14. doi: 10.2165/00003088-200140010-00001
- Eckenhoff, R. G., Olstad, C. S., and Carrod, G. (1990). Human dose-response relationship for decompression and endogenous bubble formation. *J. Appl. Physiol.* 69, 914–918.
- Eftedal, O., and Brubakk, A. O. (1997). Agreement between trained and untrained observers in grading intravascular bubble signals in ultrasonic images. *Undersea Hyperb. Med.* 24, 293–299.
- Eggleton, P., Elsdon, S. R., Fegle, J., and Hebb, C. O. (1945). A study of the effects of rapid “decompression” in certain animals. *J. Physiol.* 104, 129–150. doi: 10.1113/jphysiol.1945.sp004111
- Evans, A., Barnard, E. E. P., and Walder, D. N. (1972). Detection of gas bubbles in man at decompression. *Aerosp. Med.* 43, 1095–1096.
- Francis, T. J. R., and Simon, J. M. (2003). “Pathology of decompression sickness,” in *Bennett and Elliott’s Physiology and Medicing of Diving*, eds A. O. Brubakk and T. S. Neuman (London: Saunders), 530–556.
- Gardette, B. (1979). Correlation between decompression-sickness and circulating bubbles in 232 divers. *Undersea Biomed. Res.* 6, 99–107.
- Germonpré, P., Papadopoulou, V., Hemelryck, W., Obeid, G., Lafère, P., Eckersley, R. J., et al. (2014). The use of portable 2D echocardiography and “frame-based” bubble counting as a tool to evaluate diving decompression stress. *Diving Hyperb. Med.* 44, 5–13.
- Hrnčir, E. (1996). Importance of surface tension in therapeutic compression in decompression sickness. *Physiol. Res.* 45, 467–470.
- Hyldegaard, O., and Madsen, J. (1989). Influence of heliox, oxygen and N2O-O2 breathing on N2 bubbles in adipose-tissue. *Undersea Biomed. Res.* 16, 185–193.
- Lever, M. J., Miller, K. W., Paton, W. D. M., and Smith, E. B. (1966). Experiments on the genesis of bubbles as a result of rapid decompression. *J. Physiol.* 184, 964–969. doi: 10.1113/jphysiol.1966.sp007960
- Møllerløkken, A., Blogg, S. L., Doolette, D. J., Nishi, R. Y., and Pollock, N. W. (2016). Consensus guidelines for the use of ultrasound for diving research. *Diving Hyperb. Med.* 46, 26–32.
- Neuman, T. S., Hall, D. A., and Linaweaver, P. G. (1976). Gas phase separation during decompression in man: ultrasound monitoring. *Undersea Biomed. Res.* 3, 121–130.
- Nishi, R. Y., Brubakk, A. O., and Eftedal, O. (2003). “Bubble detection,” in *Bennett and Elliott’s Physiology and Medicine of Diving*, eds A. O. Brubakk and T. S. Neuman (London: Saunders), 501–529.
- Pollock, N. W. (2007). Use of ultrasound in decompression research. *Diving Hyperb. Med.* 37, 68–72.
- Sawatzky, K. D. (1991). *The Relationship between Intravascular Doppler-Detected Gas Bubbles and Decompression Sickness after Bounce Diving in Humans*. Toronto, ON: York University.
- Shim, S. S., Patterson, F. P., and Kendall, M. J. (1967). Hyperbaric chamber and decompression sickness: an experimental study. *Can. Med. Assoc. J.* 97, 1263–1272.
- Spencer, M. P. (1976). Decompression limits for compressed air determined by ultrasonically detected blood bubbles. *J. Appl. Physiol.* 40, 229–235.
- Su, C. L., Wu, C. P., Chen, S. Y., Kang, B. H., Huang, K. L., and Lin, Y. C. (2004). Acclimatization to neurological decompression sickness in rabbits. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287, R1214–R1218. doi: 10.1152/ajpregu.00260.2004
- Swan, J. G., Bollinger, B. D., Donoghue, T. G., Wilbur, J. C., Phillips, S. D., Alvarenga, D. L., et al. (2011). Microbubble detection following hyperbaric chamber dives using dual-frequency ultrasound. *J. Appl. Physiol.* 111, 1323–1328. doi: 10.1152/jappphysiol.01203.2010

- Swan, J. G., Wilbur, J. C., Moodie, K. L., Kane, S. A., Knaus, D. A., Phillips, S. D., et al. (2014). Microbubbles are detected prior to larger bubbles following decompression. *J. Appl. Physiol.* 116, 790–796. doi: 10.1152/jappphysiol.01156.2013
- Tanoue, K., Mano, Y., Kuroiwa, K., Suzuki, H., Shibayama, M., and Yamazaki, H. (1987). Consumption of platelets in decompression-sickness of rabbits. *J. Appl. Physiol.* 62, 1772–1779.
- Vann, R. D., Butler, F. K., Mitchell, S. J., and Moon, R. E. (2011). Decompression illness. *Lancet*, 377, 153–164. doi: 10.1016/S0140-6736(10)61085-9
- Weathersby, P. K., Homer, L. D., and Flynn, E. T. (1984). On the likelihood of decompression sickness. *J. Appl. Physiol.* 57, 815–825.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Bernaldo de Quirós, Møllerløkken, Havnes, Brubakk, González-Díaz and Fernández. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.