



Why Is there a Limit to the Changes in Myofilament Ca^{2+} -Sensitivity Associated with Myopathy Causing Mutations?

Steven B. Marston *

National Heart & Lung Institute, Imperial College London, London, UK

OPEN ACCESS

Edited by:

P. Bryant Chase,
Florida State University, USA

Reviewed by:

Murali Chandra,
Washington State University, USA
Michelle Parvatiyar,
University of California, Los Angeles,
USA

Nicolas Brunet,
SUNY Downstate Medical Center
Brooklyn, USA

*Correspondence:

Steven B. Marston
s.marston@imperial.ac.uk

Specialty section:

This article was submitted to
Striated Muscle Physiology,
a section of the journal
Frontiers in Physiology

Received: 06 July 2016

Accepted: 05 September 2016

Published: 26 September 2016

Citation:

Marston SB (2016) Why Is there a
Limit to the Changes in Myofilament
 Ca^{2+} -Sensitivity Associated with
Myopathy Causing Mutations?
Front. Physiol. 7:415.
doi: 10.3389/fphys.2016.00415

Mutations in striated muscle contractile proteins have been found to be the cause of a number of inherited muscle diseases; in most cases the mechanism proposed for causing the disease is derangement of the thin filament-based Ca^{2+} -regulatory system of the muscle. When considering the results of experiments reported over the last 15 years, one feature has been frequently noted, but rarely discussed: the magnitude of changes in myofilament Ca^{2+} -sensitivity due to myopathy-causing mutations in skeletal or heart muscle seems to be always in the range 1.5–3x EC_{50} . Such consistency suggests it may be related to a fundamental property of muscle regulation; in this article we will investigate whether this observation is true and consider why this should be so. A literature search found 71 independent measurements of HCM mutation-induced change of EC_{50} ranging from 1.15 to 3.8-fold with a mean of 1.87 ± 0.07 (sem). We also found 11 independent measurements of increased Ca^{2+} -sensitivity due to mutations in skeletal muscle proteins ranging from 1.19 to 2.7-fold with a mean of 2.00 ± 0.16 . Investigation of dilated cardiomyopathy-related mutations found 42 independent determinations with a range of EC_{50} wt/mutant from 0.3 to 2.3. In addition we found 14 measurements of Ca^{2+} -sensitivity changes due skeletal muscle myopathy mutations ranging from 0.39 to 0.63. Thus, our extensive literature search, although not necessarily complete, found that, indeed, the changes in myofilament Ca^{2+} -sensitivity due to disease-causing mutations have a bimodal distribution and that the overall changes in Ca^{2+} -sensitivity are quite small and do not extend beyond a three-fold increase or decrease in Ca^{2+} -sensitivity. We discuss two mechanism that are not necessarily mutually exclusive. Firstly, it could be that the limit is set by the capabilities of the excitation-contraction machinery that supplies activating Ca^{2+} and that striated muscle cannot work in a way compatible with life outside these limits; or it may be due to a fundamental property of the troponin system and the permitted conformational transitions compatible with efficient regulation.

Keywords: muscle regulation, Ca^{2+} -sensitivity, troponin C, HCM, DCM, myopathy, mutation

Abbreviations: HCM, hypertrophic cardiomyopathy; RCM, Restrictive cardiomyopathy; DCM, dilated cardiomyopathy; EC_{50} , Ca^{2+} concentration that gives 50% maximal activation; pCa_{50} , $-\log \text{EC}_{50}$.

Mutations in striated muscle contractile proteins have been found to be the cause of a number of inherited muscle diseases; in most cases the mechanism proposed for causing the disease is derangement of the thin filament-based Ca²⁺-regulatory system of the muscle. Hypertrophic cardiomyopathy and hypercontractile diseases of skeletal muscle, such as distal arthrogryposis and “stiff child syndrome,” have been linked to a higher myofilament Ca²⁺-sensitivity (Marston, 2011; Donkervoort et al., 2015). In contrast dilated cardiomyopathy mutations are commonly, but not exclusively, linked to decreased Ca²⁺-sensitivity. Mutations in contractile proteins that are linked to nemaline myopathy and related skeletal muscle myopathies have also been found to be associated with reduced Ca²⁺ sensitivity (Marttila et al., 2012, 2014). The causative connection between myofilament Ca²⁺-sensitivity and muscle dysfunction is a field of intensive research that is too complex to consider in this account. However, when considering the results of such experiments reported over the last 15 years, one feature has been frequently noted, but rarely discussed. The magnitude of changes in myofilament Ca²⁺-sensitivity due to myopathy-causing mutations in skeletal or heart muscle seems to be always in the range 1.5–3x EC₅₀. Such consistency suggests it may be related to a fundamental property of muscle regulation; in this article we will investigate whether this observation is true and consider why this should be so.

Most investigations have found increased Ca²⁺-sensitivity in muscle with hypertrophic cardiomyopathy (HCM) and restrictive cardiomyopathy (RCM)-causing mutations. Our literature search found 71 independent measurements of the mutation-induced change of EC₅₀ ranging from 1.15 to 3.8-fold with a mean of 1.87 ± 0.07 (sem) (Table 1). We also found 11 independent measurements of increased Ca²⁺-sensitivity due to mutations in skeletal muscle proteins ranging from 1.19 to 2.7-fold with a mean of 2.00 ± 0.16 (Table 2).

Dilated cardiomyopathy-causing mutations were initially found to decrease Ca²⁺-sensitivity but more recent studies have indicated the situation is more complex. DCM-linked mutations can both increase and decrease Ca²⁺-sensitivity depending on the individual mutations, moreover the direction of change can be different with a single mutation measured in different systems (Marston, 2011; Memo et al., 2013). This is illustrated in Table 3 where 42 independent determinations show a range of EC₅₀ wt/mutant from 0.3 to 2.3. In addition we found 14 measurements of Ca²⁺-sensitivity changes due skeletal muscle myopathy mutations ranging from 0.39 to 0.63 (Table 4).

Thus, our extensive literature search, although not necessarily complete, found that, indeed, the changes in myofilament Ca²⁺-sensitivity due to disease-causing mutations have a bimodal distribution and that the overall changes in Ca²⁺-sensitivity are quite small and do not extend beyond a 3–4-fold increase or decrease in Ca²⁺-sensitivity. Indeed when all the findings are plotted as a histogram one finds that increases in Ca²⁺-sensitivity on a log scale have an approximately normal distribution with mean increase in Ca²⁺-sensitivity (EC₅₀ wt/mutant) of 1.86-fold (corresponding to ΔpCa₅₀ = 0.255 ± 0.015), whilst the decreases in Ca²⁺ sensitivity have a mean EC₅₀ wt/mutant of 0.54-fold (corresponding to ΔpCa₅₀ of -0.286 ± 0.01; Figure 1A). It

TABLE 1 | Effect of HCM-associated mutations on myofilament Ca²⁺-sensitivity.

| Gene name | Mutation | wt/mutant EC ₅₀ ratio | Measured in | References |
|------------|-----------|----------------------------------|-----------------|-------------------------------|
| HCM | | | | |
| ACTC | E99K | 2.45 | IVMA | Song et al., 2011 |
| ACTC | E99K | 1.24 | IVMA (human) | Song et al., 2011 |
| ACTC | E99K | 1.89 | IVMA | Papadaki et al., 2015 |
| ACTC | E99K | 1.3 | Fibers TG | Song et al., 2011 |
| ACTC | E99K | 2.35 | Myofibrils TG | Song et al., 2013 |
| MYL2 | R58Q | 1.29 | Fibers X | Szczesna-Cordary et al., 2004 |
| MYL2 | D166V | 1.78 | Fibers TG | Kerrick et al., 2009 |
| MYL2 | D166V | 1.82 | Fibers TG | Yuan et al., 2015 |
| MYH7 | R403Q | 1.79 | Human fibers | Sequeira et al., 2013 |
| MYH7 | R403Q | 1.41 | Fibers TG | Blanchard et al., 1999 |
| MYH7 | R453C | 1.99 | Human fibers | Palmer et al., 2004 |
| MYBPC3 | Cat R820W | 2.01 | IVMA | Messer et al., 2016a |
| MYBPC3 | “K1” | 1.35 | Fibers TG | Frayse et al., 2012 |
| MYBPC3 | E258K | 1.80 | Human fibers | Sequeira et al., 2013 |
| TNNC1 | A8V | 2.51 | Fibers TG | Martins et al., 2015 |
| TNNC1 | A8V | 2.3 | Fibers X | Pinto et al., 2009 |
| TNNC1 | L29Q | 1.26 | Fibers X 2.3 μm | Li et al., 2013 |
| TNNC1 | L29Q | 1.17 | Fibers X 1.9 μm | Li et al., 2013 |
| TNNC1 | L29Q | 2.1 | IVMA | Schmidtman et al., 2005 |
| TNNC1 | A31S | 1.48 | Fibers X | Parvatiyar et al., 2012 |
| TNNC1 | A31S | 2.75 | ATPase | Parvatiyar et al., 2012 |
| TNNC1 | D145E | 1.74 | Fibers X | Pinto et al., 2009 |
| TNNC1 | C84Y | 1.86 | Fibers X | Pinto et al., 2009 |
| TNNI3 | R21C | 2.16 | Fibers X | Gomes et al., 2005a |
| TNNI3 | L144Q | 2.04 | Fibers X | Gomes et al., 2005b |
| TNNI3 | R145G | 3.63 | ATPase | Elliott et al., 2000 |
| TNNI3 | R145G | 2.09 | ATPase | Takahashi-Yanaga et al., 2001 |
| TNNI3 | R145G | 1.82 | IVMA | Brunet et al., 2014 |
| TNNI3 | R145G | 1.41 | IVMA | Deng et al., 2001 |
| TNNI3 | R145G | 1.35 | Fibers X | Lang et al., 2002 |
| TNNI3 | R145G | 1.15 | Fibers TG | Krüger et al., 2005 |
| TNNI3 | R145Q | 1.41 | Fibers X | Takahashi-Yanaga et al., 2001 |
| TNNI3 | R145Q | 1.70 | ATPase | Takahashi-Yanaga et al., 2001 |
| TNNI3 | R145W | 2.45 | Fibers X | Gomes et al., 2005b |
| TNNI3 | R145W | 1.15 | Human fibers | Sequeira et al., 2013 |
| TNNI3 | R162W | 1.28 | ATPase | Takahashi-Yanaga et al., 2001 |
| TNNI3 | A171T | 1.38 | Fibers X | Gomes et al., 2005b |
| TNNI3 | K178E | 2.95 | Fibers X | Gomes et al., 2005b |
| TNNI3 | ΔK182 | 1.51 | ATPase | Takahashi-Yanaga et al., 2001 |
| TNNI3 | ΔK183 | 3.8 | IVMA | Köhler et al., 2003 |
| TNNI3 | R192H | 2.29 | Fibers X | Gomes et al., 2005b |
| TNNI3 | G203S | 3.02 | IVMA | Köhler et al., 2003 |

(Continued)

TABLE 1 | Continued

| Gene name | Mutation | wt/mutant EC ₅₀ ratio | Measured in | References |
|------------|----------|----------------------------------|------------------|-------------------------------|
| HCM | | | | |
| TNNI3 | K206Q | 2.51 | IVMA | Köhler et al., 2003 |
| TNNI3 | K206Q | 1.51 | ATPase | Takahashi-Yanaga et al., 2001 |
| TNNI3 | K206I | 1.81 | ATPase | Warren et al., 2015 |
| TNNT2 | TnTΔ14 | 2.51 | Fibers X | Gafurov et al., 2004 |
| TNNT2 | TnTdel | 2.69 | ATPase | Redwood et al., 2000 |
| TNNT2 | I79N | 1.41 | Fibers X | Szczesna et al., 2000 |
| TNNT2 | I79N | 2.04 | Fibers TG | Baudenbacher et al., 2008 |
| TNNT2 | R92L | 1.65 | Fibers TG | Ford et al., 2012 |
| TNNT2 | R92Q | 1.66 | Fibers TG | Ford et al., 2012 |
| TNNT2 | R92Q | 1.74 | ATPase | Robinson et al., 2002 |
| TNNT2 | R92Q | 1.94 | IVMA | Robinson et al., 2002 |
| TNNT2 | F110I | 2.34 | Fibers TG | Szczesna et al., 2000 |
| TNNT2 | F110I | 1.32 | Fibers TG | Baudenbacher et al., 2008 |
| TNNT2 | ΔE160 | 1.41 | Fibers TG | Lu et al., 2003 |
| TNNT2 | R278C | 2.19 | Fibers TG | Szczesna et al., 2000 |
| TNNT2 | K280N | 1.64 | IVMA | Messer et al., 2016b |
| TNNT2 | K280N | 1.26 | IVMA (human Tn) | Messer et al., 2016b |
| TPM1 | E62Q | 1.21 | ATPase | Chang et al., 2005 |
| TPM1 | A63V | 1.91 | Transfected cell | Michele et al., 1999 |
| TPM1 | A63V | 1.99 | ATPase | Heller et al., 2003 |
| TPM1 | K70T | 1.58 | Transfected cell | Michele et al., 1999 |
| TPM1 | K70T | 2.13 | ATPase | Heller et al., 2003 |
| TPM1 | D175N | 1.23 | IVMA | Bing et al., 2000 |
| TPM1 | E180G | 1.30 | IVMA | Bing et al., 2000 |
| TPM1 | E180G | 1.63 | IVMA | Papadaki et al., 2015 |
| TPM1 | E180G | 1.44 | Transfected cell | Michele et al., 1999 |
| TPM1 | E180G | 2.75 | ATPase | Chang et al., 2005 |
| TPM1 | L185R | 2.51 | ATPase | Chang et al., 2005 |
| TPM1 | I284V | 1.50 | Human fibers | Sequeira et al., 2013 |

The criteria for inclusion in the table are (1) that a missense mutation has been convincingly linked to the myopathy phenotype and (2) that only direct Ca²⁺-sensitivity comparisons of mutant and "normal" are included. Seventy-one independent measurements of the HCM mutation-induced change of EC₅₀ shown as EC₅₀ WT/mutant. Values range from 1.15 to 3.8-fold with a mean of 1.87 ± 0.07 (sem). Shading indicates gene studied.

Gene names: ACTC, cardiac alpha actin; TNNI3, cardiac troponin I; TNNT2, cardiac troponin T (T3 isoform); TNINC2 cardiac troponin C; MYL2, ventricular regulatory myosin light chain; MYH7, beta myosin heavy chain; MYBPC3, cardiac myosin binding protein C; TPM1, alpha tropomyosin, Tpm1.1.

Measurement methods: IVMA, *in vitro* motility assay; Fibers TG, skinned fibers from transgenic or knock-in mouse heart; Myofibrils TG, single myofibrils from transgenic or knock-in mouse heart; Fibers X, skinned fibers with mutation protein exchanged in Human fibers, skinned fibers from human heart muscle; ATPase, reconstituted thin filament activation of myosin ATPase activity.

is also worth noting that this small Ca²⁺-sensitivity shift is observed independent of the measurement method **Figure 1B** compares the ΔpCa₅₀ distribution measured by unloaded assays (actomyosin ATPase or *in vitro* motility) and by loaded assays (force measurements in skinned muscles, cell, and isolated myofibrils). The mean magnitude of the Ca²⁺-sensitivity change is about 20% less when measured in loaded assays.

TABLE 2 | Effect of skeletal muscle gain-of-function mutations on Ca²⁺-sensitivity shown as EC₅₀ WT/mutant.

| Gene name | Mutation | wt/mutant EC ₅₀ ratio | Measured in | References |
|-----------|-----------|----------------------------------|--------------|--------------------------|
| ACTA1 | K326N | 2.50 | IVMA | Jain et al., 2012 |
| TPM2 | ΔK49 | 1.19 | IVMA | Marston et al., 2013 |
| TPM2 | ΔE139 | 1.51 | IVMA | Marston et al., 2013 |
| TPM2 | E181K | 1.58 | Human fibers | Ochala et al., 2012 |
| TPM2 | ΔK7 50% | 2.00 | IVMA | Mokbel et al., 2013 |
| TPM2 | ΔK7 | 2.70 | Human fibers | Mokbel et al., 2013 |
| TPM3 | K168E | 2.67 | IVMA | Marston et al., 2013 |
| TPM3 | K168E 50% | 1.85 | IVMA | Marston et al., 2013 |
| TPM3 | ΔE224 | 1.34 | Human fibers | Donkervoort et al., 2015 |
| TPM3 | ΔE224 | 2.2 | IVMA | Donkervoort et al., 2015 |
| TPM3 | Δ218 | 2.5 | IVMA | Donkervoort et al., 2015 |

The mean change is 1.65 ± 0.16-fold (range 1.19–2.70).

GENE NAMES: ACTA1, skeletal muscle alpha actin; TPM2, beta tropomyosin, Tpm2.2; TPM3, Tpm3.12, "gamma tropomyosin."

Shading indicates gene studied.

What could be the underlying reason for this consistent and small effect of mutations on EC₅₀? We will consider two possible mechanisms that are not necessarily mutually exclusive. Firstly, it could be that the limit is set by the capacity of the EC coupling system that supplies activating Ca²⁺ and that striated muscle cannot work in a way compatible with life outside these limits; alternatively it may be due to a fundamental property of the troponin system and the permitted conformational transitions compatible with efficient regulation.

Before attempting to discuss these mechanisms it is worthwhile considering some additional evidence on Ca²⁺-sensitivity shifts. Perhaps the most puzzling observation is that there appears to be no correlation between the Ca²⁺-sensitivity shift and disease severity. Skeletal myopathy mutations that cause life-threatening muscle weakness from birth and often require mechanical assistance in breathing (Ravenscroft et al., 2015), have the same Ca²⁺-sensitivity shifts as dilated cardiomyopathy mutations which are considerably less lethal (Hershberger et al., 2013). Whilst heart muscle has compensatory strategies not available in skeletal muscle to account for this difference, the small change in Ca²⁺-sensitivity even in the most severe skeletal muscle disease might be indicative of a fundamental structure-based limit on changes in EC₅₀.

Consideration of the Ca²⁺-sensitivity shifts in cardiomyopathies (**Tables 1, 3**) do not indicate any correlation with disease severity. Any relationship that may exist is masked by the extreme variability of Ca²⁺-sensitivity shift measurements. For instance, the "severe" TNNI3 R145G HCM/RCM-linked mutation features at both extremes of the Ca²⁺-sensitivity range (1.15x and 3.65x); for the 6 assays in the table the mean is 1.84, close to the mean of all 71 HCM measurements (1.87). The same variability can be seen with other mutations where multiple values are available: ACTC E99K, *n* = 5, 1.24–2.45 mean 1.85; TPM1 E180G, *n* = 4, 1.30–2.75, mean 1.78. The second relevant observation is that the physiological modulation of cardiac muscle myofilament Ca²⁺-sensitivity due to phosphorylation

TABLE 3 | Effect of dilated cardiomyopathy linked mutations on Ca²⁺-sensitivity.

| Gene name | Mutation | wt/mutant EC ₅₀ ratio | Measured in | References |
|-----------|-------------------|----------------------------------|-------------|--------------------------|
| ACTC | E361G | 1.05 | IVMA | Song et al., 2010 |
| ACTC | E361G skTn | 0.30 | IVMA | Song et al., 2010 |
| TNNI3 | K36Q | 0.47 | IVMA | Memo et al., 2013 |
| TNNI3 | K36Q | 0.41 | ATPase | Carballo et al., 2009 |
| TNNI3 | N185K | 0.42 | ATPase | Carballo et al., 2009 |
| TNNT2 | R131W | 0.59 | ATPase | Mirza et al., 2005 |
| TNNT2 | R131W | 0.63 | IVMA | Mirza et al., 2005 |
| TNNT2 | R134G | 0.89 | Fibers X | Hershberger et al., 2009 |
| TNNT2 | R141W | 0.69 | IVMA | Memo et al., 2013 |
| TNNT2 | R141W | 0.80 | ATPase | Mirza et al., 2005 |
| TNNT2 | R141W | 0.89 | Fibers X | Venkatraman et al., 2005 |
| TNNT2 | R151C | 0.81 | Fibers X | Hershberger et al., 2009 |
| TNNT2 | R159Q | 0.83 | Fibers X | Hershberger et al., 2009 |
| TNNT2 | R206L | 0.35 | IVMA | Mirza et al., 2005 |
| TNNT2 | R205L | 0.34 | ATPase | Mirza et al., 2005 |
| TNNT2 | R205L | 0.68 | Fibers X | Mirza et al., 2005 |
| TNNT2 | R205W | 0.83 | Fibers X | Hershberger et al., 2009 |
| TNNT2 | ΔK210 hetero | 0.63 | IVMA | Du et al., 2007 |
| TNNT2 | ΔK210 | 0.75 | Fibers X | Venkatraman et al., 2005 |
| TNNT2 | ΔK210 | 0.45 | IVMA | Du et al., 2007 |
| TNNT2 | ΔK210 recombinant | 1.54 | ATPase | Mirza et al., 2005 |
| TNNT2 | ΔK210 50% | 0.46 | IVMA | Mirza et al., 2005 |
| TNNT2 | D270N | 0.65 | IVMA | Mirza et al., 2005 |
| TNNT2 | D270N | 0.64 | ATPase | Mirza et al., 2005 |
| TNNC1 | Y5H | 0.82 | Fibers X | Pinto et al., 2011 |
| TNNC1 | D73N | 0.55 | ATPase | McConnell et al., 2015 |
| TNNC1 | D73N | 0.59 | Fibers X | McConnell et al., 2015 |
| TNNC1 | D145E | 0.52 | Fibers X | Pinto et al., 2011 |
| TNNC1 | I148V | 0.91 | Fibers X | Pinto et al., 2011 |
| TNNC1 | G159D | 0.56 | ATPase | Mirza et al., 2005 |
| TNNC1 | G159D | 0.55 | IVMA | Mirza et al., 2005 |
| TNNC1 | G159D | 1.86 | IVMA | Dyer et al., 2009 |
| TNNC1 | G159D skTn | 0.56 | IVMA | Dyer et al., 2009 |
| TNNC1 | G159D | | Fibers X | Biesiadecki et al., 2007 |
| TPM1 | E40K | 0.69 | IVMA | Memo et al., 2013 |
| TPM1 | E40K baculovirus | 0.38 | IVMA | Memo et al., 2013 |
| TPM1 | E40K | 0.64 | ATPase | Chang et al., 2005 |
| TPM1 | E54K | 0.58 | ATPase | Mirza et al., 2005 |
| TPM1 | E54K | 1.90 | Ca binding | Robinson et al., 2007 |
| TPM1 | D230N baculovirus | 2.30 | IVMA | Memo et al., 2013 |
| TPM1 | D230N bacu+skTn | 0.59 | IVMA | Memo et al., 2013 |
| TPM1 | D230N Recombinant | 0.54 | ATPase | Lakdawala et al., 2010 |

Forty-two independent measurements of the mutation-induced change of EC₅₀ shown as EC₅₀ WT/mutant.

Shading indicates gene studied.

of troponin I by protein kinase A has been known to be a 2–3-fold shift for many years (Solaro et al., 2008). **Table 5** lists a number of recent determinations of this Ca²⁺-sensitivity shift

TABLE 4 | Skeletal myopathy mutations causing a loss of function.

| Gene name | Mutation | wt/mutant EC ₅₀ ratio | Measured in | References |
|-----------|-----------|----------------------------------|-------------|-----------------------|
| TPM2 | E117K | 0.41 | IVMA | Marttila et al., 2012 |
| TPM2 | Q147P | 0.63 | IVMA | Marttila et al., 2012 |
| TPM3 | L100M | 0.52 | IVMA | Marttila et al., 2012 |
| TPM3 | R167C | 0.36 | Myofibers | Ochala et al., 2012 |
| TPM3 | R167H | 0.59 | IVMA | Marston et al., 2013 |
| TPM3 | R167H 50% | 0.58 | IVMA | Marston et al., 2013 |
| TPM3 | R244G | 0.46 | IVMA | Marston et al., 2013 |
| TPM3 | R244G 50% | 0.60 | IVMA | Marston et al., 2013 |
| TPM3 | K169E | 0.55 | Myofibers | Yuen et al., 2015 |
| TPM3 | R245G | 0.45 | Myofibers | Yuen et al., 2015 |
| TPM3 | L100M | 0.53 | Myofibers | Yuen et al., 2015 |
| TPM3 | R168G | 0.48 | Myofibers | Yuen et al., 2015 |
| TPM3 | R168H | 0.42 | Myofibers | Yuen et al., 2015 |
| TPM3 | R167C | 0.39 | Myofibers | Yuen et al., 2015 |


Fourteen independent measurements of the mutation-induced change of EC₅₀ shown as EC₅₀ WT/mutant. The mean change is 0.49 ± 0.02-fold (range 0.36–0.63).


Shading indicates gene studied.

in several species and measured by both loaded and unloaded assays illustrating its small range. **Figure 1C** shows how the magnitude and distribution of measured changes is similar to the changes induced by disease-causing mutations. It would be logical to conclude that this represents the range of achievable Ca²⁺ sensitivity shifts in cardiac muscle due to the limitations of the EC coupling system.

In principle, it should be possible to go beyond the Ca²⁺-sensitivity limits set by EC coupling in an *in vitro* system where Ca²⁺ binding affinity can be much greater or much less than the native troponin. Cardiac troponin C presents extreme examples in a single molecule. Only site II binds Ca²⁺ in the physiologically relevant range (2.5 × 10⁵ M⁻¹) and so is solely responsible for Ca²⁺-regulation (Holroyde et al., 1980). A few amino acid changes in the EF-hand motifs results in sites that do not bind Ca²⁺ (Site I) or sites that bind Ca²⁺ 200x tighter (sites III and IV) and are permanently occupied by Ca²⁺ or Mg²⁺ (Li and Hwang, 2015). Thus, it would seem that neither a very high Ca²⁺ sensitivity nor a very low one are able to participate in regulation. How much deviation of Ca²⁺ affinity from the norm is compatible with muscle regulation?

It is known that for mutations, the small Ca²⁺-sensitivity changes correlate with Ca²⁺ binding affinity to thin filaments (Robinson et al., 2007). In a study of mutations induced in skeletal muscle troponin C, Davis et al. achieved a 243-fold range of Ca²⁺ binding affinities for troponin C. However, this did not translate into such a great range when Ca²⁺-binding was measured in the presence of TnI (96-148) and caused a still smaller shift in the Ca²⁺-sensitivity of force production (Davis et al., 2004). Thus, the most extreme Ca²⁺-sensitizing mutation, V45Q increased TnC Ca²⁺ binding affinity 19-fold, but the increase was only 3.1-fold when measured in the presence of the TnI peptide and Ca²⁺-sensitivity in skinned fibers was just 2.3-fold more than wild-type. This is within the same





that Ca²⁺-activation of contraction is highly cooperative. Most measurements suggest a five-fold range in free Ca²⁺ concentration during a cardiac muscle contraction. Peak Ca²⁺ concentration is about 600 nM at rest and can be substantially higher during adrenergic stimulation, thus normally muscle is only partially activated (Negretti et al., 1995; Dibb et al., 2007).

Figure 2 shows a real life example: in a mouse model of HCM (ACTC E99K) we measured both the Ca²⁺-activation curve for myofibrils and the contractility of intact papillary muscle as well as the Ca²⁺-transient (Song et al., 2013). Under the conditions of this experiment the Ca²⁺ transient was the same in Wild-type and ACTC E99K muscle, Ca²⁺ sensitivity was 0.8 μM for wild-type and 0.34 μM for ACTC E99K with a Hill coefficient of about 4. The increase in Ca²⁺-sensitivity due to the ACTC E99K HCM mutation corresponds to an approximately four-fold increase in twitch force in the absence of a change in the Ca²⁺-transient that was actually observed.

We can use this model to consider what would happen if Ca²⁺-sensitivity changed beyond the normal range. If myofilament Ca²⁺-sensitivity was 4 times normal, maximum force would reach close to 100%, leaving no range for it to be

REFERENCES

Baudenbacher, F., Schober, T., Pinto, J. R., Sidorov, V. Y., Hilliard, F., Solaro, R. J., et al. (2008). Myofilament Ca²⁺ sensitization causes susceptibility to cardiac arrhythmia in mice. *J. Clin. Invest.* 118, 3893–3903. doi: 10.1172/jci.36642

modulated by adrenergic agents. Moreover, it is likely that the muscle would not fully relax, since, based on the five-fold range of the Ca²⁺ transient even at the lowest Ca²⁺ level force would be 5–10%, a substantial fraction of the peak force of wild-type muscle, thus the hypercontractile phenotype would impose a major defect in relaxation, much more severe than the diastolic dysfunction associated with HCM mutations with only a 1.8-fold average Ca²⁺ sensitivity increase.

If myofilament Ca²⁺-sensitivity were decreased to half the normal, contractility would be very low indeed. The fact that mutations that decrease Ca²⁺-sensitivity are not lethal and indeed in transgenic mice, may exhibit little phenotype, is probably due to a compensatory increase in the Ca²⁺-transient (Du et al., 2007). However, this compensation may not be enough to support normal contraction in the long term, leading to DCM, the phenotype commonly associated with reduced Ca²⁺-sensitivity.

CONCLUSION

The objective of this article was to confirm that Ca²⁺-sensitivity of contractility only varies within a narrow range of three-fold above and below the normal EC₅₀ at rest and to investigate why this should be. The high cooperativity of muscle activation by Ca²⁺ means there is a narrow [Ca²⁺] range between relaxed and active muscle. It would appear that the excitation-contraction coupling machinery of the cell has limited ability to change the amplitude of the Ca²⁺-transient or baseline [Ca²⁺] to compensate for changes in EC₅₀; thus increased Ca²⁺-sensitivity would be limited by inability to relax and reduced Ca²⁺-sensitivity would be limited by inability to contract. It is intriguing that the Ca²⁺-sensitivity range of the thin filament itself is independently limited. Mutations that change Ca²⁺-binding affinity to TnC by a large amount nevertheless only produce a small change in EC₅₀ for activation of loaded or unloaded contractility *in vitro*. Whether this property is an evolutionary adaptation that limits the deleterious effects of mutations in thin filaments or simply fortuitous is unknown.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and approved it for publication.

FUNDING

SM's research is funded by British Heart Foundation programme grant RG/11/20/29266.

Bayliss, C. R., Jacques, A. M., Leung, M.-C., Ward, D. G., Redwood, C. S., Gallon, C. E., et al. (2012). Myofibrillar Ca²⁺-sensitivity is uncoupled from troponin I phosphorylation in hypertrophic obstructive cardiomyopathy due to abnormal troponin T. *Cardiovasc. Res.* 97, 500–508. doi: 10.1093/cvr/cvs322

Biesiadecki, B. J., Kobayashi, T., Walker, J. S., John Solaro, R., and de Tombe, P. (2007). The troponin C G159D mutation blunts myofilament desensitization

- induced by troponin I Ser23/24 phosphorylation. *Circ. Res.* 100, 1486–1493. doi: 10.1161/01.RES.0000267744.92677.7f
- Bing, W., Knott, A., Redwood, C. S., Esposito, G., Purcell, I., Watkins, S., et al. (2000). Effect of hypertrophic cardiomyopathy mutations in human cardiac α -tropomyosin (Asp175Asn and Glu180Gly) on the regulatory properties of human cardiac troponin determined by *in vitro* motility assay. *Biochem. Biophys. Res. Commun.* 32, 1489–1498. doi: 10.1006/jmcc.2000.1182
- Blanchard, E., Seidman, C., Seidman, J. G., LeWinter, M., and Maughan, D. (1999). Altered crossbridge kinetics in the α MHC403/+ mouse model of familial hypertrophic cardiomyopathy. *Circ. Res.* 84, 475–483. doi: 10.1161/01.RES.84.4.475
- Brunet, N. M., Chase, P. B., Mihajlovic, G., and Schoffstall, B. (2014). Ca²⁺-regulatory function of the inhibitory peptide region of cardiac troponin I is aided by the C-terminus of cardiac troponin T: effects of familial hypertrophic cardiomyopathy mutations cTnI R145G and cTnT R278C, alone and in combination, on filament sliding. *Arch. Biochem. Biophys.* 552, 11–20. doi: 10.1016/j.abb.2013.12.021
- Carballo, S., Robinson, P., Otway, R., Fatkin, D., Jongbloed, J. D., de Jonge, N., et al. (2009). Identification and functional characterization of cardiac troponin I as a novel disease gene in autosomal dominant dilated cardiomyopathy. *Circ. Res.* 105, 375–382. doi: 10.1161/CIRCRESAHA.109.196055
- Chang, A. N., Harada, K., Ackerman, M. J., and Potter, J. D. (2005). Functional consequences of hypertrophic and dilated cardiomyopathy-causing mutations in α -tropomyosin. *J. Biol. Chem.* 280, 34343–34349. doi: 10.1074/jbc.M505014200
- Davis, J. P., Rall, J. A., Alionte, C., and Tikunova, S. B. (2004). Mutations of hydrophobic residues in the N-terminal domain of troponin C affect calcium binding and exchange with the troponin C-troponin I 96-148 complex and muscle force production. *J. Biol. Chem.* 279, 17348–17360. doi: 10.1074/jbc.M314095200
- Deng, Y., Schmidtmann, A., Redlich, A., Westerdorf, B., Jaquet, K., and Thieleczek, R. (2001). Effects of phosphorylation and mutation R145G on human cardiac troponin I function. *Biochemistry* 40, 14593–14602. doi: 10.1021/bi0115232
- Dibb, K. M., Eisner, D. A., and Trafford, A. W. (2007). Regulation of systolic [Ca²⁺]_i and cellular Ca²⁺ flux balance in rat ventricular myocytes by SR Ca²⁺, L-type Ca²⁺ current and diastolic [Ca²⁺]_i. *J. Physiol.* 585, 579–592. doi: 10.1113/jphysiol.2007.141473
- Donkervoort, S., Papadaki, M., de Winter, J. M., Neu, M. B., Kirschner, J., Bolduc, V., et al. (2015). TPM3 deletions cause a hypercontractile congenital muscle stiffness phenotype. *Ann. Neurol.* 78, 982–994. doi: 10.1002/ana.24535
- Du, C. K., Morimoto, S., Nishii, K., Minakami, R., Ohta, M., Tadano, N., et al. (2007). Knock-in mouse model of dilated cardiomyopathy caused by troponin mutation. *Circ. Res.* 101, 185–194. doi: 10.1161/CIRCRESAHA.106.146670
- Dyer, E. C., Jacques, A. M., Hoskins, A. C., Ward, D. G., Gallon, C. E., Messer, A. E., et al. (2009). Functional analysis of a unique troponin C mutation, Gly159Asp that causes familial dilated cardiomyopathy, studied in explanted heart muscle. *Circ. Heart Fail.* 2, 456–464. doi: 10.1161/CIRCHEARTFAILURE.108.818237
- Elliott, K., Watkins, H., and Redwood, C. S. (2000). Altered regulatory properties of human cardiac troponin I mutants that cause hypertrophic cardiomyopathy. *J. Biol. Chem.* 275, 22069–22074. doi: 10.1074/jbc.M002502200
- Ford, S. J., Mamidi, R., Jimenez, J., Tardiff, J. C., and Chandra, M. (2012). Effects of R92 mutations in mouse cardiac troponin T are influenced by changes in myosin heavy chain isoform. *J. Mol. Cell. Cardiol.* 53, 542–551. doi: 10.1016/j.yjmcc.2012.07.018
- Frayse, B., Weinberger, F., Bardswell, S. C., Cuello, F., Vignier, N., Geertz, B., et al. (2012). Increased myofilament Ca²⁺ sensitivity and diastolic dysfunction as early consequences of *Myhpc3* mutation in heterozygous knock-in mice. *J. Mol. Cell. Cardiol.* 52, 1299–1307. doi: 10.1016/j.yjmcc.2012.03.009
- Gafurov, B., Fredricksen, S., Cai, A., Brenner, B., Chase, P. B., and Chalovich, J. M. (2004). The Delta 14 mutation of human cardiac troponin T enhances ATPase activity and alters the cooperative binding of S1-ADP to regulated actin. *Biochemistry* 43, 15276–15285. doi: 10.1021/bi048646h
- Gomes, A. V., Harada, K., and Potter, J. D. (2005a). A mutation in the N-terminus of troponin I that is associated with hypertrophic cardiomyopathy affects the Ca(2+)-sensitivity, phosphorylation kinetics and proteolytic susceptibility of troponin I. *J. Mol. Cell. Cardiol.* 39, 754–765. doi: 10.1016/j.yjmcc.2005.05.013
- Gomes, A. V., Liang, J., and Potter, J. D. (2005b). Mutations in human cardiac troponin I that are associated with restrictive cardiomyopathy affect basal ATPase activity and the calcium sensitivity of force development. *J. Biol. Chem.* 280, 30909–30915. doi: 10.1074/jbc.M500287200
- Heller, M. J., Nili, M., Homsher, E., and Tobacman, L. S. (2003). Cardiomyopathic tropomyosin mutations that increase thin filament Ca²⁺ sensitivity and tropomyosin N-domain flexibility. *J. Biol. Chem.* 278, 41742–41748. doi: 10.1074/jbc.M303408200
- Hershberger, R. E., Hedges, D. J., and Morales, A. (2013). Dilated cardiomyopathy: the complexity of a diverse genetic architecture. *Nat. Rev. Cardiol.* 10, 531–547. doi: 10.1038/nrcardio.2013.105
- Hershberger, R. E., Pinto, J. R., Parks, S. B., Kushner, J. D., Li, D., Ludwigsen, S., et al. (2009). Clinical and functional characterization of TNNT2 mutations identified in patients with dilated cardiomyopathy. *Circ. Cardiovasc. Genet.* 2, 306–313. doi: 10.1161/CIRCGENETICS.108.846733
- Holroyde, M. J., Robertson, S. P., Johnson, J. D., Solaro, R. J., and Potter, J. D. (1980). The calcium and magnesium binding sites on cardiac troponin and their role in the regulation of myofibrillar adenosine triphosphatase. *J. Biol. Chem.* 255, 11688–11693.
- Jain, R. K., Jayawant, S., Squier, W., Muntoni, F., Sewry, C. A., Manzur, A., et al. (2012). Nemaline myopathy with stiffness and hypertonias associated with an ACTA1 mutation. *Neurology* 78, 1100–1103. doi: 10.1212/WNL.0b013e31824e8e8e
- Kerrick, W. G., Kazmierczak, K., Xu, Y., Wang, Y., and Szczesna-Cordary, D. (2009). Malignant familial hypertrophic cardiomyopathy D166V mutation in the ventricular myosin regulatory light chain causes profound effects in skinned and intact papillary muscle fibers from transgenic mice. *FASEB J.* 23, 855–865. doi: 10.1096/fj.08-118182
- Köhler, J., Chen, Y., Brenner, B., Gordon, A. M., Kraft, T., Martyn, D. A., et al. (2003). Familial hypertrophic cardiomyopathy mutations in troponin I (K183D, G203S, K206Q) enhance filament sliding. *Physiol. Genomics* 14, 117–128. doi: 10.1152/physiolgenomics.00101.2002
- Krüger, M., Zittrich, S., Redwood, C., Blaudeck, N., James, J., Robbins, J., et al. (2005). Effects of the mutation R145G in human cardiac troponin I on the kinetics of the contraction–relaxation cycle in isolated cardiac myofibrils. *J. Physiol.* 564, 347–357. doi: 10.1113/jphysiol.2004.079095
- Lakdawala, N. K., Dellefave, L., Redwood, C. S., Sparks, E., Cirino, A. L., Depalma, S., et al. (2010). Familial dilated cardiomyopathy caused by an α -tropomyosin mutation: the distinctive natural history of sarcomeric dilated cardiomyopathy. *J. Am. Coll. Cardiol.* 55, 320–329. doi: 10.1016/j.jacc.2009.11.017
- Lang, R., Gomes, A. V., Zhao, J., Housmans, P. R., Miller, T., and Potter, J. D. (2002). Functional analysis of a troponin I (R145G) mutation associated with familial hypertrophic cardiomyopathy. *J. Biol. Chem.* 277, 11670–11678. doi: 10.1074/jbc.M108912200
- Li, A. Y., Stevens, C. M., Liang, B., Rayani, K., Little, S., Davis, J., et al. (2013). Familial hypertrophic cardiomyopathy related cardiac troponin C L29Q mutation alters length-dependent activation and functional effects of phosphomimetic troponin I*. *PLoS ONE* 8:e79363. doi: 10.1371/journal.pone.0079363
- Li, M. X., and Hwang, P. M. (2015). Structure and function of cardiac troponin C (TNNC1): implications for heart failure, cardiomyopathies, and troponin modulating drugs. *Gene* 571, 153–166. doi: 10.1016/j.gene.2015.07.074
- Lu, Q. W., Morimoto, S., Harada, K., Du, C. K., Takahashi-Yanaga, F., Miwa, Y., et al. (2003). Cardiac troponin T mutation R141W found in dilated cardiomyopathy stabilizes the troponin T-tropomyosin interaction and causes a Ca(2+) desensitization. *J. Mol. Cell. Cardiol.* 35, 1421–1427. doi: 10.1016/j.yjmcc.2003.09.003
- Marston, S. B. (2011). How do mutations in contractile proteins cause the primary familial cardiomyopathies? *J. Cardiovasc. Transl. Res.* 4, 245–255. doi: 10.1007/s12265-011-9266-2
- Marston, S., Memo, M., Messer, A., Papadaki, M., Nowak, K., McNamara, E., et al. (2013). Mutations in repeating structural motifs of tropomyosin cause gain of function in skeletal muscle myopathy patients. *Hum. Mol. Genet.* 22, 4978–4987. doi: 10.1093/hmg/ddt345
- Martins, A. S., Parvatiyar, M. S., Feng, H.-Z., Bos, J. M., Gonzalez-Martinez, D., Vukmirovic, M., et al. (2015). *In vivo* analysis of troponin C knock-in (A8V) mice: evidence that TNNC1 is a hypertrophic

- cardiomyopathy susceptibility gene. *Circ. Cardiovasc. Genet.* 8, 653–664. doi: 10.1161/CIRCGENETICS.114.000957
- Marttila, M., Lehtokari, V.-L., Marston, S., Nyman, T. A., Barnerias, C., Beggs, A. H., et al. (2014). Mutation update and genotype-phenotype correlations of novel and previously described mutations in TPM2 and TPM3 causing congenital myopathies. *Hum. Mutat.* 35, 779–790. doi: 10.1002/humu.22554
- Marttila, M., Lemola, E., Wallefeld, W., Memo, M., Donner, K., Laing, N. G., et al. (2012). Abnormal actin binding of aberrant β -tropomyosins is a molecular cause of muscle weakness in TPM2-related nemaline and cap myopathy. *Biochem. J.* 442, 231–239. doi: 10.1042/BJ20111030
- McConnell, B. K., Singh, S., Fan, Q., Hernandez, A., Portillo, J. P., Reiser, P. J., et al. (2015). Knock-in mice harboring a Ca²⁺ desensitizing mutation in cardiac troponin C develop early onset dilated cardiomyopathy. *Front. Physiol.* 6:242. doi: 10.3389/fphys.2015.00242
- Memo, M., Leung, M.-C., Ward, D. G., dos Remedios, C., Morimoto, S., Zhang, L., et al. (2013). Mutations in thin filament proteins that cause familial dilated cardiomyopathy uncouple troponin I Phosphorylation from changes in myofibrillar Ca²⁺-sensitivity. *Cardiovasc. Res.* 99, 65–73. doi: 10.1093/cvr/cvt071
- Messer, A. (2007). *Structural and Functional Polymorphisms of Troponin in Failing Heart*. Ph.D., Thesis NHLI London, London, 343.
- Messer, A., Bayliss, C., El-Mezgueldi, M., Redwood, C., Ward, D. G., Leung, M.-C et al. (2016b). Mutations in troponin T associated with Hypertrophic Cardiomyopathy increase Ca²⁺-sensitivity and suppress the modulation of Ca²⁺-sensitivity by troponin I phosphorylation. *Arch. Biochem. Biophys.* 601, 113–120. doi: 10.1016/j.abb.2016.03.027
- Messer, A. E., Jacques, A. M., and Marston, S. B. (2007). Troponin phosphorylation and regulatory function in human heart muscle: dephosphorylation of Ser23/24 on troponin I could account for the contractile defect in end-stage heart failure. *J. Mol. Cell. Cardiol.* 42, 247–259. doi: 10.1016/j.yjmcc.2006.08.017
- Messer, A. E., Papadaki, M., Vikhorev, P. G., Sebzali, Y., El-Mezgueldi, M., Daley, A., et al. (2016a). *Primary effects of HCM mutations in humans and cats. *Biophys. J.* 110, 123a–124a. doi: 10.1016/j.bpj.2015.11.713
- Michele, D. E., Albayya, F. P., and Metzger, J. M. (1999). Direct, convergent hypersensitivity of calcium-activated force generation produced by hypertrophic cardiomyopathy mutant alpha-tropomyosins in adult cardiac myocytes. *Nat. Med.* 5, 1413–1417. doi: 10.1038/70990
- Mirza, M., Marston, S., Willott, R., Ashley, C., Mogensen, J., McKenna, W., et al. (2005). Dilated cardiomyopathy mutations in three thin filament regulatory proteins result in a common functional phenotype. *J. Biol. Chem.* 280, 28498–28506. doi: 10.1074/jbc.M412281200
- Mokbel, N., Ilkovski, B., Kreissl, M., Memo, M., Jeffries, C. M., Marttila, M., et al. (2013). K7del is a common TPM2 gene mutation associated with nemaline myopathy and raised myofibre calcium sensitivity. *Brain* 136, 494–507. doi: 10.1093/brain/aww348
- Negretti, N., Varro, A., and Eisner, D. A. (1995). Estimate of net calcium fluxes and sarcoplasmic reticulum calcium content during systole in rat ventricular myocytes. *J. Physiol.* 486(Pt 3), 581–591. doi: 10.1113/jphysiol.1995.sp020836
- Ochala, J., Gokhin, D. S., Péniisson-Besnier, I., Quijano-Roy, S., Monnier, N., Lunardi, J., et al. (2012). Congenital myopathy-causing tropomyosin mutations induce thin filament dysfunction via distinct physiological mechanisms. *Hum. Mol. Genet.* 21, 4473–4485. doi: 10.1093/hmg/dds289
- Palmer, B. M., Fishbaugher, D. E., Schmitt, J. P., Wang, Y., Alpert, N. R., Seidman, C. E., et al. (2004). Differential cross-bridge kinetics of FHC myosin mutations R403Q and R453C in heterozygous mouse myocardium. *Am. J. Physiol. Heart Circ. Physiol.* 287, H91–H99. doi: 10.1152/ajpheart.01015.2003
- Papadaki, M., Vikhorev, P. G., Marston, S. B., and Messer, A. E. (2015). Uncoupling of myofilament Ca²⁺ sensitivity from troponin I phosphorylation by mutations can be reversed by epigallocatechin-3-gallate. *Cardiovasc. Res.* 108, 99–110. doi: 10.1093/cvr/cvv181
- Parvatiyar, M. S., Landstrom, A. P., Figueiredo-Freitas, C., Potter, J. D., Ackerman, M. J., Pinto, J. R., et al. (2012). A mutation in TNNC1-encoded cardiac troponin C, TNNC1-A31S, predisposes to hypertrophic cardiomyopathy and ventricular fibrillation. *J. Biol. Chem.* 287, 31845–31855. doi: 10.1074/jbc.M112.377713
- Parvatiyar, M. S., Pinto, J. R., Liang, J., and Potter, J. D. (2010). Predicting cardiomyopathic phenotypes by altering Ca²⁺ affinity of cardiac troponin C. *J. Biol. Chem.* 285, 27785–27797. doi: 10.1074/jbc.M110.112326
- Pinto, J. R., Parvatiyar, M. S., Jones, M. A., Liang, J., Ackerman, M. J., and Potter, J. D. (2009). A functional and structural study of troponin C mutations related to hypertrophic cardiomyopathy. *J. Biol. Chem.* 284, 19090–19100. doi: 10.1074/jbc.M109.007021
- Pinto, J. R., Siegfried, J. D., Parvatiyar, M. S., Li, D., Norton, N., Jones, M. A., et al. (2011). Functional characterization of TNNC1 rare variants identified in dilated cardiomyopathy. *J. Biol. Chem.* 286, 34404–34412. doi: 10.1074/jbc.M111.267211
- Ravenscroft, G., Laing, N. G., and Bönnemann, C. G. (2015). Pathophysiological concepts in the congenital myopathies: blurring the boundaries, sharpening the focus. *Brain* 138(Pt 2), 246–268. doi: 10.1093/brain/awu368
- Redwood, C., Lohmann, K., Bing, W., Esposito, G., Elliott, K., Abdurazzak, H., et al. (2000). Investigation of a truncated troponin T that causes familial hypertrophic cardiomyopathy: Ca²⁺ regulatory properties of reconstituted thin filaments depend on the ratio of mutant to wild-type peptide. *Circ. Res.* 86, 1146–1152. doi: 10.1161/01.RES.86.11.1146
- Robinson, P., Griffiths, P. J., Watkins, H., and Redwood, C. S. (2007). Dilated and hypertrophic cardiomyopathy mutations in troponin and alpha-tropomyosin have opposing effects on the calcium affinity of cardiac thin filaments. *Circ. Res.* 101, 1266–1273. doi: 10.1161/CIRCRESAHA.107.156380
- Robinson, P., Mirza, M., Knott, A., Abdurazzak, H., Willott, R., Marston, S., et al. (2002). Alterations in thin filament regulation induced by a human cardiac troponin T mutant that causes dilated cardiomyopathy are distinct from those induced by troponin T mutants that cause hypertrophic cardiomyopathy. *J. Biol. Chem.* 277, 40710–40716. doi: 10.1074/jbc.M203446200
- Schmidtman, A., Lindow, C., Villard, S., Heuser, A., Mügge, A., Gessner, R., et al. (2005). Cardiac troponin C-L29Q, related to hypertrophic cardiomyopathy, hinders the transduction of the protein kinase A dependent phosphorylation signal from cardiac troponin I to C. *FEBS J.* 272, 6087–6097. doi: 10.1111/j.1742-4658.2005.05001.x
- Sequeira, V., Wijnker, P. J., Nijenkamp, L. L., Kuster, D. W., Najafi, A., Witjas-Paalberends, E. R., et al. (2013). Perturbed length-dependent activation in human hypertrophic cardiomyopathy with missense sarcomeric gene mutations. *Circ. Res.* 112, 1491–1505. doi: 10.1161/CIRCRESAHA.111.300436
- Solaro, R. J., Rosevear, P., and Kobayashi, T. (2008). The unique functions of cardiac troponin I in the control of cardiac muscle contraction and relaxation. *Biochem. Biophys. Res. Commun.* 369, 82–87. doi: 10.1016/j.bbrc.2007.12.114
- Song, W., Dyer, E., Stuckey, D., Copeland, O., Leung, M., Bayliss, C., et al. (2011). Molecular mechanism of the Glu99lys mutation in cardiac actin (ACTC gene) that causes apical hypertrophy in man and mouse. *J. Biol. Chem.* 286, 27582–27593. doi: 10.1074/jbc.M111.252320
- Song, W., Dyer, E., Stuckey, D., Leung, M.-C., Memo, M., Mansfield, C., et al. (2010). Investigation of a transgenic mouse model of familial dilated cardiomyopathy. *J. Mol. Cell. Cardiol.* 49, 380–389. doi: 10.1016/j.yjmcc.2010.05.009
- Song, W., Vikhorev, P. G., Kashyap, M. N., Rowlands, C., Ferenczi, M. A., Woledge, R. C., et al. (2013). Mechanical and energetic properties of papillary muscle from ACTC E99K transgenic mouse models of hypertrophic cardiomyopathy. *Am. J. Physiol. Heart Circ. Physiol.* 304, H1513–H1524. doi: 10.1152/ajpheart.00951.2012
- Szczesna, D., Zhang, R., Zhao, J., Jones, M., Guzman, G., and Potter, J. D. (2000). Altered regulation of cardiac muscle contraction by troponin T mutations that cause familial hypertrophic cardiomyopathy. *J. Biol. Chem.* 275, 624–630. doi: 10.1074/jbc.275.1.624
- Szczesna-Cordary, D., Guzman, G., Ng, S. S., and Zhao, J. (2004). Familial hypertrophic cardiomyopathy-linked alterations in Ca²⁺ binding of human cardiac myosin regulatory light chain affect cardiac muscle contraction. *J. Biol. Chem.* 279, 3535–3542. doi: 10.1074/jbc.M307092200
- Takahashi-Yanaga, F., Morimoto, S., Harada, K., Minakami, R., Shiraishi, F., Ohta, M., et al. (2001). Functional consequences of the mutations in human cardiac troponin I gene found in familial hypertrophic cardiomyopathy. *J. Mol. Cell. Cardiol.* 33, 2095–2107. doi: 10.1006/jmcc.2001.1473
- van der Velden, J., Papp, Z., Zaremba, R., Boontje, N. M., de Jong, J. W., Owen, V. J., et al. (2003). Increased Ca²⁺-sensitivity of the contractile apparatus in end-stage human heart failure results from altered phosphorylation

- of contractile proteins. *Cardiovasc. Res.* 57, 37–47. doi: 10.1016/S0008-6363(02)00606-5
- Venkatraman, G., Gomes, A. V., Kerrick, W. G., and Potter, J. D. (2005). Characterization of troponin T dilated cardiomyopathy mutations in the fetal troponin isoform. *J. Biol. Chem.* 280, 17584–17592. doi: 10.1074/jbc.M409337200
- Vikhorev, P. G., Song, W., Wilkinson, R., Copeland, O., Messer, A. E., Ferenczi, M. A., et al. (2014). The dilated cardiomyopathy-causing mutation ACTC E361G in cardiac muscle myofibrils specifically abolishes modulation of Ca^{2+} regulation by phosphorylation of troponin I. *Biophys. J.* 107, 2369–2380. doi: 10.1016/j.bpj.2014.10.024
- Warren, C. M., Karam, C. N., Wolska, B. M., Kobayashi, T., de Tombe, P. P., Arteaga, G. M., et al. (2015). A green tea catechin normalizes the enhanced Ca^{2+} sensitivity of myofilaments regulated by a hypertrophic cardiomyopathy associated mutation in human cardiac troponin I (K206I). *Circ. Cardiovasc. Genet.* 8, 765–773. doi: 10.1161/CIRCGENETICS.115.001234
- Yuan, C.-C., Muthu, P., Kazmierczak, K., Liang, J., Huang, W., Irving, T. C., et al. (2015). Constitutive phosphorylation of cardiac myosin regulatory light chain prevents development of hypertrophic cardiomyopathy in mice. *Proc. Natl. Acad. Sci. U.S.A.* 112, E4138–E4146. doi: 10.1073/pnas.1505819112
- Yuen, M., Cooper, S. T., Marston, S. B., Nowak, K. J., McNamara, E., Mokbel, N., et al. (2015). Muscle weakness in TPM3-myopathy is due to reduced Ca^{2+} -sensitivity and impaired acto-myosin cross-bridge cycling in slow fibres. *Hum. Mol. Genet.* 24, 6278–6292. doi: 10.1093/hmg/ddv334

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Marston. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.