



Carotid Body Type-I Cells Under Chronic Sustained Hypoxia: Focus on Metabolism and Membrane Excitability

Raúl Pulgar-Sepúlveda¹, Rodrigo Varas^{1,2}, Rodrigo Iturriaga³, Rodrigo Del Río^{4,5,6*} and Fernando C. Ortiz^{1*}

¹ Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad Autónoma de Chile, Santiago, Chile, ² Facultad de Ciencias de la Salud, Universidad Autónoma de Chile, Talca, Chile, ³ Laboratorio de Neurobiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile, ⁴ Laboratory of Cardiorespiratory Control, Department of Physiology, Pontificia Universidad Católica de Chile, Santiago, Chile, ⁵ Centro de Envejecimiento y Regeneración, Pontificia Universidad Católica de Chile, Santiago, Chile, ⁶ Centro de Excelencia en Biomedicina de Magallanes, Universidad de Magallanes, Punta Arenas, Chile

OPEN ACCESS

Edited by:

Eduardo Colombari,
Universidade Estadual Paulista Júlio
de Mesquita Filho (UNESP), Brazil

Reviewed by:

Thiago S. Moreira,
Universidade de São Paulo, Brazil
Davi J. A. Moraes,
Universidade de São Paulo, Brazil

*Correspondence:

Rodrigo Del Río
rdelrio@bio.puc.cl
Fernando C. Ortiz
fernando.ortiz@uautonoma.cl

Specialty section:

This article was submitted to
Integrative Physiology,
a section of the journal
Frontiers in Physiology

Received: 29 May 2018

Accepted: 24 August 2018

Published: 19 September 2018

Citation:

Pulgar-Sepúlveda R, Varas R,
Iturriaga R, Del Río R and Ortiz FC
(2018) Carotid Body Type-I Cells
Under Chronic Sustained Hypoxia:
Focus on Metabolism and Membrane
Excitability. *Front. Physiol.* 9:1282.
doi: 10.3389/fphys.2018.01282

Chronic sustained hypoxia (CSH) evokes ventilatory acclimatization characterized by a progressive hyperventilation due to a potentiation of the carotid body (CB) chemosensory response to hypoxia. The transduction of the hypoxic stimulus in the CB begins with the inhibition of K⁺ currents in the chemosensory (type-I) cells, which in turn leads to membrane depolarization, Ca²⁺ entry and the subsequent release of one- or more-excitatory neurotransmitters. Several studies have shown that CSH modifies both the level of transmitters and chemoreceptor cell metabolism within the CB. Most of these studies have been focused on the role played by such putative transmitters and modulators of CB chemoreception, but less is known about the effect of CSH on metabolism and membrane excitability of type-I cells. In this mini-review, we will examine the effects of CSH on the ion channels activity and excitability of type-I cell, with a particular focus on the effects of CSH on the TASK-like background K⁺ channel. We propose that changes on TASK-like channel activity induced by CSH may contribute to explain the potentiation of CB chemosensory activity.

Keywords: carotid body, chronic hypoxia, membrane depolarization, ion channels, TASK-like channel

INTRODUCTION

The carotid body (CB) is the main peripheral chemoreceptor in mammals. Natural stimuli such as hypoxemia, hypercapnia, and/or acidosis increase the firing rate of the petrosal ganglion (PG) sensory afferent neurons projecting to the cardiovascular and respiratory regions in the brain stem (Gonzalez et al., 1994).

The current model for hypoxic chemoreception states that hypoxia evokes a depolarization of CB type-I (glomus) cells, leading to an increment of intracellular Ca²⁺ and the subsequent release of one- or more-excitatory neurotransmitters to the nerve terminals of the PG neurons (Gonzalez et al., 1994; Iturriaga and Alcayaga, 2004; Nurse, 2005). It is well established that a key event in triggering the hypoxic response is the depolarization of the type-I cells (Buckler and Vaughan-Jones, 1994). In type-I cells from several species it has been found that hypoxia produces a fast

and reversible inhibition of K^+ currents (López-Barneo et al., 1988; López-López and González, 1992; Wyatt and Peers, 1995; Buckler, 1997) leading to a depolarization of type-I cells membrane and the consequence Ca^{2+} entry, mainly throughout L-type Ca^{2+} channels (Fieber and McCleskey, 1993; Wyatt et al., 1994). Among several molecules present in type-I cells, acetylcholine (ACh) and adenosine triphosphate (ATP) meets most of the criteria to be considered excitatory transmitters in the pathway (Varas et al., 2003; Iturriaga and Alcajaga, 2004), while dopamine (DA), nitric oxide (NO), and endothelin-1 (ET-1) modulate the chemosensory process (Iturriaga and Alcajaga, 2004).

Mammals exposed to sustained hypoxia (i.e., high altitude) develop ventilatory acclimatization featured by a progressively hyperventilation, due to an augmented CB responsiveness to hypoxia (Bisgard, 2000). In addition to the enhanced CB chemosensory responses, chronic sustained hypoxia (CSH) for weeks or months induces angiogenesis in the CB along with type-I cell hypertrophy and hyperplasia (Heath et al., 1985). Most of the studies on CB hypoxia acclimatization have been focused on putative changes in transmitters or modulators of CB chemoreception. Interestingly, DA, NO, and ET-1 are up-regulated in the CB during the first week of chronic hypoxia (Bisgard, 2000). In addition, some reports indicate that CSH increases type-I cell excitability due to changes in K^+ and Na^+ channels expression. In spite of the efforts, the mechanisms by which CSH enhances CB chemosensory responses to hypoxia remains to be established. In this review, we will examine the effects of sustained hypoxia on CB metabolism and ion channels function, focusing on the type-I cell excitability.

ELECTRICAL PROPERTIES OF TYPE-I CELLS


Type-I cells are small round-shaped cells (diameter $\sim 10 \mu\text{m}$ in the rat), with high input resistance of 5–6 $G\Omega$ (Duchen et al., 1988; Xu et al., 2005), and resting membrane potential ranging from -50 to -70 mV (Gonzalez et al., 1994). Regarding to the inward currents expressed by type-I cells, in the adult rabbit it has been reported a tetrodotoxin-sensitive, voltage-gated Na^+ current with fast activation and inactivation kinetics (Duchen et al., 1988). However, in both adult and neonatal rat type-I cells, there are contradictory results regarding the expression of functional voltage gated Na^+ channels (Fieber and McCleskey, 1993; Hempleman, 1995; Stea et al., 1995; Cáceres et al., 2007). Voltage-gated Ca^{2+} channels have been found in both rat and rabbit type-I cells (Fieber and McCleskey, 1993; Buckler and Vaughan-Jones, 1994; Wyatt et al., 1994). Most of the studies agree that L-type Ca^{2+} channel is the most abundant subtype of Ca^{2+} channel in type-I cells, however, in rabbit CB type-I cells, pharmacological evidences suggest the additional presence of N-, P/Q-type Ca^{2+} channels and a ω -conotoxin-resistant Ca^{2+} current (Overholt and Prabhakar, 1997, 1999). In addition, rat type-I cells express anionic currents (mainly Cl^-) which are involved in the chemosensory transduction of acidic and hypercapnic stimuli (Carpenter and Peers, 1997; Iturriaga et al., 1998).

Type-I cells express a wide variety of K^+ channels. Rabbit type-I cells express at least two voltage-gated, TEA-sensitive Ca^{2+} -independent K^+ currents. One of these conductances correspond to an oxygen-sensitive voltage-gated K^+ channel with a unitary conductance of ~ 40 pS and an activation threshold around -40 to -30 mV (hereafter KO_2 ; López-Barneo et al., 1988). The opening of this K^+ channel is reversibly inhibited by hypoxia with an IP_{50} (PO_2 at which 50% of maximal inhibition is reached) near to 5–10 mmHg. This inhibition by hypoxia depends on membrane potential: maximal inhibition (40% of control activity) is reached at 0 mV (López-Barneo et al., 1988). The KO_2 channel has at least four closed states (C0–C4), one open state (O) and two inactivated states (I1–I2). Hypoxia induces both, stabilization of the “C0” state and promoting the channel from “O” state to inactivation state “I1” (Ganformina and Lopez-Barneo, 1992).

Neonatal rat type-I cells express a maxi-K (BK) channel, whose opening depends on the PO_2 . This maxi-K channel has a large unitary conductance of ~ 200 pS, it is blocked by charybdotoxin and activated by both membrane depolarization (threshold ranging from -40 to -20 mV) and by a rise in intracellular $[Ca^{2+}]$ above 100 nM (Peers, 1990; Wyatt and Peers, 1995). Wyatt and Peers (1995) found that hypoxia ($PO_2 = 5$ –10 mmHg) causes a reversible inhibition of this maxi-K current in cell-attached patch-clamp recordings. Later, Williams et al. (2004) proposed that the oxygen-sensitivity of the rat type-I cells is mediated by a heme oxygenase-2 (HO-2) associated with the maxi-K channel complex. However, this proposition was challenged by Ortega-Sáenz et al. (2006), when they found that hypoxic response of type-I cells remains intact in HO-2 null mice.

Interestingly, Buckler (1997) found that voltage gated K^+ channel blockers (10 mM TEA plus 5 mM 4-aminopyridine and 20 nM charybdotoxin) failed to depolarize type-I cells and to modify intracellular $[Ca^{2+}]$. However, in the presence of these K^+ channel blockers, a hypoxic stimulus was able to evoke cell depolarization and a rise in intracellular Ca^{2+} levels. This background K^+ current was inhibited by hypoxia with an $IP_{50} \sim 10$ mmHg, reaching a maximal $\sim 70\%$ inhibition during anoxia. Hypoxia stabilized the background K^+ channel in its close state, but did not affect the opening/closing kinetics, or the open state duration (Williams and Buckler, 2004). This background K^+ channel correspond to a TWIK-related acid-sensitive K^+ channel (TASK), member of the two-pore domain K^+ channel superfamily (Buckler et al., 2000). Additional evidences suggest the presence of TASK-1, TASK-2, TASK-3, TRAAK, and TREK (Buckler, 1997; Buckler et al., 2000; Yamamoto et al., 2002; Yamamoto and Taniguchi, 2006).

The discovery of oxygen-dependent voltage-gated K^+ channels promptly leads to the suggestion that hypoxia inhibits K^+ channels, which in turn produces type-I cells depolarization. Nevertheless, the resting membrane potential of type-I cells is quite stable at least 10 mV below the threshold activation of both maxi-K and KO_2 channels. Therefore, it was hard to conciliate a possible role of these K^+ channels in initiating the depolarization in response to hypoxia in type-I cells. Since background K^+ channels (Buckler, 1997; Buckler et al., 2000)



are active at resting conditions, the closure of these channels may explain the initiation of the depolarization evoked by hypoxia. The precise mechanism by which acute hypoxia is sensed remains controversial, but is clear that TASK, maxi-K, and KO_2 channels plays a key role in the depolarization required for the neurotransmitters release from type-I cells in response to hypoxia (Figure 1).

CHRONIC SUSTAINED HYPOXIA (CSH)

Hypoxic-hypoxia (a PO_2 fall), could be classified as acute (seconds to minutes) or chronic (days to years). Acute hypoxia produced CB chemosensory excitation that evokes a reflex hyperventilation. CB increases its size in response to chronic sustained hypoxia (CSH) – to differentiate it from the *intermittent* paradigm – due to both increased number of cells and enhanced cell bodies diameter (Pequignot et al., 1984; Mills and Nurse, 1993). In this line, Pardal et al. (2007) have reported that in mice exposed to 10% O_2 for 7 days, a subpopulation of type-II cells acts like stem-cells, differentiating into new type-I cells. This mechanism may explain why CB hyperplasia is induced by CSH (Pardal et al., 2007). Likewise, several evidences suggest an important role of type-II cells in CB adaptation to different paradigms of chronic hypoxia (see Leonard et al., 2018 for a recent revision on this subject). The progressively increases in ventilation elicited by CSH, it is known as ventilatory acclimatization (Bisgard, 2000). Ventilatory acclimatization induced by CSH depends on the potentiation of the CB chemosensory response to hypoxia in several species (Bouverot et al., 1973; Olson and Dempsey, 1978; Smith et al., 1986; Vizek et al., 1987; Di Giulio et al., 2003). In this section we summarized some evidences concerning possible mechanisms behind this phenomenon.

Increased Levels of NO and ET-1 in the CB

Chronic sustained hypoxia increases the levels of NO in the CB (Ye et al., 2002) due to an augmented expression of endothelial and inducible nitric oxide synthase (Ye et al., 2002). An increased NO levels evoked by CSH may change the oxidative state and/or produce protein nitrosylation in type-I cells (Monteiro et al., 2005). It is possible that increased levels of NO induced by CSH may contribute to enhances the oxygen sensitivity in the CB. Iturriaga et al. (2000) reported that during steady chemosensory excitation induced by hypoxia, bolus injections of a NO donor transiently reduced chemosensory discharges. However, during normoxia the same concentration of NO increased chemosensory discharge, suggesting a dual role of NO (see also Iturriaga, 2001). Endothelin-1 (ET-1) is also present in the CB. Chronic hypoxia for 2 weeks increases the ET-1 levels and the expression of the ETA receptor in rat type-I cells (Chen Y. et al., 2002). ET peptides have a proliferative effect in the CB stimulating cellular proliferation in CB primary cultures trough the ETA receptor activation (Paciga et al., 1999). In the rabbit CB superfused *in vitro*, Chen et al. (2000) found that ET-1 did not modify basal CB chemosensory discharge, but potentiates the response induced by hypoxia. In addition, Chen Y. et al. (2002) found that ET-1 increases Ca^{2+} acting on L-type Ca^{2+} channels in rabbit type-I cells. Although, ET may enhance the CB response to hypoxia in a CB preparation devoid of vascular effects, the excitatory effect of ET-1 on CB chemoreception is most likely mediated by its vasoconstrictor effect (Rey and Iturriaga, 2004). Chen et al. (2007) found that the enhanced basal and hypoxic evoked chemosensory activity following CSH was significantly reduced by concurrent treatment with the ETA receptors blocker bosentan. Thus, it is plausible that CSH-induced CB acclimatization would be partially mediated by ETA receptors.

Changes in Levels and Secretion of DA, ACh, ATP, or Their Receptors

Carotid body type-I cells contains high levels of dopamine (DA), thus tyrosine hydroxylase (TH) is a conventional marker for the identification of type-I cells (Chou et al., 1998). CSH enhances both, the expression and activity of TH in CBs (Hanbauer, 1977; Czyzyk-Krzeska et al., 1992); as a consequence, there is an increased DA content. It is worth to mention that even though DA is released in response to acute hypoxia, its role in CB neurotransmission remains controversial (Fidone et al., 1990; Alcayaga et al., 2006). In most species, the evidence strongly suggests an inhibitory role for DA on CB chemoreception. It has been proposed that DA is involved in an auto-regulatory mechanism mediated by D2 dopaminergic receptors expressed in type-I cells (Fidone et al., 1990). Some evidences suggesting that the inhibitory role of DA could be attenuated in CBs exposed to CSH, leading to the increased chemoreactivity (Tatsumi et al., 1995). However, this is in sharp contrast with the increased TH activity during chronic hypoxia. Zhang et al. (2017) proposed that adenosine and DA control CB chemosensory discharge acting on petrosal neurons via their opposing actions on the hyperpolarization-activated, cyclic nucleotide-gated (HCN) cation current (I_h). By using a functional *in vitro* preparation of rat type-I cells/petrosal neurons co-cultures, they found that adenosine enhanced I_h in petrosal neurons acting on A2a receptors, while DA had the opposite effect via D2 receptors. Indeed, adenosine modified the I_h activation curve and increased firing frequency, while DA caused a hyperpolarizing shift in the curve and decreased the firing frequency (Zhang et al., 2017).

Acetylcholine and ATP have been proposed as excitatory transmitters in the CB, acting on nicotinic and P2X receptors, respectively (Zhang et al., 2000; Varas et al., 2003; Meza et al., 2012). In response to CSH, an increase of nicotinic acetylcholine receptor (nAChR) expression in chemosensory afferents has been reported (He et al., 2005). Paradoxically, nAChRs blockers does not revert the CB hyperreactivity evoked by chronic hypoxia (Bisgard, 2000). Regarding to the role played by ATP, it seems that CSH has no effect on P2X receptors expression in both, type-I cells and postsynaptic terminals of petrosal neurons. However, administration of P2X receptor blockers reduced the enhanced chemosensory response to acute hypoxia induced by CSH (He et al., 2006). Therefore, reports suggest that ACh and ATP may play a role in the hyperreactivity of the CB induced by CSH. Hence, it seems that chronic hypoxia enhances both the excitatory (purinergic and cholinergic) and inhibitory (dopaminergic) chemoreception pathways, in agreement with the proposal stating that the balance between excitatory and inhibitory transmitters is more relevant during chronically hypoxic conditions rather than normoxia (Prabhakar, 2006).

Increase in the Proportion of Na⁺ Currents and a Decrease in K⁺ Currents

In type-I cells of neonatal rats subjected to long-lasting chronic hypoxia there is an increase in the proportion of Na⁺ currents and a decrease in oxygen-sensitive K⁺ current density (Hempleman, 1995). Likewise, Cáceres et al. (2007) found

that rat CB hyperreactivity induced by CSH correlated with an increased expression of voltage-gated Na⁺ channel Nav1.1 subunit. Carpenter et al. (1998) found that exposing adult rats to 10% O₂ for 3 weeks significantly reduced K⁺ current density with no change in voltage-gated K⁺ current amplitudes. However, rabbit CBs cultured in 5% O₂ for 2 days shows a decrease in expression of the Kv3.4 subunit of the voltage-gated K⁺ channel (Käab et al., 2005), suggesting a species-specific effect on Kv-conductance.

Additionally, neonatal rat type-I cells exposed to hypoxia (9–14 days, 10% O₂) shows a decreased charybdotoxin-sensitive K⁺ current, suggesting the downregulation of the maxi-K channel (Wyatt et al., 1995). It is worth to mention that in this paradigm, the rats have “blunted” rather hyperreactive responses to acute hypoxia (Wyatt et al., 1995). In culture, human maxi-K channels show an increased calcium sensitivity after 72 h of hypoxia. This is probably due to the reported threefold increase in maxi-K β-regulatory subunit expression induced by CSH (Hartness et al., 2003). However, cultured type-I cells from neonatal rat subjected to CSH, express normal levels of maxi-K and TASK-1 channels (Nurse and Fearon, 2002). Additionally, cultured CBs cells exposed to chronic hypoxia for 2 weeks showed an increase in Na⁺ currents, but no changes in Ca²⁺ currents were reported (Stea et al., 1995), findings confirmed later in a paradigm of adult rats exposed to CSH (Peers et al., 1996).

We found an enhanced inhibition of the TASK-like current in neonatal rat type-I cells in cultured exposed to hypoxia for 2 days (Ortiz et al., 2007). While acute hypoxia (1% O₂) evokes a ~70% inhibition of the TASK-like current in control conditions, after 48 h of hypoxia current inhibition reaches ~90%. Importantly, these results were later reproduced (Ortiz et al., 2013) in an animal model for chronic *intermittent* hypoxia suggesting a common preserved mechanism of CB adaptation to hypoxic environments.

The evidence suggests that regulation of the expression and/or function of ionic currents are important adaptive mechanisms for the CB subjected to chronic hypoxia. Nevertheless, with the exception of the work of Cáceres et al. (2007), there is no physiological evidence that the CSH protocols used in above mentioned studies (Hempleman, 1995; Stea et al., 1995; Carpenter et al., 1998) really potentiated the CB chemosensory responses to acute hypoxia. On the contrary, most of them used long-term CSH exposition (>2 weeks) which is associated with an attenuation rather than a potentiation of CB chemosensory responses to acute hypoxia (Tatsumi et al., 1991). Hence, no convincing conclusion can be draw.

Increased Activity of AMP-Dependent Kinase and Protein Kinase C (PKC)

The activities of maxi-K and TASK channels are modulated by cell metabolism. Wyatt et al. (2007) found that AMP-dependent kinase (AMPK) is colocalized with both maxi-K and TASK channels in rat type-I cells. They found that AMPK activation inhibits those K⁺ currents triggering membrane depolarization, whereas AMPK-antagonist application prevents the depolarization induced by hypoxia (Wyatt et al., 2007).

TASK channel current is activated by intracellular nucleotides, such as ATP (Varas et al., 2007). At physiological concentrations (1–3 mM) ATP strongly promotes channel opening. Accordingly, in absence of ATP, TASK channel activity decreases in ~50 % in inside-out patches (Williams and Buckler, 2004; Varas et al., 2007). Thus, ATP production seems to be central in regulating the activity of TASK-like and other potassium channels such as maxi-K (Williams and Buckler, 2004; Varas et al., 2007; Holmes et al., 2018). Even though some reports postulate a null or small effect of ATP on these conductances (Xu et al., 2005) one of the earliest proposition, the “metabolic hypothesis”, states that CB oxygen sensing is mediated by oxidative phosphorylation, supported by the fact that virtually all metabolic poisons or oxidative phosphorylation inhibitors induce CB type-I cell excitation (Mulligan and Lahiri, 1982; Buerk et al., 1994; Wilson et al., 1994; Buckler and Turner, 2013). This response is normally triggered by potassium conductance(s) inhibition leading to the depolarization of the type-I cell membrane, reinforcing the notion that low PO₂ trigger potassium current blockage (Figure 1, see also Holmes et al., 2018). In this regard, it has been postulated that CB-mitochondria express a particular subtype of *cytochrome a3* – the functional core of mitochondrion complex IV – which is at least seven times more sensitive to a drop in the PO₂, further supporting a pivotal role of mitochondrial metabolism on CB-type-I cell excitation (for a full revision on this matter see Holmes et al., 2018 in this same research topic).

Therefore, in addition to changes in the expression and intrinsic function of ion channels, the fact that cellular metabolism is modified by CSH provides a potential link between CSH and the type-I cell membrane excitability. It is known that CSH increases the cytosolic AMP/ATP ratio in type-I cells. Thus, several enzymatic systems can be activated, including AMPK (Laderoute et al., 2006). Likewise, CSH induces an increase in cAMP levels in CB cells (Nurse et al., 1994). Protein kinase C (PKC) activity is enhanced by chronic hypoxia in several tissues (Zhang et al., 2004; Neckar et al., 2005), whereas the protein kinase A (PKA) activity decreases (Kobayashi et al., 1998). Increased PKC activity in response to acute hypoxia (Summers et al., 2000) was associated to the regulation of L-type Ca²⁺, maxi-K channels and background K⁺ channels in type-I cells (Peers and Carpenter, 1998; Summers et al., 2000; Zhang et al., 2003). Since PKC inhibits TASK channel activity, whereas PKA enhanced it (Zilberberg et al., 2000; Besana et al., 2004) these pathways are well-suited as potential effectors of CSH induced-changes.

Activation of Hypoxia-Inducible Transcription Factors (HIFs)

The cellular long-term response to hypoxia is driven by changes in the expression of several genes to cope with the new hypoxic environment. These cellular transcriptional responses are largely dependent on the activation of the so-called hypoxia-inducible transcription factors (Shimoda and Semenza, 2011). HIFs are composed by a HIF α subunit (mainly 1 α and 2 α), regulated by hypoxia and a constitutive β subunit (HIF-1 β). Under normoxic conditions, HIF α is continuously degraded

through hydroxylation modification catalyzed by a set of oxygen sensitive prolyl-hydroxylases enzymes. During hypoxia, the activity of prolyl-hydroxylases are dramatically reduced, allowing the translocation of HIF α to the cell nuclei where it forms an heterodimer with HIF-1 β . Following translocation and dimerization, the HIF-1 α /HIF-1 β complex bind to hypoxia response elements (HRE) present in the DNA to regulate gene expression (Shimoda and Semenza, 2011). In the CB, the contribution of HIF-1 α and HIF-2 α has been largely studied in the setting of intermittent hypoxia; much less it is known in the context of CSH. Nevertheless, it has been shown that activation of the HIF-1 α pathway promotes the up-regulation of pro-oxidant enzymes (i.e., NADPH) in the CB. On the contrary, intermittent hypoxia reduced the expression of HIF-2 α within the CB, resulting in a significant down-regulation of antioxidant enzymes (i.e., SOD) levels being the outcome an increase ROS formation within CB glomus cells, increasing both intracellular Ca²⁺ and neurotransmitter release (Prabhakar and Semenza, 2016). The specific contribution of HIF-1 α and HIF-2 α pathways on the sensitization of the CB to hypoxia under CSH remains to be fully determined. Nevertheless, it is possible to speculate that activation of HIF-1 α and inhibition of HIF-2 α pathways may occur in the CBs of animals exposed to sustained hypoxia.

We have here discussed some of the mechanisms explaining CB chemosensory hyperreactivity under CSH. This exacerbated activity has systemic consequences inducing the adaptation to CSH. In this regard, a central aspect is the CB-mediated autonomic function and even though the neural mechanism associated to autonomic responses to CSH are not completely elucidated, it has been described that CB type-I cells could be involved. During acute hypoxic exposure, excitatory inputs from the CB activate neural pathways in the brainstem increasing minute ventilation and sympathetic outflow (Barros et al., 2002; Braga et al., 2007). However, during CSH the baseline minute ventilation increases in a time-dependent manner (ventilatory acclimatization to hypoxia), probably related to the changes in oxygen sensitivity of the CB as well as the neural network in the brainstem related to sympathetic response to hypoxia (Powell et al., 2000; Powell, 2007). It has been proposed that hypoxia-mediated sympathetic activation act as a compensatory mechanism to increase oxygen supply to critical organs through regulating cardiac output and vascular conductance (Leuenberger et al., 1991; Calbet, 2003). Indeed, exposure of healthy humans to CSH produces a significant increase in circulating norepinephrine levels reflecting increases in sympathetic outflow during hypoxic exposure (Calbet, 2003; Hansen and Sander, 2003). In addition, it has been shown that rats subjected to 24 h of FiO₂ 10% displayed an increase of pre-sympathetic neurons activation located in the rostral aspect of the ventral medullary surface (RVLM). Furthermore, these set of neurons showed a further increase activation during the late expiratory phase during hypoxia stimulation suggesting a potential role on regulating the neural circuitry related to the respiratory-sympathetic coupling observed during CSH (Moraes et al., 2014). Interestingly, previous studies from our group showed that chronic activation of the CB afferent input to the brainstem using episodic hypoxic stimulation induces the activation of RVLM neurons (Del Rio et al., 2016). Therefore,

it is plausible to hypothesize that during CSH, activation of the CB may contribute to the regulation of sympathetic drive. This hypothesis remains to be determined.

CONCLUSION

Potential of CB chemosensory responses is an absolute requirement for ventilatory acclimatization. How the increased response to acute hypoxia in chronically hypoxic CBs is achieved? There are probably multiple mechanisms underlying such phenomena. In the present review we focused on evidences suggesting that type-I cells excitability and metabolism undergoes substantial changes when exposed to CSH. Interestingly, ion channels, including some involved in the generation of the chemosensory response to acute hypoxia (i.e., maxi-K and TASK channels) are tightly regulated by PKC, PKA, AMPK, and ATP. Therefore, it is possible that throughout changes in cell metabolism, in addition to modifications on ion channels expression, CSH generates changes in type-I cells excitability, ultimate leading to increased depolarization during acute

hypoxia. Importantly, increased inhibition of TASK-like currents under chronic – sustained and intermittent – hypoxia provide a novel potential mechanism to explain the CB hyperreactivity (Ortiz et al., 2007, 2013). It seems clear that there is no a single mechanism as full explanation for CB hyperreactivity during CSH. Combined studies covering from cellular/molecular studies to functional approaches will help to understand the physiology of ventilatory acclimatization during chronic sustained hypoxia.

AUTHOR CONTRIBUTIONS

All authors contributed to writing the article. RV, RI, RDR, and FO designed and edited the paper.

FUNDING

This work was supported by grants FONDECYT 11160616 to FO, FONDECYT 1180172 to RDR, and FONDECYT 1150040 to RI.

REFERENCES

- Alcayaga, J., Soto, C. R., Vargas, R. V., Ortiz, F. C., Arroyo, J., and Iturriaga, R. (2006). Carotid body transmitters actions on rabbit petrosal ganglion in vitro. *Adv. Exp. Med. Biol.* 580, 331–337. doi: 10.1007/0-387-31311-7_51
- Barros, R. C., Bonagamba, L. G., Okamoto-Canesin, R., de Oliveira, M., Branco, L. G., and Machado, B. H. (2002). Cardiovascular responses to chemoreflex activation with potassium cyanide or hypoxic hypoxia in awake rats. *Auton. Neurosci.* 97, 110–115. doi: 10.1016/S1566-0702(02)00050-4
- Besana, A., Barbuti, A., Tateyama, M. A., Symes, A. J., Robinson, R. B., and Feinmark, S. J. (2004). Activation of protein kinase C epsilon inhibits the two-pore domain K⁺ channel, TASK-1, inducing repolarization abnormalities in cardiac ventricular myocytes. *J. Biol. Chem.* 279, 33154–33160. doi: 10.1074/jbc.M403525200
- Bigard, G. E. (2000). Carotid body mechanisms in acclimatization to hypoxia. *Respir. Physiol.* 121, 237–246. doi: 10.1016/S0034-5687(00)00131-6
- Bouverot, P., Candas, V., and Libert, J. P. (1973). Role of the arterial chemoreceptors in ventilatory adaptation to hypoxia of awake dogs and rabbits. *Respir. Physiol.* 17, 209–219. doi: 10.1016/0034-5687(73)90062-5
- Braga, V. A., Soriano, R. N., Bracciali, A. L., de Paula, P. M., Bonagamba, L. G., Paton, J. F., et al. (2007). Involvement of L-glutamate and ATP in the neurotransmission of the sympathoexcitatory component of the chemoreflex in the commissural nucleus tractus solitarii of awake rats and in the working heart-brainstem preparation. *J. Physiol.* 581, 1129–1145. doi: 10.1113/jphysiol.2007.129031
- Buckler, K. (1997). A novel oxygen-sensitive potassium current in rat carotid body type I cells. *J. Physiol.* 498, 649–662. doi: 10.1113/jphysiol.1997.sp021890
- Buckler, K., Williams, B., and Honoré, E. (2000). An oxygen, acid and anaesthetic sensitive TASK-like background potassium channel in rat arterial chemoreceptor cells. *J. Physiol.* 525, 135–142. doi: 10.1111/j.1469-7793.2000.00135.x
- Buckler, K. J., and Turner, P. J. (2013). Oxygen sensitivity of mitochondrial function in rat arterial chemoreceptor cells. *J. Physiol.* 591, 3549–3563. doi: 10.1113/jphysiol.2013.257741
- Buckler, K. J., and Vaughan-Jones, R. D. (1994). Effects of hypoxia on membrane potential and intracellular calcium in rat neonatal carotid body type I cells. *J. Physiol.* 476, 423–428. doi: 10.1113/jphysiol.1994.sp020143
- Buerk, D. G., Iturriaga, R., and Lahiri, S. (1994). Testing the metabolic hypothesis of O₂ chemoreception in the cat carotid body in vitro. *J. Appl. Physiol.* 76, 1317–1323. doi: 10.1152/jappl.1994.76.3.1317
- Cáceres, A. I., Obeso, A., Gonzalez, C., and Rocher, A. (2007). Molecular identification and functional role of voltage-gated sodium channels in rat carotid body chemoreceptor cells. Regulation of expression by chronic hypoxia in vivo. *J. Neurochem.* 102, 231–245. doi: 10.1111/j.1471-4159.2007.04465.x
- Calbet, J. A. L. (2003). Chronic hypoxia increases blood pressure and noradrenaline spillover in healthy humans. *J. Physiol.* 551, 379–386. doi: 10.1113/jphysiol.2003.045112
- Carpenter, E., Bee, D., and Peers, C. (1998). Ionic currents in carotid body type I cells isolated from normoxic and chronically hypoxic adult rats. *Brain Res.* 811, 79–87. doi: 10.1016/S0006-8993(98)00962-7
- Carpenter, E., and Peers, C. (1997). Swelling- and cAMP-activated Cl⁻ currents in isolated rat carotid body type I cells. *J. Physiol.* 503, 497–511. doi: 10.1111/j.1469-7793.1997.497bg.x
- Chen, J., He, L., Dinger, B., and Fidone, S. (2000). Pharmacological effects of endothelin in rat carotid body. Activation of second messenger pathways and potentiation of chemoreceptor activity. *Adv. Exp. Med. Biol.* 475, 517–525.
- Chen, Y., Tipoe, G. L., Liong, E., Leung, S., Lam, S. Y., Iwase, R., et al. (2002). Chronic hypoxia enhances endothelin-1-induced intracellular calcium elevation in rat carotid body chemoreceptors and up-regulates ETA receptor expression. *Pflugers Arch.* 443, 565–573. doi: 10.1007/s00424-001-0728-2
- Chen, J., He, L., Liu, X., Dinger, B., Stensaas, L., and Fidone, S. (2007). Effect of the endothelin receptor antagonist bosentan on chronic hypoxia-induced morphological and physiological changes in rat carotid body. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 292, L1257–L1262. doi: 10.1152/ajplung.00419.2006
- Chou, C. L., Sham, J. S., Schofield, B., and Shirahata, M. (1998). Electrophysiological and immunocytological demonstration of cell-type specific responses to hypoxia in the adult cat carotid body. *Brain Res.* 789, 229–238. doi: 10.1016/S0006-8993(97)01472-8
- Czyzyk-Krzeska, M. F., Bayliss, D. A., Lawson, E. E., and Millhorn, D. E. (1992). Regulation of tyrosine hydroxylase gene expression in the rat carotid body by hypoxia. *J. Neurochem.* 58, 1538–1546. doi: 10.1111/j.1471-4159.1992.tb11376.x
- Del Rio, R., Andrade, D. C., Lucero, C., Arias, P., and Iturriaga, R. (2016). Carotid body ablation abrogates hypertension and autonomic alterations induced by intermittent hypoxia in rats. *Hypertension* 68, 436–445. doi: 10.1161/HYPERTENSIONAHA.116.07255
- Di Giulio, C., Huang, W. X., Mokashi, A., Roy, A., Cacchio, M., Macrí, M. A., et al. (2003). Sustained hypoxia promotes hyperactive response of carotid body in the cat. *Respir. Physiol. Neurobiol.* 134, 69–74. doi: 10.1016/S1569-9048(02)00203-3

- Duchen, M. R., Caddy, K. W., Kirby, G. C., Patterson, D. L., Ponte, J., and Biscoe, T. J. (1988). Biophysical studies of the cellular elements of the rabbit carotid body. *Neuroscience* 26, 291–313. doi: 10.1016/0306-4522(88)90146-7
- Fidone, S. J., Gonzalez, C., Obeso, A., Gomez-Nino, A., and Dinger, B. (1990). “Biogenic amine and neuropeptide transmitters in carotid body chemotransmission: experimental findings and perspectives,” in *Hypoxia: the Adaptations*, ed. J. E. Remmers (Toronto: B.C. Decker Inc), 116–126.
- Fieber, L. A., and McCleskey, E. W. (1993). L-type calcium channels in type I cells of the rat carotid body. *J. Neurophysiol.* 70, 1378–1384. doi: 10.1152/jn.1993.70.4.1378
- Ganfornina, M. D., and Lopez-Barneo, J. (1992). Gating of O₂-sensitive K⁺ channels of arterial chemoreceptor cells and kinetic modifications induced by low PO₂. *J. Gen. Physiol.* 100, 427–455. doi: 10.1085/jgp.100.3.427
- Gonzalez, C., Almaraz, L., Obeso, A., and Rigual, R. (1994). Carotid body chemoreceptors: from natural stimuli to sensory discharges. *Physiol. Rev.* 74, 829–898. doi: 10.1152/physrev.1994.74.4.829
- Hanbauer, I. (1977). Regulation of tyrosine hydroxylase in carotid body. *Adv. Biochem. Psychopharmacol.* 16, 275–280.
- Hansen, J., and Sander, M. (2003). Sympathetic neural overactivity in healthy humans after prolonged exposure to hypobaric hypoxia. *J. Physiol.* 546, 921–929. doi: 10.1113/jphysiol.2002.031765
- Hartness, M. E., Brazier, S. P., Peers, C., Bateson, A. N., Ashford, M. L., and Kemp, P. J. (2003). Post-transcriptional control of human maxiK potassium channel activity and acute oxygen sensitivity by chronic hypoxia. *J. Biol. Chem.* 278, 51422–51432. doi: 10.1074/jbc.M309463200
- He, L., Chen, J., Dinger, B., Stensaas, L., and Fidone, S. (2006). Effect of chronic hypoxia on purinergic synaptic transmission in rat carotid body. *J. Appl. Physiol.* 100, 157–162. doi: 10.1152/jappphysiol.00859.2005
- He, L., Dinger, B., and Fidone, S. (2005). Effect of chronic hypoxia on cholinergic chemotransmission in rat carotid body. *J. Appl. Physiol.* 98, 614–619. doi: 10.1152/jappphysiol.00714.2004
- Heath, D., Smith, P., and Fitch, R. (1985). Comparative pathology of the enlarged carotid body. *J. Comp. Pathol.* 95, 259–271. doi: 10.1016/0021-9975(85)90012-X
- Hempleman, S. C. (1995). Sodium and potassium current in neonatal rat carotid body cells following chronic in vivo hypoxia. *Brain Res.* 699, 42–50. doi: 10.1016/0006-8993(95)00850-P
- Holmes, A. P., Ray, C. J., Coney, A. M., and Kumar, P. (2018). Is carotid body physiological O₂ sensitivity determined by a unique mitochondrial phenotype? *Front. Physiol.* 9:562. doi: 10.3389/fphys.2018.00562
- Iturriaga, R. (2001). Nitric oxide and carotid chemoreception. *Biol. Res.* 34, 135–139. doi: 10.4067/S0716-97602001000200019
- Iturriaga, R., and Alcayaga, J. (2004). Neurotransmission in the carotid body: transmitters and modulators between glomus cells and petrosal ganglion nerve terminals. *Brain Res. Rev.* 47, 46–53. doi: 10.1016/j.brainresrev.2004.05.007
- Iturriaga, R., Mokashi, A., and Lahiri, S. (1998). Anion exchanger and chloride channel in cat carotid body chemotransduction. *J. Auton. Nerv. Syst.* 70, 23–31. doi: 10.1016/S0165-1838(98)00019-8
- Iturriaga, R., Villanueva, S., and Mosqueira, M. (2000). Dual effects of nitric oxide on cat carotid body chemoreception. *J. Appl. Physiol.* 89, 1005–1012. doi: 10.1152/jappl.2000.89.3.1005
- Kääb, S., Miguel-Velado, E., López-López, J. R., and Pérez-García, M. T. (2005). Down regulation of kv3.4 channels by chronic hypoxia increases acute oxygen sensitivity in rabbit carotid body. *J. Physiol.* 566(Pt 2), 395–408. doi: 10.1113/jphysiol.2005.085837
- Kobayashi, S., Beitner-Johnson, D., Conforti, L., and Millhorn, D. E. (1998). Chronic hypoxia reduces adenosine A_{2A} receptor-mediated inhibition of calcium current in rat PC12 cells via downregulation of protein kinase A. *J. Physiol.* 512, 351–363. doi: 10.1111/j.1469-7793.1998.351be.x
- Laderoute, K. R., Amin, K., Calaoagan, J. M., Knapp, M., Le, T., Orduna, J., et al. (2006). 5'-AMP-activated protein kinase (AMPK) is induced by low-oxygen and glucose deprivation conditions found in solid-tumor microenvironments. *Mol. Cell. Biol.* 26, 5336–5347. doi: 10.1128/MCB.00166-06
- Leonard, E. M., Salman, S., and Nurse, C. A. (2018). Sensory processing and integration at the carotid body tripartite synapse: neurotransmitter functions and effects of chronic hypoxia. *Front. Physiol.* 9:225. doi: 10.3389/fphys.2018.00225
- Leuenberger, U., Gleeson, K., Wroblewski, K., Prophet, S., Zelis, R., Zwillich, C., et al. (1991). Norepinephrine clearance is increased during acute hypoxemia in humans. *Am. J. Physiol.* 261, H1659–H1664.
- López-Barneo, J., López-López, J. R., Ureña, J., and González, C. (1988). Chemotransduction in the carotid body: current modulated by PO₂ in type I chemoreceptor cells. *Science* 241, 580–582. doi: 10.1126/science.2456613
- López-López, J. R., and González, C. (1992). Time course of K⁺ current inhibition by low oxygen in chemoreceptor cells of adult rabbit carotid body. Effects of carbon monoxide. *FEBS Lett.* 299, 251–254. doi: 10.1016/0014-5793(92)80126-2
- Meza, R. C., Ortiz, F. C., Bravo, E., Iturriaga-Vásquez, P., Eugenin, J. L., and Varas, R. (2012). Functional expression of the $\alpha 7$ and $\alpha 4$ -containing nicotinic acetylcholine receptors on the neonatal rat carotid body. *Neurochem. Int.* 60, 115–124. doi: 10.1016/j.neuint.2011.11.011
- Mills, L., and Nurse, C. (1993). Chronic hypoxia in vitro increases volume of dissociated carotid body chemoreceptors. *Neuroreport* 4, 619–622. doi: 10.1097/00001756-199306000-00004
- Monteiro, H. P., Rocha Oliveira, C. J., Curcio, M. F., Moraes, M. S., and Arai, R. J. (2005). Tyrosine phosphorylation in nitric oxide-mediated signaling events. *Methods Enzymol.* 396, 350–358. doi: 10.1016/S0076-6879(05)96029-1
- Moraes, D. J. A., Bonagamba, L. G. H., Costa, K. M., Costa-Silva, J. H., Zoccal, D. B., and Machado, B. H. (2014). Short-term sustained hypoxia induces changes in the coupling of sympathetic and respiratory activities in rats. *J. Physiol.* 592, 2013–2033. doi: 10.1113/jphysiol.2013.262212
- Mulligan, E., and Lahiri, S. (1982). Separation of carotid body chemoreceptor responses to O₂ and CO₂ by oligomycin and by antimycin A. *Am. J. Physiol.* 242, C200–C206. doi: 10.1152/ajpcell.1982.242.3.C200
- Neckar, J., Markova, I., Novak, F., Novakova, O., Szarszoi, O., Ost'adal, B., et al. (2005). Increased expression and altered subcellular distribution of PKC- δ in chronically hypoxic rat myocardium: involvement in cardioprotection. *Am. J. Physiol. Heart Circ. Physiol.* 288, H1566–H1572.
- Nurse, C. A. (2005). Neurotransmission and neuromodulation in the chemosensory carotid body. *Auton. Neurosci.* 120, 1–9. doi: 10.1016/j.autneu.2005.04.008
- Nurse, C. A., and Fearon, I. M. (2002). Carotid body chemoreceptors in dissociated cell culture. *Microsc. Res. Tech.* 59, 249–255. doi: 10.1002/jemt.10199
- Nurse, C. A., Jackson, A., and Stea, A. (1994). Plasticity in cultured arterial chemoreceptors: effects of chronic hypoxia and cyclic AMP analogs. *Adv. Exp. Med. Biol.* 360, 167–170. doi: 10.1007/978-1-4615-2572-1_20
- Olson, E. B. J. R., and Dempsey, J. A. (1978). Rat as a model for human like ventilatory adaptation to chronic hypoxia. *J. Appl. Physiol.* 44, 763–769. doi: 10.1152/jappl.1978.44.5.763
- Ortega-Sáenz, P., Pascual, A., Gómez-Díaz, R., and López-Barneo, J. (2006). Acute oxygen sensing in heme oxygenase-2 null mice. *J. Gen. Physiol.* 128, 405–411. doi: 10.1085/jgp.200609591
- Ortiz, F., Varas, R., and Iturriaga, R. (2007). Chronic hypoxia enhanced the TASK-like current inhibition evoked by acute hypoxia. *Biol. Res.* 40:135.
- Ortiz, F. C., Del Rio, R., Ebensperger, G., Reyes, V. R., Alcayaga, J., Varas, R., et al. (2013). Inhibition of rat carotid body glomus cells TASK-like channels by acute hypoxia is enhanced by chronic intermittent hypoxia. *Respir. Physiol. Neurobiol.* 185, 600–607. doi: 10.1016/j.resp.2012.11.015
- Overholt, J. L., and Prabhakar, N. R. (1997). Ca²⁺ current in rabbit carotid body glomus cells is conducted by multiple types of high-voltage-activated Ca²⁺ channels. *J. Neurophysiol.* 78, 2467–2474. doi: 10.1152/jn.1997.78.5.2467
- Overholt, J. L., and Prabhakar, N. R. (1999). Norepinephrine inhibits a toxin resistant Ca²⁺ current in carotid body glomus cells: evidence for a direct G protein mechanism. *J. Neurophysiol.* 81, 225–233. doi: 10.1152/jn.1999.81.1.225
- Paciga, M., Vollmer, C., and Nurse, C. (1999). Role of ET-1 in hypoxia-induced mitosis of cultured rat carotid body chemoreceptors. *Neuroreport* 10, 3739–3744. doi: 10.1097/00001756-199912160-00003
- Pardal, R., Ortega-sáenz, P., Durán, R., and López-barneo, J. (2007). Glia-like stem cells sustain physiologic neurogenesis in the adult mammalian carotid body. *Cell* 131, 234–236. doi: 10.1016/j.cell.2007.07.043

- Peers, C. (1990). Hypoxic suppression of K^+ currents in type I carotid body cells: selective effect on the Ca^{2+} -activated K^+ current. *Neurosci. Lett.* 119, 253–256. doi: 10.1016/0304-3940(90)90846-2
- Peers, C., and Carpenter, E. (1998). Inhibition of Ca^{2+} -dependent K^+ channels in rat carotid body type I cells by protein kinase C. *J. Physiol.* 512, 743–750. doi: 10.1111/j.1469-7793.1998.743bd.x
- Peers, C., Carpenter, E., Hatton, C. J., Wyatt, C. N., and Bee, D. (1996). Ca^{2+} channel currents in type I carotid body cells of normoxic and chronically hypoxic neonatal rats. *Brain Res.* 739, 251–257. doi: 10.1016/S0006-8993(96)00832-3
- Pequignot, J. M., Hellstrom, S., and Johannsson, H. (1984). Intact and sympathectomized carotid bodies of long-term hypoxic rats: a morphometric ultrastructural study. *J. Neurocytol.* 13, 481–493. doi: 10.1007/BF01148336
- Powell, F. L. (2007). The influence of chronic hypoxia upon chemoreception. *Respir. Physiol. Neurobiol.* 157, 154–161. doi: 10.1016/j.resp.2007.01.009
- Powell, F. L., Huey, K. A., and Dwinell, M. R. (2000). Central nervous system mechanisms of ventilatory acclimatization to hypoxia. *Respir. Physiol.* 121, 223–236. doi: 10.1016/S0034-5687(00)00130-4
- Prabhakar, N. R. (2006). O₂ sensing at the mammalian carotid body: why multiple O₂ sensors and multiple transmitters? *Exp. Physiol.* 91, 17–23.
- Prabhakar, N. R., and Semenza, G. L. (2016). Regulation of carotid body oxygen sensing by hypoxia-inducible factors. *Pflugers Arch.* 468, 71–75. doi: 10.1007/s00424-015-1719-z
- Rey, S., and Iturriaga, R. (2004). Endothelins and nitric oxide: vasoactive modulators of carotid body chemoreception. *Curr. Neurovasc. Res.* 1, 465–473. doi: 10.2174/1567202043361857
- Shimoda, L. A., and Semenza, G. L. (2011). HIF and the lung: role of hypoxia-inducible factors in pulmonary development and disease. *Am. J. Respir. Crit. Care Med.* 183, 152–156. doi: 10.1164/rccm.201109-1393PP
- Smith, C. A., Bisgard, G. E., Nielsen, A. M., Daristotle, L., Kressin, N. A., Forster, H. V., et al. (1986). Carotid bodies are required for ventilatory acclimatization to chronic hypoxia. *J. Appl. Physiol.* 60, 1003–1010. doi: 10.1152/jappl.1986.60.3.1003
- Summers, B. A., Overholt, J. L., and Prabhakar, N. R. (2000). Augmentation of calcium current by hypoxia in carotid body glomus cells. *Adv. Exp. Med. Biol.* 475, 589–599. doi: 10.1007/0-306-46825-5_57
- Stea, A., Jackson, A., Macintyre, L., and Nurse, C. A. (1995). Long-term modulation of inward currents in O₂ chemoreceptors by chronic hypoxia and cyclic AMP in vitro. *J. Neurosci.* 15, 2192–2202. doi: 10.1523/JNEUROSCI.15-03-02192.1995
- Tatsumi, K., Pickett, C. K., and Weil, J. V. (1991). Attenuated carotid body hypoxic sensitivity after prolonged hypoxic exposure. *J. Appl. Physiol.* 70, 748–755. doi: 10.1152/jappl.1991.70.2.748
- Tatsumi, K., Pickett, C. K., and Weil, J. V. (1995). Possible role of dopamine in ventilatory acclimatization to high altitude. *Respir. Physiol.* 99, 63–73. doi: 10.1016/0034-5687(94)00084-D
- Varas, R., Alcayaga, J., and Iturriaga, R. (2003). ACh and ATP mediate excitatory transmission in cat carotid identified chemoreceptor units in vitro. *Brain Res.* 988, 154–163. doi: 10.1016/S0006-8993(03)03366-3
- Varas, R., Wyatt, C. N., and Buckler, K. J. (2007). Modulation of TASK-like background potassium channels in rat arterial chemoreceptor cells by intracellular ATP and other nucleotides. *J. Physiol.* 583, 521–536. doi: 10.1111/jphysiol.2007.135657
- Vizek, M., Pickett, C. K., and Weil, J. V. (1987). Increased carotid body hypoxic sensitivity during acclimatization to hypobaric hypoxia. *J. Appl. Physiol.* 63, 2403–2410. doi: 10.1152/jappl.1987.63.6.2403
- Williams, B. A., and Buckler, K. J. (2004). Biophysical properties and metabolic regulation of a TASK-like potassium channel in rat carotid body type 1 cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 286, L221–L230. doi: 10.1152/ajplung.00010.2003
- Williams, S. E., Wootton, P., Mason, H. S., Bould, J., Iles, D. E., Riccardi, D., et al. (2004). Hemoxigenase-2 is an oxygen sensor for a calcium-sensitive potassium channel. *Science* 306, 2093–2097. doi: 10.1126/science.1105010
- Wilson, D. F., Mokashi, A., Chugh, D., Vinogradov, S., Osanai, S., and Lahiri, S. (1994). The primary oxygen sensor of the cat carotid body is cytochrome c3 of the mitochondrial respiratory chain. *FEBS Lett.* 351, 370–374. doi: 10.1016/0014-5793(94)00887-6
- Wyatt, C. N., Mustard, K. J., Pearson, S. A., Dallas, M. L., Atkinson, L., Kumar, P., et al. (2007). AMP-activated protein kinase mediates carotid body excitation by hypoxia. *J. Biol. Chem.* 282, 8092–8098. doi: 10.1074/jbc.M608742200
- Wyatt, C. N., and Peers, C. (1995). Ca^{2+} -activated K^+ channels in isolated type I cells of the neonatal rat carotid body. *J. Physiol.* 483, 559–565. doi: 10.1113/jphysiol.1995.sp020606
- Wyatt, C. N., Weir, E. K., and Peers, C. (1994). Diphenylene iodonium blocks K^+ and Ca^{2+} currents in type I cells isolated from the neonatal rat carotid body. *Neurosci. Lett.* 172, 63–66. doi: 10.1016/0304-3940(94)90663-7
- Wyatt, C. N., Wright, C., Bee, D., and Peers, C. (1995). O₂-sensitive K^+ currents in carotid body chemoreceptor cells from normoxic and chronically hypoxic rats and their roles in hypoxic chemotransduction. *Proc. Natl. Acad. Sci. U.S.A.* 92, 295–299. doi: 10.1073/pnas.92.1.295
- Xu, J., Xu, F., Tse, F., and Tse, A. (2005). ATP inhibits the hypoxia response in type I cells of rat carotid bodies. *J. Neurochem.* 92, 1419–1430. doi: 10.1111/j.1471-4159.2004.02978.x
- Yamamoto, Y., Kummer, W., Atoji, Y., and Suzuki, Y. (2002). TASK-1, TASK-2, TASK-3 and TRAAK immunoreactivities in the rat carotid body. *Brain Res.* 950, 304–307. doi: 10.1016/S0006-8993(02)03181-5
- Yamamoto, Y., and Taniguchi, K. (2006). Expression of tandem P domain K^+ channel, TREK-1, in the rat carotid body. *J. Histochem. Cytochem.* 54, 467–472. doi: 10.1369/jhc.5A6755.2005
- Ye, J. S., Tipoe, G. L., Fung, P. C., and Fung, M. L. (2002). Augmentation of hypoxia-induced nitric oxide generation in the rat carotid body adapted to chronic hypoxia: an involvement of constitutive and inducible nitric oxide synthases. *Pflugers Arch.* 444, 178–185. doi: 10.1007/s00424-002-0785-1
- Zhang, M., Fearon, I. M., Zhong, H., and Nurse, C. A. (2003). Presynaptic modulation of rat arterial chemoreceptor function by 5-HT: role of K^+ channel inhibition via protein kinase C. *J. Physiol.* 551, 825–842. doi: 10.1113/jphysiol.2002.038489
- Zhang, M., Vollmer, C., and Nurse, C. A. (2017). Adenosine and dopamine oppositely modulate a hyperpolarization-activated current I_h in chemosensory neurons of the rat carotid body in co-culture. *J. Physiol.* doi: 10.1113/JP274743 [Epub ahead of print].
- Zhang, M., Zhong, H., Vollmer, C., and Nurse, C. A. (2000). Co-release of ATP and ACh mediates hypoxic signaling at rat carotid body chemoreceptors. *J. Physiol.* 1, 143–158. doi: 10.1111/j.1469-7793.2000.t01-1-00143.x
- Zhang, Y. C., Ni, W., Zhang, Z. K., and Xu, Y. J. (2004). The effect of protein kinase C on voltage-gated potassium channel in pulmonary artery smooth muscle cells from rats exposed to chronic hypoxia. *Chin. Med. J.* 117, 19–23.
- Zilberberg, N., Ilan, N., Gonzalez-Colaso, R., and Goldstein, S. A. (2000). Opening and closing of KCNK0 potassium leak channels is tightly regulated. *J. Gen. Physiol.* 116, 721–734. doi: 10.1085/jgp.116.5.721

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Pulgar-Sepúlveda, Varas, Iturriaga, Del Rio and Ortiz. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.