



# High-Fat Diet-Induced Obesity Model Does Not Promote Endothelial Dysfunction via Increasing Leptin/Akt/eNOS Signaling

Vanessa da Silva Rocha<sup>1</sup>, Erick Roberto Gonçalves Claudio<sup>2</sup>, Vitor Loureiro da Silva<sup>1</sup>, Jóctan Pimentel Cordeiro<sup>1</sup>, Lucas Furtado Domingos<sup>1</sup>, Márcia Regina Holanda da Cunha<sup>1</sup>, Helder Mauad<sup>2</sup>, Thiago Bruder do Nascimento<sup>3</sup>, Ana Paula Lima-Leopoldo<sup>1</sup> and André Soares Leopoldo<sup>1\*</sup>

<sup>1</sup> Physiology and Biochemistry Laboratory, Department of Sports, Center of Physical Education and Sports, Federal University of Espírito Santo, Vitória, Brazil, <sup>2</sup> Department of Physiological Sciences, Health Sciences Center, Federal University of Espírito Santo, Vitória, Brazil, <sup>3</sup> Vascular Biology Center, Medical College of Georgia, Augusta University, Augusta, GA, United States

## OPEN ACCESS

### Edited by:

Tommaso Angelone,  
University of Calabria, Italy

### Reviewed by:

Antonio Colantuoni,  
University of Naples Federico II, Italy  
Antonella Amato,  
Università degli Studi di Palermo, Italy

### \*Correspondence:

André Soares Leopoldo  
andre.leopoldo@ufes.br

### Specialty section:

This article was submitted to  
Vascular Physiology,  
a section of the journal  
Frontiers in Physiology

**Received:** 12 October 2018

**Accepted:** 28 February 2019

**Published:** 20 March 2019

### Citation:

Rocha VS, Claudio ERG,  
da Silva VL, Cordeiro JP,  
Domingos LF, da Cunha MRH,  
Mauad H, Nascimento TB,  
Lima-Leopoldo AP and Leopoldo AS  
(2019) High-Fat Diet-Induced Obesity  
Model Does Not Promote Endothelial  
Dysfunction via Increasing  
Leptin/Akt/eNOS Signaling.  
Front. Physiol. 10:268.  
doi: 10.3389/fphys.2019.00268

Experimental studies show that the unsaturated high-fat diet-induced obesity promotes vascular alterations characterized by improving the endothelial L-arginine/Nitric Oxide (NO) pathway. Leptin seems to be involved in this process, promoting vasodilation via increasing NO bioavailability. The aim of this study was to test the hypothesis that unsaturated high-fat diet-induced obesity does not generate endothelial dysfunction via increasing the vascular leptin/Akt/eNOS signaling. Thirty-day-old male *Wistar* rats were randomized into two groups: control (C) and obese (Ob). Group C was fed a standard diet, while group Ob was fed an unsaturated high-fat diet for 27 weeks. Adiposity, hormonal and biochemical parameters, and systolic blood pressure were observed. Concentration response curves were performed for leptin or acetylcholine in the presence or absence of Akt and NOS inhibitor. Our results showed that an unsaturated high-fat diet promoted a greater feed efficiency (FE), elevation of body weight and body fat (BF), and an adiposity index, characterizing a model of obesity. However, comorbidities frequently associated with experimental obesity were not visualized, such as glucose intolerance, dyslipidemia and hypertension. The evaluation of the endothelium-dependent relaxation with acetylcholine showed no differences between the C and Ob rats. After NOS inhibition, the response was completely abolished in the Ob group, but not in the C group. Furthermore, Akt inhibition completely blunted vascular relaxation in the C group, but not in the Ob group, which was more sensitive to leptin-induced vascular relaxation. L-NAME incubation abolished the relaxation in both groups at the same level. Although Akt inhibitor pre-incubation reduced the leptin response, group C was more sensitive to its effect. In conclusion, the high-unsaturated fat diet-induced obesity improved the vascular reactivity to leptin and does not generate endothelial dysfunction, possibly by the increase in the vascular sensitivity to leptin and increasing NO bioavailability. Moreover, our results suggest that the increase in NO production occurs through the increase in NOS activation by leptin and is partially mediated by the Akt pathway.

**Keywords:** obesity, high-fat diet, leptin, Akt, vascular reactivity

## INTRODUCTION

Obesity, a disease characterized by excess body fat (BF), is recognized as a global epidemic since it affects virtually all age groups and social classes in both developed and developing countries (Suastika, 2006; Engin, 2017). The most common causes of the occurrence of obesity include increased caloric food intake and physical inactivity, which lead to a positive calorie balance and, consequently, increased BF (Stein and Colditz, 2004). Adipose tissue is considered not only a passive organ that stores energy, but a paracrine and endocrine organ that releases several active substances, including tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-6 (IL-6), adiponectin, resistin and leptin, which may contribute to endothelial dysfunction (Frühbeck et al., 2001).

Endothelial dysfunction associated with obesity has been considered an independent risk of developing cardiovascular diseases (Poirier et al., 2006; Acree et al., 2007; Kumanyika et al., 2008; Stapleton et al., 2008; Graupera and Claret, 2018). For instance, obese animal models induced by a high-energy diet (33% ground chow, 33% Nestle condensed milk, 7% sucrose, and 27% water) present impaired endothelium-dependent and independent relaxation in mesenteric arteries (Naderali et al., 2001). Moreover, Fatani et al. (2007) demonstrated that animals fed a high-saturated fat diet (45% fat) do not present any difference for vasoconstrictor responses to potassium chloride (KCl) or noradrenaline; however, the vascular relaxation in response to carbachol and sodium nitroprusside (SNP) is attenuated in mesenteric arteries, indicating that the high-saturated fat diet promotes vascular dysfunction. In addition, other authors found no difference in acetylcholine-induced endothelium-dependent relaxation in the aorta of mice and obese rabbits induced by a high-fat diet with 58–60% of fat calories (Mundy et al., 2007; Bruder-Nascimento et al., 2017) or standard rabbit chow with 10% added fat (Jerez et al., 2012), which apparently is sex independent.

The mechanisms and factors involved in the vascular disease associated with obesity are uncertain, especially in the vascular endothelium. Several factors, such as insulin resistance and dyslipidemia, have been cited as possible factors that are responsible for cardiovascular dysfunction in models of obesity (Antonopoulos and Tousoulis, 2017). Furthermore, hyperleptinemia induces vasodilation directly, producing and releasing nitric oxide (NO), which diffuses through cellular membranes (Maenhaut and Voorde, 2011). Therefore, this double behavior of leptin and the modulation of vascular tone may help explain why obesity is not always accompanied by high blood pressure (Lembo et al., 2000).

Leptin is an adipocyte-derived hormone, which is elevated in obese humans and rodents (Maffei et al., 1995). In addition to controlling energy stores, it is also involved in several other physiological processes, including vascular tone modulation, acting through specific leptin receptors (Ob-Rb type) located in the vascular endothelium (Leung and Kwan, 2008). Leptin binding to its receptor will activate phosphatidylinositol 3-kinase (PI3K), activating protein kinase B (Akt) and protein kinase A (PKA), which phosphorylates endothelial nitric oxide synthase

(eNOS), resulting in the production of NO (Lembo et al., 2000; Vecchione et al., 2002).

Our laboratory has revealed that obese rats induced by a high-unsaturated fat diet (20% fat) show an increase of vascular L-arginine/NO pathway in order to protect the integrity of vascular function in obesity, possibly mediated by elevated levels of plasma leptin (Bruder-Nascimento et al., 2011). Thus, the aim of this study was to test the hypothesis that unsaturated high-fat diet-induced obesity does not generate endothelial dysfunction via increasing the vascular leptin/Akt/eNOS signaling.

## MATERIALS AND METHODS

### Animals

Thirty-day-old male *Wistar* rats ( $\approx 150$  g) were obtained from the animal facility of the Federal University of Espírito Santo. Rats were individually housed, at constant room temperature ( $24 \pm 2^\circ\text{C}$ ), humidity ( $55 \pm 5\%$ ) and light cycles (12-h light/dark), and had free access to water.

All experiments and procedures were performed in accordance with the Guide for the Care and Use of Laboratory Animals, published by the United States National Institutes of Health, and were approved by the Ethics Committee on Animal Use of the Federal University of Espírito Santo (protocol 027/2014).

### Experimental Design

Rats were randomly distributed into two groups: control (C,  $n = 20$ ), which were fed standard rat chow (Presence chow, SP, Brazil); and obese (Ob,  $n = 18$ ), which were fed a high-unsaturated fat diet (RC Focus 2413, Agrocere<sup>®</sup>, Rio Claro, SP, Brazil) for 27 weeks. The dietary ingredients used to prepare the high-fat diet were sodium chloride, casein, powdered milk, soybean protein concentrate, whole corn, cracker flour, dicalcium phosphate, calcium carbonate, additives emulsifier, antioxidants, and cheese flavoring, along with a vitamins and minerals mixture (Leopoldo et al., 2011). The animals received 50 g of feed daily, and after 24 h, food that was not ingested was weighed for food intake evaluation. The components and nutritional values are displayed in **Table 1**. The high-fat diet was high in calories (high-fat diet = 3.65 kcal/g vs. standard diet = 2.95 kcal/g) due to a higher fat composition made from saturated (20%) and non-saturated (80%) fatty acids. The nutritional status was determined by food consumption (FC), caloric intake (CI) and feed efficiency (FE). Calorie intake (CI) was calculated weekly by the average weekly FC (g)  $\times$  dietary energetic density (kcal). FE was defined as the ability of animals to convert feed energy consumed in body weight and was measured by dividing body weight gain (g) by the total caloric intake (kcal) and multiplying by 100. In addition, body weight was recorded weekly.

### Determination of Obesity

At the end of the experimental protocol, the fat index was used as an indicator of obesity. After 27 weeks of developing obesity, animals were anesthetized intraperitoneally by ketamine and xylazine (50 and 10 mg/kg, respectively) and decapitated.

**TABLE 1** | Components and nutritional values of diets.

Components (%)	Experimental diets	
	Control	High-fat
Protein	23.0	20.0
Carbohydrate	42.8	26.4
Fat	4.0	20.0
Mineral and vitamins	12.1	12.1
Fiber	5.1	9.0
Humidity	13.0	12.5
<b>Nutrient composition (%)</b>		
Energy from protein	30.7	21.9
Energy from carbohydrate	57.3	28.9
Energy from fat	12.0	49.2
<b>Energy density (Kcal/g)</b>	2.99	3.65

After median thoracotomy, fat pads of adipose tissue were dissected and weighed. This method allowed the accurate and consistent assessment of the amount of BF. The adiposity index, used to assess obesity, was calculated with the following formula: adiposity index [BF/final body wt]  $\times$  100. BF was measured from the sum of the individual fat pad weights as follows: BF = epididymal fat + retroperitoneal fat + visceral fat.

## Constitution of the Control and Obese Groups

After 27 weeks of the experimental protocol, a 95% confidence interval (CI) was built for the adiposity index from the Ob and C rats and was adopted as the separation point (SP) between the groups, as previously described (Ferron et al., 2015). After the application of the criterion for determining obesity, seven animals from the control group and nine animals from the obese group were excluded. Thus, our study was carried out with 20 animals in the control group and 18 animals in the obese group.

## Comorbidities and Hormones Associated With Obesity

Obesity developed by a high-fat diet may have changes in the cardiovascular, metabolic and hormonal profile, such as hypertension, glucose intolerance, systemic insulin resistance, dyslipidemia, hyperglycemia, hyperinsulinemia, and hyperleptinemia. Thus, these were evaluated in all groups.

## Measurement of Systolic Blood Pressure (SBP)

Systolic blood pressure was determined by using the non-invasive tail-cuff plethysmography method (IITC INC/Life Science, Woodland Hills, CA, United States) in conscious rats. The rats were pre-warmed at  $35 \pm 2^\circ\text{C}$  for 5 min and placed in a cylindrical acrylic tube. At the proximal region of the animal's tail, a cuff was connected that was inflated to 200–250 mmHg; signals were sent and recorded in the computer to obtain pressure data. At least three measurements were performed for each animal to obtain an average record.

## Glucose Tolerance Test (GTT)

Following 27 weeks of treatment, all rats fasted for 6 h, and a blood sample from the tip of the tail of conscious rats was collected. The basal blood glucose level of each animal was immediately determined using a handheld glucometer (Accu-Check Go Kit; Roche Diagnostic Ltd., São Paulo, Brazil). Subsequently, an injection of glucose solution (2 g/kg body weight) dissolved in water was administered intraperitoneally, and blood glucose levels were measured after 15, 30, 60, 90, and 120 min. Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) was calculated according to the following formula: fasting glucose (mmol/l)  $\times$  fasting insulin ( $\mu\text{U/ml}$ )/22.5.

## Biochemical and Hormonal Determination

For biochemical and hormonal analysis, animals fasted for 12–15 h, then were anesthetized with chloral hydrate (1.0 ml/100 g body wt, intraperitoneally) and sacrificed by decapitation. Blood was collected in Falcon tubes, centrifuged at  $3,000 \times g$  for 15 min and stored at  $-80^\circ\text{C}$ . Serum levels of glucose, triacylglycerol (TG), total cholesterol (T-Chol), protein and high-density lipoprotein (HDL) were determined using a specific kit (Bioclin Bioquímica®, Belo Horizonte, Minas Gerais, Brazil) and analyzed by biochemical analyzer BS-200 (Mindray do Brasil – Comércio e Distribuição de Equipamentos Médicos Ltda, São Paulo, Brazil). Serum insulin and leptin levels were measured using an enzyme-linked immunosorbent assay (ELISA) using specific kits (Linco Research Inc., St. Louis, MO, United States). The reading was carried out using a microplate reader (Asys Expert Plus Microplate Reader, Cambourne, Cambridge, United Kingdom).

## Post-death Morphological Analysis

Animals were anesthetized with chloral hydrate, euthanized by decapitation and thoracotomized. At the end of the experimental protocol, the heart, ventricles and atrium, and tibia were separated, dissected, weighed and measured on a precision scale (AD 500, Mars Scientific and Industrial Instrumentation Ltd., Minas Gerais, Brazil). Cardiac remodeling at the macroscopic level, which identifies the presence or absence of cardiac hypertrophy, was determined by analyzing the following parameters: heart weight (HW), left and right ventricles (LV and RV) and atrium (AT) weights, HW, LV and RV/tibia length ratios.

## Aortic Vascular Reactivity

The animals were anesthetized with a mixture containing ketamine and xylazine (50 mg/kg and 10 mg/kg, respectively) and sacrificed by decapitation. The chest was opened, and the descending thoracic aorta artery was removed and immersed in a modified Krebs-Henseleit solution composed of (in mM/L) NaCl 118, KCl 4.7,  $\text{MgSO}_4$  1.2,  $\text{CaCl}_2$  1.6,  $\text{K}_2\text{HPO}_4$  1.2,  $\text{NaHCO}_3$  25 and glucose 11, and aerated with a carbogenic mixture containing 5%  $\text{CO}_2$  and 95%  $\text{O}_2$ . The connective and adipose tissues were removed, and the artery was divided into sections of approximately 3.5 mm in length. Each vascular ring was mounted in an isolated organ bath system (Panlab

Organ Bath System, ADInstruments, Australia) containing 5 mL of Krebs-Henseleit solution heated to  $37 \pm 0.5^\circ\text{C}$  and was continuously gassed with the carbogenic mixture. Changes in vessel diameter were recorded by a force transducer connected to a data acquisition system (Power Lab, ADInstruments, Australia). During the 45 min stabilization period, the aortic rings were subjected to a resting tension of approximately 1 g and adjusted when necessary. To test the functional integrity the rings were contracted with KCl 75 mM, and after washing and returning to the baseline values, the rings were contracted with  $10^{-7}$  M phenylephrine and relaxed with  $10^{-5}$  M acetylcholine to test the endothelial integrity.

The concentration-response constriction curve was obtained using increasing concentrations of phenylephrine ( $10^{-10}$  to  $10^{-4}$  M), and the data was shown as a percentage of contraction corrected by the maximal contraction induced by KCl. Prior to the concentration-response relaxation curves, all the rings were pre-constricted with  $10^{-7}$  M of phenylephrine. The endothelium-dependent relaxation curves with acetylcholine ( $10^{-11}$  to  $10^{-4}$  M) and leptin ( $10^{-14}$  to  $10^{-6}$  M) were obtained. To analyze the role of Akt and NO in response to leptin and acetylcholine, aortic rings were exposed to the 1L6-hydroxymethyl-chiro-inositol-2-(R)-2-Omethyl-3-O-octadecyl-*sn*-glycerocarbonate “Akt inhibitor” (5  $\mu\text{M}$ , Calbiochem, Merck, Darmstadt, Germany) and N $\omega$ -Nitro-L-arginine methyl ester (L-NAME, 100  $\mu\text{M}$ , Sigma-Aldrich, United States) 30 min prior to the concentration-response curves. To evaluate the baseline NO production in the vascular constriction, the vessel rings of control and obese rats were exposed to L-NAME (100  $\mu\text{M}$ ) before the constrictor response curve to phenylephrine.


### Statistical Analysis

Data on general characteristics, nutritional assessment and comorbidities were reported as mean  $\pm$  standard error of the mean (SEM) and submitted to the Kolmogorov-Smirnov test to determine adherence to normality. Comparisons of groups were analyzed by Student’s *t*-test for independent samples. A repeated measure of analysis of variance (ANOVA), followed by Tukey *post hoc* test, was performed to evaluate the change in body weight. In addition, the concentration-response curves were analyzed using two-way ANOVA, followed by Fisher *post hoc* tests. The *p*-value was considered significant when  $<5\%$ .

## RESULTS

### General Characteristics and Cardiac Morphology

Figure 1 illustrates the significant separation of body weight between C and Ob groups observed at week 9, which remained significantly greater during the 27 weeks of the experimental protocol. Obesity induced by a high-fat diet had a substantial elevation in adiposity parameters, FE, LV, and LVW/tibia length when compared to C rats (Table 2). In contrast, the FC was lower in Ob than in C rats. In addition, there was no difference between the groups in relation to IBW, calorie intake and



**FIGURE 1 |** Evolution of body weight at 27 week of experimental protocol of obesity. C, control; Ob, obese; BW, body weight. Data are reported as the means  $\pm$  SEM. Two-way ANOVA repeated measures for independent samples followed by Tukey *post hoc* test. \**p* < 0.05 Ob vs. C.

**TABLE 2 |** General characteristics and cardiac morphology.

Variables	Groups	
	C (n = 20)	Ob (n = 18)
IBW (g)	134 $\pm$ 3	126 $\pm$ 3
FBW (g)	504 $\pm$ 11	614 $\pm$ 19*
Weight gain (g)	370 $\pm$ 11	488 $\pm$ 21*
Visceral fat (g)	8.2 $\pm$ 0.7	15.3 $\pm$ 1.5*
Retroperitoneal fat (g)	13.2 $\pm$ 0.9	32.2 $\pm$ 3.0*
Epididymal fat (g)	6.8 $\pm$ 0.3	11.2 $\pm$ 0.6*
Body fat (g)	28.2 $\pm$ 1.8	58.6 $\pm$ 4.7*
Adiposity index (%)	5.5 $\pm$ 0.3	9.3 $\pm$ 0.5*
FC (g/day)	22.8 $\pm$ 0.6	18.6 $\pm$ 0.3*
CI (g $\times$ kcal/day)	67.3 $\pm$ 1.7	67.9 $\pm$ 1.1
FE (%)	2.91 $\pm$ 0.05	3.82 $\pm$ 0.18*
Heart (g)	1.18 $\pm$ 0.05	1.25 $\pm$ 0.04
LVW (g)	0.83 $\pm$ 0.02	0.90 $\pm$ 0.03*
Heart/Tibia-length (g/cm)	0.28 $\pm$ 0.01	0.30 $\pm$ 0.01
LVW/Tibia-length (g/cm)	0.20 $\pm$ 0.01	0.22 $\pm$ 0.01*

Values are means  $\pm$  SEM. C, control; Ob, obese; IBW, initial body weight; FBW, final body weight; FC, food consumption; CI, calorie intake; FE, feed efficiency; LVW, left ventricle weight. Student’s *t*-test for independent samples. \**p* < 0.05 Ob vs. C.

absolute heart, as well as heart/tibia length and tibia length (data not shown) (Table 2).

### Comorbidities Associated With Obesity

There were no significant differences in comorbidities associated with obesity, since the SBP, TG, Cholesterol, LDL, HDL, and protein were similar between the groups (Table 3). The glucose tolerance profile, AUC, insulin levels and HOMA-IR were also not affected by exposure to obesity (Figure 2). However, the Ob rats presented only higher levels of glucose at 90 min than the C rats (Figure 2A). In addition, the Ob group exhibited higher levels of leptin when compared to C (Table 3).



**TABLE 3** | Comorbidities associated with obesity.

Variables	Groups	
	C (n = 9)	Ob (n = 7)
SBP (mmHg)	122 ± 5	128 ± 6
TG (mg/dL)	76 ± 6	76 ± 10
Cholesterol (mg/dL)	78 ± 6	68 ± 4
LDL (mg/dL)	11.1 ± 1.4	11.3 ± 0.9
HDL (mg/dL)	20 ± 2	21 ± 2
Protein (g/dL)	5.36 ± 0.34	5.40 ± 0.28
Leptin (ng/mL)	0.77 ± 0.08	1.32 ± 0.21*

Values are reported as the means ± SEM. C, control; Ob, obese; SBP, systolic blood pressure; TG, triglycerides; LDL, low-density lipoprotein; HDL, high-density lipoprotein. Student's t-test for independent samples. \*p < 0.05 vs. C.


### Vascular Reactivity

The maximal contraction induced by 75 mM KCl stimulation was not different between groups (C: 3.25 ± 0.08 g vs. Ob: 3.05 ± 0.08 g; p > 0.05). The concentration-response curve to phenylephrine showed a significant reduction in the constrictor response obtained from vessels of Ob animals compared to C (Figure 3A). L-NAME increased the adrenergic response



in both groups compared with its respective control curve (Figures 3B,C). However, Ob rats were more sensitive to L-NAME effects, and the phenylephrine curve from Ob rats presented a shift to the left compared to the C rats, but no difference in the maximum response (Figure 3D).

The endothelium-dependent relaxation with acetylcholine showed no differences between the C and Ob rats (Figure 4A). However, NO synthesis inhibition generated a significant reduction in relaxation for both groups, compared with their respective controls (Figure 4A). In addition, L-NAME completely abolished the vascular relaxation in the Ob group, and partially in the C group (Figure 4B). On the other hand, Akt inhibition promoted reduction in the acetylcholine response only in C rats (Figure 4B).

Vasodilation-induced by leptin showed a greater response in aortic rings from Ob rats compared to that of the C rats (Figure 5A). The presence of L-NAME blunted the response of leptin-induced relaxation in both groups (Figure 5A). On the other hand, Akt inhibition resulted in reduced vascular leptin response in both groups, but this effect was less pronounced in Ob rats compared to C rats, suggesting that leptin promoted vascular relaxation partially mediated by Akt activation in obesity (Figure 5B).




**FIGURE 2** | Glucose tolerance profile at 27 week of experimental protocol of obesity. Glucose tolerance test (A), area under the curve for glucose – AUC (B), insulin serum level (C) and (D) homeostatic model assessment index (HOMA-IR) from control (C; n = 9) and obese (Ob; n = 7) groups. Data are reported as the means ± SEM. Two-way ANOVA repeated measures for independent samples followed by Tukey post hoc test. \*p < 0.05 Ob vs. C.



## DISCUSSION

The main findings of the present study were as follows: (i) the high-fat diet led to the development of obesity; (ii) a reduction in the vasoconstrictor response to phenylephrine in obese animals;

(iii) endothelial dysfunction was not observed in this model of obesity, since thoracic aorta from obese rats were still responsive to leptin effects; (iv) a significantly higher vasodilator response to leptin was observed in the obese group, which was abolished when exposed to NOS blockade in both groups; (v) there was a



reduction in the vasodilator response to leptin in both groups, compared with the respective control, when the rings were exposed to Akt inhibitor.

Experimental models that mimic the eating habits of the human population have been widely used to elucidate the mechanisms of obesity and metabolic disorders (Angelova and Boyadjiev, 2013). In this sense, our data show that rats fed the high-fat diet developed obesity characterized by an increase in the FBW, total BF and fat pads (retroperitoneal, visceral and epididymal); however, these animals had hyperleptinemia frequently related and observed in the experimental and human studies with obesity (Woods et al., 2003; Nascimento et al., 2008). In addition, obesity is often associated with metabolic and endocrine disorders, such as glucose intolerance, insulin resistance, hyperleptinemia, dyslipidemia, and hypertension. Nevertheless, in this obesity model, no changes were observed in basal glucose, insulin or alterations of the lipid profile.

In addition to absence of changes in the metabolic abnormalities, the model proposed did not lead to an elevation in blood pressure levels. This finding corroborates studies that have verified a significant reduction in SBP in obese animals (Reilly and Rader, 2003; Carroll et al., 2006; Lima-Leopoldo et al., 2011). Some studies have shown that a high-unsaturated fat diet is related to the production of eNOS stimulated by hyperleptinemia and that it may be responsible for increased production and release of NO, resulting in a reduction of peripheral resistance and, consequently, lower blood pressure (Bruder-Nascimento et al., 2011, 2017).

The results from the post-death morphological analysis revealed obesity-induced cardiac hypertrophy, evidenced by elevation of absolute LV weight, as well as LV/tibia length. These data corroborate several studies that indicate obesity induced by a high-fat diet for 15 and 30 weeks promotes cardiac hypertrophy; these authors suggest that cardiac remodeling occurs due to the long period of obesity (Lima-Leopoldo et al., 2014; Silva et al., 2014), which is considered an adaptive characteristic of obesity (Rider et al., 2009). Thus, myocyte hypertrophy is considered a mechanism of adaptation in response to several stimuli, such as increased cardiac output due to the metabolic

demand caused by obesity condition (Mill and Vassallo, 2001; Rahmouni et al., 2005).

Obesity constitutes a major cause of CVD, including hypertension, atherosclerosis, ischemic heart disease and heart failure. It is widely known that a high intake of saturated fat increases the risk of cardiovascular diseases, while diets enriched with unsaturated fatty acids seem to lower this risk (Lorente-Cebrián et al., 2013). Thus, several authors have shown beneficial effects on obesity-related vascular alterations (Bruder-Nascimento et al., 2011; Lorente-Cebrián et al., 2013; de Oliveira et al., 2017) In this sense, these alterations play a key role in the pathogenesis of these diseases, which includes the reduction of NO bioavailability as well as the elevation of components that induce vascular contractility, such as angiotensin-II and endothelin-1 (Zemel et al., 1990; Hopfner et al., 1998; Mundy et al., 2007). Furthermore, adiposity is associated with elevations in the adipocyte-derived hormone leptin, which causes a dual effect on the cardiovascular system. Centrally, it stimulates the sympathetic nervous system and increasing the arterial blood pressure, while peripherally, it modulates the release of several vasoactive factors, mainly NO, in the vascular endothelium (Kimura et al., 2000; Lembo et al., 2000).

Another effect of obesity induced by a diet rich in unsaturated fat for 27 weeks was the significant reduction in the vascular adrenergic response in Ob animals. This result was in accordance with our previously published results (Bruder-Nascimento et al., 2011, 2017). However, when the animals were submitted to the diet for 30 weeks, this difference was abolished (Bruder-Nascimento et al., 2011). In another study with obese mice induced by a high sugar diet for 8 weeks, Silva et al. (2016) verified a significant decrease in the contractile response to phenylephrine in obese animals that was associated with an increase in the vascular production of NO, which the authors attributed to the higher expression of iNOS. Moreover, in rabbits fed a hyperlipidemic diet for 12 weeks, hyporeactivity was observed in the responses stimulated by the noradrenaline and angiotensin-II (Jerez et al., 2012). Taken together, these results suggest that the time-exposure to the diet, and consequently to the obesity and hyperleptinemia, might be a determinant factor on vascular

adaptations and the vasoconstrictor responses, which is mainly related to the adrenergic system.

On the other hand, Keita et al. (2013) and Bautista et al. (2014) evaluated the aortic vascular reactivity in rats with unsaturated hyperlipidemic diet-induced obesity (with fried canola oil and margarine) for 10 and 20 weeks, respectively, and demonstrated that obesity does not change the vasoconstrictor response to phenylephrine or noradrenaline. The responses to these vasoconstrictor agonists may be related to the amount and the nature of the lipids in the diet, which are considered protective factors and/or atherogenic and are associated with the modulation of NOS activity (Santos et al., 2013).

In our study, the lowest vasoconstrictor response observed in Ob animals occurred due to a higher basal production of NO, counterbalancing and reducing the constriction caused by the stimulation of adrenergic receptors. Furthermore, the vessels of Ob animals showed a higher dependency of NO in the endothelium-mediated relaxation since the response to acetylcholine and leptin were totally abolished when NOS was inhibited with L-NAME (Figures 4A, 5A). The higher basal production of NO is possibly associated with the increase in the plasmatic concentration of leptin in obese animals. Several mechanisms have been proposed for the increase in the NO production stimulated by leptin, which might be associated with a compensatory mechanism against its sympathetic stimulation (Mitchell et al., 2001). Lembo et al. (2000) and Kimura et al. (2000) verified that the relaxation induced by leptin occurs via production of endothelial NO in the rat aorta. In this sense (Vecchione et al., 2002). showed that leptin activates the Akt pathway by a mechanism independent of PI3K phosphorylation, increasing NO production by phosphorylating eNOS at the Ser1177 residue in the aorta of *Wistar Kyoto* rats. In addition, it was verified that insulin potentiates this response by increasing the Akt phosphorylation at the sites of Ser<sup>473</sup> and Thr<sup>308</sup> (Vecchione et al., 2003). In another study that evaluated the effect of leptin in human aortic endothelial cells, a new pathway was activated by leptin that was mediated by the  $\alpha 1$  subunit of AMP-activated protein kinase (AMPK), which phosphorylates Akt at the Ser<sup>473</sup> residue, promoting the activation of eNOS (Procópio et al., 2009). Recently, a study verified that in eNOS knockout mice, leptin and a fat-rich diet induced an increase in the endothelial nNOS expression by phosphorylating JAK-2/STAT-3, thereby compensating for the reduction of NO and restoring the endothelial function in angiotensin-II-infused mice (Benkhoff et al., 2012). We can verify that among many pathways

described for leptin-induced NOS activation, most of them are triggered by Akt phosphorylation. However, according to our results, this pathway is partially activated in this model of obesity, since the Akt inhibition did not change the endothelium-mediated dilation with either acetylcholine or leptin to the same level as observed in C animals.

## CONCLUSION

In conclusion, the high-unsaturated fat (49.2 energy from fat) diet-induced obesity improved the vascular reactivity to leptin and does not generate endothelial dysfunction, possibly by the increase in the vascular sensitivity to leptin and increasing NO bioavailability. Moreover, our results suggest that the increase in NO production occurs through the increase in NOS activation by leptin and is partially mediated by the Akt pathway.

## AUTHOR CONTRIBUTIONS

AL-L, VR, and AL participated and conceived the research design. EC, Vds, JC, LD, MdC, AL-L, and AL performed the experiments and data analysis. VR, EC, MdC, HM, TBdN, AL-L, and AL interpreted and discussed the data. AL, VR, EC, MdC, and AL wrote the manuscript. HM, TBdN, AL-L, and AL refined the final draft and revised the manuscript. All authors read and approved the final version of the manuscript.

## FUNDING

This work was supported by Fundação de Amparo à Pesquisa e Inovação do Espírito Santo – FAPES (Grant Number: 67636713/15), Brazilian National Council for Scientific and Technological Development – CNPq (Grant Number: 445349/2014-0) and financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

## ACKNOWLEDGMENTS

The authors are grateful to Priscilla Spadeto Altoé for their assistance.

## REFERENCES

- Acree, L. S., Comp, P. C., Whitsett, T. L., Montgomery, P. S., Nickel, K. J., Fjeldstad, A. S., et al. (2007). The influence of obesity on calf blood flow and vascular reactivity in older adults. *Dyn. Med.* 6, 1–8. doi: 10.1186/1476-5918-6-4
- Angelova, P., and Boyadjiev, N. (2013). A review on the models of obesity and metabolic syndrome in rats. *Trakia J. Sci.* 1, 5–12.
- Antonopoulos, A. S., and Tousoulis, D. (2017). The molecular mechanisms of obesity paradox. *Cardiovasc. Res.* 113, 1074–1086. doi: 10.1093/cvr/cvx106
- Bautista, R., Carreón-Torres, E., Luna-Luna, M., Komera-Arenas, Y., Franco, M., Frago, J. M., et al. (2014). Early endothelial nitrosylation and increased abdominal adiposity after long-term consumption of frying used canola oil in Wistar rats. *Nutrition* 30, 1055–1060. doi: 10.1016/j.nut.2014.01.010
- Benkhoff, S., Loot, A. E., Pierson, I., Sturza, A., Kohlstedt, K., Fleming, I., et al. (2012). Leptin potentiates endothelium-dependent relaxation by inducing endothelial expression of neuronal NO Synthase. *Arterioscler. Thromb. Vasc. Biol.* 32, 1605–1612. doi: 10.1161/ATVBAHA.112.251140
- Bruder-Nascimento, T., Baptista, R. F. F., Pereira, P. C., Campos, D. H. S., Leopoldo, A. S., Leopoldo, A. P. L., et al. (2011). Vascular alterations in high-fat diet-obese rats: role of endothelial L-arginine/NO pathway. *Arq. Bras. Card.* 97, 40–45.



- Bruder-Nascimento, T., Ekeledo, O. J., Anderson, R., Le, H. B., and Belin de Chantemèle, E. J. (2017). Long term high fat diet treatment: an appropriate approach to study the sex-specificity of the autonomic and cardiovascular responses to obesity in mice. *Front. Physiol.* 8:32. doi: 10.3389/fphys.2017.00032
- Carroll, J. F., Zenebe, W. J., and Strange, T. B. (2006). Cardiovascular function in rat model of diet-induced obesity. *Hypertension* 48, 65–72. doi: 10.1161/01.HYP.0000224147.01024.77
- de Oliveira, P. A., Kovacs, C., Moreira, P., Magnoni, D., Saleh, M. H., and Faintuch, J. (2017). Unsaturated fatty acids improve atherosclerosis markers in obese and overweight non-diabetic elderly patients. *Obes. Surg.* 27, 2663–2671. doi: 10.1007/s11695-017-2704-8
- Engin, A. (2017). The definition and prevalence of obesity and metabolic syndrome. *Adv. Exp. Med. Biol.* 960, 1–17. doi: 10.1007/978-3-319-48382-5\_1
- Fatani, S., Pickavance, L. C., Sadler, C. J., Harrold, J. A., Cassidy, R., Wilding, J. P. H., et al. (2007). Differential vascular dysfunction in response to diets of differing macronutrient composition: a phenomenological study. *Nutr. Metab.* 4, 1–6. doi: 10.1186/1743-7075-4-15
- Ferron, A. J., Jacobsen, B. B., Sant'Ana, P. G., de Campos, D. H., de Tomasi, L. C., Luvizotto Rde, A., et al. (2015). Cardiac dysfunction induced by obesity is not related to  $\beta$ -adrenergic system impairment at the receptor-signalling pathway. *PLoS One* 10:e0138605. doi: 10.1371/journal.pone.0138605
- Frühbeck, G., Gomez-Ambrosi, J., Bal, M. J. F., and Burrell, M. A. (2001). The adipocyte: a model for integration of endocrine and metabolic signaling in energy metabolism regulation. *Am. J. Physiol. Endocrinol. Metab.* 280, E827–E847. doi: 10.1152/ajpendo.2001.280.6.E827
- Graupera, M., and Claret, M. (2018). Endothelial cells: new players in obesity and related metabolic disorders. *Trends Endocrinol. Metab.* 29, 781–794. doi: 10.1016/j.tem.2018.09.003
- Hopfner, R. L., McNeill, J. R., and Gopalakrishnan, V. (1998). Vanadate treatment normalizes exaggerated vascular smooth muscle responses in the obese Zucker rat. *Eur. J. Pharmacol.* 357, 61–65. doi: 10.1016/S0014-2999(98)00535-4
- Jerez, S., Scacchi, F., Sierra, L., Karbinder, S., and Bruno, M. P. (2012). Vascular hyporeactivity to angiotensin II and noradrenaline in a rabbit model of obesity. *J. Cardiovasc. Pharmacol.* 59, 49–57. doi: 10.1097/FJC.0b013e318235156a
- Keita, H., Ramírez-San, J. E., Paniagua-Castro, N., Garduño-Siciliano, L., and Quevedo, L. (2013). The long-term ingestion of a diet high in extra virgin olive oil produces obesity and insulin resistance but protects endothelial function in rats: a preliminary study. *Diabetol. Metab. Syndr.* 5, 1–10. doi: 10.1186/1758-5996-5-53
- Kimura, K., Tsuda, K., Baba, A., Kawabe, T., Boh-oka, S., Ibata, M., et al. (2000). Involvement of nitric oxide in endothelium-dependent arterial relaxation by leptin. *Biochem. Biophys. Res. Commun.* 273, 745–749. doi: 10.1006/bbrc.2000.3005
- Kumanyika, S. K., Obarzanek, E., Stettler, N., Bell, R., Field, A. E., Fortmann, S. P., et al. (2008). Population-based prevention of obesity. *Circulation* 118, 428–464. doi: 10.1161/CIRCULATIONAHA.108.189702
- Lembo, G., Vecchione, C., Fratta, L., Marino, G., Trimarco, V., D'amati, G., et al. (2000). Leptin induces direct vasodilation through distinct endothelial mechanisms. *Diabetes Metab. Res. Rev.* 49, 293–297. doi: 10.2337/diabetes.49.2.293
- Leopoldo, A. S., Lima-Leopoldo, A. P., Sugizaki, M. M., do Nascimento, A. F., de Campos, D. H., Luvizotto, R. A., et al. (2011). Involvement of L-type calcium channel and SERCA2A in myocardial dysfunction induced by obesity. *J. Cell. Physiol.* 2011, 2934–2942. doi: 10.1002/jcp.22643
- Leung, Y. M., and Kwan, C. Y. (2008). Dual vascular effects of leptin via endothelium: hypothesis and perspective. *Chin. J. Physiol.* 51, 1–6.
- Lima-Leopoldo, A. P., Leopoldo, A. S., da Silva, D. C., do Nascimento, A. F., de Campos, D. H., Luvizotto, R. A., et al. (2014). Long-term obesity promotes alterations in diastolic function induced by reduction of phospholamban phosphorylation at serine-16 without affecting calcium handling. *J. Appl. Physiol.* 117, 669–678. doi: 10.1152/jappphysiol.00088.2014
- Lima-Leopoldo, A. P., Leopoldo, A. S., Sugizaki, M. M., Bruno, A., Nascimento, A. F., Luvizotto, R. A., et al. (2011). Myocardial dysfunction and abnormalities in intracellular calcium handling in obese rats. *Arq. Bras. Cardiol.* 97, 232–240. doi: 10.1590/S0066-782X2011005000061
- Lorente-Cebrián, S., Costa, A. G., Navas-Carretero, S., Zabala, M., Martínez, J. A., and Moreno-Aliaga, M. J. (2013). Role of omega-3 fatty acids in obesity, metabolic syndrome, and cardiovascular diseases: a review of the evidence. *J. Physiol. Biochem.* 69, 633–651. doi: 10.1007/s13105-013-0265-4
- Maenhaut, N., and Voorde, J. V. (2011). Regulation of vascular tone by adipocytes. *BMC Med.* 9:25. doi: 10.1186/1741-7015-9-25
- Maffei, M., Halaas, J., Ravussin, E., Pratley, R. E., Lee, G. H., Zhang, Y., et al. (1995). Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat. Med.* 1, 1155–1161. doi: 10.1038/nm1195-1155
- Mill, J. G., and Vassallo, D. V. (2001). Hipertrofia cardíaca. *Rev. Bras. Hipertens.* 8, 63–75.
- Mitchell, J. L., Morgan, D. A., Correia, M. L. G., Mark, A. L., Sivitz, W. I., and Haynes, W. G. (2001). Does leptin stimulate nitric oxide to oppose the effects of sympathetic activation? *Hypertension* 38, 1081–1086.
- Mundy, A. L., Haas, E., Bhattacharya, I., Widmer, C. C., Kretz, M., Hofmann-Lehmann, R., et al. (2007). Fat intake modifies vascular responsiveness and receptor expression of vasoconstrictors: implications for diet-induced obesity. *Cardiovasc. Res.* 73, 368–375. doi: 10.1016/j.cardiores.2006.11.019
- Naderali, E. K., Brown, M. J., Pickavance, L. C., Wilding, J. P. H., Doyle, P. J., and Williams, G. (2001). Dietary obesity in the rat induces endothelial dysfunction without causing insulin resistance: a possible role for triacylglycerols. *Clin. Sci.* 101, 499–506. doi: 10.1042/cs1010499
- Nascimento, A. F., Sugizaki, M. M., Leopoldo, A. S., Lima-Leopoldo, A. P., Nogueira, C. R. N., Novelli, E. L. B., et al. (2008). Misclassification probability as obese or lean in hypercaloric and normocaloric diet. *Biol. Res.* 41, 253–259. doi: 10.4067/S0716-97602008000300002
- Poirier, P., Giles, T. D., Bray, G. A., Hong, Y., Stern, J. S., Pi-Sunyer, F. X., et al. (2006). Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss. *Arterioscler. Thromb. Vasc. Biol.* 26, 968–976. doi: 10.1161/01.ATV.0000216787.85457.f3
- Procópio, C., Andreozzi, F., Laratta, E., Cassese, A., Beguinot, F., Arturi, F., et al. (2009). Leptin-stimulated endothelial nitric-oxide synthase via an adenosine 5'-monophosphate-activated protein kinase/Akt signaling pathway is attenuated by interaction with C-reactive protein. *Endocrinology* 150, 3584–3593. doi: 10.1210/en.2008-0921
- Rahmouni, K., Correia, M. L., Haynes, W. G., and Mark, A. L. (2005). Obesity-associated hypertension: new insights into mechanisms. *Hypertension* 45, 9–14. doi: 10.1161/01.HYP.0000151325.83008.b4
- Reilly, M. P., and Rader, D. J. (2003). The metabolic syndrome: more than the sum of its parts? *Circulation* 108, 1546–1551. doi: 10.1161/01.CIR.0000088846.10655.E0
- Rider, O. J., Francis, J. M., Ali, M. K., Byrne, J., Clarke, K., Neubauer, S., et al. (2009). Determinants of left ventricular mass in obesity; a cardiovascular magnetic resonance study. *J. Cardiovasc. Magn. Reson.* 11:9. doi: 10.1186/1532-429X-11-9
- Santos, R. D., Gagliardi, A. C. M., Xavier, H. T., Magnoni, C. D., Cassani, R., Lottenberg, A. M., et al. (2013). Sociedade Brasileira de Cardiologia. I Diretriz sobre o consumo de gorduras e saúde cardiovascular. *Arq. Bras. Cardiol.* 100, 1–40.
- Silva, D. C., Lima-Leopoldo, A. P., Leopoldo, A. S., Campos, D. H., Nascimento, A. F., Oliveira Junior, S. A., et al. (2014). Influence of term of exposure to high-fat diet-induced obesity on myocardial collagen type I and III. *Arq. Bras. Cardiol.* 102, 157–163.
- Silva, J. F., Correa, I. C., Diniz, T. F., Lima, P. M., Santos, R. L., Cortes, S. F., et al. (2016). Coimbra CC, Lemos VS. Obesity, inflammation, and exercise training: relative contribution of iNOS and eNOS in the modulation of vascular function in the mouse aorta. *Front. Physiol.* 7:386. doi: 10.3389/fphys.2016.00386
- Stapleton, P. A., James, M. E., Goodwill, A. G., and Frisbee, J. C. (2008). Obesity and vascular dysfunction. *Pathophysiology.* 15, 79–89. doi: 10.1016/j.pathophys.2008.04.007
- Stein, C. J., and Colditz, G. A. (2004). The epidemic of obesity. *J. Clin. Endocrinol. Metab.* 89, 2522–2525. doi: 10.1210/jc.2004-0288
- Suastika, K. (2006). Update in the management of obesity. *Acta. Med. Indones.* 38, 231–237.
- Vecchione, C., Aretini, A., Maffei, A., Marino, G., Selvetella, G., Poulet, R., et al. (2003). Cooperation between insulin and leptin in the modulation of vascular tone. *Hypertension.* 42, 166–170. doi: 10.1161/01.HYP.0000082806.73530.68

- Vecchione, C., Maffei, A., Colella, S., Aretini, A., Poulet, R., Frati, G., et al. (2002). Leptin effect on endothelial nitric oxide is mediated through Akt-endothelial nitric oxide synthase phosphorylation pathway. *Diabetes Metab. Res. Rev.* 51, 168–173. doi: 10.2337/diabetes.51.1.168
- Woods, S. C., Seeley, R. J., Rushing, P. A., D'Alessio, D., and Tso, P. (2003). A controlled high-fat diet induces an obese syndrome in rats. *J. Nutr.* 133, 1081–1087. doi: 10.1093/jn/133.4.1081
- Zemel, M. B., Reddy, S., Shehin, S. E., Lockette, W., and Sowers, J. R. (1990). Vascular reactivity in Zucker obese rats: role of insulin resistance. *J. Vasc. Med. Biol.* 2, 81–85.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Rocha, Claudio, da Silva, Cordeiro, Domingos, da Cunha, Mauad, Nascimento, Lima-Leopoldo and Leopoldo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.