



Activation of Immune Genes in Leafhoppers by Phytoplasmas and Symbiotic Bacteria

Elena Gonella¹, Mauro Mandrioli², Rosemarie Tedeschi¹, Elena Crotti³, Marianna Pontini¹ and Alberto Alma^{1*}

¹ Dipartimento di Scienze Agrarie, Forestali e Alimentari (DISAFA), Università degli Studi di Torino, Grugliasco, Italy,

² Dipartimento di Scienze della Vita (DSV), Università degli Studi di Modena e Reggio Emilia, Modena, Italy, ³ Dipartimento di Scienze per gli Alimenti, la Nutrizione e l'Ambiente (DeFENS), Università degli Studi di Milano, Milan, Italy

OPEN ACCESS

Edited by:

Michel Cusson,
Natural Resources Canada, Canada

Reviewed by:

Arash Zibaei,
University of Guilan, Iran
Jalal Jalali Sendi,
University of Guilan, Iran
Olga Makarova,
Freie Universität Berlin, Germany

*Correspondence:

Alberto Alma
alberto.alma@unito.it

Specialty section:

This article was submitted to
Invertebrate Physiology,
a section of the journal
Frontiers in Physiology

Received: 12 December 2018

Accepted: 06 June 2019

Published: 21 June 2019

Citation:

Gonella E, Mandrioli M, Tedeschi R, Crotti E, Pontini M and Alma A (2019) Activation of Immune Genes in Leafhoppers by Phytoplasmas and Symbiotic Bacteria. *Front. Physiol.* 10:795. doi: 10.3389/fphys.2019.00795

Insect immunity is a crucial process in interactions between host and microorganisms and the presence of pathogenic, commensal, or beneficial bacteria may result in different immune responses. In Hemiptera vectors of phytoplasmas, infected insects are amenable to carrying high loads of phytopathogens, besides hosting other bacterial affiliates, which have evolved different strategies to be retained; adaptation to host response and immunomodulation are key aspects of insect-symbiont interactions. Most of the analyses published to date has investigated insect immune response to pathogens, whereas few studies have focused on the role of host immunity in microbiota homeostasis and vectorial capacity. Here the expression of immune genes in the leafhopper vector of phytoplasmas *Euscelidius variegatus* was investigated following exposure to *Asaia* symbiotic bacteria, previously demonstrated to affect phytoplasma acquisition by leafhoppers. The expression of four genes related to major components of immunity was measured, i.e., defensin, phenoloxidase, kazal type 1 serine protease inhibitor and Raf, a component of the Ras/Raf pathway. The response was separately tested in whole insects, midguts and cultured hemocytes. Healthy individuals were assessed along with specimens undergoing early- and late-stage phytoplasma infection. In addition, the adhesion grade of *Asaia* strains was examined to assess whether symbionts could establish a physical barrier against phytoplasma colonization. Our results revealed a specific activation of Raf in midguts after double infection by *Asaia* and *flavescence dorée* phytoplasma. Increased expression was observed already in early stages of phytoplasma colonization. Gut-specific localization and timing of Raf activation are consistent with the role played by *Asaia* in limiting phytoplasma acquisition by *E. variegatus*, supporting the involvement of this gene in the anti-pathogen activity. However, limited attachment capability was found for *Asaia* under *in vitro* experimental conditions, suggesting a minor contribution of physical phytoplasma exclusion from the vector gut wall. By providing evidence of immune modulation played by *Asaia*, these results contribute to elucidating the molecular mechanisms regulating interference

with phytoplasma infection in *E. variegatus*. The involvement of Raf suggests that in the presence of reduced immunity (reported in Hemipterans), immune genes can be differently regulated and recruited to play additional functions, generally played by genes lost by hemipterans.

Keywords: insect immunity, plant pathogen, symbiotic bacterium, *Asaia*, Raf

INTRODUCTION

The interactions of organisms with the surrounding environment are related to their ability in responding to damage and non-self particles. In insects, responses are carried out via innate immunity exclusively, by means of both humoral and cellular defense. A major component of humoral and cellular immune response is the production of antimicrobial peptides (AMPs), which takes place in different organs and tissues, including the digestive tract, salivary glands, and fat body (Tsakas and Marmaras, 2010; Li et al., 2016; Shao et al., 2017). Many AMPs such as defensins are highly conserved in different orders of insects, while others are not evenly distributed (Tedeschi et al., 2017). Moreover, other important components of insect immunity are commonly found, such as phenoloxidase-related machinery and other genes. Insect immune responses are differentially stimulated by either pathogenic or beneficial microorganisms (Tedeschi et al., 2017). Indeed, besides activating responses to neutralize infections by entomopathogens, insects may modulate their immunity in the presence of bacterial symbionts to regulate their density and to maintain the microbiota balance (Login et al., 2011; Eleftherianos et al., 2013; Skidmore and Hansen, 2017). Differently from what is reported for entomopathogens that induce very similar responses, the interaction with non-pathogenic microbes depends on specific traits of insect–microorganism interactions: divergent reactions have indeed been recorded even for closely related microorganisms in the same host. For example, some pathogens of animals or plants, vectored by insects, may establish different interplay with their vectors, which in turn react with different responses. Although pathogenic agents for humans are often perceived as harmful agents by infected insects (Weiss and Aksoy, 2011), in the case of plant pathogens, exclusively transmitted by Hemiptera, divergent kinds of interactions have been observed. Infection by the α -proteobacterial ‘*Candidatus Liberibacter asiaticus*’ resulted in downregulation of immune genes in nymphs of the psyllid *Diaphorina citri* Kuwayama, suggesting that the pathogen is able to modulate the vector immune response to promote its colonization of the hosts (Vyas et al., 2015). Considering plant pathogenic Mollicutes, *Spiroplasma citri* was shown to induce a specific response in the vector *Circulifer haematocaps* (Mulsant and Rey), consisting of increased phagocytosis and upregulation of a gene related to hexamerin, a protein playing a crucial role in phenoloxidase activation (Eliautout et al., 2016). However, the response is balanced by the capability of *S. citri* to inhibit phenoloxidase activity and escape phagocytosis (Eliautout et al., 2016). In phytoplasmas, diverse insect responses have been reported following infection by different strains in the same host species, i.e., immune response

or immune priming from infections (Galletto et al., 2018). Moreover, some phytoplasmas counteract the insect response by expressing genes involved in limiting the products of immunity (Makarova et al., 2015).

The leafhopper *Euscelidius variegatus* Kirschbaum (Hemiptera: Cicadellidae) is a polyphagous polyvoltine species capable of transmitting phytoplasmas belonging to different taxonomic groups, including chrysanthemum yellows phytoplasma (CYP, 16SrI group) and flavescence dorée phytoplasma (FDp, 16SrV group), under laboratory conditions. These pathogens have been shown to have opposite effects on *E. variegatus*, with CYP slightly enhancing the fitness of the leafhopper (Bosco and Marzachi, 2016) and FDp reducing its survival and fecundity (Bressan et al., 2005). Transcriptomic analysis of leafhoppers infected with either CYP or FDp has demonstrated that only infection by FDp resulted in activation of insect immune response (Galletto et al., 2018). Besides carrying phytoplasmas, *E. variegatus* harbors bacterial symbionts, like many other Auchenorrhyncha (Baumann, 2005); among these, the acetic acid bacterium *Asaia* has been experimentally documented to limit the acquisition of FDp, after oral administration (Gonella et al., 2018). Symbiont-mediated control mechanisms against phytopathogens include competitive nutrient uptake by symbiotic bacteria, erection of a physical barrier preventing gut establishment and crossing by pathogens, symbiont-mediated immune response of the insect, and the release of antagonistic compounds (Gonella et al., 2019). In *E. variegatus*, the involvement of either *Asaia*-mediated mechanisms stimulating the host immune response or physical exclusion were suggested (Gonella et al., 2018); however, little experimental evidence has been provided on the effects of symbiotic bacteria on *E. variegatus* immunity (Tedeschi et al., 2017) and no data are available on the influence of phytoplasma-symbiont multiple infection on the insect response. Additionally, interest in the molecular machinery involved in the immune response of hemipteran species is hampered by the limited immune repertoire possessed by these insects, as reported for many species (Arp et al., 2016; Skidmore and Hansen, 2017). Such a reduced response is thought to result from the need of Hemiptera to maintain stable relationships with bacterial symbionts. On the other hand, symbiotic bacteria may compensate for the reduction of immunity of their hosts by stimulating insects’ responses to protect them from enemies or directly protecting their hosts from pathogens (Eleftherianos et al., 2013). Moreover, insect immune activation is a candidate strategy used by endosymbionts to modulate the density of other bacteria (Skidmore and Hansen, 2017), including vector-transmitted disease agents, actually altering the insect transmission competence (Weiss and Aksoy, 2011; Kliot et al., 2014). The immune response may be especially crucial for those

phytopathogens that cause decreased vector fitness (Cassone et al., 2014; Nachappa et al., 2014; Alma et al., 2015; Olson and Blair, 2015), as in the case of FDp and *E. variegatus*.

In this study, we examined the question as to whether reduced phytoplasma acquisition observed after *Asaia* infection is related to stimulation of the insect immune system, and whether this mechanism is tissue-specific and related to phytoplasma infection timing. To this end, we investigated the expression pattern of four immune genes in *E. variegatus* whole adult leafhoppers, dissected midguts and cultured hemocytes, after exposure to *Asaia* strains and/or FDp. We selected genes involved in different immune pathways to explore possible peculiar regulation of immune genes in insects with reduced immunity such as Hemiptera. In particular, we analyzed phenoloxidase and kazal type 1 serine protease inhibitor, since both genes have been reported to respond to FDp infection (Galetto et al., 2018), together with the *defensin* gene that is one of the most commonly activated genes after bacterial challenges. Moreover, special attention was given to the *Raf* gene, a component of the Ras/Raf pathway, which is known in *Drosophila* as involved in the response to septic injury and in hemocyte proliferation and survival (Asha et al., 2003). The latter was selected since during the latent period (before the insect becomes infective) phytoplasmas multiply in different organs/tissues, including hemocytes, affecting their proliferation/survival (Bosco et al., 2007). Finally, an alternative hypothetical mechanism of interference with FDp acquisition was tested, involving the production of air-liquid interface (ALI) biofilm (Supplementary Table S1), i.e., masking of gut epithelial receptors through adhesion to insect gut wall.

MATERIALS AND METHODS

Insect Material and Bacterial Strains

A laboratory mass rearing of *E. variegatus* present at the DISAFA laboratories was used as a source of healthy adult leafhoppers for this work. Leafhoppers were kept on oat plants (*Avena sativa* L.) in growth chambers at 25°C with a photoperiod of 16:8 (L:D). Three groups of *E. variegatus* individuals were used for our experiments: healthy adults, specimens at the early stage of phytoplasma colonization, and individuals chronically infected by FDp. The first group was directly collected from the lab colony, while the second and the third groups were obtained by exposing insects to broad beans (*Vicia faba* L.) infected by FDp (strain FD-C). Adults at the early FD phytoplasma infection stage (EFDi) were chosen immediately after being submitted to a 5-day acquisition access period. Late FDp-infected (LFDi) *E. variegatus* were obtained by rearing nymphs on infected broad beans until adult emergence and for at least 21 days (Figure 1).

Bacterial colonization experiments in *E. variegatus* were performed using the spontaneous rifampicin-resistant mutant strains SF2.1Rif^R and SF15.14Rif^R of *Asaia*, the first being a non-ALI forming strain, and the latter an ALI-producer isolate that was reported to reduce FDp acquisition in *E. variegatus* (Gonella et al., 2018). *Escherichia coli* strain DH5 α pKan(DsRed) (Crotti et al., 2009) was also used as a non-symbiotic control. Before use, the strains of *Asaia* were cultivated overnight at 30°C in GLY

medium (Favia et al., 2007) under the selection of rifampicin (100 μ g/ml), whereas *E. coli* DH5 α pKan(DsRed) was cultured overnight at 37°C in Luria-Bertani (LB) medium under the selection of kanamycin (50 μ g/ml).


Bacterial Colonization of *E. variegatus* Individuals

Asaia strains SF2.1 Rif^R and SF15.14 Rif^R, as well as *E. coli* DH5 α pKan(DsRed), were individually administered to *E. variegatus* adults from the healthy, EFDi, and LFDi groups, following the protocol described by Crotti et al. (2009). Briefly, cultivated bacterial cells were harvested by centrifugation (10 min, 800 g), washed three times with 0.9% NaCl, and adjusted to 10⁸ cells/ml in 5% (w/v) sucrose solution in TE (10 mM Tris HCl, 1 mM EDTA, pH 8) (Crotti et al., 2009; Gonella et al., 2012). Insects were allowed to feed for 48 h on artificial feeding systems (Gonella et al., 2015) containing the cell suspensions. Specimens directly collected from the mass rearings without any further treatment were used as reference samples for the gene expression analysis, while adults maintained in the artificial feeding systems with added sterile sugar solution were taken as a control. Five insects for each group and treatment (total number of specimens: 75), corresponding to 5 biological replicates, were directly collected and stored at -80°C in RNA later[®] (Sigma-Aldrich, St. Louis, MO, United States) until RNA extraction, while further individuals were submitted to midgut dissection.

Midgut Dissection and Establishment of Hemocyte Cell Cultures

To obtain midgut samples, *E. variegatus* individuals were dissected in Phosphate Buffered Solution (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 2 mM KH₂PO₄; pH 7.4) made with Diethyl pyrocarbonate-treated water, by using sterile forceps under a stereomicroscope. Five replicates for each insect group and treatment, each one consisting of midguts from three adults, were collected and stored at -80°C in RNA later[®] (Sigma-Aldrich, St. Louis, MO, United States) until RNA extraction. Dissected midguts from *E. variegatus* adults directly collected from our lab colony without any further treatment were used as reference samples, and the midguts of insects fed with a sterile sugar diet were employed as a control.

Primary hemocyte cell cultures were established following the method described by Tedeschi et al. (2017). Specifically, groups of two adults were washed in 0.115% sodium hypochlorite, 75% ethanol and MilliQ sterile water for 10, 30, and 20 s, respectively, and then dried on a filter paper for a couple of seconds. Washed insects were placed in a single well of a sterile 24-well cell culture plate (Costar[®], Corning[®], NY, United States) containing 1 ml of Hert-Hunter 70 medium (Marutani-Hert et al., 2009), supplemented with 10 ml/L L-glutamine (Invitrogen, Carlsbad, CA, United States). Gentamicin (Sigma-Aldrich, St. Louis, MO, United States) at a final concentration of 50 μ g/ml, penicillin/streptomycin (Sigma-Aldrich, St. Louis, MO, United States) at a final concentration of 50 U/ml and 50 μ g/ml, respectively, and the antimycotic agent nystatin (Sigma-Aldrich, St. Louis, MO, United States) at a final



concentration of 100 U/ml, were also added. Plates were incubated at 24 – 26°C for 24 h, to allow cell establishment in the medium, and subsequently incubated for further 24 h with 10⁸ cells/ml bacterial suspension [*Asaia* SF2.1 Rif^R and SF15.14 Rif^R, and *E. coli* DH5α pKan(DsRed)] in the same medium deprived of antibiotics, after removal of the hemocytes culture supernatant. For each of healthy, EFDi and IFDi sample groups, five replicates were treated with bacteria, five replicates were incubated with a sterile antibiotic-free medium in the absence of bacteria as a control, and five replicates were left untreated to be used as the reference samples for gene expression analyses. At the end of treatments, cells were centrifuged at 800 g for 5 min at room temperature and immediately subjected to RNA extraction after discarding the supernatant.

RNA Extraction, cDNA Synthesis, and Quantitative Real Time PCR (qPCR)

RNA extraction was performed with the “SV Total RNA Isolation System” (Promega, Fitchburg, WI, United States), according to the supplier’s suggestions. Briefly, insect tissues were lysed and homogenized with a sterile pestle in 175 μl RNA Lysis Buffer; then samples were heated at 70°C for 3 min after adding 350 μl of RNA Dilution Buffer. Cleared lysate solutions were obtained by centrifugation, and subsequently

provided with 200 μl 95% ethanol and transferred in the supplied Spin Column Assembly. Once samples were washed with RNA Wash Solution, they were incubated for 15 min at room temperature with DNase incubation mix, then 200 μl of DNase Stop Solution were added to stop the reaction. Finally, samples were washed twice with RNA Wash Solution and resuspended in 50 μl nuclease-free water. After extraction, RNA quality and concentration were assessed with a ND-1000 spectrophotometer (NanoDrop, Wilmington, DE, United States) and by electrophoresis on a denaturing agarose gel (**Supplementary Figure S1**). First strand cDNA was synthesized by using the “Reverse Transcription System” (Promega) and Random Primers with 9 μl of RNA, following the manufacturer’s instructions. cDNA was used as a template for qPCR analysis with primer pairs specifically targeting the following genes: *defensin*, *Raf*, *phenoloxidase*, and *kazal type 1 serine protease inhibitor*. A list of primers is presented in **Table 1**. Raf-specific primers were specifically designed on conserved sequences identified by the alignment of Raf sequences of *D. citri* (XM_008488867.2), *Nilaparvata lugens* (Stål) (XM_022347672.1) and *Acyrtosiphon pisum* (Harris) (XM_001952258.4) using the on-line software Primer 3¹. A preliminary survey was conducted by amplification of cytoplasmic actin with the following parameters: initial

¹<http://bioinfo.ut.ee/primer3/>

TABLE 1 | Primers used in this study.

Primer pair	Target gene	Sequence (5' → 3')	Size (bp)	Source
EvDef F	Defensin	ATGCATTCTCCATTACTGCTG	200	Tedeschi et al., 2017
EvDef R		CAGCTGCCTCCCTTCTTGC		
Raf F	Raf	CAAGTGGAGAGGATTACAGCAG	200	This study
Raf R		GTGTGTTGGAGCCAGGTCTAT		
PO2_F1020	Phenoloxidase	CAATGTGGTCCCTCAGGAT	115	Galetto et al., 2018
PO2_R1085		CTGCGAGGTCTCATTTCTGT		
kaz1_F70	Kazal type 1 serine protease inhibitor	CTGGTTCGCAGGCCAAATACC	103	Galetto et al., 2018
kaz1_R172		GGCATGACACTCGGTACACT		
actF	Actin	AGCAGGAGATGGCCACC	300	Tedeschi et al., 2017
actR		TCCACATCTGCTGGAAGG		
fAY	16S rRNA	GCACGTAATGGTGGGCACCTT	300	Marcone et al., 1996
rEY		CGAAGTTAAGCCACTGCTTTTC		

denaturation at 95°C for 3 min, then 50 cycles consisting of denaturation at 95°C for 30 s, annealing at 58°C for 40 s and elongation at 72°C for 45 s. Even if a possible influence of immune challenge on actin expression was reported (de-Morais et al., 2005), stable actin expression was observed among sample groups and treatments (**Supplementary Table S2**), as previously observed by Capone et al. (2013) and Tedeschi et al. (2017). Consequently, *actin* was selected to be amplified as a reference gene. Moreover, samples belonging to the EFDi and LFDi groups were quantitatively checked for FDP infection by 16SrV group phytoplasma-specific PCR reactions, conducted with the fAY/rEY primer pair as described by Galetto et al. (2005); only positive samples were considered in this study. qPCRs were performed on a CFX Connect™ Real-Time PCR Detection System (Bio-Rad, Hercules, CA, United States). Reactions were conducted in clear HardShell® Low-Profile 96-Well PCR Plates (Bio-Rad) with a 25 µl mixture containing 12.5 µl of 2 × SsoFast™ EvaGreen® Supermix (Bio-Rad), 0.1 µl of each primer (100 mM), 100 ng of sample cDNA and 11.3 µl of double distilled H₂O, sealed with adhesive Microseal® PCR Plate Sealing Film (Bio-Rad); samples were analyzed in triplicate. An initial denaturation at 95°C for 3 min was followed by 50 cycles consisting of denaturation at 95°C for 30 s and annealing at 54°C (for qPCR targeting defensin and Raf) or 58°C (for qPCR targeting phenoloxidase and kazal type 1 serine protease inhibitor) for 20 s. A final step for melting curve analysis from 70 to 95°C, measuring fluorescence every 0.5°C, was added. Results were analyzed using the CFX Manager™ Software (Bio-Rad) for Ct determination. Normalization of primer efficiency was obtained by the one-point calibration (OPC) method, according to Brankatschk et al. (2012); normalized efficiencies of the target genes, with respect to the standard, ranged between 96 and 101%. Relative quantification of target genes was calculated using the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen, 2001). Statistical analyses were performed with SPSS Statistics 25 (IBM Corp. Released 2017, Armonk, NY, United States). One-way analysis of variance (ANOVA) was applied and means separated by a Tukey's test ($P < 0.05$) when variance homogeneity was satisfied (Levene test, $P < 0.05$).


Evaluation of Strain Adhesion Capacity

Adhesion capacity of strains *Asaia* SF2.1Rif^R, *Asaia* SF15.14Rif^R, *E. coli* DH5α pKan(DsRed), and the positive control *E. coli* ATCC 25404 was evaluated by crystal violet staining assay (Barbato et al., 2016). *Asaia* strains were grown overnight in GE medium (2% glucose, 0.8% yeast extract, pH 7), while *E. coli* strains were cultured overnight in LB medium [added with 50 µg/ml kanamycin in case of *E. coli* DH5α pKan(DsRed)]. Following overnight growths, 200 µl of the bacterial suspensions containing 10⁶ cell/ml of the different strains were transferred to a flat bottom, polystyrene microtiter plate. Eight replicates were inoculated for each strain and eight uninoculated controls were prepared, as well. Strains were grown at 24 or 30°C for 48 or 72 h. Following the incubation time, the optical density at 610 nm (OD 610 nm) was measured. Bacterial cultures were then removed and microtiter-adhering cells were gently washed with PBS three times. The microtiter plate was dried for 15 min and then stained with a solution of crystal violet (0.5 g/L crystal violet in 20% ethanol) for 15 min at room temperature. Finally, crystal violet solution was removed, the microtiter plate was washed twice with distilled water and let dry for 15 min. Crystal violet contained in adhering cells was then solubilized in 96% ethanol by pipetting. Absorbance (OD 610 nm) of solubilized crystal violet was measured by using a microtiter reader (Tecan Infinite F200Pro).

RESULTS

Expression of Immune Genes



Prior to measuring the expression of our target immune genes, cDNA from samples belonging to the EFDi and LFDi groups was subjected to phytoplasma quantification to confirm their infection status. We measured the average number of 16SrV phytoplasma cells per sample, which ranged between 1.04×10^2 , detected in hemocytes, and 5.54×10^4 found in midguts. No significant differences were observed between treatments within the same sample origin, according to ANOVA performed on log-transformed values (whole insects: df = 7, 32, $F = 1.466$,






$P = 0.215$; midguts: $df = 7, 32, F = 1.042, P = 0.422$; hemocytes: $df = 7, 32, F = 1.220, P = 0.321$). In the EFDi and LFDi sample groups, only the phytoplasma-positive samples were used for the immune gene expression analysis. Normalized relative quantities of immune genes considered in this study are reported in **Supplementary Table S3** and **Figures 2–4**. In the whole bodies of *E. variegatus* adults, the levels of transcripts showed little variability for all genes after supply of either bacteria, in each of the healthy, EFDi, and LFDi groups. Only in a few cases were significant differences observed in the expression values (**Supplementary Table S4**): defensin was overexpressed in healthy specimens fed with *E. coli* DH5α pKan(DsRed) (**Figure 2**), while phenoloxidase was activated after administering *Asaia* SF2.1Rif^R to individuals of the EFDi group (**Figure 3**). No differences were detected in LFDi insects (**Figure 4**). On the other hand, when considering midgut samples, significantly different transcript levels of *Raf* gene were found in the samples exposed to phytoplasmas (EFDi and LFDi groups) (**Figures 3, 4**). In both cases, *Raf* was overexpressed in the midguts belonging to specimens fed with *Asaia* SF15.14Rif^R relative to the control, consisting of leafhoppers reared in the presence of phytoplasmas without bacteria. *Kazal type 1* gene was found to be upregulated as well, although only in EFDi midguts. In this case, the transcripts related to insects fed with *E. coli* DH5α pKan(DsRed) were significantly more abundant than those detected in the presence of *Asaia* SF15.14Rif^R. In the case of hemocyte samples, most significant differences were observed for the expression levels

of *kazal type 1* gene. Strikingly, upregulation was observed in healthy (**Figure 2**) and LFDi samples (**Figure 4**), whereas *kazal type 1* transcripts from the EFDi group were not significantly different, particularly when considering hemocytes treated with either bacteria (**Figure 3**). Moreover, in samples from the healthy group, *kazal type 1* was overexpressed after exposure to *Asaia* SF2.1Rif^R relative to *E. coli* DH5α pKan(DsRed) and the control, while in samples from the LFDi group the most abundant transcripts were detected in the presence of *Asaia* SF15.14Rif^R. Other significant differences were recorded in hemocyte samples: defensin was downregulated in samples treated with *Asaia* SF2.1Rif^R relative to the control in EFDi samples, and *Raf* was upregulated in samples exposed to *Asaia* SF15.14Rif^R relative to those provided with *Asaia* SF2.1Rif^R in samples from the LFDi group (**Figures 2–4**).

Besides comparing gene expression profiles obtained for different treatments and sample groups, data were further analyzed to evaluate the temporal expression trend of each gene with respect to phytoplasma infection. Hence, normalized values obtained for single genes were examined, considering the healthy group as time 0 with respect to FDP infection, the EFDi group as time 0 + 5 days from the beginning of FDP infection, and finally the LFDi group as time 0 + 21 days from the beginning of phytoplasma infection. Statistical analysis performed to compare data from similarly treated samples from the healthy, EFDi and LFDi groups revealed a growing expression trend only for *Raf* and *kazal type 1* genes (**Figures 5, 6**). Specifically, *Raf* was







overexpressed in the whole body of *E. variegatus* fed with *Asaia* SF2.1Rif^R as a consequence of chronic FDP infection (LFDi vs. healthy + EFDi), and in the midguts of insects provided with *Asaia* SF15.14Rif^R subsequent to FDP exposure (EFDi + LFDi vs. healthy) (Figure 5 and Supplementary Table S5). Likewise, *kazal type 1* gene was upregulated in samples from the LFDi group (LFDi vs. healthy + EFDi), and precisely in midguts of insects fed with *Asaia* SF2.1Rif^R and *E. coli* DH5α pKan(DsRed), and in hemocytes following treatment with *Asaia* SF15.14Rif^R (Figure 6 and Supplementary Table S5). Conversely, we did not find any significant trend in defensin or phenoloxidase in consequence of bacterial challenge (Figures 7, 8 and Supplementary Table S5). Instead, phenoloxidase was significantly overexpressed over time in hemocytes from the control group (Figure 8).

Adhesion Test


Adhesion capacity of *Asaia* strains was evaluated in comparison to the one shown by a well-known biofilm-producer strain, i.e., *E. coli* ATCC 25404, which was considered as positive control (Wood et al., 2006). *Asaia* SF2.1Rif^R and *E. coli* DH5α pKan(DsRed) weakly adhered to the polystyrene microtiter plate (Supplementary Figure S2A and Supplementary Table S6) in comparison with the positive control strain. Moreover, following a longer incubation (72 h) *E. coli* DH5α pKan(DsRed) displayed a decreasing adhesion capacity. *Asaia* SF15.14Rif^R was not able to adhere to the microtiter wells in our experimental conditions either considering incubation at 24 and 30°C, or 48 and 72 h. In wells inoculated with *Asaia* SF15.14Rif^R bacterial biomasses were

observed on the medium surface that resembled the ALI films already described to be produced by this strain (Supplementary Figure S2B): it is likely that the shorter incubation times used in our experiments, in comparison to those described in Gonella et al. (2018) did not allow the complete film formation observed before. Conversely, strain *E. coli* ATCC 25404 showed itself to be a strong adherent strain.

DISCUSSION

Our investigation focused on the differential regulation of immune genes chosen to represent a range of elements shaping the leafhopper immune response, revealing localized regulation in *E. variegatus* adults, for all of the four genes taken into consideration. Indeed, even if gene expression variability generally appeared mitigated in whole-body samples for most of the genes, in midgut and hemocyte samples, significantly diverging gene expression levels were observed, in particular for the genes *Raf* and *kazal type 1*.

Raf is the only gene that exhibited consistent upregulation in response to exposure to a single bacterial strain, being overexpressed in the presence of *Asaia* SF15.14Rif^R. Since this strain was reported by Gonella et al. (2018) to induce a reduced FDP acquisition in *E. variegatus*, this gene might be involved in enhancing the insect response to phytoplasma infection. Specifically, *Raf* gene was more specifically activated in midgut samples when the presence of *Asaia* SF15.14Rif^R was combined with FDP infection (EFDi and LFDi groups),



while the same gene was overexpressed in the hemocytes only at the late phase of phytoplasma infection. Our experimental evidence confirms that Raf expression is relevant in the immune response and in persistent infection and/or sepsis since Raf induction causes activation of the hematopoiesis (as reported in *Drosophila*) (Asha et al., 2003; Zettervall et al., 2004). Our observations also suggest that the midgut could be a crucial site for Raf activation, supporting a role for Raf in limiting the capability of FDP to colonize the gut, and crossing this barrier to reach the hemolymph. This indication is in accordance with previous results suggesting that gut homeostasis is maintained through a balance between cell damage, due to the collateral effects of bacteria killing, and epithelial repair by stem cell division (Buchon et al., 2009) and that Raf is involved in the control of *Drosophila* gut stem cell proliferation (Jin et al., 2015). Accordingly, in the midgut of insects fed with *Asaia* SF15.14Rif^R, a significant rise of Raf expression was recorded corresponding to the early stage of FDP infection, which is a crucial stage for phytoplasma passage from the midgut to the insect hemolymph.

In the case of kazal type 1 serine protease inhibitor, a hemocyte-specific response was observed following challenge with *Asaia*, while no significant activation relative to controls was observed in whole insects or dissected midguts following exposure to these strains. Nonetheless, different strains induced kazal type 1 overexpression in the hemocytes of healthy and LFDi individuals, namely strains SF2.1Rif^R and SF15.14Rif^R in the

first and in the latter groups, respectively, whereas no significant upregulation was found in the EFDi hemocytes. Accordingly, significantly increased expression of kazal type 1 was found in late phytoplasma infection (LFDi), in hemocytes exposed to *Asaia* SF15.14Rif^R (while an increasing trend over FDP infection time was observed for strain SF2.1Rif^R in midgut samples). Hence, a response in the hemocytes could be speculated to be related to genus-specific traits rather than to the production of ALI biofilm. Previous work experimentally combing *Asaia* strains with FDP infection did not show inhibition of phytoplasma transmission by strain SF2.1Rif^R (Gonella et al., 2018), suggesting the absence of genus-specific interference with the pathogen. Moreover, although FDP itself was demonstrated to induce *kazal type 1* gene overexpression (Galletto et al., 2018), this pathogen can bloom to high concentrations in *E. variegatus* (Rashidi et al., 2014), indicating that phytoplasmas may be only partially susceptible to such a response. Furthermore, no clear temporal trend was observed for *kazal type 1* gene overexpression in the control group, in none of sample types, even though a slight increment was visible for whole-insect samples, in agreement with the upregulation reported by Galletto et al. (2018). This is indicative of limited activation of this gene in our experimental conditions. Taken together, this evidence does not support a role for kazal type 1 activation in limiting FDP acquisition by *E. variegatus*.

The expression of *defensin* and *phenoloxidase* genes was not as affected as that of *Raf* and *kazal type 1* genes in

response to bacterial infection. Defensin was overexpressed in the whole body of insects fed with *E. coli* DH5 α pKan(DsRed) only considering healthy individuals. Conversely, considering the EFDi and LFDi insect groups and single-body parts, samples exposed to this strain did not exhibit higher defensin expression than those treated with *Asaia* strains or the control, confirming the limited activation of this AMP in *E. variegatus* in response to Gram-negative bacteria (Tedeschi et al., 2017). Relative defensin transcript levels were significantly reduced in the hemocytes of insects from the EFDi group, after challenge with *Asaia* SF2.1Rif^R. However, no further data sustained the downregulation of this gene operated by the non-ALI producer strain. In previous experiments performed with hemocyte cultures of *Anopheles stephensi* Liston and *Drosophila melanogaster* Meigen, the administration of a DsRed-tagged strain of *Asaia* SF2.1 and *E. coli* DH5 α pKan(DsRed) did not induce the expression of *defensin* gene (Capone et al., 2013). However, it is worth considering that a higher load of bacterial cells (10⁹ cell/ml) and shorter incubation times (0, 4, 8, 12 h) than the ones used in our experimental set up were used by Capone et al. (2013). Taking into account the tendency of defensin expression over FDP infection-time, statistical analysis did not demonstrate any significant variation. However, an increase in the expression levels was recorded in the midgut of control insects, never exposed to bacteria other than the phytoplasma. This is in agreement with evidence reported by Yang et al. (2017), showing that defensin is activated by insect infection with *Spiroplasma melliferum*, a close relative of phytoplasmas recognized as a model for phytoplasma infection (Naor et al., 2011). On the other hand, when analyzing the FDP infection temporal trend in the control hemocytes, enhanced expression was detected in correspondence to the early stage of pathogen infection, suggesting hemocyte-specific defensin stimulation, consistently with the key role of the hemolymph for phytoplasma multiplication, resulting in a higher load of phytopathogen cells, which in turn stimulated an insect response. However, our data suggest an overall limited role of defensin in altering the capacity of *E. variegatus* to acquire FDP.

Phenoloxidase exhibited mostly irregular expression profiles, without significant diversity of transcript levels corresponding to distinct bacterial treatments according to ANOVA analysis, with the only exception being EFDi whole insects. Such an erratic response suggests the absence of a specific induction mechanism related to a challenge with any of administered bacterial strains. Limited phenoloxidase activation in response to *Asaia* infection could be expected considering its symbiotic interplay with leafhoppers (Crotti et al., 2009). On the other hand, in *Anopheles* mosquitoes, the adaptation of *Asaia* to host body environment was suggested to be related to resistance to immune response, rather than to a low immunogenicity (Capone et al., 2013); however, these authors did not investigate phenoloxidase activation. Similarly, although being a non-symbiotic strain, *E. coli* DH5 α pHM2(GFP) did not induce phenoloxidase-related responses in leafhoppers, as previously reported for *E. coli*-infected hemocytes of *C. haematoceps* (Eliautout et al., 2016). However, phenoloxidase was shown to be significantly activated in the hemocytes during the early stage of phytoplasma infection,

as a transcript peak was detected for EFDi samples in the control group. This result supports the hypothesis put forward by Galetto et al. (2018) that an increment in the expression of phenoloxidase might be evident in the early stages of FDP infection, whereas a lack of activation is typical of *E. variegatus* individuals chronically infected by phytoplasma.

Besides demonstrating that *Asaia* SF15.14Rif^R elicited in *E. variegatus* a midgut-specific immune response, this work explored an alternative process that may modulate the interference exerted by this strain on FDP, i.e., physical exclusion by means of complete adhesion to the gut wall. However, the adhesion test did not show successful adhesion to microtiter plates for *Asaia* SF15.14Rif^R. This result might be affected by the experimental conditions that were applied in terms of incubation times or medium used. The artificial substrate may indeed imperfectly mimic the insect gut epithelia. As a matter of fact, in Hemiptera the gut content is in contact with the microvilli of midgut cells, in the absence of a peritrophic membrane (Nation, 2016). Therefore, the increased contact surface provided by microvilli may offer a more suitable substrate for bacterial adhesion. Acetic acid bacteria closely related to *Asaia* were reported to be specifically located near the host gut wall (Kounatidis et al., 2009; Vacchini et al., 2017); however, our results do not support massive attachment of *Asaia* SF15.14Rif^R to the gut epithelial layer. A possible explanation of limited attachment could be the production of flocculant flake-like bacterial masses (**Supplementary Figure S2B**). These masses have previously been proposed to play a role in entrapping phytoplasma cells or erecting a barrier against them, contributing to hampering pathogen establishment in the insect body (Gonella et al., 2018). Moreover, we cannot rule out the involvement of specific host factors under *in vivo* conditions necessary to mediate the bacterial adhesion. Future work is thus required to investigate the role of bacterial entrapment or physical containment in limiting FDP acquisition by *E. variegatus*.

CONCLUSION

This work sheds light on the molecular interplay occurring among insect hosts and bacteria, focusing on symbiotic and non-symbiotic strains, as well as on leafhopper-vectored plant pathogens. Among different immune-related genes indicative of distinct response mechanisms, *Raf* gene showed midgut-specific activation in response to *Asaia* strain SF15.14Rif^R after insect infection by FDP, while the pathogen alone did not stimulate the same reaction. It can therefore be suggested that *Asaia* strain SF15.14Rif^R may elicit a basal host immune activity that appears to act against other microorganisms, including phytoplasma. The elicitation of *Raf*-mediated response is thus predicted to be a major component of the interfering effect displayed by *Asaia* SF15.14Rif^R toward FDP acquisition by *E. variegatus*. Considering the limited number of immune genes reported for *E. variegatus* and other Hemiptera (Arp et al., 2016; Skidmore and Hansen, 2017; Tedeschi et al., 2017), our results suggest a compensation of lost defensive functions by genes that have minor functions in insects exhibiting a more complete immunity.

Investigating the role of other immune genes will help to further elucidate this phenomenon, as well as exploring possible differential response profiles in different conditions, such as for example when insects are exposed to different temperatures. On the other hand, a minor contribution to phytoplasma exclusion seems to be provided by *Asaia* attachment to the vector gut wall. Further possible mechanisms of phytoplasma segregation, involving a physical barrier created by *Asaia* SF15.14Rif^R through the production of ALI biofilm, need to be investigated.

AUTHOR CONTRIBUTIONS

EG, MM, RT, EC, and AA contributed to conception and design of the study. EG, EC, and MP carried out the experiments. EG performed the statistical analysis. AA, RT, and EC provided reagents and analytical tools. EG, MM, EC, and RT wrote sections of the manuscript. All authors contributed to manuscript revision, read and approved the submitted version.

REFERENCES

- Alma, A., Tedeschi, R., Lessio, F., Picciau, L., Gonella, E., and Ferracini, C. (2015). Insect vectors of plant pathogenic mollicutes in the euro-mediterranean region. *Phytopathogen Mollicutes* 5, 53–73. doi: 10.5958/2249-4677.2015.00063.8
- Arp, A. P., Hunter, W. B., and Pelz-Stelinski, K. S. (2016). Annotation of the Asian citrus psyllid genome reveals a reduced innate immune system. *Front. Physiol.* 7:570. doi: 10.3389/fphys.2016.00570
- Asha, H., Nagy, I., Kovacs, G., Stetson, D., Ando, I., and Dearolf, C. R. (2003). Analysis of Ras-induced overproliferation in *Drosophila* hemocytes. *Genetics* 163, 203–215.
- Barbato, M., Mapelli, F., Magagnini, M., Chouaia, B., Armeni, M., Marasco, R., et al. (2016). Hydrocarbon pollutants shape bacterial community assembly of harbor sediments. *Mar. Pollut. Bull.* 104, 211–220. doi: 10.1016/j.marpolbul.2016.01.029
- Baumann, P. (2005). Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Ann. Rev. Microbiol.* 59, 155–189. doi: 10.1146/annurev.micro.59.030804.121041
- Bosco, D., Galetto, L., Leoncini, P., Saracco, P., Raccach, B., and Marzachi, C. (2007). Interrelationships between “*Candidatus* Phytoplasma asteris” and its leafhopper vectors (Homoptera: Cicadellidae). *J. Econom. Entomol.* 100, 1504–1511. doi: 10.1603/0022-0493-100.5.1504
- Bosco, D., and Marzachi, C. (2016). “Insect transmission of phytoplasmas,” in *Vector-Mediated Transmission of Plant Pathogens*, ed. J. K. Brown (St. Paul, MN: APS Press), 319–327.
- Brankatschk, R., Bodenhausen, N., Zeyer, J., and Bürgmann, H. (2012). Simple absolute quantification method correcting for quantitative PCR efficiency variations for microbial community samples. *Appl. Environ. Microbiol.* 78, 4481–4489. doi: 10.1128/AEM.07878-11
- Bressan, A., Clair, D., Sémétey, O., and Boudon-Padieu, E. (2005). Effect of two strains of flavescence dorée phytoplasma on the survival and fecundity of the experimental leafhopper vector *Euscelidius variegatus* Kirschbaum. *J. Invertebr. Pathol.* 89, 144–149. doi: 10.1016/j.jip.2005.03.001
- Buchon, N., Broderick, N. A., Poidevin, M., Pradervand, S., and Lemaitre, B. (2009). *Drosophila* intestinal response to bacterial infection: activation of host defense and stem cell proliferation. *Cell Host Microbe* 5, 200–211. doi: 10.1016/j.chom.2009.01.003
- Capone, A., Ricci, I., Damiani, C., Mosca, M., Rossi, P., Scuppa, P., et al. (2013). Interactions between *Asaia*, *Plasmodium* and *Anopheles*: new insights into mosquito symbiosis and implications in malaria symbiotic control. *Parasit. Vectors* 6:182. doi: 10.1186/1756-3305-6-182
- Cassone, B. J., Michel, A. P., Stewart, L. R., Bansal, R., Mian, M. A. R., and Redinbaugh, M. G. (2014). Reduction in fecundity and shifts in cellular processes by a native virus on an invasive insect. *Genome Biol. Evol.* 6, 873–885. doi: 10.1093/gbe/evu057
- Crotti, E., Damiani, C., Pajoro, M., Gonella, E., Rizzi, A., Ricci, I., et al. (2009). *Asaia*, a versatile acetic acid bacterial symbiont, capable of cross-colonizing insects of phylogenetically distant genera and orders. *Environ. Microbiol.* 11, 3252–3264. doi: 10.1111/j.1462-2920.2009.02048.x
- de-Morais, G. S., Vitorino, R., Domingues, R., Tomer, K., Correia, A. J., Amado, F., et al. (2005). Proteomics of immune-challenged *Drosophila melanogaster* larvae haemolymph. *Biochem. Biophys. Res. Commun.* 328, 106–115. doi: 10.1016/j.bbrc.2004.12.135
- Eleftherianos, I., Atri, J., Accetta, J., and Castillo, J. C. (2013). Endosymbiotic bacteria in insects: guardians of the immune system? *Front. Physiol.* 4:46. doi: 10.3389/fphys.2013.00046
- Eliautout, R., Dubrana, M. P., Vincent-Monégat, C., Vallier, A., Braquart-Varnier, C., Poirié, M., et al. (2016). Immune response and survival of *Circulifer haematoceps* to *Spiroplasma citri* infection requires expression of the gene hexamerin. *Dev. Comp. Immunol.* 54, 7–19. doi: 10.1016/j.dci.2015.08.007
- Favia, G., Ricci, I., Damiani, C., Raddadi, N., Crotti, E., Marzorati, M., et al. (2007). Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc. Natl. Acad. Sci. U.S.A.* 104, 9047–9051. doi: 10.1073/pnas.0610451104
- Galetto, L., Abbà, S., Rossi, M., Vallino, M., Pesando, M., Arricaus-Bouvery, N., et al. (2018). Two phytoplasmas elicit different responses in the insect vector *Euscelidius variegatus* Kirschbaum. *Infect. Immun.* 86:e042-18. doi: 10.1128/IAI.00042-18
- Galetto, L., Bosco, D., and Marzachi, C. (2005). Universal and group-specific real-time PCR diagnosis of flavescence dorée (16Sr-V), bois noir (16Sr-XII) and apple proliferation (16Sr-X) phytoplasmas from field-collected plant hosts and insect vectors. *Ann. Appl. Biol.* 147, 191–201. doi: 10.1111/j.1744-7348.2005.00030.x
- Gonella, E., Crotti, E., Mandrioli, M., Daffonchio, D., and Alma, A. (2018). *Asaia* symbionts interfere with infection by “flavescence dorée” phytoplasma in leafhoppers. *J. Pest Sci.* 91, 1033–1046. doi: 10.1007/s10340-018-0973-1
- Gonella, E., Crotti, E., Rizzi, A., Mandrioli, M., Favia, G., Daffonchio, D., et al. (2012). Horizontal transmission of the symbiotic bacterium *Asaia* sp. in the leafhopper *Scaphoideus titanus* Ball (Hemiptera: Cicadellidae). *BMC Microbiol.* 12(Suppl. 1):S4. doi: 10.1186/1471-2180-12-S1-S4
- Gonella, E., Pajoro, M., Marzorati, M., Crotti, E., Mandrioli, M., Pontini, M., et al. (2015). Plant-mediated interspecific horizontal transmission of an intracellular symbiont in insects. *Sci. Rep.* 5:15811. doi: 10.1038/srep15811

FUNDING

This work was partially supported by the “INTEFLAVI” (Un approccio integrato alla lotta contro la flavescenza dorata della vite) project. Funding was provided by Fondazione Cassa di Risparmio di Cuneo.

ACKNOWLEDGMENTS

The authors are grateful to Luca Picciau and Giuseppe Galante for insect rearing and phytoplasma maintenance.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2019.00795/full#supplementary-material>

- Gonella, E., Tedeschi, R., Crotti, E., and Alma, A. (2019). Multiple guests in a single host: interactions across symbiotic and phytopathogenic bacteria in phloem-feeding vectors – a review. *Entomol. Exp. Appl.* 167, 171–185. doi: 10.1111/eea.12766
- Jin, Y., Ha, N., Forés, M., Xiang, J., Gläßer, C., Maldera, J., et al. (2015). EGFR/Ras signaling controls *Drosophila* intestinal stem cell proliferation via capicua-regulated genes. *PLoS Genet.* 11:e1005634. doi: 10.1371/journal.pgen.1005634
- Kliot, A., Cilia, M., Csosek, H., and Ghanim, M. (2014). Implication of the bacterial endosymbiont *Rickettsia* spp. in interactions of the whitefly *Bemisia tabaci* with tomato yellow leaf curl virus. *J. Virol.* 88, 5652–5660. doi: 10.1128/JVI.00071-14
- Kounatidis, I., Crotti, E., Sapountzis, P., Sacchi, L., Rizzi, A., Chouaia, B., et al. (2009). *Acetobacter tropicalis* is a major symbiont of the olive fruit fly (*Bactrocera oleae*). *Appl. Environ. Microbiol.* 75, 3281–3328. doi: 10.1128/AEM.02933-08
- Li, Z., An, X. K., Liu, Y. D., and Hou, M. L. (2016). Transcriptomic and expression analysis of the salivary glands in white-backed planthoppers, *Sogatella furcifera*. *PLoS One* 11:e0159393. doi: 10.1371/journal.pone.0159393
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ Ct method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- Login, F. H., Balmant, S., Vallier, A., Vincent-Monegat, C., Vigneron, A., Weiss-Gayet, M., et al. (2011). Antimicrobial peptides keep insect endosymbionts under control. *Science* 334, 362–365. doi: 10.1126/science.1209728
- Makarova, O., MacLean, A., and Nicolaisen, M. (2015). Phytoplasma adapt to the diverse environments of their plant and insect hosts by altering gene expression. *Physiol. Mol. Plant Pathol.* 91, 81–87. doi: 10.1016/j.pmp.2015.06.003
- Marcone, C., Ragozzino, A., Schneider, B., Lauer, U., Smart, C. D., and Seemüller, E. (1996). Genetic characterization and classification of two phytoplasmas associated with spartium witches'-broom disease. *Plant Dis.* 80, 365–371. doi: 10.1094/PD-80-0365
- Marutani-Hert, M., Hunter, W. B., and Hall, D. G. (2009). Establishment of Asian citrus psyllid (*Diaphorina citri*) primary cultures. *In Vitro Cell. Dev. Biol. – Animal* 45, 317–320. doi: 10.1007/s11626-009-9188-3
- Nachappa, P., Levy, J., Pierson, E., and Tamborindeguy, C. (2014). Correlation between “*Candidatus Liberibacter solanacearum*” infection levels and fecundity in its psyllid vector. *J. Invertebr. Pathol.* 115, 55–61. doi: 10.1016/j.jip.2013.10.008
- Naor, V., Ezra, D., and Zahavi, T. (2011). The use of *Spiroplasma melliferum* as a model organism to study the antagonistic activity of grapevine endophytes against phytoplasma. *Bull. Insectol.* 64, S265–S266.
- Nation, J. L. (2016). *Insect Physiology and Biochemistry*. Boca Raton, FL: CRC Press.
- Olson, K. E., and Blair, C. D. (2015). Arbovirus-mosquito interactions: RNAi pathway. *Curr. Opin. Virol.* 15, 119–126. doi: 10.1016/j.coviro.2015.10.001
- Rashidi, M., D'Amelio, R., Galetto, L., Marzachi, C., and Bosco, D. (2014). Interactive transmission of two phytoplasmas by the vector insect. *Ann. Appl. Biol.* 165, 404–413. doi: 10.1111/aab.12146
- Shao, E., Lin, G., Liu, S., Ma, X., Chen, M., Lin, L., et al. (2017). Identification of transcripts involved in digestion, detoxification and immune response from transcriptome of *Empoasca vitis* (Hemiptera: Cicadellidae) nymphs. *Genomics* 109, 58–66. doi: 10.1016/j.ygeno.2016.11.006
- Skidmore, I. H., and Hansen, A. K. (2017). The evolutionary development of plant-feeding insects and their nutritional endosymbionts. *Insect Sci.* 24, 910–928. doi: 10.1111/1744-7917.12463
- Tedeschi, R., Monti, M., Gonella, E., Melchiori, G., Alma, A., and Mandrioli, M. (2017). Molecular and cellular analysis of immunity in the phytoplasma vector *Euscelidius variegatus*: exploiting immunity to improve biological control strategies. *Inv. Surv. J.* 14, 63–72.
- Tsakas, S., and Marmaras, V. J. (2010). Insect immunity and its signalling: an overview. *Inv. Surv. J.* 7, 228–238. doi: 10.1111/j.1462-2920.2009.02048.x
- Vacchini, V., Gonella, E., Crotti, E., Prosdocimi, E. M., Mazzetto, F., Chouaia, B., et al. (2017). Bacterial diversity shift determined by different diets in the gut of the spotted wing fly *Drosophila suzukii* is primarily reflected on acetic acid bacteria. *Environ. Microbiol. Rep.* 9, 91–103. doi: 10.1111/1758-2229.12505
- Vyas, M., Fisher, T. W., He, R., Nelson, W., Yin, G., Cicero, J. M., et al. (2015). Asian citrus psyllid expression profiles suggest *Candidatus liberibacter asiaticus*-mediated alteration of adult nutrition and metabolism, and of nymphal development and immunity. *PLoS One* 10:e0130328. doi: 10.1371/journal.pone.0130328
- Weiss, B., and Aksoy, S. (2011). Microbiome influences on insect host vector competence. *Trends Parasitol.* 27, 514–522. doi: 10.1016/j.pt.2011.05.001
- Wood, T. K., González Barrios, A. F., Herzberg, M., and Lee, J. (2006). Motility influences biofilm architecture in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 72, 361–367. doi: 10.1007/s00253-005-0263-8
- Yang, D., Zha, G., Li, X., Gao, H., and Yu, H. (2017). Immune responses in the haemolymph and antimicrobial peptide expression in the abdomen of *Apis mellifera* challenged with spiroplasma melliferum CH-1. *Microb. Pathog.* 112, 279–287. doi: 10.1016/j.micpath.2017.10.006
- Zettervall, C. J., Anderl, I., Williams, M. J., Palmer, R., Kurucz, E., Ando, I., et al. (2004). A directed screen for genes involved in *Drosophila* blood cell activation. *Proc. Natl. Acad. Sci. U.S.A.* 101, 14192–14197. doi: 10.1073/pnas.0403789101

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Gonella, Mandrioli, Tedeschi, Crotti, Pontini and Alma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.