



Tolerance and Reduction of Chromium(VI) by *Bacillus* sp. MNU16 Isolated from Contaminated Coal Mining Soil

Neha Upadhyay¹, Kanchan Vishwakarma¹, Jaspreet Singh¹, Mitali Mishra^{1,2}, Vivek Kumar³, Radha Rani¹, Rohit K. Mishra², Devendra K. Chauhan⁴, Durgesh K. Tripathi² and Shivesh Sharma^{1,2*}

¹ Department of Biotechnology, Motilal Nehru National Institute of Technology Allahabad, Allahabad, India, ² Centre for Medical Diagnostic and Research, Motilal Nehru National Institute of Technology Allahabad, Allahabad, India, ³ Department of Biotechnology, Himalayan Institute of Biosciences, Swami Rama Himalayan University, Dehradun, India, ⁴ D D Plant Interdisciplinary Research Laboratory, Department of Botany, University of Allahabad, Allahabad, India

OPEN ACCESS

Edited by:

Radha Prasanna,
Indian Agricultural Research Institute,
India

Reviewed by:

Ernesto Garcia-Pineda,
University System of New Hampshire,
United States
Misao Itouga,
RIKEN Center for Sustainable
Resource Science (CSRS), Japan

*Correspondence:

Shivesh Sharma
shiveshs@mnnit.ac.in;
ssnvsharma@gmail.com

Specialty section:

This article was submitted to
Plant Microbe Interactions,
a section of the journal
Frontiers in Plant Science

Received: 27 December 2016

Accepted: 25 April 2017

Published: 22 May 2017

Citation:

Upadhyay N, Vishwakarma K, Singh J, Mishra M, Kumar V, Rani R, Mishra RK, Chauhan DK, Tripathi DK and Sharma S (2017) Tolerance and Reduction of Chromium(VI) by *Bacillus* sp. MNU16 Isolated from Contaminated Coal Mining Soil. *Front. Plant Sci.* 8:778. doi: 10.3389/fpls.2017.00778

The bacterium MNU16 was isolated from contaminated soils of coal mine and subsequently screened for different plant growth promoting (PGP) activities. The isolate was further identified by 16S rRNA sequencing as *Bacillus subtilis* MNU16 with IAA concentration ($56.95 \pm 0.43 \mu\text{g/ml}$), siderophore unit ($9.73 \pm 2.05\%$), phosphate solubilization ($285.13 \pm 1.05 \mu\text{g/ml}$) and ACC deaminase activity ($116.79 \pm 0.019 \mu\text{moles } \alpha\text{-ketobutyrate/mg/24 h}$). Further, to evaluate the metal resistance profile of bacterium, the isolate was screened for multi-metal resistance (viz. 900 mg/L for Cr, 600 mg/L for As, 700 mg/L for Ni and 300 mg/L for Hg). Additionally, the resistance pattern of *B. subtilis* MNU16 against Cr(VI) (from 50 to 300 mg/L) treatments were evaluated. An enriched population was observed at 0–200 mg/L Cr(VI) concentration while slight reductions were observed at 250 and 300 mg/L Cr(VI). Further, the chromium reduction ability at 50 mg/L of Cr(VI) highlighted that the bacterium *B. subtilis* MNU16 reduced 75% of Cr(VI) to 13.23 mg/L within 72 h. The localization of electron dense precipitates was observed in the TEM images of *B. subtilis* MNU16 which is might be due to the reduction of Cr(VI) to Cr(III). The data of fluorescence microscopy and flow cytometry with respect to Cr(VI) treatments (50–300 mg/L) showed a similar pattern and clearly revealed the less toxic effect of hexavalent chromium upto 200 mg/L Cr(VI) concentration. However, toxicity effects were more pronounced at 300 mg/L Cr(VI). Therefore, the present study suggests that the plant growth promoting potential and resistance efficacy of *B. subtilis* MNU16 will go a long way in developing an effective bioremediation approach for Cr(VI) contaminated soils.

Keywords: *Bacillus subtilis*, chromium(VI), propidium iodide, flow cytometry, fluorescence microscopy

Abbreviations: CFU, colony forming unit; EDP, electron dense precipitates; FCM, flow cytometry; FM, fluorescence microscopy; MHA, Mueller Hinton Agar; MIC, minimal inhibitory concentration; PI, propidium Iodide; TEM, transmission electron microscopy.

INTRODUCTION

Soil contamination and degradation are important issues from both environmental and agricultural points of view. The continual anthropogenic activities, extensive use of chemical fertilizers, pesticides and rapid industrialization are responsible for increasing environmental pollution (Aktar et al., 2009; Singh et al., 2015, 2017; Tripathi et al., 2016a). These activities are major causes of soil degradation and contamination and thus pose a great challenge for food safety and public health (Qishlaqi and Moore, 2007; Tripathi et al., 2012, 2015, 2016b). Soils of mining sites have extremely high percentage of heavy metals such as chromium, lead, cadmium, and arsenic which cause destruction of soil properties such as pH, nutrient content and quality (Ladwani et al., 2012; Hansda et al., 2014); thereby, significantly affecting the fertility of surrounding soil, associated microbial community, vegetation status as well as pose several health risks at different levels of food chain through a well-established process of bioaccumulation (Wuana and Okieimen, 2011; Upadhyay et al., 2016).

Heavy metals are the most dominant form of contamination found in the waste dump sites of mines and their remediation is considered to be a technically difficult task due to their recalcitrant nature (Wuana and Okieimen, 2011). There are a number of traditional and conventional technologies which are widely acceptable to remediate metal contaminated soils. However, these approaches are time consuming and usually generate complex secondary waste. In this context, our major concern is to reclaim and restore soil properties through implementation of situation-specific technique(s) that permit the best possible decontamination at the lowest potential cost with modest or no side effects. The application of plant-microbe interaction for the remediation of contaminated soil is an important and well-adapted technology (Hansda et al., 2014). Plant growth is regulated by soil fertility which influences the physical (e.g., soil structure), chemical (e.g., soil pH, organic matter, C: N ratio) and biological (e.g., microbial diversity in soil) properties of soil. On the other hand, most of the time, adverse conditions of degraded soils results in the destruction of aboveground vegetation. Rhizobacteria generally colonize root and enhance plant growth in degraded soils, as they can alleviate biotic and abiotic stress by releasing phytohormones (e.g., IAA, ethylene), solubilizing minerals and producing iron chelating compounds (e.g., siderophores) (Ahemad, 2015). Several studies have reported the utilization of plant growth promoting (PGP) rhizobacteria for the bioremediation of heavy metals including *Bacillus* sp., *Pseudomonas* sp. etc. (Samuel et al., 2013; Sobariu et al., 2016; Ndeddy Aka and Babalola, 2016).

Among various heavy metals, chromium (Cr) is one of the most common pollutants originating mainly from industrial effluents, mining practices, petroleum refining industries and use of pesticides (Oliveira, 2012). In nature, chromium mainly exists in trivalent [Cr (III)] form and found naturally in environment. Hexavalent [Cr (VI)] is mainly released in environment through various anthropogenic activities and imparts toxic effects on microorganisms, human beings and soil fertility (Oliveira, 2012; Tripathi et al., 2012; Ahemad, 2015). It is well documented

that bacterial composition is greatly affected by soil type, its characteristics and stress factors which restrict the bacterial growth in soil (Upadhyay et al., 2016). Only a few bacteria may adapt to these stressed environments and proliferate. The transition into this physiological state and recovery were found to be correlated with the environmental stress equivalent to pH, temperature and bioavailability of minerals and nutrients (Olszewska et al., 2016).

The behavior of micro-organisms toward Cr(VI) was studied previously which demonstrated the significant alleviative impact of many microorganisms, including strains of *Pseudomonas*, *Enterobacter*, *Bacillus*, and *Shewanella* (Samuel et al., 2013; Ahemad, 2015). Microorganisms were found to reduce hexavalent chromium by different mechanisms either by utilizing the hexavalent chromium as final electron acceptor or by secreting certain soluble enzymes (Samuel et al., 2013; Ahemad, 2015). However, the availability of efficient hexavalent chromium reducing microorganisms is a crucial requirement for the bioremediation of soils contaminated with hexavalent chromium. With this background, the present study is envisaged to isolate and select the most resistant bacterial isolate from the contaminated soils with potential to reduce hexavalent chromium. In addition, the bacterial growth and morphological modifications were examined against various Cr(VI) treatments using modern techniques so that it can be utilized further for the effective bioremediation in associated with the enhancement in plant growth promotion when grown in contaminated soil.

MATERIALS AND METHODS

Isolation and Characterization of Bacteria for PGP Traits

Samples were collected in triplicates from the degraded soils of coal mining site of Uttar Pradesh. Nutrient agar medium was used to isolate bacterial colonies by serial dilution technique and plates were observed after 24 h of incubation. Colonies were further purified by streak plate and preserved at 4°C for further experiments.

Plant growth promoting characteristics of isolates were further identified by the standard procedure discussed below. The bacterial isolates were screened for indole-3-acetic acid (IAA) production using L-tryptophan and quantified colorimetrically by Salkowski method (Loper and Scroth, 1986). The production of siderophore was initially screened on Chrome Azurol S (CAS) agar medium (Schwyn and Neilands, 1987) and further estimated quantitatively by CAS shuttle assay as per described by Payne (1994). An equal volume of culture supernatant was mixed with CAS reagent and absorbance was measured at 630 nm against the reference containing equal volume of uninoculated medium and CAS reagent. Percentage siderophore produced was calculated by using the formula:

$$\% \text{ siderophore units} = (Ar - As/Ar) \times 100$$

Where Ar = absorbance of the reference and As = absorbance of the sample.

Bacterial isolates were further screened for phosphate solubilizing potential first qualitatively by growing isolates on Pikovskaya's medium, and isolates showing clear zone were selected as potential phosphate solubilizer (Premono et al., 1996). Bacterial isolates that have the highest solubilization index were selected for the quantitative analysis done by using Pikovskaya Broth and estimated spectrophotometrically using chlorostannous-reduced-molybdo-phosphoric acid method (Yadav and Verma, 2012).

The ACC deaminase activity of the isolates was further determined by the method of Saleh and Glick (2001). The concentration of α -ketobutyrate produced by the action of ACC deaminase was determined spectrophotometrically and calculated with the help of standard curve. The amount of α -ketobutyrate produced is expressed as μ moles α -ketobutyrate/mg/24 h.

Molecular Characterization of Potential Isolate

The genomic DNA was isolated from the bacteria by enzymatic method using phenol/chloroform (Maniatis et al., 1982). The DNA isolated was resuspended in TE buffer (pH 8.0) and amplified by PCR to obtain the 16S rRNA sequences. Universal primers, 8F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-AAGGAGGTGATCCAGCCGCA-3') were used for DNA amplification. The PCR product was purified using QIAquick PCR Purification Kit (Qiagen). Purified PCR fragments were directly sequenced with Applied Biosystems, 3500XL Genetics Analyzer using the manufacturer's instructions. Chromas Lite 2.0 software was used to read the sequences and the sequences were then aligned with the previous sequences available in the GenBank database by BLAST tool. A phylogenetic tree was created by the use of multiple sequence alignment tool, ClustalW and MEGA 6.06 software.

Screening of Bacterial Isolates for Chromium(VI) Resistance

The selected bacterium was further screened for its resistance to chromium (Cr VI) by agar well diffusion method (Hassen et al., 1998). In addition the bacterium was also screened for resistance to various metals including Iron (Fe), Copper (Cu), Arsenic (As), Mercury (Hg), Cadmium (Cd) and Nickel (Ni). Various concentrations of metals were prepared by using their salts as: $K_2Cr_2O_7$ for Cr(VI), $NaAsO_2$ for As, $HgCl_2$ for Hg, $CdCl_2$ for Cd, $FeCl_3$ for Fe and $Ni(NO_3)_2$ for Ni in distilled water and sterilized by autoclaving. Sterile MHA plates were prepared and spreaded with the overnight grown bacterial culture. After this, approximate 7 mm diameter wells were punched with a sterile borer and filled with metals (100 μ l) with various concentrations to determine the value of MIC. The plates were then incubated at 30°C for 24–48 h and zones around the wells were measured. Metal concentration which gave a clear zone of 1 mm or less than 1 mm was considered as the MIC and the bacterium resistant for that particular concentration (Rani et al., 2010).

Screening of Isolate for Cr (VI) Reduction

The ability of bacterial isolate to reduce Cr(VI) into less toxic form was analyzed using diphenylcarbazide (DPC) method by estimating the decrease in concentration of hexavalent chromium (Zahoor and Rehman, 2009). The 24 h old grown culture was inoculated in 100 ml of LB broth containing 50 mg/L of Cr(VI) as $K_2Cr_2O_7$ at 30°C to estimate the chromium reduction. The samples were collected by centrifugation at 10,000 rpm for 10 min. The remaining concentration of Cr(VI) in the supernatant was determined by measuring the absorbance of Cr(VI)-DPC complex at 540 nm using spectrophotometer at various time intervals. The percentage reduction of Cr(VI) was calculated by using the following formula:

$$\text{Cr(VI) reduction (\%)} = \frac{A-B}{B} \times 100$$

Where A- Absorbance of control; B- Absorbance of sample.

Transmission Electron Microscopy (TEM) Analysis

The bacterial strain was analyzed through the TEM to confirm the presence and accumulation of hexavalent chromium inside the cell and to detect the effect of metal on the bacterium. Samples were prepared according to the method described by Bano et al. (2013) with slight modifications. The bacterial cell without any treatment served as control and various concentrations of Cr(VI) were taken as treatments for TEM analysis. The 48 h grown bacterial culture was harvested and pellet was washed with 1 M phosphate buffer saline (PBS) buffer. The cells were fixed using fixative (2.5% glutaraldehyde and 2% paraformaldehyde in 1 M PBS) for 6 h at 4°C and then washed three times with PBS to remove the fixative solution. The pellet was then suspended in 1 M PBS buffer and sent to SAIF, AIIMS, New Delhi for TEM analysis.

Assessment of Physiological Changes in Bacteria under Different Cr(VI) Treatments

Culture Preparations

The bacteria were inoculated in nutrient broth medium at 28–30°C overnight under continuous shaking. Samples were collected from the bacterial suspension at exponential phase and the concentration was adjusted to nearly 2×10^6 bacteria/ml (Walberg et al., 1997). The pellet of bacterial cells were then collected by centrifugation at 8000 rpm for 5 min, washed thrice with 0.1 M phosphate buffered saline (PBS) and resuspend the pellet in PBS to reach an optical density of 0.1 at 600 nm.

Exposure to Chromium Treatment

The bacterial cells were analyzed for their resistance pattern against various concentrations of chromium (VI) by studying CFU pattern and performing growth profile study by UV spectrometry, FM and FCM. The metal stress was generated by method described by Boswell et al. (1998). Potassium dichromate ($K_2Cr_2O_7$) salt solution was prepared for generating chromium stress at concentration starting from 50, 100, 150, 200, 250,

and 300 mg/L and the bacterial suspension was then exposed to different concentration of Cr(VI). A control was prepared without any metal exposure. The resistance pattern and bacterial responses in the presence of various Cr(VI) treatments were studied after 12 h of incubation.

Fluorescence Spectroscopy and Flow Cytometric Measurements

The bacterial suspension was further stained with PI, a nucleic acid staining dye, and analyzed through FM and FCM. After treatment, 10 μ l PI (40 μ g/ml) was added to 200 μ l of bacterial suspension in the centrifuge tube and incubated for 15 min in dark to allow staining of the cells. The cells were immediately analyzed after the incubation period by FM and FCM.

Flow cytometry study was carried out using a BD Accuri™ C6 Flow Cytometer with a red laser of 14.7 mW output and a constant wavelength excitation of 640 nm and a blue laser of 20 mW with excitation at a wavelength of 488 nm. The FCM instrument detects forward scatter (FSC) and side scatter (SSC) and is equipped with four different types of fluorescence detectors with optical filters. The red color (FL2, 585 nm; PE/PI) fluorescence detector was used for the current flow cytometric study. The optical detectors employed to collect the scattered laser light and fluorescent emissions, and electronics digitize these signals for computational analysis. The light scatter data gives a basic idea regarding cells relative size, and morphology. The fluorescence data reveals the cells' auto fluorescence and/or labeling with fluorescent dyes, which can help characterize bacteria, resolve them from electronic noise and debris, and indicate cell viability and vitality. A detection limit of 10,000 bacteria was set for each sample and all experiments were conducted in triplicate.

Statistical Analysis

The data obtained from the study were statistically analyzed and presented with the appropriate standard deviation from the data obtained in triplicates. The data were standardized by one-way analysis of variance (ANOVA). The least significant differences among means were compared at $P \leq 0.05$ significance level. The data analysis and preparation of graphs were done by using the software OriginPro 8.3.

RESULTS

Isolation and Characterization of Bacteria to Test Plant Growth Promoting Activity

A total of 26 bacterial isolates were isolated from the degraded soils of coal mines and further characterized for their growth promoting characteristics. Bacterial isolate MNU16 was found to be positive for all screened PGP characteristics and proved to be the most effective producer of maximum growth promoting substances and therefore, selected for further studies. Five isolates were recorded as being efficient IAA producers, out of which highest IAA production was shown by isolate MNU16 with an

TABLE 1 | Details of plant growth promoting traits produced by *Bacillus subtilis* MNU16 strain.

Plant growth promoting characteristics	Results
IAA production (μ g/ml)	56.95 \pm 0.436 ^a
Siderophore production (%)	9.73 \pm 2.05 ^b
Phosphate solubilization (μ g/ml)	285.13 \pm 1.05 ^c
ACC deaminase activity (μ moles α -ketobutyrate/mg/24 h)	116.79 \pm 0.019 ^e
Cr(VI) reduction %	75.39% \pm 0.97 ^{dc}

IAA, Indole 3-acetic acid. Results are average of data in triplicates \pm SD (standard deviation). Mean with different letters in the same column differ significantly at $P \leq 0.05$ (One-way ANOVA) significance level.

IAA concentration of 56.95 \pm 0.436 μ g/ml (Table 1) after 48 days of incubation.

Siderophore production was further checked by plate assay and the isolates that were able to produce an orange halo zone around the bacterial colony were selected as siderophore producers. The production of siderophore by isolate MNU16 was first observed in plate assay and then further quantified for siderophore production. On quantifying, it was observed that the isolate MNU16 produced 9.73 \pm 2.05% siderophore in succinate medium (Table 1).

The bacterial strain MNU16 was also assessed for ACC deaminase activity, and 116.79 \pm 0.019 μ moles α -ketobutyrate/mg/24 h (Table 1) activity was recorded. The phosphate solubilization activity of the bacterial strain MNU16 was observed by the presence of clear halo zone around the Pikovskaya's agar plate. It was also observed that the phosphate solubilization zone increased with increasing incubation time around the bacterial colony. The bacterial isolate was used to further quantified for phosphate solubilization potential. A significant increase in phosphate solubilization was observed concomitant with the decrease in pH, as the period of incubation increased from 24 to 120 h (Figures 1A,B). The quantitative estimation of phosphate solubilization is presented in Figure 1A. The bacterium was found to solubilize 81.75 μ g/ml of phosphate after 24 h of incubation and maximum solubilization (285.13 \pm 1.05 μ g/ml) was recorded after 120 h of incubation. A continuous decline in the medium pH was observed from 6.5 to 4.1 which is presented in Figure 1B. It was clearly observed that the initial pH of the medium was 6.5 which decreased to 5.5 in 24 h and after 120 h, the pH of the medium was found to be 4.1, which is due to the production of several organic acids that are responsible for phosphate solubilization.

Identification of Isolate through Molecular Characterization

On the basis of 16S rRNA gene sequence, isolate MNU16 showed maximum similarity with *Bacillus* spp. The *B. subtilis* MNU16 sequences were submitted to Genbank database with Accession No. KY012360. A phylogenetic tree (Figure 1C) was constructed using MEGA 5.0 by comparing it with other 16S rRNA sequences of related species and the isolate showed maximum similarity with *Bacillus* sp. (Accession No. KT150178.1).




FIGURE 1 | (A) Quantitative solubilization of phosphate by *Bacillus* sp. MNU16. **(B)** Lowering of pH in the broth due to P-solubilizing activity of *Bacillus* sp. MNU16. Data are average \pm standard deviation of experiments in triplicate. **(C)** Phylogenetic tree, based on 16S ribosomal RNA gene sequences, showing relationships between MNU16 and other *Bacillus* sp. **(D)** Chromium [Cr (VI)] reduction by *Bacillus subtilis* MNU16 at 120 rpm and 30°C. **(E)** Growth of *B. subtilis* MNU16 in the presence of various concentration of chromium at various time intervals. Data are average \pm standard deviation of experiments in triplicate.

Identification of Minimal Inhibitory Concentration of Cr(VI)

The bacterial isolate was screened for chromium (Cr) and resistance to other heavy metals (As, Fe, Hg, Ni, Cd, and Cu) by agar well diffusion method (Supplementary Figure S1). The bacterial isolate MNU16 was found to be resistant at 900 mg/L chromium concentration. The MIC for Cd was recorded upto 150 mg/L, 300 mg/L for Hg, 600 mg/L for As, 700 mg/L for Ni, 1000 mg/L for Fe and 500 mg/L for Cu, respectively (Supplementary Figure S1). The order of toxicity of metals for the strain MNU16 was observed as highest for cadmium > mercury > arsenic > nickel > chromium > iron.

Bacillus subtilis MNU16 under Different Chromium Treatments

The growth profile of bacterium *B. subtilis* MNU16 was performed at different Cr(VI) treatments to study the growth pattern of isolate at different time intervals. In addition, the

isolate *B. subtilis* MNU16 was also studied for its potential to reduce hexavalent chromium [Cr(VI)] in nutrient broth medium having initial chromium concentration of 50 mg/L (Figure 1D). Further, it was observed that the isolate reduced the concentration of chromium upto 13.23 mg/L within 72 h of incubation at 30°C and shaking at 120 rpm (Figure 1D). The growth kinetics of bacterial isolate in the presence of chromium is shown in Figure 1E which illustrated that the bacterium shows higher growth at chromium concentrations 50, 100, 150, and 200 mg/L. However, the absorbance reduced at higher concentration and bacterial growth got affected at 250 and 300 mg/L as shown in Figure 1E.

Physiological Modulations in *Bacillus subtilis* MNU16 under Different Cr Treatments

Bacterial growth was observed in the presence of Cr stress by studying the plate count and growth profile of isolate (Figure 1E).

The overnight grown bacterial culture was inoculated into the medium having 0 mg/L (control), 50, 100, 150, 200, 250, and 300 mg/L Cr(VI) concentrations, respectively, and was plated onto the nutrient agar medium after 12 h of incubation. The plate count assessment is considered as an initial screening of bacterial viability and the result showed that some morphological changes in bacterial cells were observed upon spreading the bacterial culture after metal exposure. The CFU count was also recorded highest in medium without metal concentration (control) and found to decrease with increasing metal concentration. The results showed that cells have lost their viability in presence of metal concentration which is in agreement with the results of FCM. The bacterial colony was small, pin point, circular at 0, 50, and 100 mg/L chromium concentration, whereas at 150, 200, 250, and 300 mg/L chromium concentration, the colonies were observed to be little elevated and whitish-yellow. It was also concluded from the study that less toxic effect of metal was observed at 50 and 100 mg/L resulting in alteration in colony morphology of bacteria at higher metal concentration. In addition, the physiological changes in bacteria were also studied by TEM (Figure 2), FM (Figure 3), and FCM in the presence of different chromium treatments (Figures 4, 5).

Transmission Microscopy of *Bacillus subtilis* MNU16

To identify the effect of hexavalent chromium inside cells of *B. subtilis* MNU16, TEM analysis was performed. The modifications due to Cr treatments (0, 100, 200, and 300 mg/L) in bacterial cells have been indicated by pointing out the arrow in the Figures 2A–D. The results signified that in control where no Cr treatment was provided, the bacterial cells have a smooth surface (Figure 2A). Modification in cell surface were recorded with increasing concentration of Cr(VI) (Figures 2B–D). Results further suggested that, once the bacterial cells get exposed in the presence of Cr(VI), the size of bacterium increased and an irregular shape and rough surface were observed as compared to control cells. The effect of Cr(VI) at high concentration (300 mg/L) initiates cell lysis which might be due to the deposition of chromium in the cytoplasm and on the surface of the cells (Figures 2C,D).

Cell Viability Assessment by Fluorescence Microscopy

The bacterium *B. subtilis* MNU16 was subjected to different chromium treatments (Supplementary Figure S2). The bacterial viability was assessed qualitatively using FM by staining bacterial cells with PI. PI is most often used as cell death marker due to its exclusion by the cell membrane of live cells; therefore, the fluorescence bestowed by the dye is usually related to cells whose membrane integrity was damaged. The PI inclusion by bacterial cells shows the percentage of bacterial death at different stress conditions. When the cells were stained with PI (red fluorescence), dead cells acquired the stain and appeared as red color spots when observed by FM (Figure 3). Results shows the images of bacterial isolates exposed to various concentrations of chromium ranged from 50 to 300 mg/L, as observed by FM. It

is clearly observed from the result that very few populations of bacterial cells were stained in control; however, the population of stained cells increased with increasing concentrations of chromium (Figure 3).

Effect of Cr Treatments on Cell Size and Granularity of *B. subtilis*

Chromium is used to generate stress in bacterial cells and the effect of chromium at various concentrations on cell size and granularity is shown in Figure 4. It was observed that by increasing concentration of metal, cell size and granularity of bacteria increases as shown in the figure below (Figure 4). The dot plot of *B. subtilis* MNU16 in the presence of chromium(VI) treatment at various concentrations shows that with increasing the concentration of metal, a shift in population was observed and clearly visible at concentration 100, 150, 200, 250, and 300 mg/L defining the changes in FSC and SSC; therefore, it is concluded that the relative size and granularity of the bacterial cells was affected due to the chromium treatment (Figures 4A–G).

Effect of Cr Treatments on Membrane Integrity and Cell Viability of *B. subtilis*

The cell membrane of all the microorganisms allows the cells to interact with the surrounding environment. The cell membrane is selectively permeable and controls the transport of molecules within the cell. To quantify or visualize dead cells by studying the membrane integrity of cells, nucleic acid stains are most commonly used because cell contains relatively high percentage of nucleic acids. Therefore, dead cells acquires the stain by binding with nucleic acids and shows enhancement in fluorescence which aids in quantifying the cell viability during stressed environment. The membrane integrity of *B. subtilis* MNU16 was checked by capturing fluorescently stained cells with PI. In current study, PI was used to discriminate between healthy stable cells and cells with ruptured membrane. PI, a small hydrophilic molecule, is impermeable to the membrane of viable cells, therefore excluded by viable cells and penetrates the membrane of dead cells having damaged membranes and intercalates double stranded DNA and emits fluorescence. It was observed that the mean fluorescence intensity is higher for the cells having ruptured membrane which is proved to be beneficial for a proper discrimination of dead and viable cells in FCM.

In the present study, cells when treated with chromium give high PI fluorescence intensities indicating the percentage cell death in the presence of chromium. Respective dot plots showing the percentage cell death in bacterial cells after Cr(VI) treatment are presented in Figure 5A. When bacterial cells are treated with chromium salt, the cytoplasmic membrane potential (represents the difference of electric potential between inside and outside of cell) may get affected and the cytoplasmic membrane permeability gets destroyed. Passive and active transport systems across the cytoplasmic membrane of healthy bacterial cells generate an electrochemical gradient, giving a measure of cellular metabolic activity. Very less intensity was observed at 50 mg/L concentration, whereas a significant increase was recorded with increasing metal concentration. However, at concentration




FIGURE 2 | Transmission electron microscopy (TEM) images of cross sectioned *B. subtilis* MNU16 cells at (A) control (without treatment). (B) *B. subtilis* MNU16 cells at 100 mg/L of Cr(VI) (C) *B. subtilis* MNU16 cells at 200 mg/L of Cr(VI) and (D) at 300 mg/L of Cr(VI) concentration. Arrows signifies SCS, smooth cell surface; RF, rough surface; BF, binary fission; EDP, electron dense precipitates; CL, cell lysis.

300 mg/L, the intensity was reduced a bit as compared to low metal concentration. The percentage of cell death (Figure 5A), the relative PI fluorescence (Figure 5B) and cell count versus relative PI fluorescence graph (Figure 5C) recorded in the cells in control and treatments are shown and it is concluded that resistance pattern of bacteria increases with increasing chromium concentration. In control, gaiting was done in a way so that no cell death is recorded and in comparison with the control, least cell death (2.8%) (Figure 5A) was recorded in 50 mg/L Cr(VI) concentration and cells that were exposed to highest concentration of chromium gave maximum fluorescence with highest percentage of cell death (38.1%) (Figure 5A) indicating the maximum toxic effect of chromium at 300 mg/L; however, percentage cell death was approximately low in the concentration 50, 100, and 200 mg/L (Figure 5A) and increases above 200 mg/L Cr(VI) concentration.

DISCUSSION

Contamination of soil with chromium can modulate the soil microbial diversity and metabolic activity. In addition,

nutrient deficiency was also reported in metal contaminated soils which often causes stress for plants (Qishlaqi and Moore, 2007; Ladwani et al., 2012). Till now, several PGP rhizobacteria have been identified such as *Azotobacter*, *Bacillus*, *Pseudomonas*, *Azospirillum*, and *Serratia* for their potential to enhance plant growth by solubilizing minerals, producing iron chelating compounds (siderophores), phytohormones and biocontrol compounds and enhancing phytoremediation of metal contaminants (Hansda et al., 2014; Ndeddy Aka and Babalola, 2016; Sarathambal et al., 2017). Chromium [Cr(VI)] exerts toxic effect on microorganisms by different mechanisms. Chromium resistant bacteria can overcome toxic effects of Cr(VI) and have been utilized for bioremediation of contaminated sites (Samuel et al., 2013; Sobariu et al., 2016). It has been reported that several metal resistant PGP rhizobacteria enhance metal translocation and improve metal uptake efficiency (Hansda et al., 2014).

Isolation and utilization of PGP rhizobacteria for rhizoremediation of heavy metal contaminated soils have gained worldwide attention because it is an environmental friendly and cost-effective approach. Therefore, the current study deals with the isolation and characterization of chromium-resistant-PGP




FIGURE 3 | Fluorescence microscopy images of *B. subtilis* stained with PI incubated at different concentration of chromium.

bacteria capable of reducing Cr(VI) to Cr(III). The PGP characteristics were screened and the potential PGP isolate was further screened for chromium resistance and tolerance potential. The reduction of Cr(VI) to Cr(III) was also studied and subsequently the morphological and physiological responses were checked to study the resistant pattern of bacterial strain in the presence of various chromium concentrations to evaluate its impact on bioremediation potential.

The results of the present study confirm those from previously reported literature on *Bacillus* sp. regarding its ability to produce PGP substances and promote plant growth (Shukla et al., 2012; Cherif-Silini et al., 2013; Ndeddy Aka and Babalola, 2016). Production of a plant growth regulator, such as IAA, is an important characteristic of PGP rhizobacteria (Saharan and Nehra, 2011). In the present study, production of IAA was reported by the test isolate and the concentration of IAA (Table 1) produced was in agreement with other studies (Karnwal, 2009; Shukla et al., 2012). However, the intrinsic potential of bacteria to produce growth hormone such as IAA in the plant rhizosphere greatly relies on the accessibility of precursors and uptake of microbial IAA by plant (Arshad and Frankenberger, 1991).

The potential of the bacteria to solubilize the insoluble tricalcium phosphate *in vitro* is an additional significant method for accomplishing plant growth promotion (Saharan and Nehra, 2011). The present study shows that the isolate MNU16 solubilizes maximum phosphate ($285.13 \pm 1.05 \mu\text{g/ml}$) after 120 h of incubation, and with increasing incubation time, the pH of the medium declined to 4.1 from an initial pH of 6.4 as shown in Figures 1A,B. The result draws support from the previous studies which highlighted the solubilization of phosphate by producing several types of organic acids by

different *Bacillus* species and other bacterial species along with the subsequent drop in medium pH (Shukla et al., 2012; Cherif-Silini et al., 2013). Besides, phosphate solubilization and IAA production, the isolate possess another important characteristic of PGP activity, i.e., ACC deaminase activity, in the bacterial isolates which are known to mitigate plant stress by reducing ethylene concentration in plants (Karthikeyan et al., 2012). ACC deaminase activity is an important property that diminishes plant stress caused by adverse environmental stresses such as presence of high metal concentration. ACC deaminase converts ACC, which serves as an ethylene precursor in plants, to ammonia and α -ketobutyrate and thus reduces the plant stress caused by higher ethylene concentration (Karthikeyan et al., 2012). A significant concentration of α -ketobutyrate was produced by the selected isolate in the present study (Table 1). It proves that bacteria possess the attribute of phytoremediation by facilitating plants to attain higher and dense roots in chromium contaminated soils.

Siderophore molecules exert both growth promotion and biocontrol mechanisms and have significantly high affinity for iron (Saharan and Nehra, 2011). Consequently, the low availability of iron in the surrounding environment of plant rhizosphere could suppress the growth of pathogenic microorganisms (Patil et al., 2014). In the present study, the bacterium isolated shows a highly orange halo zone in plate assay which confirms siderophore production. Therefore, it was studied quantitatively for estimation of siderophore, which was significantly lower than the values previously reported for PGP strains (Patil et al., 2014; Dutta et al., 2015).

The MIC of the bacteria for chromium was obtained by agar well diffusion method in Muller Hinton agar medium with Cr(VI) concentration ranging from 50 to 1000 mg/L




FIGURE 4 | Representative dot-plots of cell size (FSC) and cell complexity (SSC) at various concentration of chromium (A) 0 mg/L; (B) 50 mg/L; (C) 100 mg/L; (D) 150 mg/L; (E) 200 mg/L; (F) 250 mg/L; (G) 300 mg/L; and (H) relative size and (I) granularity of *B. subtilis* MNU16 at different Cr(VI) concentrations. Data are average \pm standard deviation of experiments in triplicate. Bars followed by various letter(s) showed significant difference at $P < 0.05$ significance level according to single factor ANOVA.

and the bacterial isolate MNU16 was found to be highly resistant for chromium with a concentration of 900 mg/L (Supplementary Figure S1). In addition, the present isolate was also found to be resistant for various other metals (Supplementary Figure S1). The previous studies reported the tolerance range of chromium resistant microorganism from 100 to 4000 mg/L Cr(VI) (Thacker et al., 2007; Zhu et al., 2008). The isolated *Bacillus* sp. MNU16 falls within the above reported range of tolerance. From the molecular characterization and phylogenetic analysis of the test isolate, it was found that the test isolate MNU16 showed maximum similarity with the *B. subtilis* species (Figure 1C). Not many isolates from the genus *Bacillus* have been known to tolerate and reduce Cr (VI) (Ibrahim et al., 2011; Mangaiyarkarasi et al., 2011). This difference in resistance profile may be because of the variation in medium composition used for studying the chromium toxicity which influences the toxicity via 'masking' effect (Caravelli et al., 2008).

In the present study, *B. subtilis* MNU16 shows chromium reduction to a concentration of 13.23 mg/L (Figure 1D) from

an initial concentration of 50 mg/L within 72 h of incubation. The reduction efficiency of Cr(VI) for *Bacillus* sp. MNU16 was evaluated and shown in Figure 1D and the percentage (%) reduction of Cr(VI) was recorded upto 75% by the bacterium *Bacillus* sp. MNU16 (Table 1). The results are in accordance with the previous study which also showed reduction of chromium by *Bacillus* sp. (Samuel et al., 2013) and the Cr(VI) reduction percentages lies within the previously reported studies (Maqbool et al., 2015). This inherent ability of many bacterial species to resist and reduce chromate both under field and laboratory conditions have been previously studied for remediation of chromium contaminated environments (Ibrahim et al., 2011; Maqbool et al., 2015). However, the exact mechanism that is responsible for the reduction of Cr(VI) to Cr(III) is not completely known, and various explanations have been given previously (Ahemad, 2015). The possible mechanism behind the conversion of Cr(VI) to Cr(III) can be either when chromate serve as a final electron acceptor to acquire energy, or the bacteria may secrete some waste product which can possibly reduce Cr(VI) to Cr(III) or by the secretion of several enzymes that




FIGURE 5 | (A) Graph represents dot plots of mean FL2-A which shows the PI inclusion (% cell death) at various chromium (VI) concentrations ranged from 0 mg/L (control), 50, 100, 150, 200, 250, and 300 mg/L. **(B)** Overview of changes in cell membrane permeability (PI fluorescence) with increasing Cr(VI) concentration **(C)** Overlay graph of cell count versus PI fluorescence with increasing Cr(VI) concentration. Data are average \pm standard deviation of experiments in triplicate. Bars followed by various letter(s) showed significant difference at $P < 0.05$ significance level according to single factor ANOVA.

might act on Cr(VI) and reduce its toxicity by converting it into Cr(III) (Ahemad, 2015).

Microorganisms when subjected to heavy metals become susceptible to the toxicity caused by reactive oxygen species. Due to the PGP and metal resistance properties, the isolates can be used to remediate metal contaminated soil. The PGP activities of microorganisms are one of the reasons behind the survival of microorganism in metal contaminated soils. The siderophore producing microorganisms have the potential to bind with metals present in surrounding and therefore enhance the bioavailability of metals in plant rhizosphere (Sarathambal et al., 2017). In general, the phytoremediation process of contaminated soils will enhance with the increase in plant growth and metal bioavailability (Ahemad, 2015).

Before studying the physiological response of *Bacillus* sp. MNU16 by FCM, the bacterial response in the presence of Cr(VI) was also studied in the liquid medium by studying the growth pattern of isolate under various chromium concentrations. The growth profile of *Bacillus* sp. MNU16 is shown in **Figure 1E**.

It was clearly observed from growth profile (**Figure 1E**) that the absorbance at 300 mg/L chromium concentration was less which indicated that chromium showed highest toxicity at concentration 300 mg/L as compared to other concentrations. The toxic effect of metal was effective in the liquid medium (Shakoori et al., 2000) as the mobility of metals is comparatively more in liquid medium as compared to solid medium. The reduced growth observed in the presence of higher concentrations of Cr(VI) was attributed to the reduction of metabolic rate of bacterium by Cr(VI) (Pal and Paul, 2004).

The application of electron microscopy to examine the morphology of cells and modification in cell structure due to exposure to various contaminants can provide novel dimensions. The results obtained from TEM analysis of *B. subtilis* MNU16 demonstrated the allocation of FDP inside and outside cells which might be due to the exposure of hexavalent chromium treatments. Similarly, study conducted by Batool et al. (2014) reported similar pattern of FDP and cell disruption in cells of *P. aeruginosa* and *Ochrobactrum intermedium* treated with

different concentrations of Cr(VI). Further, in the present study, the deposition of EDP inside the *B. subtilis* MNU16 might be due to the accumulation and intracellular reduction of Cr(VI) as shown in **Figures 2C,D**. The study of TEM also showed that the toxicity of Cr(VI) is maximum at 300 mg/L Cr(VI) concentration as shown in **Figure 2D**. The intracellular reduction mechanism of Cr(VI) was reported previously on the strains of *Acinetobacter* sp. and *Shewanella oneidensis* which explains the deposition of EDP inside the cells (Daulton et al., 2007; Srivastava and Thakur, 2007). Thus the presence of these EDP inside cells clearly revealed the modification mechanisms of Cr(VI) into Cr(III) in the form of hydroxyl and carboxyl groups (Bruins et al., 2000; Bencheikh et al., 2007).

The resistance pattern of the test isolate *Bacillus* sp. MNU16 was further studied by the flow cytometric technique in the presence of different chromium concentrations. FCM method has been previously studied for analyzing bacterial physiology when exposed to various stresses (Ionescu et al., 2015; Olszewska et al., 2016) as well as for screening antibiotic susceptibility in bacteria and fungi (Wright et al., 2006). This technique is more suitable for studying physical and chemical characteristics of bacteria due to its ability to rapidly and efficiently analyze large number of microorganisms with better accuracy. The current study reports the use of PI, an exclusion dye used for the analysis of cell viability and complexity by FCM analysis of *B. subtilis* MNU16, during chromium treatments to evaluate its effect on the membrane permeability and eventually on cell viability. The relative sizes and internal complexity of bacterial cells were estimated by flow cytometric analysis by estimating the forward scatter (FSC) and side scatter (SSC) of light, respectively, and shown in **Figures 4A–I**. The results showed that with increasing chromium concentration, a shift in FSC and SSC of bacteria was observed (**Figure 4**). The forward scatter (FSC) is affected by the size of bacteria and the intensity of scatter depends on the refractive index of the particle and flow medium; although, for particles having size in the range of bacteria were found to be correlated with the total volume of bacterial cell (Baatout et al., 2007). The side scatter of light (SSC) is proportional to cell granularity or internal complexity. The reflected and refracted light was measured at several interfaces of bacterial cell where alteration in refractive index was observed and also aids in detecting the presence of inclusion bodies generated due to the environmental stresses (Baatout et al., 2007).

The mean FL2 graph showing the relative fluorescence due to PI staining of *B. subtilis* MNU16 grown in the presence of various chromium concentration given in **Figure 5B**, describing the comparison of cell count and fluorescence intensity at various Cr(VI) concentrations. It is clear from the **Figure 5C** that high cell count and less fluorescence intensity was observed in the control sample where no stress was provided and eventually a decrease in cell count was recorded when growing the cells in the presence of different Cr(VI) concentrations.

The study shows that a significant population of bacterial species is present in the medium at high concentration of chromium which shows *B. subtilis* MNU16 has chromium resistance potential. The noticeable increase in PI fluorescence

intensity with increasing Cr(VI) concentration (**Figure 5**) shows the enhancement in loss of membrane integrity with the increasing concentration of chromium. Furthermore, in the presence of chromium treatments, variation in cell death was observed by varying the concentration of chromium with the help of FM. The presence of maximum number of cells stained with PI showed the highest cell death in the sample which is correlated with the data obtained from the PI fluorescence of FCM as shown in **Figure 5**. These results are confirmed by the data obtained by fluorescence imaging of bacteria that gives fluorescence when PI intercalates with bacterial DNA which shows the increase in membrane permeability due to membrane damage.

Furthermore, the results obtained from the study confirm the fact that the most crucial target for any environment stress is cell membrane whether it is pH, heavy metal or temperature (Mangaiyarkarasi et al., 2011; Olszewska et al., 2016). This suggests that differences in properties of cell membrane can possibly be an essential factor in establishing the metal tolerance potential between various bacterial strains. On the other hand, the nature of Cr(VI) toxicity as a function of membrane composition and structure in determining tolerance of metal and other environmental stresses still needs to be clarified.

CONCLUSION

The main focus of the study was to isolate an efficient chromium resistant PGP rhizobacterium having the potential to tolerate chromium stress which can be utilized for bioremediation purpose. The selected *B. subtilis* MNU16 strain shows significant PGP characteristics highlighting resistance to high concentration of chromium and able to decrease the toxicity of Cr(VI) by reducing it to Cr(III), a less toxic form. The resistance pattern against chromium was studied by analyzing the cell viability percentage by FCM and FM. FCM presents the probability of performing both qualitative and quantitative evaluation established on the simultaneous measurements of structural and practical parameters of single cell. The study concluded that the isolated PGP rhizobacterial strain with significant Cr(VI) reduction potential provides new possibilities to design strategies for improving the efficacy of rhizoremediation (microbial assisted phytoremediation) of contaminated soils.

AUTHOR CONTRIBUTIONS

NU, KV, JS, RM, SS, MM, and DT designed experiments. NU, KV, RM, and JS performed experiments. NU, KV, VK, RR, DC, RM, and DT, analyzed data and wrote the manuscript. DT, DC, and SS critically evaluated the manuscript.

ACKNOWLEDGMENTS

Authors are thankful to Department of Biotechnology, Govt. of India for providing financial assistance and Director MNNIT Allahabad for providing necessary facilities. The services and

facilities provided by Centre for Medical Diagnostic and Research, MNNIT Allahabad and MHRD sponsored project, “Design Innovation Centre” are also acknowledged. The authors are also thankful to AIIMS, New Delhi for providing the TEM facility.

REFERENCES

- Ahemad, M. (2015). Enhancing phytoremediation of chromium-stressed soils through plant-growth-promoting bacteria. *J. Genet. Eng. Biotechnol.* 13, 51–58. doi: 10.1016/j.jgeb.2015.02.001
- Aktar, W., Sengupta, D., and Chowdhury, A. (2009). Impact of pesticides use in agriculture: their benefits and hazards. *Interdiscip. Toxicol.* 2, 1–12. doi: 10.2478/v10102-009-0001-7
- Arshad, M., and Frankenberger, W. T. (1991). Microbial production of plant hormones. *Plant Soil* 133, 1–8. doi: 10.1007/BF00011893
- Baatout, S., Leys, N., Hendrickx, L., Dams, A., and Mergeay, M. (2007). Physiological changes induced in bacteria following pH stress as a model for space research. *Acta Astronaut.* 60, 451–459. doi: 10.1016/j.actastro.2006.09.012
- Bano, L., Kumar, R. V., and Hameed, A. (2013). In vitro bactericidal and TEM study of the interaction of Ag-modified titania with coliform bacteria. *Int. J. Environ. Health Res.* 23, 407–418. doi: 10.1080/09603123.2012.743115
- Batool, R., Hasnain, S., Batool, R., and Yrjälä, K. (2014). Impact of environmental stress on biochemical parameters of bacteria reducing chromium. *Braz. J. Microbiol.* 45, 573–583. doi: 10.1590/S1517-83822014000200028
- Bencheikh, L. R., Obratsova, A., Mackey, M., Ellisman, M., and Tebo, B. (2007). Toxicity of Cr(III) to *Shewanella* sp. strain MR-4 during Cr(VI) reduction. *Environ. Sci. Technol.* 41, 214–220. doi: 10.1021/es0622655
- Boswell, C. D., Hewitt, C. J., and Macaskie, L. E. (1998). An application of bacterial flow-cytometry - evaluation of the toxic effects of 4 heavy-metals on *Acinetobacter* sp. with potential for bioremediation of contaminated wastewaters. *Biotechnol. Lett.* 20, 857–863. doi: 10.1023/A:1005363525882
- Bruins, M. R., Kapil, S., and Oehme, W. (2000). Microbial resistance to metals in the environment. *Ecotoxicol. Environ. Saf.* 45, 198–207. doi: 10.1006/eesa.1999.1860
- Caravelli, A. H., Giannuzzi, L., and Zaritzky, N. E. (2008). Reduction of hexavalent chromium by *Sphaerotilus natans* a filamentous micro-organism present in activated sludges. *J. Hazard. Mater.* 156, 214–222. doi: 10.1016/j.jhazmat.2007.12.014
- Cherif-Silini, H., Silini, A., Ghoul, M., Yahiaoui, B., and Arif, F. (2013). Solubilization of phosphate by the *Bacillus* under salt stress and in the presence of osmoprotectant compounds. *Afr. Microbiol. Res.* 7, 4562–4571.
- Daulton, T. L., Little, B. J., Jones-Meehan, J., Blom, D. A., and Lawrence, F. (2007). Microbial reduction of chromium from the hexavalent to divalent state. *Geochim. Cosmochim. Acta* 71, 556–565. doi: 10.1016/j.gca.2006.10.007
- Dutta, J., Handique, P. J., and Thakur, D. (2015). Assessment of culturable tea rhizobacteria isolated from tea estates of Assam, India for growth promotion in commercial tea cultivars. *Front. Microbiol.* 6:1252. doi: 10.3389/fmicb.2015.01252
- Hansda, A., Kumar, V., Anshumali, A., and Usmani, Z. (2014). Phytoremediation of heavy metals contaminated soil using plant growth promoting rhizobacteria (PGPR): a current perspective. *Recent Res. Sci. Technol.* 6, 131–134.
- Hassen, A., Saidi, N., Cherif, M., and Boudabous, A. (1998). Resistance of environmental bacteria to heavy metal. *Bioresour. Technol.* 64, 7–15. doi: 10.1016/S0960-8524(97)00161-2
- Ibrahim, A. S. S., El-Tayeb, M. A., Elbadawi, Y. B., and Al-Salamah, A. A. (2011). Bioreduction of Cr (VI) by potent novel chromate resistant alkaliphilic *Bacillus* sp strain KSUCr5 isolated from hypersaline Soda lakes. *Afr. J. Biotechnol.* 10, 7207–7218.
- Ionescu, R., Măruțescu, L., Tănase, A.-M., Chiciudean, I., Csutak, O., Pelinescu, D., et al. (2015). Flow cytometry based method for evaluation of biodegradative potential of *Pseudomonas fluorescens*. *Agric. Agric. Sci. Procedia* 6, 567–578. doi: 10.1016/j.aaspro.2015.08.088
- Karnwal, A. (2009). Production of indole acetic acid by fluorescent *Pseudomonas* in the presence of L-tryptophan and rice root exudates. *J. Plant Pathol.* 91, 61–63.
- Karthikeyan, B., Joe, M. M., Islam, M. R., and Sa, T. (2012). ACC deaminase containing diazotrophic endophytic bacteria ameliorate salt stress in *Catharanthus roseus* through reduced ethylene levels and induction of antioxidative defense systems. *Symbiosis* 56, 77–86. doi: 10.1007/s13199-012-0162-6
- Ladwani, K. D., Manik, V. S., and Ramteke, D. S. (2012). Assessment of heavy metal contaminated soil near coal mining area in Gujarat by toxicity characteristics. *Int. J. Life Sci. Biotechnol. Pharma Res.* 1, 73–80.
- Loper, J. E., and Scroth, M. N. (1986). Influence of bacterial sources of indole-3-acetic acid on root elongation of sugar beet. *Phytopathology* 76, 386. doi: 10.1094/Phyto-76-386
- Mangaiyarkarasi, M. M., Vincent, S., Janarthanan, S., Rao, T. S., and Tata, B. V. R. (2011). Bioreduction of Cr(VI) by alkaliphilic *Bacillus subtilis* and interaction of the membrane groups. *Saudi J. Biol. Sci.* 18, 157–167. doi: 10.1016/j.sjbs.2010.12.003
- Maniatis, T., Fritsch, E. F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. New York, NY: Cold Spring Harbor Laboratory.
- Maqbool, Z., Asghar, H. N., Shahzad, T., Hussain, S., Riaz, M., Ali, S., et al. (2015). Isolating, screening and applying chromium reducing bacteria to promote growth and yield of okra (*Hibiscus esculentus* L.) in chromium contaminated soils. *Ecotoxicol. Environ. Saf.* 114, 343–349. doi: 10.1016/j.ecoenv.2014.07.007
- Ndeddy Aka, R. J., and Babalola, O. O. (2016). Effect of bacterial inoculation of strains of *Pseudomonas aeruginosa*, *Alcaligenes faecalis* and *Bacillus subtilis* on germination, growth and heavy metal (cd, cr, and ni) uptake of *Brassica juncea*. *Int. J. Phytoremediation* 18, 200–209. doi: 10.1080/15226514.2015.1073671
- Oliveira, H. (2012). Chromium as an environmental pollutant: insights on induced plant toxicity. *J. Bot.* 2012, 1–8. doi: 10.1155/2012/375843
- Olaszewska, M. A., Kocot, A. M., Nynca, A., and Łaniewska-Trokenheim, Ł. (2016). Utilization of physiological and taxonomic fluorescent probes to study Lactobacilli cells and response to pH challenge. *Microbiol. Res* 192, 239–246. doi: 10.1016/j.micres.2016.07.011
- Pal, A., and Paul, A. K. (2004). Aerobic chromate reduction by chromium-resistant bacteria isolated from serpentine soil. *Microbiol. Res.* 159, 347–354. doi: 10.1016/j.micres.2004.08.001
- Patil, S., Bheemaraddi, M. C., and Shivannavar, C. T. (2014). Biocontrol activity of siderophore producing *Bacillus subtilis* CTS-G24 against wilt and dry root rot causing fungi in Chickpea. *J. Agric. Vet. Sci.* 7, 63–68.
- Payne, S. (1994). Detection, isolation and characterization of siderophores. *Methods Enzymol.* 235, 329–344. doi: 10.1016/0076-6879(94)35151-1
- Premono, M. E., Moawad, A. M., and Vlek, P. L. G. (1996). Effect of phosphate-solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere. *Indones J. Crop Sci.* 11, 13–23.
- Qishlaqi, A., and Moore, F. (2007). Statistical analysis of accumulation and sources of heavy metals occurrence in agricultural soils of Khoshk River Banks, Shiraz, Iran. *Am. Euras. J. Agric. Environ. Sci.* 2, 565–573.
- Rani, J., Hemambika, B., Hemapriya, J., and Rajeshkannan, V. (2010). Comparative assessment of heavy metal removal by immobilized and dead bacterial cells: a biosorption approach. *Glob. J. Environ. Res.* 4, 23–30.
- Saharan, B. S., and Nehra, V. (2011). Plant growth promoting rhizobacteria?: a critical review. *Life Sci. Med. Res.* 21, 1–30.
- Saleh, S. S., and Glick, B. R. (2001). Involvement of *gacS* and *rpoS* in enhancement of the plant growth-promoting capabilities of *Enterobacter cloacae* CAL2 and UW4. *Can. J. Microbiol.* 47, 698–705. doi: 10.1139/cjm-47-8-698
- Samuel, J., Paul, M. L., Ravishankar, H., Mathur, A., Saha, D. P., Natarajan, C., et al. (2013). The differential stress response of adapted chromite mine isolates *Bacillus subtilis* and *Escherichia coli* and its impact on bioremediation potential. *Biodegradation* 24, 829–842. doi: 10.1007/s10532-013-9631-8
- Sarathambal, C., Khankhane, P. J., Gharde, Y., Kumar, B., Varun, M., and Arun, S. (2017). The effect of Plant growth promoting rhizobacteria on the growth,

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2017.00778/full#supplementary-material>

- physiology, and Cd uptake of *Arundo donax* L. *Int. J. Phytoremediation* 19, 360–370. doi: 10.1080/15226514.2016.1225289
- Schwyn, B., and Neilands, J. B. (1987). Universal chemical assay for the detection and determination of siderophores. *Anal. Biochem.* 160, 47–56. doi: 10.1016/0003-2697(87)90612-9
- Shakoori, A. R., Makhdoom, M., and Haq, R. U. (2000). Hexavalent chromium reduction by a dichromate-resistant gram-positive bacterium isolated from effluents of tanneries. *Appl. Microbiol. Biotechnol.* 53, 348–351. doi: 10.1007/s002530050033
- Shukla, K. P., Sharma, S., Singh, N. K., and Singh, V. (2012). Deciphering rhizosphere soil system for strains having plant growth promoting and bioremediation traits. *Agric. Res.* 1, 251–257. doi: 10.1007/s40003-012-0028-4
- Singh, S., Srivastava, P. K., Kumar, D., Tripathi, D. K., Chauhan, D. K., and Prasad, S. M. (2015). Morpho-anatomical and biochemical adapting strategies of maize (*Zea mays* L.) seedlings against lead and chromium stresses. *Biocatal. Agric. Biotechnol.* 4, 286–295. doi: 10.1016/j.bcab.2015.03.004
- Singh, S., Tripathi, D. K., Singh, S., Sharma, S., Dubey, N. K., Chauhan, D. K., et al. (2017). Toxicity of aluminium on various levels of plant cells and organism: a review. *Environ. Exp. Bot.* 137, 177–193. doi: 10.1016/j.envexpbot.2017.01.005
- Sobariu, D. L., Fertu, D. I. T., Diaconu, M., Pavel, L. V., Hlihor, R.-M., Drăgoi, E. N., et al. (2016). Rhizobacteria and plant symbiosis in heavy metal uptake and its implications for soil bioremediation. *N. Biotechnol.* doi: 10.1016/j.nbt.2016.09.002 [Epub ahead of print].
- Srivastava, S., and Thakur, I. S. (2007). Evaluation of biosorption potency of *Acinetobacter* sp. for removal of hexavalent chromium from tannery effluent. *Biodegradation* 18, 637–646. doi: 10.1007/s10532-006-9096-0
- Thacker, U., Parikh, R., Shouche, Y., and Madamwar, D. (2007). Reduction of chromate by cell-free extract of *Brucella* sp. isolated from Cr (VI) contaminated sites. *Bioresour. Technol.* 98, 1541–1547. doi: 10.1016/j.biortech.2006.06.011
- Tripathi, A., Tripathi, D. K., Chauhan, D. K., and Kumar, N. (2016a). Chromium (VI)-induced phytotoxicity in river catchment agriculture: evidence from physiological, biochemical and anatomical alterations in *Cucumis sativus* (L.) used as model species. *Chem. Ecol.* 32, 12–33. doi: 10.1080/02757540.2015.1115841
- Tripathi, D. K., Singh, V. P., Kumar, D., and Chauhan, D. K. (2012). Impact of exogenous silicon addition on chromium uptake, growth, mineral elements, oxidative stress, antioxidant capacity, and leaf and root structures in rice seedlings exposed to hexavalent chromium. *Acta Physiol. Plant.* 34, 279–289. doi: 10.1007/s11738-011-0826-5
- Tripathi, D. K., Singh, V. P., Prasad, S. M., Chauhan, D. K., Dubey, N. K., and Rai, A. K. (2015). Silicon-mediated alleviation of Cr(VI) toxicity in wheat seedlings as evidenced by chlorophyll fluorescence, laser induced breakdown spectroscopy and anatomical changes. *Ecotoxicol. Environ. Saf.* 113, 133–144. doi: 10.1016/j.ecoenv.2014.09.029
- Tripathi, D. K., Singh, V. P., Prasad, S. M., Dubey, N. K., Chauhan, D. K., and Rai, A. K. (2016b). LIB spectroscopic and biochemical analysis to characterize lead toxicity alleviative nature of silicon in wheat (*Triticum aestivum* L.) seedlings. *J. Photochem. Photobiol. B* 154, 89–98. doi: 10.1016/j.jphotobiol.2015.11.008
- Upadhyay, N., Verma, S., Singh, A. P., Devi, S., Vishwakarma, K., Kumar, K., et al. (2016). Soil ecophysiological and microbiological indices of soil health: a study of coal mining site in Sonbhadra, Uttar Pradesh. *J. Soil Sci. Plant Nutr.* 6, 778–800. doi: 10.4067/s0718-95162016005000056
- Walberg, M., Gaustad, P., and Steen, H. B. (1997). Rapid assessment of ceftazidime, ciprofloxacin, and gentamicin susceptibility in exponentially-growing *E. coli* cells by means of flow cytometry. *Cytometry* 27, 169–178. doi: 10.1002/(SICI)1097-0320(19970201)27:2<169::AID-CYTO9>3.0.CO;2-B
- Wright, M. S., Peltier, G. L., Stepanauskas, R., and McArthur, J. V. (2006). Bacterial tolerances to metals and antibiotics in metal-contaminated and reference streams. *FEMS Microbiol. Ecol.* 58, 293–302. doi: 10.1111/j.1574-6941.2006.00154.x
- Wuana, R. A., and Okieimen, F. E. (2011). Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. *ISRN Ecol.* 2011, 1–20. doi: 10.5402/2011/402647
- Yadav, B. K., and Verma, A. (2012). *Phosphate Solubilization and Mobilization in Soil Through Microorganisms Under Arid Ecosystems*. Rijeka: INTECH Open Access Publisher, 93–108.
- Zahoor, A., and Rehman, A. (2009). Isolation of Cr(VI) reducing bacteria from industrial effluents and their potential use in bioremediation of chromium containing wastewater. *J. Environ. Sci.* 21, 814–820. doi: 10.1016/S1001-0742(08)62346-3
- Zhu, W., Yang, Z., Ma, Z., and Chai, L. (2008). Reduction of high concentrations of chromate by *Leucobacter* sp. CRB1 isolated from Changsha, China. *World J. Microbiol. Biotechnol.* 24, 991–996. doi: 10.1007/s00253-012-4361-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Upadhyay, Vishwakarma, Singh, Mishra, Kumar, Rani, Mishra, Chauhan, Tripathi and Sharma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.