



Crassulacean Acid Metabolism Abiotic Stress-Responsive Transcription Factors: a Potential Genetic Engineering Approach for Improving Crop Tolerance to Abiotic Stress

Atia B. Amin^{1†}, Kumudu N. Rathnayake^{1†}, Won C. Yim², Travis M. Garcia², Beate Wone¹, John C. Cushman² and Bernard W. M. Wone^{1*}

¹ Department of Biology, University of South Dakota, Vermillion, SD, United States, ² Department of Biochemistry and Molecular Biology, University of Nevada, Reno, Reno, NV, United States

OPEN ACCESS

Edited by:

Wolfram Weckwerth,
University of Vienna, Austria

Reviewed by:

Dong-Ha Oh,
Louisiana State University,
United States
Douglas S. Domingues,
São Paulo State University (UNESP),
Brazil

*Correspondence:

Bernard W. M. Wone
bernie.wone@usd.edu

[†]Co-first authors

Specialty section:

This article was submitted to
Plant Systems and Synthetic Biology,
a section of the journal
Frontiers in Plant Science

Received: 29 September 2018

Accepted: 25 January 2019

Published: 22 February 2019

Citation:

Amin AB, Rathnayake KN,
Yim WC, Garcia TM, Wone B,
Cushman JC and Wone BWM (2019)
Crassulacean Acid Metabolism
Abiotic Stress-Responsive
Transcription Factors: a Potential
Genetic Engineering Approach
for Improving Crop Tolerance
to Abiotic Stress.
Front. Plant Sci. 10:129.
doi: 10.3389/fpls.2019.00129

This perspective paper explores the utilization of abiotic stress-responsive transcription factors (TFs) from crassulacean acid metabolism (CAM) plants to improve abiotic stress tolerance in crop plants. CAM is a specialized type of photosynthetic adaptation that enhances water-use efficiency (WUE) by shifting CO₂ uptake to all or part of the nighttime when evaporative water losses are minimal. Recent studies have shown that TF-based genetic engineering could be a useful approach for improving plant abiotic stress tolerance because of the role of TFs as master regulators of clusters of stress-responsive genes. Here, we explore the use of abiotic stress-responsive TFs from CAM plants to improve abiotic stress tolerance and WUE in crops by controlling the expression of gene cohorts that mediate drought-responsive adaptations. Recent research has revealed several TF families including *AP2/ERF*, *MYB*, *WRKY*, *NAC*, *NF-Y*, and *bZIP* that might regulate water-deficit stress responses and CAM in the inducible CAM plant *Mesembryanthemum crystallinum* under water-deficit stress-induced CAM and in the obligate CAM plant *Kalanchoe fedtschenkoi*. Overexpression of genes from these families in *Arabidopsis thaliana* can improve abiotic stress tolerance in *A. thaliana* in some instances. Therefore, we propose that TF-based genetic engineering with a small number of CAM abiotic stress-responsive TFs will be a promising strategy for improving abiotic stress tolerance and WUE in crop plants in a projected hotter and drier landscape in the 21st-century and beyond.

Keywords: abiotic stress response, crassulacean acid metabolism, drought tolerance, extremophytes, genetic engineering, transcription factor

INTRODUCTION

Formidable challenges facing humankind include a burgeoning global human population (Godfray et al., 2010; Gerland et al., 2014) and the increasing frequency and intensity of droughts related to global warming (Cook et al., 2014; Singh et al., 2015). In addition, abiotic stresses including high salinity, temperature extremes, increased UV radiation, heavy metals, and high light intensities

are and will continue to be major constraints for global crop production and food security (Lesk et al., 2016). Among these abiotic stresses, drought is of major concern as it has dire effects on crop productivity (Fahad et al., 2017), plant growth, and development (Yordanov et al., 2000). By the end of the 21st century, rapid changes in the global climate will likely increase the frequencies of drought by more than 20% compared to current rates (Lobell et al., 2011; Cook et al., 2014). Indeed, Daryanto et al. (2016) showed that an approximately 40% decrease in water availability can decrease wheat (*Triticum aestivum* L.) and maize (*Zea mays* L.) yields by 20.6% and 39.3%, respectively. High salinity, another abiotic stress that is harmful to crop production, affects 20% of total cultivated and 33% of irrigated agricultural lands worldwide (Shrivastava and Kumar, 2015). Similarly, low temperatures and heatwaves cause significant reduction in crop yields across the world (Sanghera et al., 2011; Challinor et al., 2014; Hatfield and Prueger, 2015). Thus, novel approaches to mitigate the negative impacts of abiotic stresses on crop yields must be explored and developed to avoid socio-economic collapse due to climate change.

Approaches to enhance sustainable bioenergy production by engineering the CAM pathway into C₃ crops to enhance their water-use efficiency (WUE) on marginal lands are already underway (Borland et al., 2014, 2015; Yang et al., 2015; Liu et al., 2018). Another approach to enhance abiotic stress tolerance is to modulate the expression of transcription factors (TFs) or the functions of abiotic stress-adaptive genes that might already be present, but that are not normally expressed in unstressed or C₃ plants (Hussain et al., 2011; Rabara et al., 2014; Joshi et al., 2016; Wang et al., 2016; Bechtold, 2018). This approach would involve bioengineering a small number of regulatory genes with potentially global effects made possible by the role of TFs in gene regulation (Rabara et al., 2014; Joshi et al., 2016).

Transcription factor-based genetic engineering could direct such regulatory TFs to modulate a large number of downstream abiotic stress-responsive genes (Rabara et al., 2014). Stress tolerance in plants is generally under polygenic control (Tran et al., 2010) and some of the genes regulating stress-tolerance responses happen to code for TFs (Villalobos et al., 2004; Qiu and Yu, 2009; Zhang et al., 2011; Yang et al., 2012; Cai et al., 2014; Chen et al., 2014; Swain et al., 2017). Therefore, TFs might be ideal candidate regulators for improving abiotic stress tolerance in crop plants. To date, TF-based genetic engineering has mainly repurposed TFs from *Arabidopsis thaliana*, *Glycine max*, *Oryza sativa*, and *T. aestivum* (Rabara et al., 2014; Joshi et al., 2016; Wang et al., 2016). As far as we know, no reports have analyzed the effects of overexpressing TFs from crassulacean acid metabolism (CAM) plants which have greater abiotic stress tolerances than mesophytes. Most CAM plants are naturally adapted to low-water environments and many other abiotic stresses compared to the agronomically important C₃ plants (Borland et al., 2009). Here, we consider using a “next-generation TF-based” approach to exploit abiotic stress-responsive TFs from CAM plants to improve abiotic stress tolerance in crop plants. The current technologies for TF-based approaches to improve plant abiotic stress tolerance have been extensively discussed

and reviewed by Rabara et al. (2014), Joshi et al. (2016), and Wang et al. (2016), and those details are therefore only briefly summarized below.

TRANSCRIPTION FACTOR-BASED APPROACH

Plants are sessile organisms that exhibit various biochemical, physiological, and molecular adaptations to extreme environments (Joshi et al., 2016). For instance, water-deficit stress activates the expression of stress-responsive genes encoding enzymes that synthesize compatible protective sugars, antioxidants, and proteins, including heat shock proteins and some classes of late embryogenesis abundant (LEA) proteins (Tran et al., 2010; Joshi et al., 2016; Wang et al., 2016). In addition to the stress-induced upregulation of the above proteins, the expression of various regulatory proteins including TFs, protein kinases, and protein phosphatases is also activated (Wang et al., 2016).

Transcription factors are master regulators of many cellular processes and can also interact with other transcriptional regulators (Joshi et al., 2016). Importantly, they play a pivotal role in different abiotic stress responses by binding to the upstream *cis*-regions of promoters in many stress-responsive genes (Yamaguchi-Shinozaki and Shinozaki, 2006). Many studies have been conducted to identify and characterize families of TFs including *AP2/ERF*, *MYB*, *WRKY*, *NAC*, *NF-Y*, and *bZIP* that are involved in abiotic stress responses (Umezawa et al., 2006; Gollack et al., 2011; Leyva-González et al., 2012; Wang et al., 2016; Swain et al., 2017; Zanetti et al., 2017). Several TFs have already been overexpressed in crop plants and *A. thaliana* to improve abiotic stress tolerance (Qiu and Yu, 2009; Zhang et al., 2011; Yang et al., 2012; Cai et al., 2014; Chen et al., 2014). For example, the *NAC* family is one of the largest TF families in plants and is involved not only in plant growth and development, but also in transcriptional reprogramming associated with plant stress responses (Tran et al., 2010; Nakashima et al., 2012). Mao et al. (2012) reported that overexpression of the *TaNAC2* gene from wheat can enhance tolerance to drought, salt, and freezing stresses in *A. thaliana*. In addition, functional characterization of the *NAC045* (Zheng et al., 2009) and *SNAC1* genes from *O. sativa* enhanced drought and salt tolerance in rice (Hu et al., 2006). Furthermore, overexpression of either *GmMYB76* or *GmMYB177* from soybean significantly enhanced salt and freezing tolerance in *A. thaliana* (Liao et al., 2008).

A relatively less explored yet high-potential approach is to discover novel abiotic stress-adaptive regulatory genes in extremophytes (i.e., CAM xerophytes and halophytes, desiccation-tolerant plants, or resurrection plants) to use for bioengineering abiotic stress tolerance in crop plants (Inan et al., 2004; Shi et al., 2013; Joshi et al., 2016; Bechtold, 2018). For example, the overexpression of the TF *CpMYB10* from the resurrection plant *Craterostigma plantagineum* in *A. thaliana* led to desiccation and salt tolerance in transgenic lines (Villalobos et al., 2004).

CRASSULACEAN ACID METABOLISM AND ABIOTIC STRESS TOLERANCE

CAM plants have evolved a specialized type of photosynthetic adaptation that allows them to live under conditions of severe water deficit and in semi-arid and arid regions of the world including deserts. These plants have shifted all or part of their primary CO₂ uptake and fixation to the nighttime, when evaporative water losses are minimal, and perform C₃ carboxylation reactions when stomata are closed during the daytime. This temporal separation of carbon fixation leads to the formation of the four-carbon organic acid malate, which is stored in the vacuole during the night and subsequently undergoes decarboxylation to release CO₂ for re-fixation during the day to produce carbohydrates (Borland et al., 2009). Because of this temporal separation of carbon fixation and inverted stomatal behavior, CAM plants can reduce water loss due to transpiration. These characteristics also allow CAM plants to fix net CO₂ 15% more efficiently than C₃ plants (Nobel, 1991) resulting in increased biomass of CAM plants while using less water than C₃ plants. Additionally, CAM plants can produce similar amounts of biomass using 80% less water in comparison to C₃ plants (Nobel, 1996; Borland et al., 2009). Thus, CAM plants have between 3- and 6-fold higher WUE than C₄ and C₃ plants, respectively (Garcia et al., 2014; Yang et al., 2015).

In addition to their higher WUE and associated drought tolerance (Yang et al., 2017), CAM plants can tolerate high temperature up to 70°C, whereas C₃ plants can tolerate only 50–55°C (Borland et al., 2009). CAM halophytes can also adapt to high salinity, as during the induction of CAM by salt stress in *Mesembryanthemum crystallinum* (Winter and Holtum, 2014). Moreover, CAM plants can better tolerate higher light intensities (>1000 μmol m⁻² s⁻¹) and UV-B irradiation levels than can agronomically important C₃ plants (Borland et al., 2009). Furthermore, CAM plants can increase daily net CO₂ uptake under increased atmospheric CO₂ concentrations, which might be advantageous in global climate change scenarios (Nobel, 1996). Some CAM plants such as *Agave salmiana*, *Opuntia ficus-indica*, and *Stenocereus queretaroensis* can also survive in subzero temperatures, and *Agave utahensis* Engelm. can tolerate temperatures as low as –18°C (Nobel, 1996).

The abiotic stress-adaptive characteristics of CAM plants will be particularly beneficial for adapting to the consequences of anthropogenic climate change, such as droughts and heatwaves, high soil salinity, temperature extremes, and high light or UV-B irradiation. Many of the stress-adaptive responses involve abscisic acid (ABA)-dependent and -independent response pathways (Song et al., 2016). ABA-dependent and independent signaling events likely participate in the stress-activation of CAM in *M. crystallinum* (Chu et al., 1990; Taybi and Cushman, 1999, 2002; Abdin et al., 2002; Cushman and Borland, 2002), suggesting that the CAM pathway likely has evolved in response to abiotic stress (Reyes-García and Andrade, 2009; Heyduk et al., 2018; Yin et al., 2018).

Crassulacean acid metabolism is thought to have evolved independently multiple times from ancestral C₃ plants (Silvera et al., 2010) because no unique metabolic pathways are required,

although some CAM-specific variant enzymes apparently evolved in some instances (Ermolova et al., 2003; Gehrig et al., 2005; Vaasen et al., 2006). However, temporal changes in gene expression of the CAM enzyme variants likely occurred because of alterations in their regulation compared to their orthologs in C₃ plants (Hermans and Westhoff, 1990; Lepiniec et al., 1993; Cushman et al., 2008; Heyduk et al., 2018; Yin et al., 2018). Furthermore, the ABA-dependent stress response pathway is involved in CAM activation not only in *M. crystallinum* (Taybi and Cushman, 2002), but also in other CAM species (Taybi et al., 1995; Rodrigues et al., 2016; Yin et al., 2018). Although CAM is found in over 36 families of vascular plants (Silvera et al., 2010), we rely on only a few major CAM species such as pineapple (*Ananas comosus*), *Agave*, and *Opuntia* as agricultural crops to provide food, forage, fiber, and biofuels (Cushman et al., 2015). Well characterized CAM model species also provide abundant resources for the identification and selection of candidate TFs involved in abiotic stress adaptations (Hartwell et al., 2016). Hence, identification of candidate CAM pathway regulators (i.e., TFs) that are expressed or activated under water-deficit stress or CAM should be prioritized to exploit the molecular and regulatory machinery of abiotic stress adaptation in CAM plants as a vital resource for applications in C₃ crop species (Yang et al., 2015; Fernie, 2016; Yin et al., 2018).

CAM ABIOTIC STRESS-RESPONSIVE TF-BASED APPROACH

Bioengineering a TF that can confer desirable traits such as increased drought tolerance (Villalobos et al., 2004) or increased biomass (Lim et al., 2018) into C₃ *A. thaliana* will be crucial as a proof of concept for the CAM abiotic stress-responsive TF-based approach to increase abiotic stress tolerance in C₃ plants. Fortunately, genetic resources (i.e., genome and transcriptome sequences) for CAM plants are now available for *Agave* (Abraham et al., 2016), *Kalanchoe* spp. (Yang et al., 2017), pineapple (Ming et al., 2015, 2016; Wai et al., 2017), and *M. crystallinum* (Chiang et al., 2016). Although genetic resources for CAM plants are becoming readily available, the underlying regulatory basis of CAM is still not completely understood. Many TFs of unknown function have been identified during recent genome and transcriptome sequencing efforts; thus, there are now many opportunities to analyze the functions of TFs involved in water-deficit-stress response or CAM function and to exploit the potential of bioengineering using CAM plant TFs to improve abiotic stress tolerance in crop plants. Indeed, candidate CAM TFs involved in C₃ to CAM transition in obligate CAM species of *Agave* (Heyduk et al., 2018; Huang et al., 2018; Yin et al., 2018), *Kanlanchoe* (Moseley et al., 2018), and *Manfreda* (Heyduk et al., 2018), and weak CAM species of *Polianthes* and *Beschorneria* (Heyduk et al., 2018), or the induction of CAM in *Tralimum triangulare* (Brilhaus et al., 2016) have been identified. Not surprisingly, a number of these candidate TFs are involved in the ABA stress responsive pathway (Heyduk et al., 2018; Yin et al., 2018). More importantly though, many of these candidate TFs are involved in the rewiring of the phase shift from C₃

to CAM transition in the evolution of CAM photosynthesis (Heyduk et al., 2018; Moseley et al., 2018; Yin et al., 2018). Although it would be interesting to attempt to reprogram a C₃ plant such that it becomes CAM performing, we are not suggesting to shift gene expression patterns of CAM pathway genes that might be present in extant C₃ plants, or to regulate the engineered CAM pathway in C₃ plants (Yang et al., 2015; Fernie, 2016; Heyduk et al., 2018; Yin et al., 2018), but rather identify and exploit the TFs involved in abiotic stress responses

from obligate and inducible CAM plants to modulate the expression of appropriate genes in C₃ plants to improve their abiotic stress tolerance.

Eight most abundant candidate TFs under water-deficit stress diel and zeitgeber time have been identified that might regulate the CAM state, water-deficit stress response, or both in *M. crystallinum* (Garcia et al., 2014; Cushman, unpubl. data; **Table 1**). *Mesembryanthemum crystallinum* switches from C₃ to CAM

TABLE 1 | List of the top eight candidate transcription factors (TFs) from the inducible CAM plant *Mesembryanthemum crystallinum* and top eight candidate TFs from the obligate CAM plant *Kalanchoe fedtschenkoi* hypothesized to regulate the CAM state or water-deficit stress responses in CAM plants and their corresponding orthologs in *Arabidopsis thaliana*.

TF Name	TF Family	<i>A. thaliana</i> Locus ID	Functional annotation of <i>A. thaliana</i> ortholog at TAIR
<i>McERF74</i>	AP2/ERF/CRF	AT1G53910	Detection of hypoxia, ethylene-activated signaling pathway, regulation of root development, response to hypoxia
<i>McNAC29</i>	NAC	AT1G69490	Embryo development ending in seed dormancy, flower development, fruit ripening, leaf senescence, multicellular organism development, multidimensional cell growth, regulation of transcription
<i>McBLH1</i>	HB/Homeodomain	AT2G35940	Polar nuclei fusion, response to abscisic acid, response to continuous far red-light stimulus by the high-irradiance response system, response to symbiotic fungus
<i>McbZIP2</i>	bZIP	AT2G18160	Positive regulation of transcription
<i>McAGL8</i>	MADS/AGAMOUS-LIKE 8	AT5G60910	Cell differentiation, developmental growth involved in morphogenesis, positive regulation of flower development, fruit development, maintenance of inflorescence meristem identity
<i>McAP2-12</i>	AP2/ERF	AT1G53910	Detection of hypoxia, ethylene-activated signaling pathway, regulation of root development, response to hypoxia
<i>McbZIP44</i>	bZIP	AT1G75390	Positive regulation of transcription, seed germination
<i>McHB7</i>	HB/Homeobox	AT2G46680	Abscisic acid-activated signaling pathway, positive regulation of transcription, response to water deprivation
<i>KfMYB59</i>	MYB	AT5G59780	Cell differentiation, response to cadmium ions, response to ethylene, response to gibberellin, response to NaCl
<i>KfLHY1</i>	Homeodomain	AT1G01060	Circadian rhythm, long-day photoperiodism, flowering, response to abscisic acid, response to auxin, response to NaCl
<i>KfBZIP29</i>	bZIP	AT4G38900	Regulation of transcription, reproductive shoot system and development
<i>KfNF-YB3</i>	NF-Ys	AT4G14540	Regulation of transcription, protein heterodimerization
<i>KfNAC83</i>	NAC	AT5G13180	Lignin biosynthetic process, plant-type secondary cell wall biogenesis, fruit dehiscence
<i>KfAP2</i>	AP2/ERF/CRF	AT4G11140	Cotyledon development cytokinin-activated signaling pathway, ethylene-activated signaling pathway, leaf development
<i>KfCOL3</i> (010148t1)	Zinc Finger CONSTANS-like 4	AT5G24930	Red light signaling pathway, regulator of flower development, regulation of photomorphogenesis
<i>KfCOL5</i> (00914t1)	Zinc Finger CONSTANS-like 5	AT5G57660	Regulation of flower development, regulation of transcription, response to light stimulus

We are only reporting functions related to transcriptional activation and abiotic stress for the *A. thaliana* ortholog from the *Arabidopsis* Information Resource (TAIR) databases (<https://www.arabidopsis.org/>) in the table.

TABLE 2 | Results from T₃ transgenic *A. thaliana* lines overexpressing candidate CAM TFs from the obligate CAM plant *K. fedtschenkoi* hypothesized to regulate water-deficit stress response or CAM activation.

CAM-related transcription factor	Integrated WUE	Drought tolerance	NaCl tolerance	Heavy metal tolerance	Biomass	Timing of bolting	Lateral roots	Root hairs
<i>KfMYB59</i>	Enhanced	NC	100 mM NaCl	50 μM Na ₂ SeO ₄	Increased	Delayed	NC	Increased
<i>KfNAC83</i>	Enhanced	Enhanced	150 mM	TBD	Increased	NC	Increased	Increased

NC – No change, same as WT; TBD – To be determined.

when salt or water-deficit stressed (Cushman, 2001; Cushman and Borland, 2002; Taybi and Cushman, 2002). *Kalanchoe fedtschenkoi* orthologs of these top candidate TFs were also highly expressed during CAM induction in older leaf pairs of *K. fedtschenkoi* plants (Garcia et al., 2014; Cushman, unpubl. data). It is known that young leaves are C₃ performing, whereas mature leaves are CAM performing in *K. fedtschenkoi* (Cushman, 2001). These TFs share a base mean expression level of >100 FPKM and at least two-fold induction during a transition from C₃ to CAM or imposition of water-deficit stress in *K. fedtschenkoi* and *M. crystallinum*, respectively (Garcia et al., 2014; Cushman, unpubl. data). These candidate CAM TFs also belong to the families of TFs reported to be involved in abiotic stress responses (Rabara et al., 2014; Joshi et al., 2016; Roy, 2016; Wang et al., 2016). Orthologs of these candidate CAM TFs in *A. thaliana* also have several reported functions in plant development and abiotic stress tolerance (Zheng et al., 2009; Zhang et al., 2011; Yang et al., 2012; Chen et al., 2014; Swain et al., 2017; Zanetti et al., 2017). Intriguingly, recent functional characterization of two putative CAM regulators of water-deficit stress response or CAM activation *via* overexpression in *A. thaliana* strongly suggest that CAM TFs have high potential to increase tolerance to drought and other abiotic stresses in C₃ plants.

One of these candidate CAM TFs is a myeloblastosis (MYB59, closest ortholog in *Arabidopsis*) TF whose transcripts are 20-fold more abundant in CAM-performing older leaf pairs relative to C₃-performing younger leaves in *K. fedtschenkoi* (Hartwell et al. unpubl. data). Results from four, third-generation (T₃) transgenic lines carrying the *KlMYB59* indicate increased rosette size and biomass at 4-week-old juvenile stage, and increased shoot length at 8-week-old mature stage in transgenic plants compared to WT (Table 2; Wone et al., unpubl. data; full results being presented in a separate publication). However, transgenic lines show delayed flowering in long-day photoperiod compared to WT plants (16 h light/8 h dark). In addition, these transgenic lines exhibit increased integrated WUE compared to WT plants. Furthermore, transgenic lines have longer primary roots despite exposure to 50 μM selenium compared to WT plants. In addition to MYB59, transcripts of the NAC83 TF (closest ortholog in *Arabidopsis*) were also more highly expressed in CAM-performing leaves of *K. laxiflora* and *K. fedtschenkoi* relative to C₃-performing leaves (Cushman et al., unpubl. data). The function of this *K. fedtschenkoi* NAC83 TF (*KfNAC83*) is not known in CAM- or C₃-performing CAM plants, but its *A. thaliana* ortholog suggests roles in abiotic stress responses and development (Table 1). Functional characterization of *KfNAC83* shows enhanced water-deficit stress tolerance and increased integrated WUE in four independent transgenic T₃-generation *A. thaliana* lines compared to WT plants (Table 2; Wone et al., unpubl. data; full results being presented in a separate publication). Furthermore, *KfNAC83*-overexpressing lines show significantly increased rosette size, leaves in the mature rosette, shoot biomass, number of siliques, and lateral roots compared to WT. Interestingly, these transgenic lines also showed tolerance to 150 mM NaCl. Collectively, our characterization results strongly suggest that at least two of these candidate CAM TFs have functions in abiotic stress responses and CAM photosynthesis.

CONCLUSION

The abiotic stress-adaptive features of CAM plants provide a wealth of genetic resources, specifically TFs, that are now available for functional testing and possible improvement of WUE and abiotic stress responses in C₃ photosynthesis plants. Our recent findings strongly suggest that a bioengineering approach using CAM abiotic stress-responsive TFs has the potential to increase abiotic stress tolerance in *A. thaliana* and possibly in C₃ crop plants. Our results indicate that CAM abiotic stress-responsive gene expression can be modulated by the appropriate CAM TFs to generate stress-adaptive phenotypes in *A. thaliana* and likely other C₃ plants because these CAM abiotic stress-responsive genes are apparently conserved and present in C₃ plants (Heyduk et al., 2018). Furthermore, although *K. fedtschenkoi* is distantly related to *A. thaliana*, transgenic *A. thaliana* lines carrying the obligate CAM plants' TFs showed favorable features for translational applications. We are optimistic that overexpressing TFs from the inducible CAM halophyte, *M. crystallinum* will have similar favorable responses in *A. thaliana* lines. Co-overexpression of only a small number of obligate and/or inducible CAM plant abiotic stress-responsive TFs with demonstrated abiotic stress-adaptive or -responsive functions would provide a facile approach for bioengineering desirable responses to abiotic stress (Song et al., 2016). Such an approach could open the door to potentially transformative applications to ensure long-term sustainable food, fiber, feed, and fuel production in a projected hotter and drier landscape in the 21st century and beyond.

AUTHOR CONTRIBUTIONS

BWMW conceived the CAM abiotic stress-responsive TF-based approach to improve C₃ crop plant abiotic stress responses. AA, KR, and BWMW wrote the manuscript. AA, KR, and BW conducted the experiments and provided the CAM plant TF overexpression data. WY, TG, and JC provided the CAM plant TF sequences. All authors reviewed the final manuscript.

FUNDING

This work was funded by South Dakota Board of Regents Competitive Research Grant #A18-0015-001 to BWMW and was supported, in part, by the Department of Energy, Office of Science, Genomic Science Program under Award Number DE-SC0008834 to JC.

ACKNOWLEDGMENTS

We thank Mary Ann Cushman for her helpful and clarifying comments on the manuscript. We also thank Austin Hacck and Savannah Aanderud for their assistance in maintaining the transgenic lines of *Arabidopsis* plants.

REFERENCES

- Abdin, M., Rehman, R., Israr, M., Srivastava, P. S., and Bansal, K. C. (2002). Abiotic stress related genes and their role in conferring resistance in plants. *Indian J. Biotech.* 1, 225–244.
- Abraham, P. E., Yin, H., Borland, A. M., Weighill, D., Lim, S. D., De Paoli, H. C., et al. (2016). Transcript, protein and metabolite temporal dynamics in the CAM plant agave. *Nat. Plants* 2:16178. doi: 10.1038/nplants.2016.178
- Bechtold, U. (2018). Plant life in extreme environments: how do you improve drought tolerance? *Front. Plant Sci.* 9:543. doi: 10.3389/fpls.2018.00543
- Borland, A. M., Griffiths, H., Hartwell, J., and Smith, J. A. C. (2009). Exploiting the potential of plants with crassulacean acid metabolism for bioenergy production on marginal lands. *J. Exp. Bot.* 60, 2879–2896. doi: 10.1093/jxb/erp118
- Borland, A. M., Hartwell, J., Weston, D. J., Schlauch, K. A., Tschaplinski, T. J., Tuskan, G. A., et al. (2014). Engineering crassulacean acid metabolism to improve water-use efficiency. *Trends Plant Sci.* 19, 327–338. doi: 10.1016/j.tplants.2014.01.006
- Borland, A. M., Wullschlegler, S. D., Weston, D. J., Tuskan, G. A., Hartwell, J., Yang, X., et al. (2015). Climate-resilient agroforestry: physiological responses to climate change and engineering of crassulacean acid metabolism (CAM) as a mitigation strategy. *Plant Cell Environ.* 38, 1833–1849. doi: 10.1111/pce.12479
- Brilhaus, D., Bräutigam, A., Mettler-Altmann, T., Winter, K., and Weber, A. P. M. (2016). Reversible burst of transcriptional changes during induction of crassulacean acid metabolism in *Talinum triangulare*. *Plant Physiol.* 170, 102–122. doi: 10.1104/pp.15.01076
- Cai, R., Zhao, Y., Wang, Y., Lin, Y., Peng, X., Li, Q., et al. (2014). Overexpression of a maize WRKY58 gene enhances drought and salt tolerance in transgenic rice. *Plant Cell Tissue Organ. Cult.* 119, 565–577. doi: 10.1007/s11240-014-0556-7
- Challinor, A. J., Watson, J., Lobell, D. B., Howden, S. M., Smith, D. R., and Chhetri, N. (2014). A meta-analysis of crop yield under climate change and adaptation. *Nat. Clim. Chang.* 4, 287–291. doi: 10.1038/nclimate2153
- Chen, X., Wang, Y., Lv, B., Li, J., Luo, L., Lu, S., et al. (2014). The NAC family transcription factor OsNAP confers abiotic stress response through the aba pathway. *Plant Cell Physiol.* 55, 604–619. doi: 10.1093/pcp/pct204
- Chiang, C.-P., Yim, W. C., Sun, Y.-H., Ohnishi, M., Mimura, T., Cushman, J. C. et al. (2016). Identification of ice plant (*Mesembryanthemum crystallinum* L.) MicroRNAs using RNA-Seq and their putative roles in high salinity responses in seedlings. *Front. Plant Sci.* 7:1143. doi: 10.3389/fpls.2016.01143
- Chu, C., Dai, Z., Ku, M. S. B., and Edwards, G. E. (1990). Induction of crassulacean acid metabolism in the facultative halophyte *Mesembryanthemum crystallinum* by abscisic acid. *Plant Physiol.* 93, 1253–1260. doi: 10.1104/pp.93.3.1253
- Cook, B. I., Smerdon, J. E., Seager, R., and Coats, S. (2014). Global warming and 21st century drying. *Clim. Dyn.* 43, 2607–2627. doi: 10.1007/s00382-014-2075-y
- Cushman, J. C. (2001). Crassulacean acid metabolism. A plastic photosynthetic adaptation to arid environments. *Plant Physiol.* 127, 1439–1448. doi: 10.1104/pp.010818
- Cushman, J. C., Agarie, S., Albion, R. L., Elliot, S. M., Taybi, T., and Borland, A. M. (2008). Isolation and characterization of mutants of common ice plant deficient in crassulacean acid metabolism. *Plant Physiol.* 147, 228–238. doi: 10.1104/pp.108.116889
- Cushman, J. C., and Borland, A. M. (2002). Induction of Crassulacean acid metabolism by water limitation. *Plant Cell Environ.* 25, 295–310. doi: 10.1046/j.0016-8025.2001.00760.x
- Cushman, J. C., Davis, S. C., Yang, X., and Borland, A. M. (2015). Development and use of bioenergy feedstocks for semi-arid and arid lands. *J. Exp. Bot.* 66, 4177–4193. doi: 10.1093/jxb/erv087
- Daryanto, S., Wang, L., and Jacinthe, P. A. (2016). Global synthesis of drought effects on maize and wheat production. *PLoS One* 11:e0156362. doi: 10.1371/journal.pone.0156362
- Ermolova, N. V., Ann Cushman, M., Taybi, T., Condon, S. A., Cushman, J. C., and Chollet, R. (2003). Expression, purification, and initial characterization of a recombinant form of plant PEP-carboxylase kinase from CAM-induced *Mesembryanthemum crystallinum* with enhanced solubility in *Escherichia coli*. *Protein Expr. Purif.* 29, 123–131. doi: 10.1016/S1046-5928(03)00014-7
- Fahad, S., Bajwa, A. A., Nazir, U., Anjum, S. A., Farooq, A., Zohaib, A., et al. (2017). Crop production under drought and heat stress: plant responses and management options. *Front. Plant Sci.* 8:1147. doi: 10.3389/fpls.2017.01147
- Fernie, A. R. (2016). Systems biology: a new CAM era. *Nat. Plants* 2:16190. doi: 10.1038/nplants.2016.190
- Garcia, T. M., Heyduk, K., Kuzmick, E., and Mayer, J. A. (2014). Crassulacean acid metabolism biology. *New Phytol.* 204, 738–740. doi: 10.1111/nph.13127
- Gehrig, H. H., Wood, J., Ann Cushman, M., Virgo, A. C., Cushman, J., and Winter, K. (2005). Research note: large gene family of phosphoenolpyruvate carboxylase in the crassulacean acid metabolism plant *Kalanchoe pinnata* (Crassulaceae) characterised by partial cDNA sequence analysis. *Funct. Plant Biol.* 32, 467–472. doi: 10.1071/FP05079
- Gerland, P., Raftery, A. E., Ševčíková, H., Li, N., Gu, D., Spoorenberg, T., et al. (2014). World population stabilization unlikely this century. *Science* 346, 234–237. doi: 10.1126/science.1257469
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., et al. (2010). Food security: the challenge of feeding 9 billion people. *Science* 327, 812–818. doi: 10.1126/science.1185383
- Gollack, D., Luking, L., and Yang, O. (2011). Plant tolerance to drought and salinity: stress regulating transcription factors and their functional significance in the cellular transcriptional network. *Plant Cell Rep.* 30, 1383–1391. doi: 10.1007/s00299-011-1068-0
- Hartwell, J., Dever, L. V., and Boxall, S. F. (2016). Emerging model systems for functional genomics analysis of crassulacean acid metabolism. *Curr. Opin. Plant Biol.* 31, 100–108. doi: 10.1016/j.pbi.2016.03.019
- Hatfield, J. L., and Prueger, J. H. (2015). Temperature extremes: effect on plant growth and development. *Weather Clim. Extrem.* 10, 4–10. doi: 10.1016/j.wace.2015.08.001
- Hermans, J., and Westhoff, P. (1990). Analysis of expression and evolutionary relationships of phosphoenol-pyruvate carboxylase genes in *Flaveria trinervia* (C4) and *F. pringlei* (C3). *Mol. Gen. Genet.* 224, 459–468. doi: 10.1007/BF00262441
- Heyduk, K., Ray, J. N., Ayyampalayam, S., and Leebens-Mack, J. (2018). Shifts in gene expression profiles are associated with weak and strong crassulacean acid metabolism. *Am. J. Bot.* 105, 587–601. doi: 10.1002/ajb2.1017
- Hu, H., Dai, M., Yao, J., Xiao, B., Li, X., Zhang, Q., et al. (2006). Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice. *Proc. Natl. Acad. Sci. U.S.A.* 103, 12987–12992. doi: 10.1073/pnas.0604882103
- Huang, X., Wang, B., Xi, J., Zhang, Y., He, C., Zheng, J., et al. (2018). Transcriptome comparison reveals distinct selection patterns in domesticated and wild agave species, the important cam plants. *Int. J. Genomics* 2018, 1–12. doi: 10.1155/2018/5716518
- Hussain, S. S., Kayani, M. A., and Amjad, M. (2011). Transcription factors as tools to engineer enhanced drought stress tolerance in plants. *Biotechnol. Prog.* 27, 297–306. doi: 10.1002/btpr.514
- Inan, G., Zhang, Q., Li, P., Wang, Z., Cao, Z., Zhang, H., et al. (2004). Salt stress. A halophyte and cryophyte Arabidopsis relative model system and its applicability to molecular genetic analyses of growth and development of extremophiles. *Plant Physiol.* 135, 1718–1737. doi: 10.1104/pp.104.041723
- Joshi, R., Wani, S. H., Singh, B., Bohra, A., Dar, Z. A., Lone, A. A., et al. (2016). Transcription factors and plants response to drought stress: current understanding and future directions. *Front. Plant Sci.* 7:1029. doi: 10.3389/fpls.2016.01029
- Lepiniec, L., Keryer, E., Philippe, H., Gadal, P., and Cretin, C. (1993). Sorghum phosphoenolpyruvate carboxylase gene family: structure, function and molecular evolution. *Plant Mol. Biol.* 21, 487–502. doi: 10.1007/BF00028806
- Lesk, C., Rowhani, P., and Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature* 529, 84–87. doi: 10.1038/nature16467
- Leyva-González, M. A., Ibarra-Laclette, E., Cruz-Ramírez, A., and Herrera-Estrella, L. (2012). Functional and transcriptome analysis reveals an acclimatization strategy for abiotic stress tolerance mediated by Arabidopsis NF-YA family members. *PLoS One* 7:e48138. doi: 10.1371/journal.pone.0048138

- Liao, Y., Zou, H. F., Wang, H. W., Zhang, W. K., Ma, B., Zhang, J. S., et al. (2008). Soybean GmMYB76, GmMYB92, and GmMYB177 genes confer stress tolerance in transgenic Arabidopsis plants. *Cell Res.* 18, 1047–1060. doi: 10.1038/cr.2008.280
- Lim, S. D., Yim, W. C., Liu, D., Hu, R., Yang, X., and Cushman, J. C. (2018). A *Vitis vinifera* basic helix-loop-helix transcription factor enhances plant cell size, vegetative biomass and reproductive yield. *Plant Biotechnol. J.* 16, 1595–1615. doi: 10.1111/pbi.12898
- Liu, D., Palla, K. J., Hu, R., Moseley, R. C., Mendoza, C., Chen, M., et al. (2018). Perspectives on the basic and applied aspects of crassulacean acid metabolism (CAM) research. *Plant Sci.* 274, 394–401. doi: 10.1016/j.plantsci.2018.06.012
- Lobell, D. B., Schlenker, W., and Costa-Roberts, J. (2011). Climate trends and global crop production since 1980. *Science* 333, 616–620. doi: 10.1126/science.1204531
- Mao, X., Zhang, H., Qian, X., Li, A., Zhao, G., and Jing, R. (2012). TaNAC2, a NAC-type wheat transcription factor conferring enhanced multiple abiotic stress tolerances in Arabidopsis. *J. Exp. Bot.* 63, 2933–2946. doi: 10.1093/jxb/err462
- Ming, R., VanBuren, R., Wai, C. M., Tang, H., Schatz, M. C., Bowers, J. E., et al. (2015). The pineapple genome and the evolution of CAM photosynthesis. *Nat. Genet.* 47, 1435–1442. doi: 10.1038/ng.3435
- Ming, R., Wai, C. M., and Guyot, R. (2016). Pineapple genome: a reference for monocots and CAM photosynthesis. *Trends Genet.* 32, 690–696. doi: 10.1016/j.tig.2016.08.008
- Moseley, R. C., Mewalal, R., Motta, F., Tuskan, G. A., Haase, S., and Yang, X. (2018). Conservation and diversification of circadian rhythmicity between a model crassulacean acid metabolism plant *Kalanchoë fedtschenkoi* and a model C3 photosynthesis plant *Arabidopsis thaliana*. *Front. Plant Sci.* 9:1757. doi: 10.3389/fpls.2018.01757
- Nakashima, K., Takasaki, H., Mizoi, J., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2012). NAC transcription factors in plant abiotic stress responses. *Biochim. Biophys. Acta* 1819, 97–103. doi: 10.1016/j.bbagr.2011.10.005
- Nobel, P. S. (1991). Achievable productivities of certain CAM plants: basis for high values compared with C3 and C4 plants. *New Phytol.* 119, 183–205. doi: 10.1111/j.1469-8137.1991.tb01022.x
- Nobel, P. S. (1996). Responses of some North American CAM plants to freezing temperatures and doubled CO₂ concentrations: implications of global climate change for extending cultivation. *J. Arid Environ.* 34, 187–196. doi: 10.1006/jare.1996.0100
- Qiu, Y., and Yu, D. (2009). Over-expression of the stress-induced OsWRKY45 enhances disease resistance and drought tolerance in Arabidopsis. *Environ. Exper. Bot.* 65, 35–47. doi: 10.1016/j.envexpbot.2008.07.002
- Rabara, R. C., Tripathi, P., and Rushton, P. J. (2014). The potential of transcription factor-based genetic engineering in improving crop tolerance to drought. *Omic* 18, 601–614. doi: 10.1089/omi.2013.0177
- Reyes-García, C., and Andrade, J. (2009). Crassulacean acid metabolism under global climate change. *New Phytol.* 181, 754–757. doi: 10.1111/j.1469-8137.2009.02762.x
- Rodrigues, M. A., Hamachi, L., Mioto, P. T., Purgatto, E., and Mercier, H. (2016). Implications of leaf ontogeny on drought-induced gradients of CAM expression and ABA levels in rosettes of the epiphytic tank bromeliad *Guzmania monostachia*. *Plant Physiol. Biochem.* 1, 400–411. doi: 10.1016/j.plaphy.2016.08.010
- Roy, S. (2016). Function of MYB domain transcription factors in abiotic stress and epigenetic control of stress response in plant genome. *Plant Signal. Behav.* 11:e1117723. doi: 10.1080/15592324.2015.1117723
- Sanghera, G. S., Wani, S. H., Hussain, W., and Singh, N. B. (2011). Engineering cold stress tolerance in crop plants. *Curr. Genomics* 12, 30–43. doi: 10.2174/138920211794520178
- Shi, Y., Yan, X., Zhao, P., Yin, H., Zhao, X., Xiao, H., et al. (2013). transcriptomic analysis of a tertiary relict plant, extreme xerophyte *Reaumuria soongorica* to identify genes related to drought adaptation. *PLoS One* 8:e63993. doi: 10.1371/journal.pone.0063993
- Shrivastava, P., and Kumar, R. (2015). Soil salinity: a serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi J. Biol. Sci.* 22, 123–131. doi: 10.1016/j.sjbs.2014.12.001
- Silvera, K., Neubig, K., Whitten, W., Williams, N., Winter, K., and Cushman, J. C. (2010). Evolution along the crassulacean acid metabolism continuum. *Funct. Plant Biol.* 37, 995–1010. doi: 10.1071/FP10084
- Singh, B., Bohra, A., Mishra, S., Joshi, R., and Pandey, S. (2015). Embracing new-generation 'omics' tools to improve drought tolerance in cereal and food-legume crops. *Biol. Plant* 59, 413–428. doi: 10.1007/s10535-015-0515-0
- Song, L., Huang, S. C., Wise, A., Castanon, R., Nery, J. R., Chen, H., et al. (2016). A transcription factor hierarchy defines an environmental stress response network. *Science* 354, aag1550. doi: 10.1126/science.aag1550
- Swain, S., Myers, Z. A., Siriwardana, C. L., and Holt, B. F. III (2017). The multifaceted roles of NUCLEAR FACTOR-Y in *Arabidopsis thaliana* development and stress responses. *Biochim. Biophys. Acta Gene Regul. Mech.* 1860, 636–644. doi: 10.1016/j.bbagr.2016.10.012
- Taybi, T., and Cushman, J. C. (1999). Signaling events leading to crassulacean acid metabolism (CAM) induction in the common ice plant, *Mesembryanthemum crystallinum*. *Plant Physiol.* 121, 545–555. doi: 10.1104/pp.121.2.545
- Taybi, T., and Cushman, J. C. (2002). Abscisic acid signaling and protein synthesis requirements for phosphoenolpyruvate carboxylase transcript induction in the common ice plant. *J. Plant Physiol.* 159, 1235–1243. doi: 10.1078/0176-1617-00834
- Taybi, T., Sotta, B., Gehrig, H., Güclü, S., Kluge, M., and Brulfert, J. (1995). Differential effects of abscisic acid on phosphoenolpyruvate carboxylase and CAM operation in *Kalanchoë blossfeldiana*. *Botanica Acta* 108, 240–246. doi: 10.1111/j.1438-8677.1995.tb00856.x
- Tran, L. S., Nishiyama, R., Yamaguchi-Shinozaki, K., and Shinozaki, K. (2010). Potential utilization of NAC transcription factors to enhance abiotic stress tolerance in plants by biotechnological approach. *GM Crops* 1, 32–39. doi: 10.4161/gmcr.1.1.10569
- Umezawa, T., Fujita, M., Fujita, Y., Yamaguchi-Shinozaki, K., and Shinozaki, K. (2006). Engineering drought tolerance in plants: discovering and tailoring genes to unlock the future. *Curr. Opin. Biotechnol.* 17, 113–122. doi: 10.1016/j.copbio.2006.02.002
- Vaasen, A., Begerow, D., and Hampp, R. (2006). Phosphoenolpyruvate carboxylase genes in C₃, crassulacean acid metabolism (CAM) and C₃/CAM intermediate species of the genus *Clusia*: rapid reversible C₃/CAM switches are based on the C₃ housekeeping gene. *Plant Cell Environ.* 29, 2113–2123. doi: 10.1111/j.1365-3040.2006.01583.x
- Villalobos, M. A., Bartels, D., and Iturriaga, G. (2004). Stress tolerance and glucose insensitive phenotypes in Arabidopsis overexpressing the CpMYB10 transcription factor gene. *Plant Physiol.* 135, 309–324. doi: 10.1104/pp.103.034199
- Wai, C. M., VanBuren, R., Zhang, J., Huang, L., Miao, W., Edger, P. P., et al. (2017). Temporal and spatial transcriptomic and microRNA dynamics of CAM photosynthesis in pineapple. *Plant J.* 92, 19–30. doi: 10.1111/tpj.13630
- Wang, H., Wang, H., Shao, H., and Tang, X. (2016). Recent advances in utilizing transcription factors to improve plant abiotic stress tolerance by transgenic technology. *Front. Plant Sci.* 7:67. doi: 10.3389/fpls.2016.00067
- Winter, K., and Holtum, J. A. (2014). Facultative crassulacean acid metabolism (CAM) plants: powerful tools for unravelling the functional elements of CAM photosynthesis. *J. Exp. Bot.* 65, 3425–3441. doi: 10.1093/jxb/eru063
- Yamaguchi-Shinozaki, K., and Shinozaki, K. (2006). Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu. Rev. Plant Biol.* 57, 781–803. doi: 10.1146/annurev.arplant.57.032905.105444
- Yang, A., Dai, X., and Zhang, W.-H. (2012). A R2R3-type MYB gene, OsMYB2, is involved in salt, cold, and dehydration tolerance in rice. *J. Exp.* 63, 2541–2556. doi: 10.1093/jxb/err431
- Yang, X., Cushman, J. C., Borland, A. M., Edwards, E. J., Wulschleger, S. D., Tuskan, G. A., et al. (2015). A roadmap for research on crassulacean acid metabolism (CAM) to enhance sustainable food and bioenergy production in a hotter, drier world. *New Phytol.* 207, 491–504. doi: 10.1111/nph.13393
- Yang, X., Hu, R., Yin, H., Jenkins, J., Shu, S., Tang, H., et al. (2017). The *Kalanchoë* genome provides insights into convergent evolution and building blocks of crassulacean acid metabolism. *Nat. Commun.* 8:1899. doi: 10.1038/s41467-017-01491-7

- Yin, H., Guo, H. B., Weston, D. J., Borland, A. M., Ranjan, P., Abraham, P. E., et al. (2018). Diel rewiring and positive selection of ancient plant proteins enabled evolution of CAM photosynthesis in Agave. *BMC Genomics* 19:588. doi: 10.1186/s12864-018-4964-7
- Yordanov, I., Velikova, V., and Tsonev, T. (2000). Plant responses to drought, acclimation, and stress tolerance. *Photosynthetica* 38, 171–186. doi: 10.1023/a:1007201411474
- Zanetti, M. E., Ripodas, C., and Niebel, A. (2017). Plant NF-Y transcription factors: key players in plant-microbe interactions, root development and adaptation to stress. *Biochim. Biophys. Acta Gene Regul. Mech.* 1860, 645–654. doi: 10.1016/j.bbagr.2016.11.007
- Zhang, X., Wang, L., Meng, H., Wen, H., Fan, Y., and Zhao, J. (2011). Maize ABP9 enhances tolerance to multiple stresses in transgenic Arabidopsis by modulating ABA signaling and cellular levels of reactive oxygen species. *Plant Mol. Biol.* 75, 365–378. doi: 10.1007/s11103-011-9732-x
- Zheng, X., Chen, B., Lu, G., and Han, B. (2009). Overexpression of a NAC transcription factor enhances rice drought and salt tolerance. *Biochem. Biophys. Res. Commun.* 379, 985–989. doi: 10.1016/j.bbrc.2008.12.163

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Amin, Rathnayake, Yim, Garcia, Wone, Cushman and Wone. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.