



Deep in the Jelly: Histochemical and Functional Aspects of Mucilage-Secreting Floral Colleters in the Orchids *Elleanthus brasiliensis* and *E. crinipes*

Fábio Cassola^{1,2*}, Carlos Eduardo Pereira Nunes^{1,3}, Makeli Garibotti Lusa^{1,4}, Vera Lúcia Garcia² and Juliana Lischka Sampaio Mayer^{1*}

OPEN ACCESS

Edited by:

Kevin Davies,
The New Zealand Institute for Plant
and Food Research Ltd.,
New Zealand

Reviewed by:

Clay Carter,
University of Minnesota, United States
Simone Maria Teixeira
Sabóia-Morais,
Universidade Federal de Goiás, Brazil

*Correspondence:

Fábio Cassola
fcassola3@gmail.com
Juliana Lischka Sampaio Mayer
mjimayer@yahoo.com.br

Specialty section:

This article was submitted to
Plant Metabolism
and Chemodiversity,
a section of the journal
Frontiers in Plant Science

Received: 25 December 2018

Accepted: 04 April 2019

Published: 24 April 2019

Citation:

Cassola F, Nunes CEP, Lusa MG,
Garcia VL and Mayer JLS (2019)
Deep in the Jelly: Histochemical
and Functional Aspects
of Mucilage-Secreting Floral Colleters
in the Orchids *Elleanthus brasiliensis*
and *E. crinipes*.
Front. Plant Sci. 10:518.
doi: 10.3389/fpls.2019.00518

¹ Institute of Biology, State University of Campinas, Campinas, Brazil, ² Department of Organic and Pharmaceutical Chemistry, Chemical, Biological and Agricultural Pluridisciplinary Research Center, Paulínia, Brazil, ³ Department of Biological and Environmental Sciences, University of Stirling, Stirling, Scotland, ⁴ Center of Biological Sciences, Federal University of Santa Catarina, Florianópolis, Brazil

Colleters are trichomes or emergencies that produce a sticky exudate consisting of a mixture of mucilage, lipids, terpenes, and phenolic compounds. Colleters occur in at least 60 families of angiosperms; however, reports of them are scarce for the Orchidaceae. *Elleanthus brasiliensis* is distinguished by the presence of an abundant gelatinous secretion that covers almost all of its inflorescences. We aimed to describe the histology of colleters in inflorescences of *E. brasiliensis* and *Elleanthus crinipes*, and to analyze the chemical composition of their secretion to better understand the functions of these secretory structures. Due to the low frequency of colleters and lack of visible secretion in *E. crinipes*, histochemical tests and chemical analyses were not performed for this species. Colleters are of a brush type and their secretion has, at the same time, hydrophilic and lipophilic components. Histochemical tests further revealed the presence of pectin, mucilage, lipids, terpenes, phenolic compounds, and proteins. The GC-MS analysis confirmed the presence of γ -sitosterol and palmitic, linoleic, and stearic acids in the secretion of *E. brasiliensis*. Infrared analysis indicated the possible presence of polysaccharides in the secretion. The occurrence of colleters in both species studied and in other orchids described in the literature suggests that these structures are common in the inflorescences of tropical orchids. In these environments, the hydrated polysaccharides in the secretion form a dense matrix that can act as a physical barrier, and terpenes may help to protect against herbivores and pathogenic microorganisms. This information broadens our knowledge of the morphological and chemical diversity of the secretions produced by orchid colleters.

Keywords: Atlantic Forest, Epidendroideae, histochemistry analysis, microstructure, plant anatomy and morphology, secretory structure, Orchidaceae

Abbreviations: a.s.l., above sea level; kV, kilovolts; MM, molecular mass.

INTRODUCTION


Plant secretions are synthesized and eliminated by specific cells, which can occur in isolation or form differentiated glandular structures, such as trichomes, emergences, canals, cavities, and laticifers (Castro and Demarco, 2008; Castro and Machado, 2013). Among such differentiated glandular structures, colleters are emergences (Leitão and Cortelazzo, 2008) formed of epidermal and subepidermal tissues or of trichomes (Mayer et al., 2011; Machado and Rodrigues, 2013) originating only from the protoderm, which produce a sticky exudate composed of mucilage and/or lipid-like substances (Fahn, 1979; Mayer et al., 2013; Machado et al., 2015; Capelli et al., 2017). In general, this secretion is associated with vegetative and reproductive organs in the process of differentiation, and can protect them against dehydration or attacks by herbivores and microorganisms (Whittier and Peterson, 1984; Thomas, 1991; Mayer et al., 2013; Coutinho et al., 2015; Lusa et al., 2015).

In eudicot plants, the most common types of colleters are composed of secretory palisade epidermal cells, with the central axis formed by parenchyma and in some cases including vascular bundles (Miguel et al., 2006; Castro and Machado, 2013). In monocots plants, these structures can be trichomes, bulky cells with a dense cytoplasm and atrophied nucleus, hairs, or epidermal appendages (Leitão and Cortelazzo, 2008;

Mayer et al., 2011). Despite this distinction, many aspects of the secretion, physiology, anatomy, and ultrastructure of colleters are poorly known due to a lack of comprehensive and in-depth studies on them (Fahn, 1979; Macêdo et al., 2016).

The genus *Elleanthus* C. Presl contains 150 described species, with representatives in Central and South America, and its greatest diversity occurs in the Andes (Dudek et al., 2017). *Elleanthus brasiliensis* (Lindl.) Rchb. f. is found in the humid forests of eastern Brazil and the Guianas, while *Elleanthus crinipes* Rchb. f. is endemic to southeastern Brazil, where it is found in highland forests in different physiognomies of the Atlantic Forest (Nunes et al., 2016). *E. brasiliensis* stands out from other species due to the large amount of mucilaginous secretion that covers its inflorescences, refracting the reddish color of the bracts and giving the globular inflorescence a bright appearance (Figures 1A,B). In *E. crinipes* (Figure 1C), this secretion is conspicuously scarce (Figures 1D,E), although these two species may occur in the same environments and share the same pollinators (Nunes et al., 2013, 2016).

Knowledge of the secretory structures of plants and the chemical constitution of their secretions can help elucidate the relationships between patterns and processes in the ecology of interactions involving plants. Thus, this study aimed to identify and describe the secretory structures present in the inflorescences of *E. brasiliensis* and *E. crinipes* and analyze the chemical






FIGURE 2 | Distribution of the colleters (arrows) and secretion in the floral organs of *E. brasiliensis*. **(A,C,E)** Light microscopy. **(B,D,F–H)** Scanning electron microscopy. **(A–D)** Bracteole covered by a large amount of secretion (arrowheads). **(C,D)** Magnified detail of the colleters (“Col” and white arrows) involved in secretion. **(E,F)** Sepals presenting colleters and a lower amount of secretion. **(G)** Ovarian wall showing colleters. **(H)** Floral column with colleters on the surface. BLE, bracteole; COL, floral column; OVA, ovary; PET, petal; SEP, sepal.

composition of their secretion to estimate its functional role in the analyzed species and provide useful data for ecological, taxonomic, and chemosystematic purposes.

MATERIALS AND METHODS

Plant Material

Elleanthus brasiliensis and *E. crinipes* are epiphytic or rupicolous herbs that occur in environments with high air humidity (Nunes et al., 2016). *E. brasiliensis* has pendent stems (Figure 1A), while *E. crinipes* has erect stems (Figure 1C), at the apex of which reproductive buds give rise to racemose inflorescences (Figures 1B,D,E). Due to the low frequency of colleters and the lack of visible or abundant secretion in *E. crinipes*, histochemical

tests were not performed for this species. In despite of belonging to the same genus, these species are not closely related inside *Elleanthus* and are classified in distinct sections within it (Dressler, 2006).

Samples were collected during the flowering and fruiting seasons from 2012 to 2015. Sampling was carried out at two sites, a lowland site (less than 100 m a.s.l.) and a highland one (800–1000 m a.s.l.), both in areas of the Atlantic Forest (Ombrophilous Dense Forest; Veloso et al., 1991) in Serra do Mar State Park (SMSP), southeastern Brazil. Inflorescences of *E. brasiliensis* were collected in the lowland area where this species is more abundant, in the municipality of Ubatuba, São Paulo state (23°20'021.9' S, 44°50'014.5' W). Inflorescences of *E. crinipes* were collected in the highland area, between the municipalities of São Luiz do Paraitinga, Cunha, and Natividade da Serra, São Paulo state




FIGURE 3 | Distribution of the colleters (arrows) and secretion in the floral organs of *E. crinipes*. **(A,B)** Scanning electron microscopy. **(C–F)** Light microscopy. **(A)** Floral bud with bracteole showing colleters (white arrow) and sepals surrounding the bud. Inset: colleter on the surface of the sepal. **(B)** Magnified detail of the bracteole, showing colleters, and secretion (arrowheads). **(C)** Ovary with colleters on outer surface, as viewed from above the ovary axis of a lateral flower surrounded by a bract. **(D)** Detail of the ovary, showing colleter on the surface and the recess of the wall (white arrow). **(E)** Bracteole with a colleter. **(F)** Detail of the bracteole's colleters surrounded by their secretion. BLE, bracteole; OVA, ovary; SEP, sepal.

(23°26008' S, 45°13022.5' W and 23°19055' S, 45°05049' W, respectively). Vouchers (12/02/2010, *E. brasiliensis*, C.E.P. Nunes 01; *E. crinipes*, C.E.P. Nunes 02) are deposited in the herbarium of the University of Campinas “Prof. Dr. João Semir” (UEC).

Light and Scanning Electron Microscopy

To analyze the general structure of colleters, tissue samples of flowers and floral bracts at different developmental stages were fixed in a formalin–acetic acid–alcohol (FAA) solution for 24 h (Johansen, 1940) and in Karnovsky's solution for 48 h (Karnovsky, 1965), and were then subjected to reduced pressure to allow adequate penetration by the fixative. Samples were subsequently stored in 70% (v/v) ethanol. The material was then dehydrated through a tertiary butanol series (Johansen, 1940). One part of the samples was embedded in plastic resin (Leica Histo-resin®, Heraeus Kulzer, Hanau, Germany), while another part was embedded in Paraplast®X-tra (Fisher, cat. n° 23-021-401) (Johansen, 1940). The embedded samples

were longitudinally and transversely sectioned with a rotary microtome (Leica®) equipped with a type C blade. For the samples in Paraplast®, sections were cut at a thickness of 12 μm, had the paraffin removed, and were then stained with safranin O and astra blue (Srebotnik and Messner, 1994). For the samples in Histo-resin®, sections were cut at a thickness of 5–7 μm and stained with 0.05% toluidine blue at in a citrate-phosphate buffer with a pH of 4.5 (Sakai, 1973). After staining, the glass slides were mounted with the synthetic resin Entellan®(Merck®). The serial sections were examined microscopically (Olympus BX51) under polarized light to verify the occurrence of starch grains, crystals, and lignified cell walls. Images of inflorescences were taken in the field with a digital camera (Canon EOS20D).

For scanning electron microscopy (SEM), flower samples were fixed as described by Karnovsky (1965) for 24 h (modified by preparation in pH 7.2 phosphate buffer), dehydrated in a graded ethanol series, and subjected to critical point drying with CO₂ (Horridge and Tamm, 1969). Samples were then attached to

aluminum stubs and coated with gold (30–40 nm). Finally, the samples were examined under a LEO model VP 435 scanning electron microscope (SEM) at 10 kV.

Histochemical Analysis

Several different histochemical procedures were carried out to detect the main classes of chemical compounds typically produced by plant secretory structures. The histochemical reactions used comprised the following: reaction with coriphosphine under fluorescence to test for pectins (Ueda and Yoshioka, 1976); ruthenium red for mucilage and pectic substances (Johansen, 1940); Sudan III, Sudan IV (Jensen, 1962), and Sudan black B (Pearse, 1968) for total lipids; Nile blue sulfate for neutral lipids (Cain, 1947); aniline blue black for proteins (Fisher, 1968); Nadi reagent for terpenes (David and Carde, 1964); and ferric chloride for phenolic compounds (Johansen, 1940). Sections were examined immediately after each reaction under an Olympus®BX 51 microscope. Photomicrographs were taken of the samples under the Olympus®BX 51 microscope, which was equipped with an Olympus DP 71 camera. For the analysis of the reaction with coriphosphine, the same microscope was used, and was equipped for epifluorescence illumination with a U-LH100HG mercury lamp to provide excitation (bandpass filter: 450–490 nm) and suppression (long-pass filter: 515 nm). Control sections were prepared simultaneously to the

histochemical tests, in accordance with standard procedures. To verify the natural appearances of organs and secretions, untreated sections were prepared and observed. Images were recorded from light and epifluorescence microscopy by capturing images of the slides using an Olympus DP71 video camera, coupled to the abovementioned microscope.

Ethanollic Extract Preparation

The secretion from the collectors was collected directly from the surfaces of inflorescences of *E. brasiliensis* (the species with abundant secretion) containing floral buds using a Pasteur pipette, and was then frozen. Four to five inflorescences (13.2 mg of secretion) were used from three different individuals. Due to the high-volume reduction of the material when dry, it was necessary to join several individuals for these analyses. To obtain the ethanollic extract of the secretion, each sample was thawed, 15 mL of ethanol was added to it, and it was then placed in an ultrasonic bath for 20 min. The solution was dried under reduced pressure, which yielded 2.96 mg of the ethanollic extract.

Gas Chromatography Coupled to Mass Spectrometry (GC-MS)

The GC-MS analysis were performed on an Agilent®6890N chromatograph, with a 5975-mass detector and a 7683B automatic injector coupled to a HP5MS capillary column




FIGURE 4 | Characterization of the collectors of *E. brasiliensis* (A–E,G,H) and *E. crinipes* (F). (A) Scanning electron microscopy. (B–H) Light microscopy. (A) Brush-type collector with short basal axis and elongated terminal cells. (B–E) Collector with short cup-shaped basal axis composed of one (B), two (C,D), or three cells (E), and presenting lignified and suberified secondary wall deposition, as indicated by double-staining with afstra blue and safranin. Elongated cells have only primary cell walls. (F) Detail of the basal axis of the collector showing the cup-shaped cell with a secondary wall. (G) Collector with elongated cells releasing secretion (arrowheads) that covers the surface of the bracteole. (H) Histochemical test with Sudan IV indicating suberification in the cells of the collector's axis and in the cell walls of the elongated cells that connect with the cells of the axis. Ax, axis of collector; El, elongated cell; Sw, secondary wall; Su, suberized wall.

(30 m × 0.25 mm × 0.25 μm). The conditions of this analysis were as follows: the temperature of the injector was set at 220°C, the detector was set at 280°C, and the column was set at 60°C and increased by 3°C/min up to 240°C, and then maintained at 240°C for 7 min.; and He super dry was added at 1.0 mL/min as the carrier gas. The mass spectra obtained for each signal observed in the chromatogram were compared to the fragmentation patterns of the NIST 2005 library compounds of the equipment with a searching similarity of ≥90%.

Methylation of the Extract

Two milliliters of dichloromethane and 0.5 mL of diazomethane solution in ethyl ether were added to 1.4 mg of the ethanolic extract of the mucilaginous secretion. After evaporation of all of the solvent, the sample was resuspended with 1 mL of ethyl acetate and injected into the GC-MS system.

Infrared Spectroscopic Analysis

Infrared reflectance spectra were obtained in an infrared spectrometer with Fourier Transform (IR-TF) in the region of 4000–450 cm⁻¹. An attenuated total reflectance accessory was used in the Cary 630 – FTIR Spectrometer (Agilent Equipment). To make the sample suitable for this analysis, a potassium bromide (KBr) tablet was prepared from 7.9 mg of a dry sample of the jelly secretion.

RESULTS

Structural and Histochemical Characterization of the Collecters

The axis of each inflorescence was found to be surrounded by fibrous bracts, and each floral bud is in turn encased by equally rigid bracts (Figures 1B,D,E). In the reproductive buds and mature inflorescences of both species, the collectors produce a sticky secretion, which envelopes the floral buds and external parts of the flowers (Figures 1B,E). These secretory structures are present in the both surfaces of the bracts, bracteoles, sepals and on ovary wall, and floral column in *E. brasiliensis* (Figures 2A–H) and are very abundant, especially in the bracteoles (Figures 2A–D), which would explain the

more abundant secretion by this species (Figures 2A–D). In *E. crinipes*, collectors are also present in the bracts, bracteoles (Figures 3A,B,E,F), sepals (Figure 3A), and ovary wall (Figures 3C,D). However, their frequency is remarkably lower when the median region is analyzed. On five bracts of each species in an area of 141 mm² were found on average of 2.4 collectors in *E. crinipes* and 18 collectors in *E. brasiliensis*. In *E. crinipes* the collectors occur only in the median region, whereas in *E. brasiliensis* these structures are present throughout the bract. This characterizes the difference in the amount of secretion produced by these species (Figures 1E, 3B,E,F).

The collectors of *E. brasiliensis* are of a brush type (Figures 4A–E,G,H), each containing a short, cup-shaped shaft at the base composed of one to three cells (Figures 4B–E). In the terminal part, these structures bear three to seven elongated cells (Figures 4A,B,E). The axillary cells present lignified and suberified secondary wall deposition (Figures 4B–F,H). However, the elongated cells present only primary cell walls (Figures 4B–E) and constitute the site of secretion synthesis (Figure 4G). On the other hand, the walls of the elongated cells in connection with the axis cells also exhibit suberification (Figure 4H). The structure of collectors of *E. crinipes* is similar to that described for the collectors of *E. brasiliensis*, as seen in the Figures 3B,E,F. Nevertheless, the basal shaft is more elongated in *E. crinipes* (Figure 3B).

The content of the collectors of *E. brasiliensis* and the secretion produced by these structures are hydrophilic and lipophilic in nature (Table 1 and Figures 5A–H). In general, lipophilic substances and phenolic compounds are evident in the secretory cell vacuoles (Figures 5C,E,G, black arrows), and the hydrophilic substances occupy the entire protoplast (Figure 5B). The exudate is a mixture of these hydrophilic and lipophilic substances (Figure 5, arrowheads), containing pectic substances (Figure 5A), mucilage (Figure 5B, black arrow), lipids (Figures 5D,E), terpenoids (Figure 5F), phenolic compounds (Figure 5G), and proteins (Figure 5H).


Chemical Analysis of the Secretion From the Collecters

The GC-MS analysis confirmed the presence of the triterpene γ -sitosterol (MM = 414 g/mol) (Figure 6A), and of palmitic (MM = 270 g/mol), linoleic (MM = 292 g/mol), and stearic (MM = 298 g/mol) acids (Figure 6B). The analysis of the ethanolic extract of the inflorescence secretion by IR-TF qualitatively revealed the presence of functional groups characteristic of polysaccharides in the region of 4000–500 cm⁻¹ (Figure 6C). A broad and intense band between 3500 and 3000 cm⁻¹ was attributed to the presence of a hydroxyl (OH) group (Hammami et al., 2018). A less intense band at 2918.91 cm⁻¹ characterized the stretch between C–H bonds (Hou et al., 2018). At 1721.93 cm⁻¹, a low-intensity band attributed to the axial deformation of the C = O of enols (Silverstein et al., 2007) was detected (Figure 6C). The presence of medium-intensity absorption at 1413.87, 1378.88, and 1251.88 cm⁻¹ and high-intensity absorption at 1039.63 cm⁻¹ were also observed (Figure 6C), which are characteristic of the S = O bonds of sulfated esters (Silverstein et al., 2007; Webber et al., 2012).

TABLE 1 | Classes of substances evidenced, and their reagents used in *E. brasiliensis* collectors and its secretion.

Substance class	Reagents	Result of reaction*
Pectins	Coriophosphine	+++
Mucilage and pectic substances	Ruthenium red	+++
Total lipids	Sudan III	++
Total lipids	Sudan black B	+++
Neutral lipids	Nile blue sulfate	+++
Terpenes	Nadi reagent	++
Phenolic compounds	Ferric chloride	++
Proteins	Aniline blue black	+


*The intensity of reactions was indicated by: +++ (very intense reaction); ++ (intense reaction); + (little intense reaction).



DISCUSSION

Secretory structures are involved in the production of different substances, both in the vegetative and reproductive organs of plants. In flowers and inflorescences, these include such structures as idioblasts, glandular trichomes (Leite et al., 2018), collectors (Lacchia et al., 2016), laticifers (Marinho et al., 2018), osmophores, and floral and extra-floral nectaries (De Souza et al., 2005). Trichomes are classified into glandular and non-glandular trichomes. Glandular trichomes present specialized cells in the glandular head with the ability to produce, store and secrete

various substances (Fahn, 1979). According to the constituents of the secretion, the glandular trichomes receive a functional denomination like the nectaries (production of nectar) and the collectors (mucilage and/or lipid-like substances secretion) (Frões et al., 2015; Tian et al., 2017). In the Orchidaceae, researchers have previously observed the presence of nectaries (Leitão et al., 2014; Pansarin et al., 2015; Solano-Gómez et al., 2016), osmophores (Pansarin and Amaral, 2009; Millner and Baldwin, 2016; Caballero-Villalobos et al., 2017), idioblasts, and collectors (Leitão and Cortelazzo, 2008; Mayer et al., 2011; Cardoso-Gustavson et al., 2014).



Collecters have been described in at least 60 families of angiosperms (Thomas, 1991), including Apocynaceae (Apezatto-da-Glória and Estelita, 2000; Martins et al., 2013; Ribeiro et al., 2017), Bromeliaceae (Ballego-Campos and Paiva, 2018a,b), Euphorbiaceae (Vitarelli et al., 2015, 2016;

Feio et al., 2016; Martins et al., 2016), Fabaceae (Paiva and Machado, 2006; Oliveira and Isaias, 2010), and Rubiaceae (Judkevich et al., 2017; Paiva-Pinheiro et al., 2019). These secretory structures are located in the vegetative and floral buds, and remain in these plant organs throughout their life.

Usually, collectors produce a sticky secretion composed of a mixture of terpenes and mucilage (complex polymers of acidic or neutral polysaccharides of high molecular weight), which helps to retain water, preventing the desiccation of the meristem (Castro and Machado, 2013).

Our results confirmed the presence of collectors in the two studied species of *Elleanthus*. However, although the type of collector was very similar in both species, the frequency of occurrence of these structures differed between the species. In *E. crinipes*, these structures are scarcer (**Figure 3A**) and apparently have much less secretory activity (**Figure 3B**), if compared with the same region in *E. brasiliensis* (**Figure 2D**), as could be perceived in the SEM images obtained. Histochemical analyses found evidence of the presence of mucilage and pectin in the secretion of *E. brasiliensis*. These findings confirmed the function of these structures in the studied species. Further, together with recent findings in several species from distinct lineages of Epidendroideae, the occurrence of collectors in these two tropical orchids suggests that these structures are common features of the inflorescences of tropical orchids in general (Leitão and Cortelazzo, 2008; Mayer et al., 2011; Cardoso-Gustavson et al., 2014).

In *E. brasiliensis*, the conspicuousness of the mucilaginous secretion, significant amounts of which accumulate on the plant and within the collectors, also suggests that the secretion functions as a physical barrier enclosing the flower buds and external parts of the bases of the flower tubes. Such a physical barrier would exclude nectar thieves and robbers attempting illegitimate visits to the nectaries, as well as feeding by insect herbivores.

The results of the present study revealed the presence of the triterpene γ -sitosterol and different fatty acids in the secretion of *E. brasiliensis* and the apparent absence of fungi on the surface of the flower parts of this species in SEM images. Terpenes are molecules with great structural diversity and are produced in the leaves, stems, flowers, and occasionally in the roots (Dudareva et al., 2003). In addition to playing a role in growth and development (Logan et al., 2000), as well as in pollinator attraction to plants (Pichersky and Gershenzon, 2002), these substances protect them against attacks by herbivores and pathogenic microorganisms (Paré and Tumlinson, 1999; Tholl, 2006; Cheng et al., 2007). Fatty acids constitute virtually all plant tissues and also play a role in protecting the plant organs from attack by microorganisms (McGaw et al., 2002; Kachroo and Kachroo, 2009). The recognition of pathogens by plants results in the triggering of defensive responses (Ramirez-Prado et al., 2018). These molecules and their derivatives help in these responses both in protection against bacteria (16C fatty acids) and fungi (16C and 18C fatty acids) (Kachroo and Kachroo, 2009). These findings support the idea that this secretion also functions in the protection of the inflorescences against microorganisms. In this case, the hydrated polysaccharides form a dense matrix that may act as a physical barrier, while other chemical components, such as terpenes or the fatty acids, may help to protect the inflorescence against herbivores and pathogenic microorganisms.

Through the IR-TF analysis performed, the presence of absorption bands that characterized the possible presence of polysaccharides in the secretion was observed. The presence of polysaccharides was also confirmed by the reaction with ruthenium red. However, other chromatographic analyses will be required to characterize which polysaccharide(s) are included in the secretion. Polysaccharides help in the retention of water and provide viscosity to the secretion (Toneli et al., 2005). In this way, the secretion remains adhered to the inflorescence even if it occurs in a pendent way. Although the results aided in understanding the chemical composition of the secretion, it is important to note that the analyses were performed with the secretion of several individuals. This was necessary because of the extremely low concentrations of the dry material in the individual samples, which may hinder the reproduction of these analyses in the study with this species.

Indeed, previous observations of floral visitors did not record any insect herbivore feeding on the flower parts of these species, but rather only signals of herbivory by vertebrates (e.g., birds) on the inflorescences have been seen. Therefore, this secretion is likely one of the mechanisms used to direct floral resources to the main pollinators of these orchid species, hummingbirds (Nunes et al., 2013, 2016), rather than nuisance insects. Additionally, in *E. brasiliensis* the bright and translucent mucilage in the secretion refracts the color signal of the reddish floral bracts and sepals, increasing the apparent volume of the inflorescence and reinforcing the signal to bird pollinators without advertising as much to others, such as bee pollinators (Lunau et al., 2011).

CONCLUSION

Anatomical and SEM analyses revealed the presence of collectors in the inflorescences of both *Elleanthus* species studied. Although the type of collectors is very similar between these species, the low frequency of these structures' occurrence together with an apparently reduced secretory activity results in a decrease in the production of secretion by *E. crinipes*. The presence of polysaccharides, fatty acids, and terpenes implies the role of the secretion on the hydration and protection of the inflorescences of *E. brasiliensis*. This information will contribute to the characterization of species of the family Orchidaceae, both in terms of their morphological and anatomical aspects, as well as possible plant defenses against herbivores and pathogens.

AUTHOR CONTRIBUTIONS

FC carried out the chemical experiments. CN was responsible for collecting the material. ML carried out the anatomical and histochemical analyses. VG assisted and interpreted in the chemical analyses. JM designed, assisted, interpreted the anatomic analysis, and supervised the work. FC wrote the manuscript and CN, ML, and JM reviewed it.

ACKNOWLEDGMENTS

We would like to thank the Pró-Reitoria de Pós-Graduação (PRPG/UNICAMP) for the research support. FC and VG thank Sinéio Boaventura Júnior for the help in methylation analysis. We thank the Instituto Florestal (Parque Estadual da Serra do Mar, Núcleo Santa Virginia and Núcleo Picinguaba) for the development of the study on protected

public land. CN thank CNPQ (131934/2009-0), FAPESP (03/12595-7), COTEC/IF (41.065/2005), and IBAMA/CGEN (093/2005) the funding support. ML and FC would like to thank CAPES and FAPEX/UNICAMP for granting the scholarship. JM thank FAPESP (2015/26479-6) for funding support. We also thank the access to equipment and assistance provided by the Electron Microscope Laboratory (LME/UNICAMP).

REFERENCES

- Appezato-da-Glória, B., and Estelita, E. M. (2000). Development, structure and distribution of collectors in *Mandevilla illustris* and *M. velutina* (Apocynaceae). *Rev. Bras. Bot.* 23, 113–120. doi: 10.1590/S0100-84042000000200001
- Ballego-Campos, I., and Paiva, E. A. S. (2018a). Collecters in the vegetative axis of *Aechmea blanchetiana* (Bromeliaceae): anatomical, ultrastructural and functional aspects. *Aust. J. Bot.* 66, 379–387. doi: 10.1071/BT18095
- Ballego-Campos, I., and Paiva, E. A. S. (2018b). Mucilage secretion in the inflorescences of *Aechmea blanchetiana*: evidence of new functions of scales in Bromeliaceae. *Flora* 246, 1–9. doi: 10.1016/j.flora.2018.06.003
- Caballero-Villalobos, L., Silva-Arias, G. A., Buzatto, C. R., Nervo, M. H., and Singer, R. B. (2017). Generalized food-deceptive pollination in four *Cattleya* (Orchidaceae: Laeliinae) species from Southern Brazil. *Flora* 234, 195–206. doi: 10.1016/j.flora.2017.07.014
- Cain, A. J. (1947). The use of Nile Blue in the examination of lipoids. *Q. J. Microsc. Sci.* 88, 383–392. doi: 10.3109/10520296809115077
- Capelli, N. V., Rodrigues, B. A., and Demarco, D. (2017). Stipules in Apocynaceae: an ontogenetic perspective. *AoB Plants* 9, 1–11. doi: 10.1093/aobpla/plw083
- Cardoso-Gustavson, P., Campbell, L. M., Mazzoni-Viveiros, S. C., and de Barros, A. F. (2014). Floral collectors in Pleurothallidinae (Epidendroideae: Orchidaceae). *Am. J. Bot.* 101, 587–597. doi: 10.3732/ajb.1400012
- Castro, M. M., and Demarco, D. (2008). Phenolic compounds produced by secretory structures in plants: a brief review. *Nat. Prod. Commun.* 3, 1273–1284.
- Castro, M. M., and Machado, S. R. (2013). “Células e Tecidos Secretores,” in *Anatomia Vegetal*, eds B. Appezato-da-Glória and S. M. Carmello-Guerreiro (Viçosa: Editora UFV), 169.
- Cheng, A. X., Lou, Y. G., Mao, Y. B., Lu, S., Wang, L. J., and Chen, X. Y. (2007). Plant terpenoids: biosynthesis and ecological functions. *J. Integr. Plant Biol.* 49, 179–186. doi: 10.1111/j.1744-7909.2007.00395.x
- Coutinho, Í. A. C., Francino, D. M. T., and Meira, R. M. S. A. (2015). New records of collectors in *Chamaecrista* (Leguminosae, Caesalpinioideae s.l.): structural diversity, secretion, functional role, and taxonomic importance. *Int. J. Plant Sci.* 176, 72–85. doi: 10.1086/679016
- David, R., and Carde, J. P. (1964). Coloration différentielle des inclusions lipidique et terpeniques des pseudophylles du *Pin maritime* au moyen du reactif Nadi. *Comptes Rendus L. Acad. Des. Sci.* 258, 1338–1340.
- De Souza, R. C. O. S., Toni, K. L. G., Andreato, R. H. P., and Costa, C. G. (2005). Anatomia e vascularização das flores estaminadas e pistiladas de *Smilax fluminensis* Steudel (Smilacaceae). *Rodriguésia* 56, 107–121. doi: 10.1590/2175-78602005568708
- Dressler, R. L. (2006). “Eleanthus,” in *Genera Orchidacearum Volume 4: Epidendroideae (Part 1)*, eds A. M. Pridgeon, P. J. Cribb, M. W. Chase, and F. N. Rasmussen (Oxford: Oxford University Press), 598.
- Dudareva, N., Martin, D., Kish, C. M., Kolosova, N., Gorenstein, N., Fäldt, J., et al. (2003). (E)-beta-ocimene and myrcene synthase genes of floral scent biosynthesis in Snapdragon: function and expression of three terpene synthase genes of a new terpene synthase subfamily. *Plant Cell* 15, 1227–1241. doi: 10.1105/tpc.011015
- Dudek, M., Baranow, P., Kolanowska, M., and Rykaczewski, M. (2017). *Eleanthus albiflorus* (Orchidaceae) a new, white-flowered species from Peru. *Phytotaxa* 312, 256–262. doi: 10.11646/phytotaxa.312.2.8
- Fahn, A. (1979). *Secretory Tissues in Plants*. London: Academic Press.
- Feio, A. C., Riina, R., and Meira, R. M. S. A. (2016). Secretory structures in leaves and flowers of two dragon's blood *Croton* (Euphorbiaceae): new evidence and interpretations. *Int. J. Plant Sci.* 177, 511–522. doi: 10.1086/685705
- Fisher, D. B. (1968). Protein staining of ribboned epon sections for light microscopy. *Histochemie* 16, 92–96. doi: 10.1007/BF00306214
- Fróes, F. F. P. C., Gama, T. S. S., Feio, A. C., Demarco, D., and Aguiar-Dias, A. C. A. (2015). Structure and distribution of glandular trichomes in three species of Bignoniaceae. *Acta Amaz.* 45, 347–354. doi: 10.1590/1809-4392201404393
- Hammami, N., Gara, A. B., Bargougui, K., Ayedi, H., Abdalleh, F., Ben, et al. (2018). Improved *in vitro* antioxidant and antimicrobial capacities of polysaccharides isolated from *Salicornia arabica*. *Int. J. Biol. Macromol.* 120, 2123–2130. doi: 10.1016/j.ijbiomac.2018.09.052
- Horridge, G. A., and Tamm, S. L. (1969). Critical point drying for scanning electron microscopy study of ciliary motion. *Science* 163, 817–818. doi: 10.1126/science.163.3869.817
- Hou, G., Chen, X., Li, J., Ye, Z., Zong, S., and Ye, M. (2018). Physicochemical properties, immunostimulatory activity of the *Lachnum* polysaccharide and polysaccharide-dipeptide conjugates. *Carbohydr. Polym.* 73, 228–238. doi: 10.1016/j.carbpol.2018.09.067
- Jensen, W. A. (1962). *Botanical Histochemistry: Principles and Practices*. San Francisco, CA: Freeman and Company.
- Johansen, D. A. (1940). *Plant Microtechnique*. New York, NY: McGraw-Hill Book Co.
- Judkevich, M. D., Salas, R. M., and Gonzalez, A. M. (2017). Collecters in american *Spermacoceae* genera (Rubiaceae): morphoanatomical and evolutionary aspects. *Int. J. Plant Sci.* 178, 378–397. doi: 10.1086/691165
- Kachroo, A., and Kachroo, P. (2009). Fatty acid-derived signals in plant defense. *Annu. Rev. Phytopathol.* 47, 153–176. doi: 10.1146/annurev-phyto-080508-081820
- Karnovsky, M. J. (1965). A formaldehyde-glutaraldehyde fixative of high osmolarity of use in electron microscopy. *J. Cell Biol.* 27, 137–138.
- Lacchia, A. P. S., Tölke, E. E. A. D., Carmello-Guerreiro, S. M., Ascensão, L., and Demarco, D. (2016). Foliar collectors in Anacardiaceae: first report for the family. *Botany* 94, 1–10. doi: 10.1139/cjb-2015-0236
- Leitão, C. A. E., and Cortelazzo, A. L. (2008). Structural and histochemical characterisation of the collectors of *Rodriguezia venusta* (Orchidaceae). *Aust. J. Bot.* 56, 161–165. doi: 10.1071/BT07114
- Leitão, C. A. E., Dolder, M. A. H., and Cortelazzo, A. L. (2014). Anatomy and histochemistry of the nectaries of *Rodriguezia venusta* (Lindl.) Rchb. f. (Orchidaceae). *Flora* 209, 233–243. doi: 10.1016/j.flora.2014.03.002
- Leite, V. G., Mansano, V. F., and Teixeira, S. P. (2018). Floral development of Moraceae species with emphasis on the perianth and androecium. *Flora* 240, 116–132. doi: 10.1016/j.flora.2018.01.009
- Logan, B. A., Monson, R. K., and Potosnak, M. J. (2000). Biochemistry and physiology of foliar isoprene production. *Trends Plant Sci.* 5, 477–481. doi: 10.1016/S1360-1385(00)01765-9
- Lunau, K., Papiorek, S., Eltz, T., and Sazima, M. (2011). Avoidance of achromatic colours by bees provides a private niche for hummingbirds. *J. Exp. Biol.* 214, 1607–1612. doi: 10.1242/jeb.052688
- Lusa, M. G., Cardoso, E. C., Machado, S. R., and Appezato-da-Glória, B. (2015). Trichomes related to an unusual method of water retention and protection of the stem apex in an arid zone perennial species. *AoB Plants* 7, 1–10. doi: 10.1093/aobpla/plu088
- Macêdo, T. P., Cortez, P. A., and Costa, L. C. B. (2016). First record of collectors in *Zanthoxylum* Linn. species (*Rutaceae* Juss., *Sapindales*): structural, functional and taxonomic considerations. *Flora* 224, 66–74. doi: 10.1016/j.flora.2016.07.007
- Machado, S. R., Palarci, L. M., Paiva, E. A. S., and Rodrigues, T. M. (2015). Collecters on the inflorescence axis of *Croton glandulosus* (Euphorbiaceae): structural

- and functional characterization. *Int. J. Plant Sci.* 176, 86–93. doi: 10.1086/678469
- Machado, S. R., and Rodrigues, T. M. (2013). “Estruturas secretoras externas,” in *Anatomia das plantas de Esau – meristemas, células e tecidos do corpo da planta: sua estrutura, função e desenvolvimento*, ed. E. R. Franklin (São Paulo: Blucher), 548–549.
- Marinho, C. R., Pereira, R. A. S., Peng, Y.-Q., and Teixeira, S. P. (2018). Laticifer distribution in fig inflorescence and its potential role in the fig-fig wasp mutualism. *Acta Oecol.* 90, 160–167. doi: 10.1016/j.actao.2017.10.005
- Martins, F. M., Cunha-Neto, I. L., and Pereira, T. M. (2016). Floral morphology and anatomy of *Dalechampia alata* Klotzsch ex Baill. (Euphorbiaceae), with emphasis on secretory structures. *Braz. J. Biol.* 76, 233–244. doi: 10.1590/1519-6984.19514
- Martins, F. M., Mascarenhas, A. A. S., Macedo, T. P., and Cunha Neto, I. L. (2013). Estruturas secretoras em órgãos vegetativos e florais de *Secondatia densiflora* A.DC. (Apocynaceae - Apocynoideae - Odontadenieae). *Rev. Bras. Plantas Med.* 15, 13–24. doi: 10.1590/S1516-05722013000100002
- Mayer, J. L. S., Cardoso-Gustavson, P., and Appezzato-da-Glória, B. (2011). Colleters in monocots: new record for orchidaceae. *Flora* 206, 185–190. doi: 10.1016/j.flora.2010.09.003
- Mayer, J. L. S., Carmello-Guerreiro, S. M., and Mazzafera, P. (2013). A functional role for the colleters of coffee flowers. *AoB Plants* 5, 1–13. doi: 10.1093/aobpla/plt029
- McGaw, L. J., Jäger, A. K., van Staden, J., and Houghton, P. J. (2002). Antibacterial effects of fatty acids and related compounds from plants. *S. Afr. J. Bot.* 68, 417–423. doi: 10.1016/S0254-6299(15)30367-7
- Miguel, E. C., Gomes, V. M., Oliveira, M. A., and Cunha, M. (2006). Colleters in *Bathysa nicholsonii* K. Schum. (Rubiaceae): ultrastructure, secretion protein composition, and antifungal activity. *Plant Biol.* 8, 715–722. doi: 10.1055/s-2006-924174
- Millner, H. J., and Baldwin, T. C. (2016). Floral micromorphology of the genus *Restrepia* (Orchidaceae) and the potential consequences for pollination. *Flora* 225, 10–19. doi: 10.1016/j.flora.2016.09.007
- Nunes, C., Castro, M. M., Galetto, L., and Sazima, M. (2013). Anatomy of the floral nectary of ornithophilous *Elleanthus brasiliensis* (Orchidaceae: Sobralieae). *Bot. J. Linn. Soc.* 171, 764–772. doi: 10.1111/boj.12024
- Nunes, C. E. P., Amorim, F. W., Mayer, J. L. S., and Sazima, M. (2016). Pollination ecology of two species of *Elleanthus* (Orchidaceae): novel mechanisms and underlying adaptations to hummingbird pollination. *Plant Biol.* 18, 15–25. doi: 10.1111/plb.12312
- Oliveira, D. C., and Isaias, R. M. S. (2010). Redifferentiation of leaflet tissues during midrib gall development in *Copaifera langsdorffii* (Fabaceae). *S. Afr. J. Bot.* 76, 239–248. doi: 10.1016/j.sajb.2009.10.011
- Paiva, E. A. S., and Machado, S. R. (2006). Ontogenesis, structure and ultrastructure of *Hymenaea stigonocarpa* (Fabaceae: Caesalpinioideae) colleters. *Rev. Biol. Trop.* 54, 943–950.
- Paiva-Pinheiro, S. K., Teófilo, F. B. S., Lima, A. K. M., Cordoba, B. V., Miguel, T. B. A. R., and Castro-Miguel, E. (2019). Ontogenesis and secretion mechanism of *Morinda citrifolia* L. (Rubiaceae) colleters. *S. Afr. J. Bot.* 121, 26–33. doi: 10.1016/j.sajb.2018.10.015
- Pansarin, E. R., and Amaral, M. C. E. (2009). Reproductive biology and pollination of southeastern Brazilian *Stanhopea* Frost ex Hook. (Orchidaceae). *Flora* 204, 238–249. doi: 10.1016/j.flora.2008.01.014
- Pansarin, E. R., Pansarin, L. M., and Alves-dos-Santos, I. (2015). Floral features, pollination biology, and breeding system of *Comparettia coccinea* (Orchidaceae: Oncidiinae). *Flora* 217, 57–63. doi: 10.1016/2015.09.008
- Paré, P. W., and Tumlinson, J. H. (1999). Plant volatiles as a defense against insect herbivores. *Plant Physiol.* 121, 325–331. doi: 10.1104/pp.121.2.325
- Pearse, A. G. E. (1968). *Histochemistry, Theoretical and Applied*, 3rd Edn. London: Churchill Livingstone.
- Pichersky, E., and Gershenzon, J. (2002). The formation and function of plant volatiles: perfumes for pollinator attraction and defense. *Curr. Opin. Plant Biol.* 5, 237–243. doi: 10.1016/S1369-5266(02)00251-0
- Ramirez-Prado, J. S., Abulfaraj, A. A., Rayapuram, N., Benhamed, M., and Hirt, H. (2018). Plant immunity: from signaling to epigenetic control of defense. *Trends Plant Sci.* 23, 833–844. doi: 10.1016/j.tplants.2018.06.004
- Ribeiro, J. C., Ferreira, M. J. P., and Demarco, D. (2017). Colleters in Asclepiadoideae (Apocynaceae): protection of meristems against desiccation and new functions assigned. *Int. J. Plant Sci.* 178, 465–477.
- Sakai, W. S. (1973). Simple method for differential staining of paraffin embedded plant material using toluidine blue O. *Stain Technol.* 48, 247–249. doi: 10.3109/10520297309116632
- Silverstein, R. M., Webster, F. X., and Kiemle, D. J. (2007). “Espectrometria no Infravermelho,” in *Identificação Espectrométrica de Compostos Orgânicos*, eds R. M. Silverstein, F. X. Webster, and D. J. Kiemle (Rio de Janeiro: LTC), 70–104.
- Solano-Gómez, R., Martínez-Ovando, E., Martínez-Feria, A., and Gutiérrez-Caballero, J. A. (2016). New records in the Orchidaceae family from Oaxaca, Mexico. *Rev. Mex. Biodivers.* 87, 1348–1351. doi: 10.1016/j.rmb.2016.09.012
- Srebotnik, E., and Messner, K. (1994). A simple method that uses differential staining and light microscopy to assess the selectivity of wood delignification by white rot fungi. *Appl. Environ. Microbiol.* 60, 1383–1386. doi: 10.1038/244060a0
- Tholl, D. (2006). Terpene synthases and the regulation, diversity and biological roles of terpene metabolism. *Curr. Opin. Plant Biol.* 9, 1–8. doi: 10.1016/j.pbi.2006.03.014
- Thomas, V. (1991). Structural, functional and phylogenetic aspects of the colleter. *Ann. Bot.* 68, 287–305. doi: 10.1093/oxfordjournals.aob.a088256
- Tian, N., Liu, F., Wang, P., Zhang, X., Li, X., and Wu, G. (2017). The molecular basis of glandular trichome development and secondary metabolism in plants. *Plant Gene* 12, 1–12. doi: 10.1016/j.plgene.2017.05.010
- Toneli, J. T. C. L., Murr, F. E. X., and Park, K. J. (2005). Estudo da reologia de polissacarídeos utilizados na indústria de alimentos. *Rev. Bras. Prod. Agroindustriais* 7, 181–204. doi: 10.15871/1517-8595/rbpa.v7n2p181-204
- Ueda, K., and Yoshioka, S. (1976). Cell wall development of *Micrasterias americana*, especially in isotonic and hypertonic solutions. *J. Cell Sci.* 21, 617–631.
- Veloso, H. P., Rangel-Filho, A. L. R., and Lima, J. C. A. (1991). *Classificação da Vegetação Brasileira, Adaptada a um Sistema Universal*. Rio de Janeiro: Fundação Instituto Brasileiro de Geografia e Estatística – IBGE.
- Vitarelli, N. C., Riina, R., Caruzo, M. B. R., Cordeiro, I., Fuertes-Aguilar, J., and Meira, R. M. S. A. (2015). Foliar secretory structures in Crotonaeae (Euphorbiaceae): diversity, anatomy, and evolutionary significance. *Am. J. Bot.* 102, 833–847. doi: 10.3732/ajb.1500017
- Vitarelli, N. C., Riina, R., Cassino, M. F., and Meira, R. M. S. A. (2016). Trichome-like emergences in *Croton* of Brazilian highland rock outcrops: evidences for atmospheric water uptake. *Perspect. Plant Ecol. Evol. Syst.* 22, 23–35. doi: 10.1016/j.ppees.2016.07.002
- Webber, V., de Carvalho, S. M., Ogliari, P. J., Hayashi, L., and Barreto, P. L. M. (2012). Optimization of the extraction of carrageenan from *Kappaphycus alvarezii* using response surface methodology. *Food Sci. Technol.* 32, 812–818. doi: 10.1590/S0101-20612012005000111
- Whittier, D. P., and Peterson, R. L. (1984). Gametophytes of *Botrychium lunarioides* and their mucilage-coated rhizoids. *Can. J. Bot.* 62, 2854–2860. doi: 10.1139/b84-380

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Cassola, Nunes, Lusa, Garcia and Mayer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.