



The Immune Receptor Roq1 Confers Resistance to the Bacterial Pathogens *Xanthomonas*, *Pseudomonas syringae*, and *Ralstonia* in Tomato

Nicholas C. Thomas^{1,2}, Connor G. Hendrich³, Upinder S. Gill^{4,5}, Caitilyn Allen³, Samuel F. Hutton⁴ and Alex Schultink^{1,2*}

¹ Fortiphyte Inc., Berkeley, CA, United States, ² Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA, United States, ³ Department of Plant Pathology, University of Wisconsin–Madison, Madison, WI, United States, ⁴ IFAS, Gulf Coast Research and Education Center, University of Florida, Wimauma, FL, United States, ⁵ Department of Plant Pathology, North Dakota State University, Fargo, ND, United States

OPEN ACCESS

Edited by:

Marc Valls,
University of Barcelona, Spain

Reviewed by:

Nemo Peeters,
Institut National de la Recherche
Agronomique (INRA), France
Kee Hoon Sohn,
Pohang University of Science
and Technology, South Korea
Kouhei Ohnishi,
Kochi University, Japan

*Correspondence:

Alex Schultink
alex.schultink@fortiphyte.com

Specialty section:

This article was submitted to
Plant Microbe Interactions,
a section of the journal
Frontiers in Plant Science

Received: 24 December 2019

Accepted: 30 March 2020

Published: 23 April 2020

Citation:

Thomas NC, Hendrich CG,
Gill US, Allen C, Hutton SF and
Schultink A (2020) The Immune
Receptor Roq1 Confers Resistance
to the Bacterial Pathogens
Xanthomonas, *Pseudomonas*
syringae, and *Ralstonia* in Tomato.
Front. Plant Sci. 11:463.
doi: 10.3389/fpls.2020.00463

Xanthomonas species, *Pseudomonas syringae* and *Ralstonia* species are bacterial plant pathogens that cause significant yield loss in many crop species. Generating disease-resistant crop varieties can provide a more sustainable solution to control yield loss compared to chemical methods. Plant immune receptors encoded by nucleotide-binding, leucine-rich repeat (NLR) genes typically confer resistance to pathogens that produce a cognate elicitor, often an effector protein secreted by the pathogen to promote virulence. The diverse sequence and presence/absence variation of pathogen effector proteins within and between pathogen species usually limits the utility of a single NLR gene to protecting a plant from a single pathogen species or particular strains. The NLR protein Recognition of XopQ 1 (Roq1) was recently identified from the plant *Nicotiana benthamiana* and mediates perception of the effector proteins XopQ and HopQ1 from *Xanthomonas* and *P. syringae* respectively. Unlike most recognized effectors, alleles of XopQ/HopQ1 are highly conserved and present in most plant pathogenic strains of *Xanthomonas* and *P. syringae*. A homolog of XopQ/HopQ1, named RipB, is present in most *Ralstonia* strains. We found that Roq1 confers immunity to *Xanthomonas*, *P. syringae*, and *Ralstonia* when expressed in tomato. Strong resistance to *Xanthomonas perforans* was observed in three seasons of field trials with both natural and artificial inoculation. The *Roq1* gene can therefore be used to provide safe, economical, and effective control of these pathogens in tomato and other crop species and reduce or eliminate the need for traditional chemical controls.

Keywords: plant immunity, *Ralstonia*, *Xanthomonas*, *Pseudomonas*, tomato

INTRODUCTION

Bacterial pathogens from the species *Pseudomonas syringae* and the genera *Ralstonia* and *Xanthomonas* can infect many different crop species and inflict significant yield losses when environmental conditions favor disease. *Xanthomonas* and *P. syringae* tend to enter plant stem, leaf, or flower tissue through wounds or natural openings, such as stomata or hydathodes, whereas

Ralstonia is soilborne, entering roots through wounds and natural openings before colonizing xylem tissue (Vasse et al., 1995; Gudesblat et al., 2009). Once inside the host these bacteria manipulate host metabolism and suppress plant immunity using multiple strategies, including effector proteins delivered by the type III secretion system (Kay and Bonas, 2009; Peeters et al., 2013; Xin et al., 2018). This enables the pathogens to multiply to high titers while the plant tissue is still alive and showing few or no visual symptoms. Once the bacteria reach high populations, they typically cause necrosis of infected leaf tissue or wilting and eventual death of the plant.

Effective control measures for bacterial pathogens are relatively limited, particularly once plants become infected (Davis et al., 2013). Soil fumigation can reduce *Ralstonia* populations in the soil but this is expensive, potentially hazardous to workers and the environment, and of limited efficacy (Yuliar et al., 2015). Copper sulfate and antibiotics such as streptomycin have been used to control *Xanthomonas* species and *P. syringae* but have adverse environmental impacts and many strains have evolved tolerance to these chemicals (Kennelly et al., 2007; Griffin et al., 2017). Applying chemicals that induce systemic acquired resistance, such as acibenzolar-S-methyl, can provide partial control but increases production cost and can depress crop yields when used repeatedly (de Pontes et al., 2016).

The most effective, economical, and safe way to control bacterial pathogens is to plant crop varieties that are immune to the target pathogen (Jones et al., 2014; Vincelli, 2016). Such immunity is often mediated by plant immune receptor genes. Plants have large families of cell surface and intracellular immune receptor proteins that surveil for the presence of invading pathogens (Zipfel, 2014; Jones et al., 2016). Effector proteins delivered by the bacterial type III secretion system are common elicitors of intracellular plant immune receptors encoded by nucleotide-binding domain and leucine-rich repeat containing (NLR) genes (Li et al., 2015; Jones et al., 2016; Kapos et al., 2019). While effector proteins contribute to virulence on a susceptible host, an immune response is activated in the plant if that plant has the cognate receptor to recognize the effector. NLR genes typically confer strong, dominant resistance to pathogens that deliver the cognate recognized effector protein (Tai et al., 1999; Jones and Dangl, 2006; Boller and He, 2009; Deslandes and Rivas, 2012; Li et al., 2015). Disease-resistant plants can be generated by identifying the appropriate plant immune receptor genes and transferring them into the target crop species (Dangl et al., 2013).

We recently identified the *Nicotiana benthamiana* immune receptor gene named *Recognition of XopQ 1 (Roq1)*, which appears to be restricted to the genus *Nicotiana* and contributes to resistance against *Xanthomonas* spp. and *P. syringae* (Schultink et al., 2017). The Roq1 protein is a Toll/Interleukin-1 Receptor (TIR) NLR immune receptor that mediates recognition of the *Xanthomonas* effector protein XopQ and the homologous effector HopQ1 from *P. syringae*. XopQ is present in most species and strains of *Xanthomonas* (Ryan et al., 2011) and HopQ1 is present in 62% (290 of 467) sequenced putative pathogenic *P. syringae* strains (Dillon et al., 2019). XopQ/HopQ1 has homology to nucleoside hydrolases and has been shown to enhance virulence on susceptible hosts (Ferrante and Scortichini, 2009;

Li et al., 2013), possibly by altering cytokinin levels or interfering with the activity of host 14-3-3 proteins (Giska et al., 2013; Li et al., 2013; Hann et al., 2014; Teper et al., 2014). The conservation of XopQ/HopQ1 and their importance in virulence suggests that *Roq1* has widespread potential to confer resistance to these pathogens in diverse crop species. Indeed, transient expression assays demonstrated that *Roq1* can recognize XopQ/HopQ1 alleles from *Xanthomonas* and *P. syringae* pathogens of tomato, pepper, rice, citrus, cassava, brassica, and bean (Schultink et al., 2017). However, it was not known if *Roq1* can confer disease resistance when expressed in a crop plant.

Tomato is one of the most important vegetable crops and is highly susceptible to several bacterial diseases. Bacterial spot, bacterial speck, and bacterial wilt of tomato are caused by *Xanthomonas* species, *P. syringae* pv. *tomato* and *Ralstonia*, respectively. These diseases are difficult to control, especially if the pathogens become established in a field and environmental conditions favor disease (Rivard et al., 2012; Potnis et al., 2015). Tomato breeding germplasm has only limited resistance against these diseases and in some cases linkage drag has complicated introgression of resistance genes from wild relatives (Sharma and Bhattarai, 2019). *Ralstonia* contains a homolog of XopQ/HopQ1 called RipB. Roq1 is able to mediate the perception of RipB (Staskawicz and Schultink, 2019), and silencing *Roq1* in *N. benthamiana* resulted in severe wilting phenotypes upon *Ralstonia* infection (Nakano and Mukaihara, 2019). This suggests that expressing *Roq1* in tomato could also confer resistance to bacterial wilt. Like XopQ/HopQ1 in *Xanthomonas* and *P. syringae*, RipB is highly conserved and is present in approximately 90% of sequenced *Ralstonia* isolates (Sabbagh et al., 2019). Here we present data showing that expression of *Roq1* in tomato confers resistance against *Xanthomonas*, *Pseudomonas*, and *Ralstonia* upon recognition of the cognate pathogen effector.

MATERIALS AND METHODS

Generation of Tomato Expressing Roq1

The Roq1 coding sequence was amplified from *N. benthamiana* cDNA and cloned into the pORE E4 binary plasmid (Coutu et al., 2007). The expression of *Roq1* was driven by the constitutive PENTCUP2 promoter, which was derived from tobacco and has been reported to drive expression in leaf, root, and stem tissue (Malik et al., 2002). *Agrobacterium tumefaciens* co-cultivation was used to transform *Roq1* into the tomato variety Fla. 8000 at the University of Nebraska Plant Transformation Core Research Facility. Transformed plants were selected by resistance to kanamycin, confirmed by genotyping, and selfed to obtain homozygous lines.

Bacterial Leaf Spot and Leaf Speck Disease Assays

Xanthomonas cultures were grown in NYG broth (0.5% peptone, 0.3% yeast extract, 2% glycerol) with rifampicin (100 µg/mL) overnight at 30°C. *P. syringae* cultures were grown in KB broth (1% peptone, 0.15% K₂HPO₄, 1.5% glycerol, 5 mM

MgSO₄, pH 7.0) with rifampicin (100 µg/mL) overnight at 28°C. Bacterial cultures were spun down at 5200 g, washed once with 10 mM MgCl₂, and then diluted to the appropriate infiltration density with 10 mM MgCl₂. Leaf tissue of tomato plants (approximately 4 weeks old) was infiltrated with bacterial solution using a needleless syringe. To quantify bacterial growth, leaf punches were homogenized in water, serially diluted and plated on NYG (for *Xanthomonas* spp.) or KB (for *P. syringae*) plates supplemented with 100 µg/mL rifampicin and 50 µg/mL cycloheximide to measure colony forming units. Three biological replicates were performed for each condition and the reported results are representative of at least three independent experiments. *Xanthomonas perforans* strain 4B, *Xanthomonas euvesicatoria* strain 85-10, and *P. syringae* strain DC3000 and the corresponding XopQ/HopQ1 deletion mutants and complemented strains were described previously (Wei et al., 2007; Schwartz et al., 2015; Schultink et al., 2017). The *P. syringae* pv. *tomato* Race 1 strain was isolated from a field of tomatoes with the PTO resistance gene in 1993 in California.

Transient Expression of RipB and XopQ

Alleles of RipB from *Ralstonia* strains GMI1000 and MolK2 (NCBI Genbank accessions CAD13773.2 and WP_003278485) were synthesized and cloned into a *Bsa*I-compatible version of the pORE E4 vector (Coutu et al., 2007). This plasmid was transformed into *A. tumefaciens* strain C58C1. *A. tumefaciens* cultures were grown on a shaker overnight at 30°C in LB broth with rifampicin (100 µg/mL), tetracycline (10 µg/mL), and kanamycin (50 µg/mL). The cells were collected by centrifugation and resuspended in infiltration buffer [10 mM 2-(*N*-morpholino)ethanesulfonic acid, 10 mM MgCl₂, pH 5.6], and diluted to an OD₆₀₀ of 0.5 for infiltration into *Nicotiana tabacum* leaf tissue. Each experiment was performed on multiple leaves and multiple plants with the selected images being representative of the observed result.

N. tabacum roq1 Mutant Lines

Nicotiana tabacum roq1 mutant lines were generated by transforming *N. tabacum* with a construct coding for CAS9 and a guide RNA targeting the *Roq1* gene with the sequence GATGATAAGGAGTTAAAGAG. This construct was also used for the generation of *N. benthamiana* roq1 mutants published in Qi et al. (2018). Transformed *N. tabacum* plants were generated by *Agrobacterium* co-cultivation and selected for using kanamycin. Transformed plants were genotyped for the presence of mutations at the target site by PCR and Sanger sequencing (Supplementary Table S1).

Bacterial Wilt Virulence Assays

Ralstonia virulence on tomato was measured as previously described (Khokhani et al., 2018). Briefly, cells of *Ralstonia* strains GMI1000 and UW551 grown overnight in CPG (0.1% casein hydrolysate, 1% peptone, 0.5% glucose, pH 7.0) at 28°C were collected by centrifugation and diluted to an OD₆₀₀ of 0.1 in water (1 × 10⁸ CFU/mL). 50 mL of this suspension was poured on the soil around 17-day old tomato plants. Disease was rated daily for two weeks on a 0–4 disease index scale, where 0 is no

leaves wilted, 1 is 1–25% wilted, 2 is 26–50% wilted, 3 is 51–75% wilted, and 4 is 76–100% wilted. Data represent a total of four biological replicates with 10 plants per replicate. Virulence data were analyzed using repeated measures ANOVA (Khokhani et al., 2018). For petiole infection, the petiole of the first true leaf was cut with a razor blade horizontally approximately 1 cm from the stem. A drop of bacterial solution (2 µL, OD₆₀₀ = 0.001) was pipetted onto the exposed cut petiole surface.

Field Trial Disease Assays


Three field trials were conducted at the University of Florida Gulf Coast Research and Education Center in Balm during the spring seasons of 2018 and 2019 and the fall season of 2018 and under the notification process of the United States Department of Agriculture. Large-fruited, fresh market tomato lines were used in these trials and included the inbred line, Fla. 8000, and nearly isogenic lines containing either *Roq1* (event 316.4) or *Bs2* (Kunwar et al., 2018). The *Roq1* tomato line selected for the field trial was the same line used in the experiments shown in Figures 1, 2, 5. For each trial, seeds were sown directly into peat-lite soilless media (Speedling, Sun City, FL, United States) in 128-cell trays (38 cm³ cell size). Transplants were grown in a greenhouse until 5 or 6 weeks, then planted to field beds that had been fumigated and covered with reflective plastic mulch. Field trials were conducted using a randomized complete block design with four blocks and 10-plant plots. Field plants were staked and tied, and irrigation was applied through drip tape beneath the plastic mulch of each bed. A recommended fertilizer and pesticide program were followed throughout the growing season, excluding the use of plant defense inducers, copper, or other bactericides (Freeman et al., 2018). Fruits were harvested from the inner eight plants of each plot at the breaker stage and beyond and graded for marketability according to USDA specifications with block considered a random effect.

Field trials were inoculated with *X. perforans* race T4 (strain mixture of GEV904, GEV917, GEV1001, and GEV1063). Bacterial strains were grown on nutrient agar medium (BBL, Becton Dickinson and Co., Cockeysville, MD, United States) and incubated at 28°C for 24 h. Bacterial cells were removed from the plates and suspended in a 10 mM MgSO₄ solution, and the suspension was adjusted to OD₆₀₀ = 0.3, which corresponds to 10⁸ CFU/mL. The suspension for each strain was then diluted to 10⁶ CFU/mL, mixed in equal volume, and applied along with polyoxyethylene sorbitan monolaurate (Tween 20; 0.05% [vol/vol]) for field inoculation. Field trial plants were inoculated approximately 3 weeks after transplanting.

Bacterial spot disease severity was recorded three to eight weeks after inoculation using the Horsfall-Barratt scale (Horsfall and Barratt, 1945), and ratings were converted to midpoint percentages for statistical analysis. Blocks were considered random effects.

Generation of the *Ralstonia* Δ ripB Mutant

An unmarked Δ ripB mutant was created using *sacB* selection with the vector pUFR80 (Castañeda et al., 2005). Briefly, the



regions upstream and downstream of *ripB* were amplified using the primers *ripBupF/R* and *ripBdwnF/R* (**Supplementary Table S1**). These fragments were inserted into pUFR80 digested with *HindIII* and *EcoRI* using Gibson Assembly (Gibson et al., 2009) (New England Biolabs, Ipswich, MA, United States) and this construct was incorporated into the genome of strain GMI1000 using natural transformation, with successful integrants selected on CPG + kanamycin (Coupat et al., 2008). Plasmid loss was then selected for on CPG plates containing 5% w/v sucrose. Correct deletions were confirmed using PCR and sequencing.

Phylogenetic Analysis of *XopQ*, *HopQ1*, and *RipB* Alleles

RipB alleles were identified by BLAST search of the NCBI protein database. Clustal Omega (Sievers et al., 2011) was used to generate a multiple sequence alignment with *XopQ* and *HopQ1* alleles. To span the diversity of *RipB* alleles without having many redundant sequences, only a single sequence was retained if there were multiple identical or nearly identical sequences identified. A maximum likelihood tree was generated using PhyML (Guindon et al., 2010). The phylotype calls of the strains were obtained from previously published work (Liu et al., 2009; Mukaiharu and Tamura, 2009; Safni et al., 2014).

RESULTS

Tomatoes Expressing *Roq1* Are Resistant to *Xanthomonas* and *P. syringae*

We generated homozygous tomato plants expressing the *Roq1* gene from *N. benthamiana* and tested them for resistance to *Xanthomonas* and *P. syringae* by measuring bacterial growth *in planta*. Population sizes of wild-type *X. perforans* strain 4B and *X. euvesicatoria* strain 85-10 were approximately 100-fold smaller in tomato expressing *Roq1* compared to wild-type tomato at 6 days post inoculation (**Figure 1**). In contrast, *XopQ* deletion mutants multiplied equally well in leaves of both wild-type and *Roq1* tomato. Disease symptoms begin as small water-soaked lesions and progress to necrosis of infected tissue. Wild-type *X. perforans* and *X. euvesicatoria* caused severe disease symptoms on wild-type tomato plants but failed to cause visible symptoms on *Roq1* plants (**Figure 2**). The *XopQ* mutants caused similar disease symptoms on both wild-type and *Roq1* tomato. Similar results were observed for *P. syringae* DC3000, and its *HopQ1* mutant (**Figures 1, 2**) and a Race 1 isolate of *P. syringae* pv. *tomato* (**Supplementary Figure S1**). Tomatoes expressing *Roq1* were resistant to *Xanthomonas* and *Pseudomonas* *XopQ/HopQ1* mutants complemented with a wild-type copy of *XopQ/HopQ1* (**Supplementary Figure S2**). A second tomato line expressing *Roq1*, derived from an independent transformation event, also showed resistance to wild-type *X. euvesicatoria* and *X. perforans* but not to the *XopQ* deletion strains (**Supplementary Figure S3**).




FIGURE 2 | Bacterial disease symptoms on *Roq1* tomato. *Xanthomonas perforans* 4B (*Xp*), *Xanthomonas euvesicatoria* 85-10 (*Xe*), and *Pseudomonas syringae* DC3000 (*Ps*) wild-type and *XopQ*/*HopQ1* knockout strains were infiltrated into tomato leaf tissue at low inoculum and disease symptoms were imaged at 12, 13, and 4 days post infiltration for *Xe*, *Xp*, and *Ps*, respectively. The infiltration was performed using a needleless syringe and circular wounds from the infiltration are visible. The distal part of region of each leaf was infiltrated and the proximal part was left untreated. *Xe* and *Xp* were infiltrated at an OD_{600} of 0.0001 whereas *Ps* was infiltrated at an OD_{600} of 0.00005. The images shown here are representative of at least three independent experiments.

Expression of *Roq1* Confers Resistance to *Xanthomonas perforans* in the Field

To determine if the resistance observed in growth chamber experiments would hold up under commercial tomato production conditions, we tested the ability of *Roq1* tomatoes to resist *X. perforans* infection in the field. *Roq1* tomatoes were grown along with the Fla. 8000 wild-type parent as well as a Fla. 8000 variety expressing the *Bs2* gene from pepper as a resistant control (Kunwar et al., 2018). For each of the three growing seasons, both *Roq1* and the resistant *Bs2* control tomatoes showed significantly lower disease severity than the parental Fla. 8000 variety (Table 1) ($p < 0.05$). The total marketable yield of the *Roq1* plants was not significantly different from that of the susceptible parent for any of the three seasons ($p > 0.05$). No obvious difference in growth morphology was observed between *Roq1* and wild-type tomato plants (Supplementary Figure S4).

The *Ralstonia* Effector RipB, a Homolog of *XopQ*/*HopQ1*, Is Recognized by *Roq1*


RipB, considered a “core” *Ralstonia* effector, is present in approximately 90% of sequenced strains (Sabbagh et al., 2019) making it an attractive target ligand for engineering crop plants to be resistant to this pathogen. *Roq1* perceives diverse alleles of *XopQ* and *HopQ1* and we hypothesized that it can also recognize different alleles of RipB. We constructed a phylogenetic tree using a subset of RipB alleles identified by BLAST search to approximately span the diversity of this effector in *Ralstonia* (Figure 3). The *Ralstonia* genus contains many diverse strains that have been divided into four phylotypes based on based on sequence analysis of the internal transcribed spacer region of the 16S–23S rRNA gene (Poussier et al., 2000; Prior and Fegan, 2004; Safni et al., 2014). We selected RipB alleles from *Ralstonia* strains GMI1000 and MolK2, from phylotypes I and II, respectively, for subsequent

analysis. These two RipB alleles share 71% amino acid identity with each other and approximately 52% identity with *XopQ* excluding the divergent N terminus containing the putative type III secretion signal. An alignment of these two RipB proteins with *XopQ* and *HopQ1* is shown in Supplementary Figure S5. To test for *Roq1*-dependent recognition of RipB, we used *Agrobacterium* to transiently express RipB from GMI1000 and MolK2 in leaf tissue of wild-type and *roq1* mutant *N. tabacum*. The *N. tabacum roq1-1* mutant was generated using a CRISPR/CAS9 construct targeting exon 1 of the *Roq1* gene (Supplementary Figure S6). Both RipB alleles

TABLE 1 | Field trial results.

Season/Genotype	Disease severity	Marketable yield (kg/ha)
Spring 2018		
Fla. 8000	86 ± 5	54, 655 ± 9, 450
Fla. 8000 <i>Roq1</i>	1 ± 1	52, 656 ± 3, 810
Fla. 8000 <i>Bs2</i>	1 ± 1	66, 270 ± 10, 309
Fall 2018		
Fla. 8000	25 ± 7	19, 576 ± 11, 038
Fla. 8000 <i>Roq1</i>	5 ± 1	18, 538 ± 5, 901
Fla. 8000 <i>Bs2</i>	0 ± 0	33, 770 ± 13, 176
Spring 2019		
Fla. 8000	84 ± 2	73, 009 ± 15, 243
Fla. 8000 <i>Roq1</i>	11 ± 7	92, 837 ± 11, 072
Fla. 8000 <i>Bs2</i>	5 ± 1	80, 516 ± 14, 531


A field trial was conducted in Florida with disease pressure from *Xanthomonas perforans*. Disease severity presented as percent infected tissue, converted from field ratings that were scored using the Horsfall–Barratt scale. Harvested tomatoes were graded and sized by USDA specifications to calculate the total marketable yield. The values shown are means ± standard deviation from at least four replicate plots of 10 plants each. Tomato plants expressing the *Bs2* immune receptor gene were included as a resistant control.



triggered a strong hypersensitive/cell death response in wild-type *N. tabacum*, indicating immune activation. This response was absent in the *roq1-1* mutant but could be restored by transiently expressing Roq1 along with XopQ, RipB^{GMI1000}, or RipB^{MolK2} (Figure 4).

Roq1 Tomatoes Are Resistant to *Ralstonia* Containing RipB


Our observation that Roq1 can recognize RipB in leaf transient expression assays suggested that Roq1 can mediate resistance to bacterial wilt caused by *Ralstonia* in tomato. We tested this hypothesis by challenging wild-type and *Roq1*-expressing tomato plants with *Ralstonia* strain GMI1000 using a soil soak inoculation disease assay. Wild-type plants developed severe wilting approximately 7 days after inoculation, whereas *Roq1* tomato plants remained mostly healthy over the 2-week time course (Figure 5A and Supplementary Figure S7). The *Roq1* tomato plants were susceptible to a deletion mutant lacking RipB (GMI1000 Δ ripB). We also challenged plants by introducing bacteria directly to the xylem by placing bacteria on the surface of



a cut petiole. Wild-type plants were wilted by eight days whereas *Roq1* plants remained healthy (Figure 5B). Tomatoes expressing *Roq1* were also resistant to *Ralstonia* strain UW551, which is a race 3 biovar 2 potato brown rot strain from phylogroup II that has a RipB allele (Figure 3 and Supplementary Figure S8).

Distribution of RipB Alleles in *Ralstonia*

To investigate the potential for using *Roq1* to protect plants from *Ralstonia*, we investigated the occurrence of RipB alleles in select strains. Table 2 lists some *Ralstonia* strains with their known hosts along with their respective phylogroup and RipB allele accession information. Table 2 illustrates that strains lacking putative functional RipB alleles correlate with strains that are virulent on tobacco, which contains a native *Roq1* gene. All strains in Table 2 except for tobacco pathogenic strains K60, Y45, BK1002, and OE1-1 contain putative full-length and functional RipB alleles. Relative to other RipB alleles, the K60 RipB allele is truncated after residue 437 and missing approximately 65 C-terminal residues and the OE1-1 allele is truncated after residue 417, missing approximately 77 residues based on a published genome sequence (Hayes et al., 2017) (Supplementary Figure S5). Y45 does not have a predicted RipB allele based on a draft genome sequence (Li et al., 2011). Published gene models for RipB disagree on which start codon is the correct one, leading some RipB alleles to look like they are missing part of the N terminus or have N-terminal extensions. Analysis of the DNA sequence of diverse RipB alleles showed that out of three possible in-frame start codons, only a single putative start codon is conserved among *Ralstonia* strains from all four



phylotypes (Supplementary Figure S9), suggesting that this is the true start codon and eliminating the N-terminal discrepancy between different RipB alleles.

DISCUSSION

Roq1 expression in tomato confers strong resistance to *X. perforans*, *X. euvesicatoria*, and *P. syringae* pv. *tomato*. Its effectiveness is dependent on the presence of the recognized effector protein XopQ/HopQ1 (Figures 1, 2). Field trials revealed that tomatoes expressing Roq1 were less susceptible to *X. perforans* than wild-type tomatoes in conditions approximating commercial production (Table 1). Roq1 conferred a similar level of resistance as the Bs2-containing resistant check

variety in one season and was slightly weaker in the other two. Bacterial spot caused by *X. perforans* can cause lesions on fruits, making them unsuitable for commercial sale, and also reduce plant productivity by damaging leaf tissue. The onset of fruit lesions requires high disease pressure during a particular phase of fruit development. Environmental conditions did not favor the development of fruit lesions and we did not observe significant fruit lesion formation on any of the genotypes in any of the three seasons. Despite showing a strong reduction in foliar disease symptoms, the Roq1 line did not have a significantly greater yield than the susceptible parental variety. A possible explanation for this finding is that bacterial spot did not appear to be a major constraint on yield in any of the three seasons. In spring 2018, weather conditions promoted the development of disease only late in the season after much of the yield was already set. Fall 2018 was unseasonably hot and dry for most of the season resulting in low disease pressure and very poor yield for all genotypes. Of the three seasons, spring 2019 had weather conditions expected to be most conducive for observing an impact of bacterial spot on marketable yield with mid-season rain promoting the development of disease symptoms. The average marketable yield for the Roq1 tomatoes was 27% higher than wild-type in this season, although a relatively small sample size (four replicate plots of 10 plants each) and a large variability of yield between plots resulted in a p -value of 0.08 by Student's t -test. Notably the yield of the resistant check variety expressing Bs2 was not significantly higher than the susceptible control in this season, though it was previously reported to give a yield increase of 1.5–11x relative to susceptible varieties under high disease pressure (Horvath et al., 2012). This suggests that bacterial spot was not severe enough to have a strong impact on yield in this season and that Roq1 may result in an increase in marketable yield under stronger disease pressure.

It was unclear if Roq1 could confer resistance to *Ralstonia* because it colonizes different tissues than *Xanthomonas* and *P. syringae*. While *Xanthomonas* and *P. syringae* colonize tomato leaf tissue, *Ralstonia* enters through the roots and colonizes xylem vessels. Although the type III secretion system is essential for virulence in *Ralstonia*, it is not clear when and where the pathogen delivers effectors into host cells. It was therefore not clear if Roq1 would be able to confer resistance to this pathogen in tomato. Here we demonstrated that tomato plants expressing Roq1 had strong resistance to *Ralstonia* expressing RipB as measured by both soil soak and cut-petiole inoculation assays (Figure 5). In addition to conferring resistance to the phylogroup I strain GMI1000, Roq1 also confers resistance to *Ralstonia* race 3 biovar 3 strain UW551, a phylogroup II strain that can overcome other known sources of bacterial wilt resistance in tomato (Milling et al., 2011). Some but not all of the Roq1 tomatoes inoculated with GMI1000 by soil soak were colonized by a moderate or low population of *Ralstonia* (Supplementary Figure S10). This observation suggests that Roq1-mediated immune responses may act to both restrict the establishment of vascular colonization and separately reduce bacterial titers if colonization does occur. Activation of immune receptors, including Roq1, is known to induce many defense-associated genes with different putative activities (Sohn et al., 2014; Qi

TABLE 2 | RipB occurrence and host range in *Ralstonia*.

<i>Ralstonia</i> strain	Host(s)	Origin	Phylotype	RipB allele	RipB accession
GMI1000	Tomato, Pepper, Arabidopsis	French Guyana	I	Present	CAD13773.2
RS1000	Tomato	Japan	I	Present	BAD42390.1
OE1-1	Tobacco	Japan	I	Truncated	RSOE_09080**
BK1002	Tobacco	Japan	not available	Truncated	BBI29704.1
Y45	Tobacco	China	I	Absent	None
K60	Tomato, Tobacco	United States	IIA	Truncated	B7R77_07450**
CFBP2957	Tomato	Martinique	IIA	Present	CBJ44424.1
Po82	Tomato, Banana, Potato	Mexico	IIB	Present	WP_014618158.1
IPO1609*	Potato	Netherlands	IIB	Present	WP_020956993.1
MolK2	Banana	Philippines	IIB	Present	WP_003278485.1
UW551*	Geranium	Kenya	IIB	Present	WP_020956993.1
CMR15	Tomato	Cameroon	III	Present	WP_020749919.1
PSI07	Tomato	Indonesia	IV	Present	WP_013213770.1
BDB R229	Banana	Indonesia	IV	Present	WP_078222314.1
R24	Clove	Indonesia	IV	Present	CCA88514.1

The published host range of select *Ralstonia* strains is listed along with the identified RipB allele. Truncated indicates that the identified allele is missing conserved residues at the C terminus and is putatively non-functional. * Indicates a race 3 biovar 2 strain. ** Indicates NCBI locus tag from genome accession CP009764.1 for strain OE1-1 and genome NCTK01000001.1 for strain K60.

et al., 2018), presumably acting to inhibit pathogen virulence by distinct mechanisms. The observation that Roq1 inhibits both colonization establishment and population growth suggests that at least two independent downstream defense responses mediate the observed resistance phenotype.

The *Roq1* tomatoes were susceptible to a *Ralstonia* mutant lacking *RipB*, indicating that the resistance depends on the interaction between RipB and Roq1. This is consistent with the observation that several naturally occurring *Ralstonia* strains that can infect tobacco have a truncated or are missing the RipB effector (Table 2) (Nakano and Mukaihara, 2019), suggesting that losing RipB can allow the pathogen to overcome the native *Roq1* gene present in *N. tabacum*. Tobacco-infecting strains K60 and OE1-1 contain independently truncated RipB alleles (Figure 3 and Supplementary Figure S5) and there have likely been multiple independent gene loss events which enable strains to evade Roq1-mediated resistance. Similarly, HopQ1 has been lost in strains of *P. syringae* that can infect tobacco (Denny, 2006; Ferrante and Scortichini, 2009; Li et al., 2011). This suggests that this effector is not essential for virulence in all circumstances and it would therefore be prudent to deploy *Roq1* in combination with other disease resistance traits to avoid resistance breakdown due to pathogens losing XopQ/HopQ1/RipB. Although minor foliar symptoms were observed on the *Roq1* tomatoes, particularly in spring 2019 (Table 1), we do not believe this was due to a naturally occurring XopQ mutant arising during the trial. Instead, we think that these low disease scores may have been the result of fungal diseases, which can cause foliar symptoms that look similar to bacterial spot, or by the *Roq1* tomatoes supporting a low level of bacterial growth.

No other known NLR immune receptor confers resistance against such a broad range of bacterial pathogenic genera as Roq1. Effectors that are recognized by NLR proteins act as avirulence factors and are under strong evolutionary pressure to diversify or be lost to evade immune activation. Therefore the effector repertoires of pathogens are often quite diverse,

with relatively few “core” effectors conserved within a species and even fewer shared between different genera (Grant et al., 2006). Effectors recognized by plant NLRs are typically narrowly conserved within a single bacterial genus (Kapos et al., 2019). One such effector is AvrBs2, recognized by the Bs2 receptor from pepper, which is present in many *Xanthomonas* strains but is absent from *P. syringae* and *Ralstonia*. In contrast, XopQ/HopQ1/RipB is highly conserved in most *Xanthomonas*, *P. syringae*, and *Ralstonia* strains that cause disease in crop plants including kiwi (*P. syringae* pv. *actinidae*), banana (*Ralstonia* and *X. campestris* pv. *musacearum*), stone fruit (*P. syringae*), pepper (*X. euvesicatoria*), citrus (*X. citri*), strawberry (*X. fragariae*), brassica (*X. campestris*), rice (*X. oryzae*), potato (*Ralstonia*), and others. *Ralstonia* race 3 biovar 2 strains are of particular concern because they are cold tolerant and potentially threaten potato cultivation in temperate climates. As a result, *Ralstonia* race 3 biovar 2 strains are strictly regulated quarantine pathogens in Europe and North America and is on the United States Select Agent list. The ability of *Roq1* to protect tomato from the race 3 biovar 2 strain UW551 (Supplementary Figure S8) suggests that *Roq1* can also protect potato from this high-concern pathogen. This work demonstrates the widespread potential of using naturally occurring plant immune receptors to safely, sustainably, and economically manage diverse and difficult to control pathogen species.

DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/Supplementary Material.

AUTHOR CONTRIBUTIONS

NT and AS wrote the manuscript and performed *Pseudomonas* and *Ralstonia* petiole infection assays. AS carried out

Xanthomonas infection and *Agrobacterium* transient expression experiments. UG and SH performed *Xanthomonas* field experiments. CH constructed the *Ralstonia* knockout and performed *Ralstonia* soil soak assays, supervised by CA. All authors analyzed the results and edited and approved the manuscript.

FUNDING

This work was supported in part by the National Institute of Food and Agriculture, US Department of Agriculture under award number 2016-67012-25106, and the UC Berkeley Innovative Genomics Institute. CH was supported by an NSF Predoctoral Fellowship.

REFERENCES

- Boller, T., and He, S. Y. (2009). Innate immunity in plants: an arms race between pattern recognition receptors in plants and effectors in microbial pathogens. *Science* 324, 742–744. doi: 10.1126/science.1171647
- Castañeda, A., Reddy, J. D., El-Yacoubi, B., and Gabriel, D. W. (2005). Mutagenesis of all eight avr genes in *Xanthomonas campestris* pv. *campestris* had no detected effect on pathogenicity, but one avr gene affected race specificity. *Mol. Plant. Microbe Interact.* 18, 1306–1317.
- Coupat, B., Chaumeille-Dole, F., Fall, S., Prior, P., Simonet, P., Nesme, X., et al. (2008). Natural transformation in the *Ralstonia solanacearum* species complex: number and size of DNA that can be transferred. *FEMS Microbiol. Ecol.* 66, 14–24. doi: 10.1111/j.1574-6941.2008.00552.x
- Coutu, C., Brandle, J., Brown, D., Brown, K., Miki, B., Simmonds, J., et al. (2007). pORE: a modular binary vector series suited for both monocot and dicot plant transformation. *Transgenic Res.* 16, 771–781. doi: 10.1007/s11248-007-9066-2
- Dangl, J. L., Horvath, D. M., and Staskawicz, B. J. (2013). Pivoting the plant immune system from dissection to deployment. *Science* 341, 746–751. doi: 10.1126/science.1236011
- Davis, R. M., Miyao, G., Subbarao, K. V., Stapleton, J. J., and Aegerter, A. B. J. (2013). *UC IPM: UC Management Guidelines - TOMATO:Diseases*. Available at: <http://ipm.ucanr.edu/PMG/r783101611.html> (Accessed June 17, 2019).
- de Pontes, N. C., Nascimento, A. D. R., Golynski, A., Maffia, L. A., and Rogério de Oliveira, J. (2016). Intervals and number of applications of acibenzolar-S-methyl for the control of bacterial spot on processing tomato. *Plant Dis.* 100, 2126–2133. doi: 10.1094/pdis-11-15-1286-re
- Denny, T. (2006). “Plant pathogenic *Ralstonia* species,” in *Plant-Associated Bacteria*, ed. S. S. Gnanamanickam (Dordrecht: Springer), 573–644. doi: 10.1007/1-4020-4538-7_16
- Deslandes, L., and Rivas, S. (2012). Catch me if you can: bacterial effectors and plant targets. *Trends Plant Sci.* 17, 644–655. doi: 10.1016/j.tplants.2012.06.011
- Dillon, M. M., Almeida, R. N. D., Laflamme, B., Martel, A., Weir, B. S., Desveaux, D., et al. (2019). Molecular evolution of *Pseudomonas syringae* Type III secreted effector proteins. *Front. Plant Sci.* 10:418.
- Ferrante, P., and Scortichini, M. (2009). Identification of *Pseudomonas syringae* pv. *actinidiae* as causal agent of bacterial canker of yellow kiwifruit (*Actinidia chinensis* Planchon) in Central Italy. *J. Phytopathol.* 157, 768–770. doi: 10.1111/j.1439-0434.2009.01550.x
- Freeman, J. H., McAvoy, E. J., Boyd, N. S., Kaniserry, R., Smith, H. A., Desaegeer, J., et al. (2018). “Tomato production,” in *Vegetable Production Handbook of Florida 2018–2019*, eds H. A. Smith, J. H. Freeman, P. J. Dittmar, M. L. Paret, and G. E. Vallad (Lincolnshire, IL: Vance Publishing Corporation), 349–393.
- Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A. III, and Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* 6, 343–345. doi: 10.1038/nmeth.1318
- Giska, F., Lichočka, M., Piechocki, M., Dadlez, M., Schmelzer, E., Hennig, J., et al. (2013). Phosphorylation of HopQ1, a type III effector from *Pseudomonas*

ACKNOWLEDGMENTS

This manuscript has been released as a pre-print at bioRxiv (Thomas et al., 2019). We thank Shirley Sato and Tom Clemente of the University of Nebraska Plant Transformation Core Research Facility for the transformation of tomato. We thank Myeong-Je Cho and Julie Pham of the UC Berkeley Innovative Genomics Institute for transformation of *Nicotiana tabacum*.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2020.00463/full#supplementary-material>

- syringae*, creates a binding site for host 14-3-3 proteins. *Plant Physiol.* 161, 2049–2061. doi: 10.1104/pp.112.209023
- Grant, S. R., Fisher, E. J., Chang, J. H., Mole, B. M., and Dangl, J. L. (2006). Subterfuge and manipulation: type III effector proteins of phytopathogenic bacteria. *Annu. Rev. Microbiol.* 60, 425–449. doi: 10.1146/annurev.micro.60.080805.142251
- Griffin, K., Gambley, C., Brown, P., and Li, Y. (2017). Copper-tolerance in *Pseudomonas syringae* pv. *tomato* and *Xanthomonas* spp. and the control of diseases associated with these pathogens in tomato and pepper. A systematic literature review. *Crop Protoc.* 96, 144–150. doi: 10.1016/j.cropro.2017.02.008
- Gudesblat, G. E., Torres, P. S., and Vojnov, A. A. (2009). *Xanthomonas campestris* overcomes *Arabidopsis* stomatal innate immunity through a DSF cell-to-cell signal-regulated virulence factor. *Plant Physiol.* 149, 1017–1027. doi: 10.1104/pp.108.126870
- Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321. doi: 10.1093/sysbio/syq010
- Hann, D. R., Dominguez-Ferreras, A., Motyka, V., Dobrev, P. I., Schornack, S., Jehle, A., et al. (2014). The *Pseudomonas* type III effector HopQ1 activates cytokinin signaling and interferes with plant innate immunity. *New Phytol.* 201, 585–598. doi: 10.1111/nph.12544
- Hayes, M. M., MacIntyre, A. M., and Allen, C. (2017). Complete genome sequences of the plant pathogens *Ralstonia solanacearum* Type Strain K60 and *R. solanacearum* race 3 biovar 2 strain UW551. *Genome Announc.* 5:e01088-17. doi: 10.1128/genomeA.01088-17
- Horsfall, J. G., and Barrat, R. W. (1945). An improved grading system for measuring plant diseases. *Phytopathology* 35:655.
- Horvath, D. M., Stall, R. E., Jones, J. B., Pauly, M. H., Vallad, G. E., Dahlbeck, D., et al. (2012). Transgenic resistance confers effective field level control of bacterial spot disease in tomato. *PLoS One* 7:e42036. doi: 10.1371/journal.pone.0042036
- Jones, J. D. G., and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329.
- Jones, J. D. G., Vance, R. E., and Dangl, J. L. (2016). Intracellular innate immune surveillance devices in plants and animals. *Science* 354, doi: 10.1126/science.aaf6395
- Jones, J. D. G., Witek, K., Verweij, W., Jupe, F., Cooke, D., Dorling, S., et al. (2014). Elevating crop disease resistance with cloned genes. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 369:20130087. doi: 10.1098/rstb.2013.0087
- Kapos, P., Devendrakumar, K. T., and Li, X. (2019). Plant NLRs: from discovery to application. *Plant Sci.* 279, 3–18. doi: 10.1016/j.plantsci.2018.03.010
- Kay, S., and Bonas, U. (2009). How *Xanthomonas* type III effectors manipulate the host plant. *Curr. Opin. Microbiol.* 12, 37–43. doi: 10.1016/j.mib.2008.12.006
- Kennelly, M. M., Cazorla, F. M., de Vicente, A., Ramos, C., and Sundin, G. W. (2007). *Pseudomonas syringae* diseases of fruit trees: progress toward understanding and control. *Plant Dis.* 91, 4–17. doi: 10.1094/pd-91-0004

- Khokhani, D., Tran, T. M., Lowe-Power, T. M., and Allen, C. (2018). Plant assays for quantifying *Ralstonia solanacearum* virulence. *Bio Protocol* 8:3028.
- Kunwar, S., Iriarte, F., Fan, Q., Evaristo da Silva, E., Ritchie, L., Nguyen, N. S., et al. (2018). Transgenic expression of *EFR* and *Bs2* genes for field management of bacterial wilt and bacterial spot of tomato. *Phytopathology* 108, 1402–1411. doi: 10.1094/phyto-12-17-0424-r
- Li, W., Yadeta, K. A., Elmore, J. M., and Coaker, G. (2013). The *Pseudomonas syringae* effector HopQ1 promotes bacterial virulence and interacts with tomato 14-3-3 proteins in a phosphorylation-dependent manner. *Plant Physiol.* 161, 2062–2074. doi: 10.1104/pp.112.211748
- Li, X., Kapos, P., and Zhang, Y. (2015). NLRs in plants. *Curr. Opin. Immunol.* 32, 114–121. doi: 10.1016/j.coi.2015.01.014
- Li, Z., Wu, S., Bai, X., Liu, Y., Lu, J., Liu, Y., et al. (2011). Genome sequence of the tobacco bacterial wilt pathogen *Ralstonia solanacearum*. *J. Bacteriol.* 193, 6088–6089. doi: 10.1128/jb.06009-11
- Liu, Y., Kanda, A., Kiba, A., Hikichi, Y., and Ohnishi, K. (2009). Distribution of avirulence genes *avrA* and *popP1* in 22 Japanese phylotype I strains of *Ralstonia solanacearum*. *J. Gen. Plant Pathol.* 75, 362–368. doi: 10.1007/s10327-009-0189-6
- Malik, K., Wu, K., Li, X. Q., Martin-Heller, T., Hu, M., Foster, E., et al. (2002). A constitutive gene expression system derived from the tCUP cryptic promoter elements. *Theor. Appl. Genet.* 105, 505–514. doi: 10.1007/s00122-002-0926-0
- Milling, A., Babujee, L., and Allen, C. (2011). *Ralstonia solanacearum* extracellular polysaccharide is a specific elicitor of defense responses in wilt-resistant tomato plants. *PLoS One* 6:e15853. doi: 10.1371/journal.pone.0015853
- Mukaihara, T., and Tamura, N. (2009). Identification of novel *Ralstonia solanacearum* type III effector proteins through translocation analysis of *hrpB*-regulated gene products. *Microbiology* 155, 2235–2244. doi: 10.1099/mic.0.027763-0
- Nakano, M., and Mukaihara, T. (2019). The type III effector RipB from *Ralstonia solanacearum* RS1000 acts as a major avirulence factor in *Nicotiana benthamiana* and other *Nicotiana* species. *Mol. Plant Pathol.* 20, 1237–1251. doi: 10.1111/mpp.12824
- Peeters, N., Carrère, S., Anisimova, M., Plener, L., Cazalé, A.-C., and Genin, S. (2013). Repertoire, unified nomenclature and evolution of the Type III effector gene set in the *Ralstonia solanacearum* species complex. *BMC Genomics* 14:859. doi: 10.1186/1471-2164-14-859
- Potnis, N., Timilsina, S., Strayer, A., Shantharaj, D., Barak, J. D., Paret, M. L., et al. (2015). Bacterial spot of tomato and pepper: diverse *Xanthomonas* species with a wide variety of virulence factors posing a worldwide challenge. *Mol. Plant Pathol.* 16, 907–920. doi: 10.1111/mpp.12244
- Poussier, S., Prior, P., Luisetti, J., Hayward, C., and Fegan, M. (2000). Partial sequencing of the *hrpB* and *endoglucanase* genes confirms and expands the known diversity within the *Ralstonia solanacearum* species complex. *Syst. Appl. Microbiol.* 23, 479–486. doi: 10.1016/s0723-2020(00)80021-1
- Prior, P., and Fegan, M. (2004). “Recent developments in the phylogeny and classification of *Ralstonia solanacearum*,” in *Proceedings of the 1 International Symposium on Tomato Diseases 695* (Leuven: International Society for Horticultural Science), 127–136. doi: 10.17660/actahortic.2005.695.14
- Qi, T., Seong, K., Thomazella, D. P. T., Kim, J. R., Pham, J., Seo, E., et al. (2018). NRG1 functions downstream of EDS1 to regulate TIR-NLR-mediated plant immunity in *Nicotiana benthamiana*. *Proc. Natl. Acad. Sci. U.S.A.* 115, E10979–E10987.
- Rivard, C. L., O’Connell, S., Peet, M. M., Welker, R. M., and Louws, F. J. (2012). Grafting tomato to manage bacterial wilt caused by *Ralstonia solanacearum* in the Southeastern United States. *Plant Dis.* 96, 973–978. doi: 10.1094/pdis-12-10-0877
- Ryan, R. P., Vorhölter, F.-J., Potnis, N., Jones, J. B., Van Sluys, M.-A., Bogdanove, A. J., et al. (2011). Pathogenomics of *Xanthomonas*: understanding bacterium-plant interactions. *Nat. Rev. Microbiol.* 9, 344–355. doi: 10.1038/nrmicro2558
- Sabbagh, C. R. R., Carrère, S., Lonjon, F., Vaillieu, F., Macho, A. P., Genin, S., et al. (2019). Pangenomic type III effector database of the plant pathogenic *Ralstonia* spp. *PeerJ* 7:e7346. doi: 10.7287/peerj.preprints.27726v1
- Safni, I., Cleenwerck, I., De Vos, P., Fegan, M., Sly, L., and Kappler, U. (2014). Polyphasic taxonomic revision of the *Ralstonia solanacearum* species complex: proposal to emend the descriptions of *Ralstonia solanacearum* and *Ralstonia syzygii* and reclassify current *R. syzygii* strains as *Ralstonia syzygii* subsp. *syzygii* subsp. nov., *R. solanacearum* phylotype IV strains as *Ralstonia syzygii* subsp. *indonesiensis* subsp. nov., banana blood disease bacterium strains as *Ralstonia syzygii* subsp. *celesbensis* subsp. nov. and *R. solanacearum* phylotype I and III strains as *Ralstonia pseudosolanacearum* sp. nov. *Int. J. Syst. Evol. Microbiol.* 64, 3087–3103. doi: 10.1099/ijs.0.066712-0
- Schultink, A., Qi, T., Lee, A., Steinbrenner, A. D., and Staskawicz, B. (2017). Roq1 mediates recognition of the *Xanthomonas* and *Pseudomonas* effector proteins XopQ and HopQ1. *Plant J.* 92, 787–795. doi: 10.1111/tj.13715
- Schwartz, A. R., Potnis, N., Timilsina, S., Wilson, M., Patané, J., Martins, J., et al. (2015). Phylogenomics of *Xanthomonas* field strains infecting pepper and tomato reveals diversity in effector repertoires and identifies determinants of host specificity. *Front. Microbiol.* 6:535.
- Sharma, S., and Bhattarai, K. (2019). Progress in developing bacterial spot resistance in tomato. *Agronomy* 9:26. doi: 10.3390/agronomy9010026
- Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7:539. doi: 10.1038/msb.2011.75
- Sohn, K. H., Segonzac, C., Rallapalli, G., Sarris, P. F., Woo, J. Y., Williams, S. J., et al. (2014). The nuclear immune receptor RPS4 is required for RRS1SLH1-dependent constitutive defense activation in *Arabidopsis thaliana*. *PLoS Genet.* 10:e1004655. doi: 10.1371/journal.pgen.1004655
- Staskawicz, B. J., and Schultink, A. C. (2019). *Roq1 Provides Resistance to Both Xanthomonas and Pseudomonas in Plants*. World Patent WO2019040483. Geneva: WIPO.
- Tai, T. H., Dahlbeck, D., Clark, E. T., Gajiwala, P., Pasion, R., Whalen, M. C., et al. (1999). Expression of the *Bs2* pepper gene confers resistance to bacterial spot disease in tomato. *Proc. Natl. Acad. Sci. U.S.A.* 96, 14153–14158. doi: 10.1073/pnas.96.24.14153
- Teper, D., Salomon, D., Sunitha, S., Kim, J.-G., Mudgett, M. B., and Sessa, G. (2014). *Xanthomonas euvesicatoria* type III effector XopQ interacts with tomato and pepper 14-3-3 isoforms to suppress effector-triggered immunity. *Plant J.* 77, 297–309. doi: 10.1111/tj.12391
- Thomas, N. C., Hendrich, C. G., Gill, U. S., Allen, C., Hutton, S. F., and Schultink, A. (2019). Roq1 confers resistance to *Xanthomonas*, *Pseudomonas syringae* and *Ralstonia solanacearum* in tomato. *bioRxiv* [Preprint]. doi: 10.1101/813758
- Vasse, J., Frey, P., and Trigalet, A. (1995). *Microscopic Studies of Intercellular Infection and Protoxylem Invasion of Tomato Roots by Pseudomonas Solanacearum*. Available at: <https://pubag.nal.usda.gov/catalog/1443867> (Accessed July 1, 2019).
- Vincelli, P. (2016). Genetic engineering and sustainable crop disease management: opportunities for case-by-case decision-making. *Sustain. Sci. Pract. Policy* 8:495. doi: 10.3390/su8050495
- Wei, C. F., Kvitko, B. H., Shimizu, R., Crabill, E., Alfano, J. R., Lin, N. C., et al. (2007). A *Pseudomonas syringae* pv. *tomato* DC3000 mutant lacking the type III effector HopQ1-1 is able to cause disease in the model plant *Nicotiana benthamiana*. *Plant J.* 51, 32–46. doi: 10.1111/j.1365-313x.2007.03126.x
- Xin, X.-F., Kvitko, B., and He, S. Y. (2018). *Pseudomonas syringae*: what it takes to be a pathogen. *Nat. Rev. Microbiol.* 16, 316–328. doi: 10.1038/nrmicro.2018.17
- Yuliar, Nion, Y. A., and Toyota, K. (2015). Recent trends in control methods for bacterial wilt diseases caused by *Ralstonia solanacearum*. *Microbes Environ.* 30, 1–11. doi: 10.1264/jsme2.me14144
- Zipfel, C. (2014). Plant pattern-recognition receptors. *Trends Immunol.* 35, 345–351.

Conflict of Interest: AS and NT are employees of and have a financial stake in Fortiphyte Inc., which has intellectual property rights related to the Roq1 resistance gene.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Thomas, Hendrich, Gill, Allen, Hutton and Schultink. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.