



Virtual Reality as a Potential Tool to Face Frailty Challenges

Silvia Serino^{1,2}, Serena Barello¹, Francesca Miraglia^{3,4}, Stefano Triberti¹ and Claudia Repetto^{1*}

¹ Department of Psychology, Catholic University of the Sacred Heart, Milan, Italy, ² Applied Technology for Neuropsychology Lab, Istituto Auxologico Italiano, Milan, Italy, ³ Department of Geriatrics, Neuroscience and Orthopedics, Institute of Neurology, Catholic University of the Sacred Heart, Rome, Italy, ⁴ Brain Connectivity Laboratory, IRCCS San Raffaele Pisana, Rome, Italy

Keywords: virtual reality, frailty, rehabilitation, executive functions, gait, aging in place, patient engagement

OPEN ACCESS

Edited by:

Albert Rizzo,
USC Institute for Creative
Technologies, United States

Reviewed by:

Pascual Gonzalez,
Universidad de Castilla-La Mancha,
Spain

*Correspondence:

Claudia Repetto
claudia.repetto@unicatt.it

Specialty section:

This article was submitted to
Human-Media Interaction,
a section of the journal
Frontiers in Psychology

Received: 14 June 2017

Accepted: 24 August 2017

Published: 05 September 2017

Citation:

Serino S, Barello S, Miraglia F,
Triberti S and Repetto C (2017) Virtual
Reality as a Potential Tool to Face
Frailty Challenges.
Front. Psychol. 8:1541.
doi: 10.3389/fpsyg.2017.01541

The aging population and the corresponding increase in age-related diseases present scientific community and public health authorities with imminent challenges. One of these challenges deals with a deeper understanding of functional status of elderly in order to prevent and/or delay the onset of late-life disability (Rodríguez-Artalejo and Rodríguez-Mañas, 2014). The syndrome of “frailty” has been recently introduced in literature to specifically characterize the health of older individuals who deserve special attention because of their increased vulnerability to adverse health outcomes (Afilalo et al., 2010). Although there is not a unique definition of frailty (Morley et al., 2013), the majority of studies refers to the five operational criteria (Fried et al., 2001): decreased gait speed, reduced grip strength, prolonged and unmotivated exhaustion, low physical activity, unintended weight loss. The problem of different definitions leads also to a large variation in reported prevalence rates, which range approximately from 5 to 60% (Collard et al., 2012). However, this multifaceted decline in different physiological systems make frail older individuals progressively more exposed to stressors (Clegg et al., 2013), making urgent the need for better care interventions.

In parallel, some authors suggested to introduce also the phenotype of “cognitive frailty” to refer to older individuals who manifest a concurrent weakness in both physical and cognitive domains (Kelaiditi et al., 2013). Other than the presence of physical frailty, Kelaiditi and colleagues (Kelaiditi et al., 2013) proposed that the key criteria of cognitive frailty is the presence of mild cognitive impairment, in the absence of dementia. A very recent large study (Delrieu et al., 2016), involving 1.617 participants, clarified that cognitive frailty individuals showed a specific weakness in executive domain (i.e., a wide range of high-level cognitive abilities including problem-solving, planning, monitoring). These findings were in line with studies suggesting a crucial link between an early cognitive decline in frontal areas and gait deficits (Montero-Odasso et al., 2011), since they share common brain networks.

Different treatment approaches have been investigated in clinical trials for reducing the functional decline of frail individuals: exercise interventions (Forster et al., 2009), nutritional programs (Fiatarone et al., 1994), and integrated approaches (Looman et al., 2016). Three main issues emerged so far: the need for a more precise identification of markers of frailty; a call for innovative therapeutic strategies; the importance to develop personalized and integrated care models and intervention approaches aimed at improving independence, preferably delivered in their home setting (de Vos et al., 2012). In this context, we suggest that Virtual Reality can be an innovative tool potentially able to address the aforementioned issues.

VIRTUAL REALITY FOR FRAIL PATIENTS: HOW AND WHY

Virtual Reality (VR) is a combination of technological devices that allows users to navigate into and interact with tridimensional computer-created environments, having the subjective sensation to be there (Sanchez-Vives and Slater, 2005; Triberti and Riva, 2015). In the last decades, VR has been extensively used in neuroscience, and recent studies investigated the role of VR on brain modulation via neuroimaging methods. These studies aimed to characterize brain activity in virtual environments in order to understand the neurophysiologic correlates of the virtual navigation (Pacheco et al., 2017). Neuroimaging evidence suggested that medial temporal lobe structures, the hippocampus in particular, as well as parietal and frontal regions have been implicated in spatial navigation in humans (Iaria et al., 2008). Furthermore, it was observed an elevated theta and gamma power associated with VR navigation and increased theta coherence between right parietal and temporal regions (Cornwell et al., 2008). Theta and gamma oscillatory activity and cortico-hippocampal communication are part of a brain mechanism involved in the information transfer of different spatial representations required for successful navigation (White et al., 2012); parietal lobe is involved in numerous aspects of visuo-spatial cognition; theta and gamma activity in this region likely reflects the mechanism by which medial temporal and parietal brain regions communicate during navigation (Moser et al., 2008).

Therefore, experimental evidence underlined that during navigation in a virtual environment the brain activity is modulated (Weidemann et al., 2009), and paved the way to consider VR a potential tool for diagnosis and rehabilitation of motor and cognitive deficits.

Along this line of reasoning, in the following paragraphs we want to clarify how VR can provide interesting opportunities in order to deal with the challenges prompted by frailty: early symptoms identification, motor and cognitive rehabilitation, home-setting interventions.

As far as motor symptoms are concerned, an increasing number of studies have developed novel paradigms to deeply model the gait characteristics (e.g., Martens et al., 2017), especially for those ones difficult to analyze in clinical setting (i.e., intermittent gait freezing in Parkinson's Disease—PD, or—more related to frailty—a decrease in gait speed). For instance, Shine and co-workers developed a VR-based environment for investigating gait features (Shine et al., 2013). In this paradigm, the participants, seated in front of a computer screen, were asked to navigate in a realistic virtual environment (i.e., a corridor) via footpedals. In addition to the “simple command” (such as “WALK” or “STOP” that appeared on the screen), they were trained to more complex stimuli, entailing executive functions: the congruent color-word (“BLUE” written in blue that means “WALK”), or the incongruent color-word (“RED” written in green that means “STOP”). Shine and co-workers (Shine et al., 2013) found that PD patients with freezing of gait had a large frequency of motor arrests in comparison with “non-freezers” PD patients on this task, making it suitable for modeling the gait behavior. However, this dual-task paradigm implemented

in VR appears also particularly useful for the evaluation of motor aspects of gait speed and related cognitive deficits. As far as cognitive symptoms are concerned, traditional paper-and-pencil tests are not reliable to capture the “complexity” of executive functioning emerging in real-life situations (Shallice and Burgess, 1991; Goldstein, 1996). One attempt to overcome this issue is the development of tests evaluating the executive functioning in real-life scenarios, such as the Multiple Errands Test (a shopping task in a supermarket, Shallice and Burgess, 1991) or the Executive Function Performance Test (simple cooking, telephone use, and medication management, Baum et al., 2008). Given the difficulties in reproducing these tests in real life situations (i.e., time consuming, high economic costs, safety of the patients, poor controllability of experimental conditions), VR technology has been increasingly used for the assessment of executive functions. Indeed, VR permits to develop scenarios reproducing daily-life situations, allowing a secure and ecologically valid assessment of executive functions (Parsons, 2011). For example, Nir-Hadad and co-workers (Nir-Hadad et al., 2017) recently developed and tested in a sample of 19 post-stroke patients a virtual version of the original Four Item Shopping Task, which requires budget management as a functional test of executive functioning. Also the virtual version of the Multiple Errands Test has been developed and tested in different clinical populations (Raspelli et al., 2012; Cipresso et al., 2014).

Second, VR could be a promising tool to enhance neuroplasticity in neurorehabilitation (Ng et al., 2013). The concept underlying VR-based therapy as a treatment for motor and cognitive dysfunction is to improve neuroplasticity of the brain by engaging users in multisensory training. VR-based intervention effectiveness was demonstrated in several chronic stroke patients (Lloréns et al., 2015), in vestibular (Alahmari et al., 2014), in sensori-motor (Fluet and Deutsch, 2013) and cognitive rehabilitation of neurological patients (Slobounov et al., 2015).

A recent systematic review found that VR-based trainings were more effective than conventional therapies in enhancing balance and gait ability in post-stroke patients (de Rooij et al., 2016). The advantages offered by VR over conventional approaches were multiple: within virtual environments, it is possible to develop repetitive and personalized motor training that are enriched by different feedbacks (proprioceptive, visual, auditory) able to maximize motor learning. In particular, the use of VR in combination with haptic devices (i.e., robotic systems able to give users tactile and force feedbacks when interacting with virtual objects) can enhance the environment realism (Hoffman et al., 1998), thus improving the efficacy of a rehabilitation program (Teruel et al., 2015). Moreover, VR-based stimulation can provide frail individuals with engaging and enriching environments, helping them to repeat the exercises harder and longer, thus exploiting the principle of motor learning (Kitago and Krakauer, 2013). Furthermore, although VR is currently in the developmental phase in terms of treatment of frailty (Mugueta-Aguinaga and Garcia-Zapirain, 2017), VR-based rehabilitation protocols have been already tested for training executive functions in other clinical populations (Faria et al., 2016). It is worthy to underline that cognitive

training might be particularly demanding for elderly, especially in case of cognitive impairment. As previously explained, VR offers the chance to set-up cognitive exercises within meaningful environments (Riva et al., 2006). Moreover, in virtual environments it is possible to reproduce real-life situation in a safer and more controlled setting: the ecological validity is an important feature in neuropsychological assessment and remediation, even more so for the executive functions trainings.

Finally, as the old and frail population continues to grow, a great deal of attention has been dedicated to find organizational solutions aimed at promoting *aging-in-place* policies, in order to facilitate individuals in living independently in one's own home as long as possible (Stones and Gullifer, 2016). Indeed, *aging-in-place* is recognized as a crucial strategy to improve the quality of life of elderly citizens as well as the sustainability of social and welfare systems. Early evidence is supportive of the advantages of structured program to enable *aging-in-place* as it enhances patient engagement in their own medical and rehabilitation processes (Kim et al., 2017), which is a crucial predictor of patients' quality of life and medication adherence (Barello and Graffigna, 2015). However, *aging-in-place* requires a reframing of the care models in terms of seamless transitions between hospitals, the welfare system and territory care, along with all other physical and social contexts in an elderly citizens life. However, accomplishing this will require substantial innovation in the incorporation of advanced technologies in the process of care and cure. In this context, VR-based technologies might be a powerful tool to make the *aging-in-place* imperative a concrete reality (Lange et al., 2010). Indeed, this tool might guarantee elderly people to follow the rehabilitation process directly at home. Moreover, VR has the potential to sustain elderly people active engagement in the medical course due to their high level of customization according the patient's unique expectations and care needs (Graffigna et al., 2014). According to these

reflections, *aging-in-place* using VR-based technologies may be a promising solution for the upcoming aging society. However, the implementation of these solutions at home should consider also the introduction of some specific systems that allow patients' monitoring (e.g., intelligent systems for teletherapy, Rodríguez et al., 2016, or wearable devices for unobtrusive monitoring, Patel et al., 2012) to make VR-based training as controlled as in clinical settings.

Beside the numerous advantages VR offers for facing frailty challenges, potential limitations should also be taken into account. At a basic research level, it should be acknowledged that there are still few evidences about of VR brain modulation effects in several domains: long-term outcomes, direct comparisons between commercial and customized modules, immersive vs. non-immersive VR, and augmented vs. fully virtual systems. At the applicative level, frail older adults could show low degrees of technology acceptance, due to a general diffidence toward the technological devices or to the discomfort elicited by the specific VR set up proposed. These limitations, thought, should not prevent researchers and clinicians from carrying on projects that test the use of VR with elderly and frail patients; on the contrary, these issues should stimulate to move forward in basic and applicative research in order to better exploit in the future the VR capabilities with frail individuals.

AUTHOR CONTRIBUTIONS

SS, SB, FM, and CR conceived the work. SS, SB, ST, and FM drafted the paper. CR and ST revised critically the work.

ACKNOWLEDGMENTS

This work was partially supported by the Fondazione Cariplo within the project "Active Aging and Healthy Living."

REFERENCES

- Afilalo, J., Eisenberg, M. J., Morin, J. F., Bergman, H., Monette, J., Noisieux, N., et al. (2010). Gait speed as an incremental predictor of mortality and major morbidity in elderly patients undergoing cardiac surgery. *J. Am. Coll. Cardiol.* 56, 1668–1676. doi: 10.1016/j.jacc.2010.06.039
- Alahmari, K. A., Sparto, P. J., Marchetti, G. F., Redfern, M. S., Furman, J. M., and Whitney, S. L. (2014). Comparison of virtual reality based therapy with customized vestibular physical therapy for the treatment of vestibular disorders. *IEEE Trans. Neural Syst. Rehabil. Eng.* 22, 389–399. doi: 10.1109/TNSRE.2013.2294904
- Barello, S., and Graffigna, G. (2015). Engaging patients to recover life projectuality: an Italian cross-disease framework. *Quality Life Res.* 24, 1087–1096. doi: 10.1007/s11136-014-0846-x
- Baum, C. M., Connor, L. T., Morrison, T., Hahn, M., Dromerick, A. W., and Edwards, D. F. (2008). Reliability, validity, and clinical utility of the executive function performance test: a measure of executive function in a sample of people with stroke. *Am. J. Occup. Ther.* 62, 446–455. doi: 10.5014/ajot.62.4.446
- Cipresso, P., Albani, G., Serino, S., Pedrolì, E., Pallavicini, F., Mauro, A., et al. (2014). Virtual multiple errands test (VMET): a virtual reality-based tool to detect early executive functions deficit in Parkinson's disease. *Front. Behav. Neurosci.* 8:405. doi: 10.3389/fnbeh.2014.00405
- Clegg, A., Young, J., Iliffe, S., Rikkert, M. O., and Rockwood, K. (2013). Frailty in elderly people. *Lancet* 381, 752–762. doi: 10.1016/S0140-6736(12)62167-9
- Collard, R. M., Boter, H., Schoevers, R. A., and Oude Voshaar, R. C. (2012). Prevalence of frailty in community-dwelling older persons: a systematic review. *J. Am. Geriatr. Soc.* 60, 1487–1492. doi: 10.1111/j.1532-5415.2012.04054.x
- Cornwell, B. R., Johnson, L. L., Holroyd, T., Carver, F. W., and Grillon, C. (2008). Human hippocampal and parahippocampal theta during goal-directed spatial navigation predicts performance on a virtual Morris water maze. *J. Neurosci.* 28, 5983–5990. doi: 10.1523/JNEUROSCI.5001-07.2008
- de Rooij, I. J. M., van de Port, I. G. L., and Meijer, J. W. G. (2016). Effect of virtual reality training on balance and gait ability in patients with stroke: systematic review and meta-analysis. *Phys. Ther.* 96, 1905–1918. doi: 10.2522/ptj.20160054
- de Vos, A. J. B. M., Asmus-Szepesi, K. J. E., Bakker, T. J. E. M., de Vreede, P. L., van Wijngaarden, J. D. H., Steyerberg, E. W., et al. (2012). Integrated approach to prevent functional decline in hospitalized elderly: the prevention and reactivation care program (PReCaP). *BMC Geriatr.* 12:7. doi: 10.1186/1471-2318-12-7
- Delrieu, J., Andrieu, S., Pahor, M., Cantet, C., Cesari, M., Ousset, P. J., et al. (2016). Neuropsychological profile of "cognitive frailty" subjects in MAPT study. *J. Prev. Alzheimers Dis.* 3, 151–159. doi: 10.14283/jpad.2016.94
- Faria, A. L., Andrade, A., Soares, L., and Badia, S. B. I. (2016). Benefits of virtual reality based cognitive rehabilitation through simulated activities of daily living: a randomized controlled trial with stroke patients. *J. Neuroeng. Rehabil.* 13:96. doi: 10.1186/s12984-016-0204-z
- Fiatrone, M. A., O'Neill, E. F., Ryan, N. D., Clements, K. M., Solares, G. R., Nelson, M. E., et al. (1994). Exercise training and nutritional supplementation

- for physical frailty in very elderly people. *N. Engl. J. Med.* 330, 1769–1775. doi: 10.1056/NEJM199406233302501
- Fluet, G. G., and Deutsch, J. E. (2013). Virtual reality for sensorimotor rehabilitation post-stroke: the promise and current state of the field. *Curr. Phys. Med. Rehab. Rep.* 1, 9–20. doi: 10.1007/s40141-013-0005-2
- Forster, A., Lambley, R., Hardy, J., Young, J., Smith, J., Green, J., et al. (2009). Rehabilitation for older people in long-term care. *Cochrane Database Syst. Rev.* doi: 10.1002/14651858.CD004294.pub2
- Fried, L. P., Tangen, C. M., Walston, J., Newman, A. B., Hirsch, C., Gottdiener, J., et al. (2001). Frailty in older adults evidence for a phenotype. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 56, M146–M157. doi: 10.1093/gerona/56.3.M146
- Goldstein, G. (1996). “Functional considerations in neuropsychology,” in *Ecological Validity of Neuropsychological Testing*, eds R. J. Sbordone and C. J. Long (Delray Beach: FL:GsRPress/St.LuciePress), 75–89.
- Graffigna, G., Barello, S., Triberti, S., Wiederhold, B. K., Bosio, A. C., and Riva, G. (2014). Enabling eHealth as a pathway for patient engagement: a toolkit for medical practice. *Stud. Health Technol. Inform.* 199, 13–21. doi: 10.3233/978-1-61499-401-5-13
- Hoffman, H. G., Hollander, A., Schroder, K., Rousseau, S., and Furness, T. (1998). Physically touching and tasting virtual objects enhances the realism of virtual experiences. *Virtual Real.* 3, 226–234. doi: 10.1007/BF01408703
- Iaria, G., Fox, C. J., Chen, J. K., Petrides, M., and Barton, J. J. S. (2008). Detection of unexpected events during spatial navigation in humans: bottom-up attentional system and neural mechanisms. *Eur. J. Neurosci.* 27, 1017–1025. doi: 10.1111/j.1460-9568.2008.06060.x
- Kelaiditi, E., Cesari, M., Canevelli, M., Van Kan, G. A., Ousset, P. J., Gillette-Guyonnet, S., et al. (2013). Cognitive frailty: rational and definition from an (IANA/IAGG) international consensus group. *J. Nutr. Health Aging* 17, 726–734. doi: 10.1007/s12603-013-0367-2
- Kim, K., Gollamudi, S. S., and Steinhilbl, S. (2017). Digital technology to enable aging in place. *Exp. Gerontol.* 88, 25–31. doi: 10.1016/j.exger.2016.11.013
- Kitago, T., and Krakauer, J. W. (2013). Motor learning principles for neurorehabilitation. *Handb. Clin. Neurol.* 110, 93–103. doi: 10.1016/B978-0-444-52901-5.00008-3
- Lange, B. S., Requejo, P., Flynn, S. M., Rizzo, A. A., Valero-Cuevas, F. J., Baker, L., et al. (2010). The potential of virtual reality and gaming to assist successful aging with disability. *Phys. Med. Rehabil. Clin. N. Am.* 21, 339–356. doi: 10.1016/j.pmr.2009.12.007
- Lloréns, R., Gil-Gómez, J. A., Alcañiz, M., Colomer, C., and Noé, E. (2015). Improvement in balance using a virtual reality-based stepping exercise: a randomized controlled trial involving individuals with chronic stroke. *Clin. Rehabil.* 29, 261–268. doi: 10.1177/0269215514543333
- Looman, W. M., Fabbriotti, I. N., de Kuyper, R., and Huijsman, R. (2016). The effects of a pro-active integrated care intervention for frail community-dwelling older people: a quasi-experimental study with the GP-practice as single entry point. *BMC Geriatr.* 16:43. doi: 10.1186/s12877-016-0214-5
- Martens, K. A. E., Shine, J. M., and Lewis, S. J. (2017). “Using virtual reality to advance the understanding and rehabilitation of gait impairments in Parkinson’s Disease,” in *Locomotion and Posture in Older Adults*, eds F. A. Barbieri and R. Vitorio (Cham: Springer International Publishing), 397–416.
- Montero-Odasso, M., Muir, S. W., Hall, M., Doherty, T. J., Kloseck, M., Beauchet, O., et al. (2011). Gait variability is associated with frailty in community-dwelling older adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 66, 568–576. doi: 10.1093/gerona/66/5/568
- Morley, J. E., Vellas, B., van Kan, G. A., Anker, S. D., Bauer, J. M., Bernabei, R., et al. (2013). Frailty consensus: a call to action. *J. Am. Med. Dir. Assoc.* 14, 392–397. doi: 10.1016/j.jamda.2013.03.022
- Moser, E. I., Kropff, E., and Moser, M. B. (2008). Place cells, grid cells, and the brain’s spatial representation system. *Annu. Rev. Neurosci.* 31, 69–89. doi: 10.1146/annurev.neuro.31.061307.090723
- Mugueta-Aguinaga, I., and Garcia-Zapirain, B. (2017). Is technology present in frailty? Technology a back-up tool for dealing with frailty in the elderly: a systematic review. *Aging Dis.* 8, 176–195. doi: 10.14336/AD.2016.0901
- Ng, Y. S., Chew, E., Samuel, G. S., Tan, Y. L., and Kong, K. H. (2013). Advances in rehabilitation medicine. *Singapore Med. J.* 54, 538–551. doi: 10.11622/smedj.2013197
- Nir-Hadad, S. Y., Weiss, P. L., Waizman, A., Schwartz, N., and Kizony, R. (2017). A virtual shopping task for the assessment of executive functions: validity for people with stroke. *Neuropsychol. Rehabil.* 27, 808–833. doi: 10.1080/09602011.2015.1109523
- Pacheco, T. B. F., Oliveira Rego, I. A., Campos, T. F., and Cavalcanti, F. A. D. C. (2017). Brain activity during a lower limb functional task in a real and virtual environment: a comparative study. *NeuroRehabilitation* 40, 391–400. doi: 10.3233/NRE-161426
- Parsons, T. D. (2011). “Neuropsychological assessment using virtual environments: enhanced assessment technology for improved ecological validity,” in *Advanced Computational Intelligence Paradigms in Healthcare 6. Virtual Reality in Psychotherapy, Rehabilitation, and Assessment*, eds S. Brahmam and L. C. Jain (Springer), 271–289.
- Patel, S., Park, H., Bonato, P., Chan, L., and Rodgers, M. (2012). A review of wearable sensors and systems with application in rehabilitation. *J. Neuroeng. Rehabil.* 9:21. doi: 10.1186/1743-0003-9-21
- Raspelli, S., Pallavicini, F., Carelli, L., Morganti, F., Pedrolì, E., Cipresso, P., et al. (2012). Validating the neuro vr-based virtual version of the multiple errands test: preliminary results. *Presence* 21, 31–42. doi: 10.1162/PRES_a_00077
- Riva, G., Castelnuovo, G., and Mantovani, F. (2006). Transformation of flow in rehabilitation: the role of advanced communication technologies. *Behav. Res. Methods* 38, 237–244. doi: 10.3758/BF03192775
- Rodríguez-Artalejo, F., and Rodríguez-Mañas, L. (2014). The frailty syndrome in the public health agenda. *J. Epidemiol. Commun. Health* 68, 703–704. doi: 10.1136/jech-2014-203863
- Rodríguez, A. C., Roda, C., Montero, F., González, P., and Navarro, E. (2016). An interactive fuzzy inference system for teletherapy of older people. *Cognit. Comput.* 8, 318–335. doi: 10.1007/s12559-015-9356-6
- Sanchez-Vives, M. V., and Slater, M. (2005). From presence to consciousness through virtual reality. *Nat. Rev. Neurosci.* 6, 332–339. doi: 10.1038/nrn1651
- Shallice, T., and Burgess, P. W. (1991). Deficits in strategy application following frontal lobe damage in man. *Brain* 114, 727–741. doi: 10.1093/brain/114.2.727
- Shine, J. M., Matar, E., Bolitho, S. J., Dilda, V., Morris, T. R., Naismith, S. L., et al. (2013). Modeling freezing of gait in Parkinson’s disease with a virtual reality paradigm. *Gait Posture* 38, 104–108. doi: 10.1016/j.gaitpost.2012.10.026
- Slobounov, S. M., Ray, W., Johnson, B., Slobounov, E., and Newell, K. M. (2015). Modulation of cortical activity in 2D vs. 3D virtual reality environments: an EEG study. *Int. J. Psychophysiol.* 95, 254–260. doi: 10.1016/j.ijpsycho.2014.11.003
- Stones, D., and Gullifer, J. (2016). ‘At home it’s just so much easier to be yourself’: older adults’ perceptions of ageing in place. *Ageing Soc.* 36, 449–481. doi: 10.1017/S0144686X14001214
- Teruel, M. A., Oliver, M., Montero, F., Navarro, E., and González, P. (2015). “Multisensory treatment of the hemispatial neglect by means of virtual reality and haptic techniques,” in *International Work-Conference on the Interplay Between Natural and Artificial Computation* (Cham: Springer), 469–478.
- Triberti, S., and Riva, G. (2015). Being present in action: a theoretical model about the “interlocking” between intentions and environmental affordances. *Front. Psychol.* 6:2052. doi: 10.3389/fpsyg.2015.02052
- Weidemann, C. T., Mollison, M. V., and Kahana, M. J. (2009). Electrophysiological correlates of high-level perception during spatial navigation. *Psychon. Bull. Rev.* 16, 313–319. doi: 10.3758/PBR.16.2.313
- White, D. J., Congedo, M., Ciorciari, J., and Silberstein, R. B. (2012). Brain oscillatory activity during spatial navigation: theta and gamma activity link medial temporal and parietal regions. *J. Cogn. Neurosci.* 24, 686–697. doi: 10.1162/jocn_a_00098

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Serino, Barello, Miraglia, Triberti and Repetto. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.