



Differential T Cell Levels of Tumor Necrosis Factor Receptor-II in Children With Autism

Paul Ashwood*

Department of Medical Microbiology and Immunology, and The Medical Investigation of Neurodevelopmental Disorders Institute, University of California, Davis, CA, United States

OPEN ACCESS

Edited by:

Richard Eugene Frye,
Phoenix Children's Hospital,
United States

Reviewed by:

Emily L. Casanova,
School of Medicine Greenville,
University of South Carolina,
United States
David Quentin Beversdorf,
University of Missouri, United States
Robert Hendren,
University of California, San Francisco,
United States

*Correspondence:

Paul Ashwood
pashwood@ucdavis.edu

Specialty section:

This article was submitted to
Child and Adolescent Psychiatry,
a section of the journal
Frontiers in Psychiatry

Received: 24 August 2018

Accepted: 11 October 2018

Published: 20 November 2018

Citation:

Ashwood P (2018) Differential T Cell
Levels of Tumor Necrosis Factor
Receptor-II in Children With Autism.
Front. Psychiatry 9:543.
doi: 10.3389/fpsy.2018.00543

Autism spectrum disorders (ASD) are characterized by impairments in verbal and non-verbal communication, in social interactions, and often accompanied by stereotypical interests and behaviors. A role for immune dysfunction has long been implicated in ASD pathophysiology, behavioral severity, and co-morbidities. The pro-inflammatory cytokine tumor necrosis factor alpha (TNF α) has been associated with ASD in some studies but little is known about its receptors. There are two receptors for TNF α , with TNFR1 relaying many of the signals from TNF α , especially those that are rapid, whilst TNFR2 relays later more long-term effects of TNF α . Proteolytic cleavage can lead to the soluble versions of these receptors which can neutralize the effects of TNF α . Here, we determined levels of TNF α and its receptors in 36 children with a confirmed diagnosis of ASD and 27 confirmed typically developing (TD) controls, 2–5 years-of-age. Children with ASD had higher levels of TNFR2 on T cells compared to controls following cell stimulation. Levels of sTNFR2 were decreased in cell supernatants following stimulation in ASD. Overall these data corroborate the role of inflammatory events in ASD and align with previous studies that have shown differential changes in cellular adaptive immunity in children with ASD. Future longitudinal analyzes of cellular immune function and downstream signaling from immune receptors will help further delineate the role of inflammation in ASD.

Keywords: autism, lymphocyte, T cells, TNF-tumor necrosis factor, behavior, cytokine receptors

INTRODUCTION

Autism spectrum disorders (ASD) are neurodevelopmental disorders affecting 1 in 59 children whom are characterized by significant deficits in communication, social interactions and frequently accompanied by stereotyped or restricted behaviors and interests. The etiology of ASD is complex and largely unknown; however, potential genetic candidates linked with ASD include many genes that regulate immune responses, for instance human leukocyte antigen (HLA)-DR, phosphatase and tensin homolog (PTEN), macrophage migration inhibitory factor (MIF), complement C4B, MET tyrosine receptors, interleukin (IL)-4 receptor, and reelin [Reviewed in (1, 2)]. Significant immune dysfunction is also seen in children with ASD, including prominent neuroinflammation in brain specimens, and alterations in adaptive and innate immune responses in the periphery (3–15).

Tumor necrosis factor- α (TNF α) is a pleiotropic and highly regulated pro-inflammatory cytokine secreted by a number of different cells that plays an important role in coordinating early inflammatory processes, cell proliferation and apoptosis (16). Biologically the activities of TNF α are mediated via two different ubiquitously expressed TNF α receptors, TNF receptor type I (TNFRI; also known as CD120a), and type II (TNFRII; CD120b) (17, 18). TNFRI seems to be the main mediator of TNF α rapid signaling and is found on most tissues, whereas TNFRII mediates later more long-term effects of TNF α and are more commonly expressed on immune cells (17, 19). Proteolytic cleavage of these receptors from the cell surface results in soluble forms (sTNFRI and sTNFRII) that can neutralize TNF α and thus modulate its biological activity (20). The measurement of levels of sTNFRs and TNF α together may be more useful and reliable markers of the inflammatory response and TNF α bioactivity than just TNF α alone.

Although many studies have documented increased expression of TNF α in different neuropsychiatric diseases, for example schizophrenia, depressive disorder, and Alzheimer's disease, only a few have evaluated TNF α in serum/plasma or following stimulation of immune cells in ASD, and the results are often conflicting (3, 4, 21–36). Information on the levels of sTNFRs are even more scant in ASD. Finally, no previous study has assessed both levels of TNFRs on the cell surface of immune cells and production of sTNFRs in ASD.

The present study sought to evaluate levels of TNFRs on T cells and sTNFRs in supernatants following immune challenge in ASD patients compared to TD control subjects.

METHODS

Subjects

This study examined 63 participants enrolled through the Childhood Autism Risk from Genetics and Environment (CHARGE) study at U.C. Davis (37). Full details regarding behavioral measures/assessments and recruitment in the CHARGE study protocols have previously been described (37). Children were consecutively assessed. Participants were free of medication and without chronic clinically defined illness or fever at time of blood draw. The participants were 27 typically developing (TD) controls median age 3.9 years [(interquartile range 2.2–6.1), 4 females] and 36 children with ASD [median age 3.6 years (interquartile range 2.5–4.8), 5 females]. Diagnoses of ASD was based on Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria, and defined as autistic disorder. Further evaluation was confirmed using the Autism Diagnostic Interview-Revised (ADI-R) and the Autism Diagnostic Observation Schedule (ADOS) assessments. Children from the TD groups were screened for autism traits using the Social Communication Questionnaire (SCQ). This study was approved by the UC Davis institutional review board and complied with all requirements regarding human subjects. Parents gave both written and informed consent.

Cell Isolation, Stimulation, and Biochemical Measures

Peripheral blood mononuclear cells (PBMC) were separated from the whole blood by centrifugation over Histopaque-1077 HybridMax lymphocyte separation medium (Sigma; St. Louis, MO) before washing twice in Hanks Balanced Salt Solution (HBSS; VWR; Brisbane, CA). PBMC were either cultured in media alone, or stimulated with PHA (10 μ g/mL; Sigma), for 24 h at 37°C in 5% CO₂. Following culture, plates were centrifuged before supernatants were harvested and stored at –80°C until cytokine analysis and cells processed for flow cytometry.

The quantification of TNF α and its soluble receptors (sTNFRI and sTNFRII) were assessed by ELISA using standard procedures recommended by the manufacturer (Quantikine, R&D Systems, Minneapolis, Minn., USA). All samples were on unstimulated and stimulated cell culture supernatants, in duplicates. The detection limits for the kits were <0.5 pg/mL. Concentrations obtained below the sensitivity limit of detection (LOD) of the method were calculated as LOD/2 for statistical comparisons. Culture supernatants had not undergone any previous freeze/thaws cycle.

Cells were harvested after culture and were washed three times in FACS buffer (PBS, 1% fetal bovine serum albumin (VWR, USA) and 0.1 % sodium azide (Sigma), before being resuspended and stained in 100 μ l FACS buffer containing either the following monoclonal antibodies fluorescein isothiocyanate (FITC)-conjugated mouse anti-human TNFRI (CD120a); phycoerythrin (PE)-conjugated mouse anti-human TNFRII (CD120b); (PE)-Cy5-conjugated mouse anti-human CD3; and allophycocyanin (APC)-conjugated mouse anti-human CD4, CD8 (all antibodies were from BD Biosciences, CA, USA). Appropriate IgG isotype controls (BD bioscience) were used to correct for compensation issues. Cells were incubated at 4°C for 30 min before being spun down and washed with staining buffer. Cells were then analyzed on a LSR II flow cytometer and the data acquired analyzed with FlowJo software (BD Immunocytometry Systems). Lymphocytes were gated using forward scatter and side scatter parameters and CD3⁺ cells for analysis of cell surface TNFRI and TNFRII expression, with further analysis of CD4 and CD8 expression where each parameter was measured separately on CD3⁺ populations, CD3⁺CD4⁺ populations, and CD3⁺CD8⁺ populations.

Statistical Analysis

In primary analyses, induced TNF α and soluble receptors and cell surface markers levels (outcome) were compared by group (predictor) and statistical significance was determined using a parametric Student's *t*-test, following confirmation of normal distribution, with a *p*-value of <0.05 considered significant. Multiple comparisons were adjusted for by using the Benjamini-Hochberg False Discovery Rate. Using answers to questions regarding loss of language (Q11) and loss of social skills (Q25) of the ADI-R, the autism population was further divided into two groups based on the clinical onset of autistic symptoms; namely, children who regressed in acquired language or social skills after initial typical development, and secondly, children who did not

TABLE 1 | Comparison of the TNF α and soluble receptors (pg/ml) following cell culture in media alone or stimulation with PHA in children with autism ($n = 36$) and typically developing controls ($n = 27$). Data are presented as mean \pm standard error of means (SEM).

		TNF α	sTNFRI	sTNFRII
Media	ASD	34.38 \pm 2.81	57.80 \pm 2.13	147.7 \pm 13.07
	TD	33.09 \pm 3.844	58.23 \pm 2.70	182.4 \pm 22.80
PHA	ASD	54.50 \pm 1.89	63.83 \pm 2.48	1,060 \pm 77.72
	TD	54.71 \pm 1.96	66.17 \pm 3.49	1,766 \pm 213.61*


* $p = 0.014$ stimulated levels of sTNFRII were decreased in ASD children compared with PHA stimulated levels in typically developing controls.

regress. We also compared ADOS scores with immune outcomes using Pearson correlations. There were too few subjects with clinical co-morbid features such as gastrointestinal symptoms or sleep disorders to determine differences between cytokine production or cell surface receptor expression. All analyses were carried out using SAS version 9.1 (SAS Inc.; Cary, NC) and graphed with Prism 5 Software (GraphPad Software; San Diego, CA).

RESULTS

Cytokine and soluble receptors were measured in harvested supernatants from unstimulated and PHA-stimulated, PBMC cultures. No difference between children with ASD and TD controls were observed in TNF α or its receptors in unstimulated media alone conditions (Table 1). Activation with PHA led to an increase in all analytes measured across both groups. Observed levels of sTNFRII after PHA-stimulation were significantly less in ASD group compared to the TD group ($p = 0.014$; Table 1). When comparisons were made among children with ASD who had regressed compared to those that had not, unstimulated sTNFRII levels were lower in children with ASD who had not regressed, compared to those that had and to TD controls ($p < 0.04$). However, following PHA-stimulation, both groups were decreased for sTNFRII compared with TD controls and were not significantly different from each other. No other differences were observed in ASD children based on regression. Levels of sTNFRII were significantly negatively correlated with impairments in ADOS social interactions ($r = -0.383$, $p < 0.025$), suggesting lower levels were associated with worse behaviors.

No significant differences in the frequencies of CD3 $^+$, CD4 $^+$, and CD8 $^+$ T cells between TD controls and children with ASD, with or without PHA stimulation were observed. The frequency of TNFRI expressing or TNFRII expressing T cell subsets was not different between groups in unstimulated or stimulated conditions (data not shown). Following immune stimulation both the frequency of cells expressing TNFRI (CD120a) and TNFRII (CD120b) on the cell surface were similar in both groups. However, after PHA stimulation, the amount of cell surface TNFRII (CD120b) receptors, as measured by mean fluorescence intensity (MFI), was significantly increased in children with



ASD compared to controls on CD3 $^+$ T cells as a whole, and both the T helper CD3 $^+$ CD4 $^+$ CD120b $^+$ subsets and cytolytic CD3 $^+$ CD8 $^+$ CD120b $^+$ T cells ($p < 0.03$; Figure 1). No differences were seen between those children with ASD who had regression and those children with ASD that did not have regression. In the ASD group frequency of CD3 $^+$ CD120b $^+$ were associated with worse social behavior on ADOS ($r = 0.238$, $p < 0.035$).

DISCUSSION

To the best of our knowledge this is the first study to assess levels of TNFRs on immune cells and release of sTNFR into cell culture supernatants following immune challenge in ASD. Our results demonstrated increased levels of TNFRII on T cells and T cell subsets following stimulation, and decreased sTNFRII in supernatants in children with ASD, but no differences in TNF α and cell surface or sTNFRI levels were found.

Moreover, we found associations between TNFRII levels on cells or in the supernatants and more impairments in behavior. It is currently unclear how TNF α can affect neurodevelopmental outcomes and behaviors during childhood in ASD, and the data should be treated with caution. However, of note, numerous studies have shown that impairments in core ASD behaviors and associated co-morbid and aberrant behaviors, are strongly correlated with altered immune profiles (38). Further validation of the link between observed behavioral severity and cytokine and cytokine receptors is warranted.

Multiple studies have demonstrated that TNF α levels are increased in serum of individuals with ASD when compared to controls (21, 23, 25, 26, 34–36); however, studies on plasma have provided conflicting results (4, 28, 35). These discordant results may reflect the matrix used as well as methodological issues, including different assessment instruments, and clinical and demographic characteristics of ASD populations studied. In addition, it is possible that TNF α is produced at affected tissues and degraded shortly after its production. As TNFRs can be induced by TNF α , the cellular expression may relate to TNF α activity. Our data suggest an increase in TNFRII on T cells, with subsequent decreases in supernatants presumably due to decreased shedding. Previous studies by Jyonouchi and colleagues corroborate our findings showing decreased sTNFRII in cell culture supernatants after T cell mitogen stimulation with PHA (29–32); however, cell surface expression of TNFR was not previously determined. Soluble TNFRII is generated by proteolytic cleavage by the metalloproteinase TNF α converting enzyme (TACE, also known as ADAM17) which despite this enzyme being increased in blood of ASD children (39) does not result in increased sTNFRII in our study, perhaps suggesting proteolytic cleavage is not the underlying mechanism and that receptors are retained on cells. Another possibility is that cell signaling after TNF α binding is weak and that proteolysis does not occur due to altered signaling pathways (40). Soluble TNFRII may lead to the inactivation of circulating TNF α by the generation of high affinity complexes. These complexes reduce the binding of TNF α to any cell membrane target receptors thus downregulating the activity and response to TNF α (41). As soluble TNFRs neutralize TNF α , measuring both receptor and cytokine may be more reflective of the net effect of cytokines, in that even though we did not see an increase in TNF α after stimulation, the increased presence of receptors on the cell surface and decreased levels of neutralizing sTNFRII in supernatants may indicate that more TNF α -receptor ligand binding and signaling occurred. Future studies should determine TNF α -TNR responses in immune cells from children with ASD, including other immune cell types not just T cells. Further

studies should also address whether cytokines such as TNF α are involved in pro-apoptotic signaling or pro-life signaling and the net balance of those signals induced by inflammatory responses in ASD.

Differential expression of markers of T cell activation have been shown in children with ASD following immune stimulation including increased CD26, CD38, CD69, HLA-DR, but decreases in CD25 (42–46). Another member of the TNF receptor family, CD137 is a co-stimulatory molecule expressed by activated T cells and enhances T cell proliferation, effector functions, and survival was also increased on T cells in ASD children (43). TNFRII is predominantly found on immune cells and is primarily associated with lymphocyte proliferation (17, 19). The TNFRs differ in their intracellular domains which induce distinct intracellular signals. We recently showed that T cell signaling was altered in children with ASD (40) and may suggest that certain downstream signals such as the mTOR pathway may lead to preferential immune activation. Taken together these results may suggest potential differential activation of T cell subsets in ASD when compared to controls.

In summary, the results presented here are in agreement with prior studies and suggest that inflammatory processes, as evinced by alterations of pro-inflammatory cytokine signaling, could contribute to ASD pathophysiology. It is also possible that other aspects of ASD might provide additional feedback that influences or compounds the altered immune state in ASD. Further, future studies including longitudinal measurements of the same participants might further reveal the putative role for levels of TNFRs as bio signatures of ASD development and progression. More studies will be required to characterize the changes in TNF α and the TNRII and their temporal relationship with ASD development, as well as assessment of cytokine and cognate receptors in the brain.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and has approved it for publication.

FUNDING

This study was funded by the NIEHS Children's Center grant (P01 ES011269), US EPA STAR program grant (R833292 and R829388), NIEHS CHARGE study (R01ES015359), NICHD (HD086669, HD090214, and U54 HD079125), Autism Speaks Foundation, HEDCO Foundation, The Boler Company Foundation, NARSAD Foundation, and the Johnson Foundation.

ACKNOWLEDGMENTS

We would like to thank all staff CHARGE study staff, and Angela Tarver, Charity Onore, Paula Krakowiak, and Amanda Enstrom for technical support. The dedication and commitment of the families who took part in these studies is gratefully acknowledged.

REFERENCES

- Careaga M, Ashwood P. Autism spectrum disorders: from immunity to behavior. *Methods Mol Biol.* (2012) 934:219–40. doi: 10.1007/978-1-62703-071-7_12
- Mead J, Ashwood P. Evidence supporting an altered immune response in ASD. *Immunol Lett.* (2015) 163:49–55. doi: 10.1016/j.iml.2014.11.006
- Ashwood P, Wakefield AJ. Immune activation of peripheral blood and mucosal CD3+ lymphocyte cytokine profiles in children with autism and gastrointestinal symptoms. *J Neuroimmunol.* (2006) 173:126–34. doi: 10.1016/j.jneuroim.2005.12.007
- Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah I, Van de Water J. Elevated plasma cytokines in autism spectrum disorders provide evidence of immune dysfunction and are associated with impaired behavioral outcome. *Brain Behav Immun.* (2011) 25:40–5. doi: 10.1016/j.bbi.2010.08.003
- Breece E, Paciotti B, Nordahl CW, Ozonoff S, Van de Water JA, Rogers SJ, et al. Myeloid dendritic cells frequencies are increased in children with autism spectrum disorder and associated with amygdala volume and repetitive behaviors. *Brain Behav Immun.* (2013) 31:69–75. doi: 10.1016/j.bbi.2012.10.006
- Careaga M, Rogers S, Hansen RL, Amaral DG, Van de Water J, Ashwood P. Immune endophenotypes in children with autism spectrum disorder. *Biol Psychiatry* (2017) 81:434–41. doi: 10.1016/j.biopsych.2015.08.036
- Enstrom AM, Lit L, Onore CE, Gregg JP, Hansen RL, Pessah IN, et al. Altered gene expression and function of peripheral blood natural killer cells in children with autism. *Brain Behav Immun.* (2009) 23:124–33. doi: 10.1016/j.bbi.2008.08.001
- Enstrom A, Onore C, Van de Water J, Ashwood P. Differential monocyte responses to TLR ligands in children with autism spectrum disorders. *Brain Behav Immun.* (2010) 24:64–71. doi: 10.1016/j.bbi.2009.08.001
- Garbett K, Ebert PJ, Mitchell A, Lintas C, Manzi B, Mirnics K, et al. Immune transcriptome alterations in the temporal cortex of subjects with autism. *Neurobiol Dis.* (2008) 30:303–11. doi: 10.1016/j.nbd.2008.01.012
- Gupta S, Aggarwal S, Rathanavaran B, Lee T. Th1- and Th2-like cytokines in CD4+ and CD8+ T cells in autism. *J Neuroimmunol.* (1998) 85:106–9. doi: 10.1016/S0165-5728(98)00021-6
- Heuer L, Ashwood P, Schauer J, Goines P, Krakowiak P, Hertz-Picciotto I, et al. Reduced levels of immunoglobulin in children with autism correlates with behavioral symptoms. *Autism Res.* (2008) 1:275–83. doi: 10.1002/aur.42
- Jyonouchi H, Geng L, Cushing-Ruby A, Quraishi H. Impact of innate immunity in a subset of children with autism spectrum disorders: a case control study. *J Neuroinflamm.* (2008) 5:52–9. doi: 10.1186/1742-2094-5-52
- Li X, Chauhan A, Sheikh AM, Patil S, Chauhan V, Li X-M, et al. Elevated immune response in the brain of autistic patients. *J Neuroimmunol.* (2009) 207:111–6. doi: 10.1016/j.jneuroim.2008.12.002
- Molloy CA, Morrow AL, Meinen-Derr J, Schleifer K, Dienger K, Manning-Courtney P, et al. Elevated cytokine levels in children with autism spectrum disorder. *J Neuroimmunol.* (2005) 172:198–205. doi: 10.1016/j.jneuroim.2005.11.007
- Vargas DL, Nascimbene C, Krishnan C, Zimmerman AW, Pardo CA. Neuroglial activation and neuroinflammation in the brain of patients with autism. *Ann Neurol.* (2005) 57:67–81. doi: 10.1002/ana.20315
- McCoy MK, Tansey MG. TNF signaling inhibition in the CNS: implications for normal brain function and neurodegenerative disease. *J Neuroinflamm.* (2008) 45:1–13. doi: 10.1186/1742-2094-5-45
- Hohmann HP, Remy R, Brockhaus M, van Loon AP. Two different cell types have different major receptors for human tumor necrosis factor (TNF alpha). *J Biol Chem.* (1989) 264:14927–34.
- Brockhaus M, Schoenfeld HJ, Schlaeger EJ, Hunziker W, Lesslauer W, Loetscher H. Identification of two types of tumor necrosis factor receptors on human cell lines by monoclonal antibodies. *Proc Natl Acad Sci USA.* (1990) 87:3127–31. doi: 10.1073/pnas.87.8.3127
- Grell M, Douni E, Wajant H. The transmembrane form of tumor necrosis factor is the prime activating ligand of the 80 kDa tumor necrosis factor receptor. *Cell* (1995) 83:793–802. doi: 10.1016/0092-8674(95)90192-2
- Mohler KM, Torrance DS, Smith CA, Goodwin RG, Stremmler KE, Fung VP, et al. Soluble tumor necrosis factor (TNF) receptors are effective therapeutic agents in lethal endotoxemia and function simultaneously as both TNF carriers and TNF antagonists. *J Immunol.* (1993) 151:1548–61.
- Al-Ayadhi LY. Pro-inflammatory cytokines in autistic children in central Saudi Arabia. *Neurosciences* (2005) 10:155–8.
- Ashwood P, Anthony A, Torrente F, Wakefield AJ. Spontaneous mucosal lymphocyte cytokine profiles in children with autism and gastrointestinal symptoms: mucosal immune activation and reduced counter regulatory interleukin-10. *J Clin Immunol.* (2004) 24:664–73. doi: 10.1007/s10875-004-6241-6
- Chez MG, Dowling T, Patel PB, Khanna P, Kominsky M. Elevation of tumor necrosis factor-alpha in cerebrospinal fluid of autistic children. *Pediatr Neurol.* (2007) 36:361–5. doi: 10.1016/j.pediatrneurol.2007.01.012
- Croonenberghs J, Bosmans E, Deboutte D, Kenis G, Maes M. Activation of the inflammatory response system in autism. *Neuropsychobiology* (2002) 45:1–6. doi: 10.1159/000048665
- Ferguson BJ, Marler S, Altstein LL, Lee EB, Mazurek MO, McLaughlin A, et al. Associations between cytokines, endocrine stress response, and gastrointestinal symptoms in autism spectrum disorder. *Brain Behav Immun.* (2016) 58:57–62. doi: 10.1016/j.bbi.2016.05.009
- Ghaffari MA, Mousavinejad E, Riahi F, Mousavinejad M, Afsharmanesh MR. Increased serum levels of tumor necrosis factor-alpha, resistin, and visfatin in the children with autism spectrum disorders: a case-control study. *Neurol Res Int.* (2016) 2016:9060751. doi: 10.1155/2016/9060751
- Goines PE, Ashwood P. Cytokine dysregulation in autism spectrum disorders (ASD): possible role of the environment. *Neurotoxicol Teratol.* (2013) 36:67–81. doi: 10.1016/j.ntt.2012.07.006
- Hu CC, Xu X, Xiong GL, Xu Q, Zhou BR, Li CY, et al. Alterations in plasma cytokine levels in chinese children with autism spectrum disorder. *Autism Res.* (2018) 11:989–999. doi: 10.1002/aur.1940
- Jyonouchi H, Sun S, Le H. Proinflammatory and regulatory cytokine production associated with innate and adaptive immune responses in children with autism spectrum disorders and developmental regression. *J Neuroimmunol.* (2001) 120:170–9. doi: 10.1016/S0165-5728(01)00421-0
- Jyonouchi H, Geng L, Ruby A, Reddy C, Zimmerman-Bier B. Evaluation of an association between gastrointestinal symptoms and cytokine production against common dietary proteins in children with autism spectrum disorders. *J Pediatr.* (2005) 146:605–10. doi: 10.1016/j.jpeds.2005.01.027
- Jyonouchi H, Geng L, Ruby A, Zimmerman-Bier B. Dysregulated innate immune responses in young children with autism spectrum disorders: their relationship to gastrointestinal symptoms and dietary intervention. *Neuropsychobiology* (2005) 51:77–85. doi: 10.1159/000084164
- Jyonouchi H, Geng L, Streck DL, Dermody JJ, Toruner GA. MicroRNA expression changes in association with changes in interleukin-1β/interleukin10 ratios produced by monocytes in autism spectrum disorders: their association with neuropsychiatric symptoms and comorbid conditions (observational study). *J Neuroinflamm.* (2017) 14:229. doi: 10.1186/s12974-017-1003-6
- Rose D, Ashwood P. Potential cytokine biomarkers in autism spectrum disorders. *Biomark Med.* (2014) 8:1171–81. doi: 10.2217/bmm.14.39
- Steeb H, Ramsey JM, Guest PC, Stocki P, Cooper JD, Rahmoune H, et al. Serum proteomic analysis identifies sex-specific differences in lipid metabolism and inflammation profiles in adults diagnosed with Asperger syndrome. *Mol Autism.* (2014) 5:4. doi: 10.1186/2040-2392-5-4
- Suzuki K, Matsuzaki H, Iwata K, Kamenoy Y, Shimmura C, Kawai S, et al. Plasma cytokine profiles in subjects with high-functioning autism spectrum disorders. *PLoS ONE* (2011) 6:e20470. doi: 10.1371/journal.pone.0020470
- Xie J, Huang L, Li X, Li H, Zhou Y, Zhu H, et al. Immunological cytokine profiling identifies TNF-α as a key molecule dysregulated in autistic children. *Oncotarget* (2017) 8:82390–8. doi: 10.18632/oncotarget.19326
- Hertz-Picciotto I, Croen LA, Hansen R, Jones CR, Van de Water J, Pessah IN. The CHARGE study: an epidemiologic investigation of genetic and environmental factors contributing to autism. *Environ Health Perspect.* (2006) 114:1119–25. doi: 10.1289/ehp.8483
- Onore C, Careaga M, Ashwood P. The role of immune dysfunction in the pathophysiology of autism. *Brain Behav Immun.* (2012) 26:383–92. doi: 10.1016/j.bbi.2011.08.007

39. Corbett BA, Kantor AB, Schulman H, Walker WL, Lit L, Ashwood P, et al. A proteomic study of serum from children with autism showing differential expression of apolipoproteins and complement proteins. *Mol Psychiatry* (2007) 12:292–306. doi: 10.1038/sj.mp.4001943
40. Onore C, Yang H, Van de Water J, Ashwood P. Dynamic Akt/mTOR signaling in children with autism spectrum disorder. *Front Pediatr*. (2017) 5:43. doi: 10.3389/fped.2017.00043
41. Bemelmans MH, Gouma DJ, Buurman WA. LPS-induced sTNF receptor release *in vivo* in a murine model. investigation of the role of tumor necrosis factor, IL-1, leukemia inhibiting factor, and IFN-gamma. *J Immunol*. (1993) 151:5554–62.
42. Ashwood P, Corbett BA, Kantor A, Schulman H, Van de Water J, Amaral DG. In search of cellular immunophenotypes in the blood of children with autism. *PLoS ONE* (2011) 6:e19299. doi: 10.1371/journal.pone.0019299
43. Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah IN, Van de Water J. Altered T cell responses in children with autism. *Brain Behav Immun*. (2011) 25:840–9. doi: 10.1016/j.bbi.2010.09.002
44. Denney DR, Frei BW, Gaffney GR. Lymphocyte subsets and interleukin-2 receptors in autistic children. *J Autism Dev Disord*. (1996) 26:87–97. doi: 10.1007/BF02276236
45. Plioplys AV, Greaves A, Kazemi K, Silverman E. Lymphocyte function in autism and Rett syndrome. *Neuropsychobiology* (1994) 29:12–6. doi: 10.1159/000119056
46. Mostafa GA, Al Shehab A, Fouad NR. Frequency of CD4+CD25high regulatory T cells in the peripheral blood of Egyptian children with autism. *J Child Neurol*. (2010) 25:328–35. doi: 10.1177/0883073809339393

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Ashwood. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.