



# The Role of Neuroinflammation in Postoperative Cognitive Dysfunction: Moving From Hypothesis to Treatment

Seyed A. Safavynia<sup>1\*</sup> and Peter A. Goldstein<sup>1,2,3</sup>

<sup>1</sup> Department of Anesthesiology, Weill Cornell Medical College, New York, NY, United States, <sup>2</sup> Department of Medicine, Weill Cornell Medical College, New York, NY, United States, <sup>3</sup> Neuroscience Graduate Program, Weill Cornell Medical College, New York, NY, United States

## OPEN ACCESS

### Edited by:

Richard Eugene Frye,  
Phoenix Children's Hospital,  
United States

### Reviewed by:

Caroline Menard,  
Laval University, Canada  
Xinhong Zhu,  
Southern Medical University, China

### \*Correspondence:

Seyed A. Safavynia  
sas9204@med.cornell.edu

### Specialty section:

This article was submitted to  
Molecular Psychiatry,  
a section of the journal  
Frontiers in Psychiatry

**Received:** 16 October 2018

**Accepted:** 19 December 2018

**Published:** 17 January 2019

### Citation:

Safavynia SA and Goldstein PA (2019)  
The Role of Neuroinflammation in  
Postoperative Cognitive Dysfunction:  
Moving From Hypothesis to  
Treatment. *Front. Psychiatry* 9:752.  
doi: 10.3389/fpsy.2018.00752

Postoperative cognitive dysfunction (POCD) is a common complication of the surgical experience and is common in the elderly and patients with preexisting neurocognitive disorders. Animal and human studies suggest that neuroinflammation from either surgery or anesthesia is a major contributor to the development of POCD. Moreover, a large and growing body of literature has focused on identifying potential risk factors for the development of POCD, as well as identifying candidate treatments based on the neuroinflammatory hypothesis. However, variability in animal models and clinical cohorts makes it difficult to interpret the results of such studies, and represents a barrier for the development of treatment options for POCD. Here, we present a broad topical review of the literature supporting the role of neuroinflammation in POCD. We provide an overview of the cellular and molecular mechanisms underlying the pathogenesis of POCD from pre-clinical and human studies. We offer a brief discussion of the ongoing debate on the root cause of POCD. We conclude with a list of current and hypothesized treatments for POCD, with a focus on recent and current human randomized clinical trials.

**Keywords:** postoperative cognitive dysfunction, cognitive decline, neuroinflammation, central nervous system, microglia, anesthesia

## INTRODUCTION

Disordered neurocognitive function following surgery is a heterogeneous set of conditions, which includes both the fluctuating and typically transient postoperative delirium and the more protracted problem of postoperative cognitive dysfunction (POCD). POCD is a well-known risk of the surgical experience, having been described as a consequence of anesthesia as early as 1887 (1), and a common complication of cardiac surgery since the 1950's (2). More than 60 years following its modern description, it is only just now that clearly articulated guidelines have been suggested for identifying POCD (3). POCD has been loosely defined as a significant reduction in cognitive performance from baseline following surgery, and diagnosed as subtle deficits in multiple core neurocognitive domains, including executive function, attention, verbal memory, psychomotor speed, and visuospatial abstraction (4, 5). Given that the literature thus far has used the term POCD to describe these deficits, we will also use the term here, but recognize going forward the nomenclature will likely evolve so as to conform with new guidelines (3). Since the 1950's, advanced age has been shown to be one of the strongest associations for development of POCD: the incidence

of POCD is reported to be anywhere between 9 and 54% 1 week after surgery in adults over age 65 (6), with no difference in rates based on the type of surgery and/or anesthetic (7). POCD itself can persist long after surgery, with an incidence between 10 and 17% at 3 months following surgery (7, 8) and 3% at 12 months following surgery (9). Moreover, POCD can contribute to severe cognitive deficits over the long term, affecting overall morbidity and mortality, with increased hospital costs (10, 11). The health and economic burdens of POCD are likely to increase over the next several years: Life expectancy is increasing, and more than 30% of individuals over age 65 have surgery annually (12).

At the epidemiological level, a handful of risk factors for the development of POCD have emerged from population studies; controversy exists, however, in the interpretation of these data and their clinical implications. Risk factors for POCD were initially identified in patients undergoing cardiac surgery, and included advanced age, aortic valve replacement, and prolonged (mean 70 min) cardiopulmonary bypass (CPB) time (13). While advanced age (>65 years) has been consistently identified as a risk factor for POCD (8, 14), the evidence is less convincing with other potential risk factors due to differences in populations and neurocognitive testing modalities (4, 7). For example, it has long been thought that preexisting frailty in general (15–20), and neurocognitive frailty in particular (9, 21, 22) may be a risk factor for POCD as these patients may be vulnerable to cognitive insults at baseline. Indeed, observational studies have shown that surgery may precipitate further cognitive decline in patients with neurodegenerative disorders such as Alzheimer's disease (AD) (23), and biomarkers of AD such as the apolipoprotein E4 (APOE-4) genotype have also been associated with development of POCD in elderly patients (24, 25). However, a long-term retrospective analysis did not show an accelerated progression to dementia in patients with AD after non-cardiac surgery (26). More recent data in humans show that while the CSF tau/ $\beta$ -amyloid ratio increases following surgery, the increase is independent of the type of anesthetic (i.e., propofol vs. isoflurane) (27), further calling into question the predictive value of these biomarker studies. These discrepancies may be in part due to confounders such as temperature regulation; hypothermia rather than anesthesia *per se* seems to be the driver behind the observed tauopathy (28, 29), with dexmedetomidine as a possible exception (30). Chronic inflammatory states such as diabetes, metabolic syndrome, and atherosclerosis have all been proposed as potential risk factors for POCD (31–33), while pro-cognitive activities such as sleep, exercise, and level of education seem to be protective (34). Despite these data, the heterogeneous populations and study paradigms used inherently limit the clinical interpretation of these risk factors.

At the cellular level, data from animal and human studies suggest that neuroinflammation from either surgery or anesthesia is a major contributor to the development of POCD, yet the specific relationship between inflammation and POCD remains unknown. Multiple rodent models of surgery have shown upregulation of pro-inflammatory cytokines and inflammatory mediators in both peripheral tissues and the central nervous system (CNS) (35, 36). Similarly in rats, inflammation in the form of prior infection can also increase the incidence and severity

of POCD (37, 38). In human studies, patients who develop POCD also show increases in serum and cerebrospinal fluid (CSF) pro-inflammatory cytokines, irrespective of the type of surgery (39–42), which has been corroborated in meta-analyses (43, 44). However, there seems to be little relationship between the magnitude of the neuroinflammation and the development of POCD. For example, while CPB was thought to be a strong initiator of peripheral and subsequent neuroinflammation (45), the rates of POCD in cardiac and non-cardiac surgery are similar (7), as well as in pulsatile vs. non-pulsatile CPB (46) and on-pump and off-pump cardiac surgery (45). Meta-regressions show a slight relationship with plasma levels of interleukin-6 (IL-6) and S100 calcium-binding protein  $\beta$  (S100 $\beta$ ) and POCD, but no other cytokines studied have shown any correlation (43). While inflammation always occurs with surgery, POCD does not, and it remains unclear what specific risk factors and triggers are responsible for this conversion.

Despite the advances in research, fundamental barriers exist to understanding POCD in a generalized context, limiting the ability to predict patients at risk for POCD and develop appropriate therapies for such patients. Firstly, POCD has been broadly defined, with no historical formal clinical definition (5, 47, 48). Similarly, animal models of POCD are defined using a variety of metrics, each testing different cognitive domains as a proxy for POCD (49). Without a formal definition, it is difficult to accurately and consistently identify patients with POCD and construct appropriate animal models, thereby limiting a generalized understanding of the epidemiology and pathogenesis of the disorder. Secondly, determining the root causes of POCD is difficult as surgery and anesthesia occur almost invariably in tandem (48), with larger and more high-risk surgery often necessitating longer anesthetic times. Thirdly, proposed treatments showing promise in animal studies are often not as effective when tested in clinical trials, revealing a need for a more nuanced understanding of POCD.

We present a broad topical overview of the current state of the literature regarding the effects of neuroinflammation on the development of POCD. We will review the proposed cellular mechanisms underlying the pathogenesis of POCD in pre-clinical and human studies. We will present the evidence underlying the debate on the etiologic contributions of neuroinflammation and POCD in both animal models and human studies, whether surgical, anesthetic, or both. Lastly, we will discuss proposed treatments for POCD, with a focus on recent and current human randomized clinical trials.

While POCD is often grouped with postoperative delirium (POD) in the literature, we limit the discussion in this review to POCD and not POD. POD and POCD are distinct disorders: Delirium is defined in the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) as a disorder of reduced attention and orientation to the environment, accompanied by cognitive disturbances in an acute and fluctuating course with lucid intervals (50). By contrast, POCD is described as an objectively measured decline in cognition in the postoperative state compared to the preoperative state (48). Unlike delirium, the time course of POCD does not fluctuate with lucid intervals, and some patients never recover

from the initial insult (51, 52). Nevertheless, there is a growing body of evidence suggesting that neuroinflammation contributes to POD; for a detailed review on the role of neuroinflammation on POD, please see Maldonado (53).


## PROPOSED MECHANISMS FOR PATHOGENESIS OF POCD

Taken together, data from animal and human studies have fueled the **hypothesis** that peripheral surgical trauma causes CNS inflammation via disruption of the blood-brain barrier (BBB), which then causes a functional disruption in neural activity, leading to POCD. Each component of this hypothesis is regulated by a variety of inflammatory mediators discussed below. This sequence of events can persist long after surgery and resolution of neuroinflammation, and can accelerate neurocognitive decline in neurocognitively frail populations.

### Peripheral Initiation of Inflammation

It is well-known that aseptic surgical trauma causes inflammation at the surgical site, which is amplified *via* peripheral pro-inflammatory cytokines. In response to surgical trauma, damaged cells at the site of injury passively release small biomolecules known as damage-associated molecular patterns (or danger-associated molecular patterns; DAMPs) (4, 54). In particular, the DAMP known as high molecular group box 1 protein (HMGB1) is released following surgical trauma and binds to Toll-like receptors (TLRs) and the receptor for advanced glycosylation end products (RAGE) on the cell membrane of peripherally circulating bone marrow derived monocytes (BMDMs) (55) (Figure 1). In rats, surgery and anesthesia have been associated with increased hippocampal HMGB1 expression (56); similarly, human studies have shown that plasma HMGB1 levels are correlated with the level of inflammation in both non-cardiac surgery and non-surgical inflammatory states (57). In rodents, elevations of HMGB1 are associated with cognitive deficits (58), which can be mitigated in the presence of HMGB1 inhibitors (4, 59). These results are corroborated by evidence that HMGB1 levels are elevated in patients with POCD following gastrointestinal surgery (60).

When bound by HMGB1, both TLR-4 and RAGE activate nuclear factor kappa B (NF- $\kappa$ B), a transcription factor which regulates the expression of pro-inflammatory cytokines (Figure 1). Normally, cytosolic NF- $\kappa$ B is bound to the NF- $\kappa$ B inhibitor I $\kappa$ B in an inactive state; however, when I $\kappa$ B is phosphorylated by I $\kappa$ B kinase (IKK), NF- $\kappa$ B is released and enters the nucleus, causing pro-inflammatory cytokine upregulation (55). Once activated by NF- $\kappa$ B, the pro-inflammatory cytokines interleukin-1 beta (IL-1 $\beta$ ), IL-6, and tumor necrosis factor alpha (TNF $\alpha$ ) cause further release of HMGB1 in a positive feedback loop, amplifying the inflammatory response (57). Additionally, IL-1 and TNF $\alpha$  can cause further activation of NF- $\kappa$ B, resulting in cyclooxygenase 2 isozyme (COX-2) upregulation (34). There is a strong association between elevations in serum pro-inflammatory cytokines and POCD in both animal models (61, 62) and human studies (41, 44). Moreover, in rats, inhibition of NF- $\kappa$ B and pro-inflammatory cytokines has been associated



**FIGURE 1** | Signaling pathways involved in peripheral initiation of inflammation. Injured cells release damage-associated molecular patterns (DAMPs) including high mobility group box-1 protein (HMGB1) in response to surgical trauma. HMGB1 activates nuclear factor-kappa B (NF- $\kappa$ B) signaling pathways in bone marrow derived monocytes (BMDMs), causing nuclear translocation of NF- $\kappa$ B, increased expression of cyclooxygenase 2 isozyme (COX-2) upregulation, and expression of pro-inflammatory cytokines interleukin-1 beta (IL-1 $\beta$ ), interleukin 6 (IL-6), and tumor necrosis factor alpha (TNF $\alpha$ ). These pro-inflammatory cytokines can act back on BMDMs in positive feedback loops (solid curved lines) as well as promote further release of HMGB1 from injured cells by unknown mechanisms (dashed curved lines). IKK, I $\kappa$ B kinase; IL-6R, IL-6 receptor; P, phosphate group; RAGE, receptor for advanced glycosylation end products; TLR-4, Toll-like receptor 4; TNF $\alpha$ R, TNF $\alpha$  receptor.

with a reduction in POCD using various metrics (including Morris water maze, elevated plus maze, fear conditioning, and passive avoidance test) (63–65).

### Blood-Brain Barrier Breakdown

Peripheral pro-inflammatory cytokines disrupt BBB permeability *via* COX-2 upregulation and matrix metalloproteinases (MMPs), allowing pro-inflammatory cytokines to enter the CNS

(**Figure 2**). Normally, the BBB is made up of tight junctions held together by transmembrane proteins (i.e., occludins, claudins, junctional adhesion molecules) between neurovascular endothelial cells (66). This structure only allows for the passive diffusion of water, gases, and small lipid-soluble molecules (67). However, pro-inflammatory cytokines IL-1 and TNF $\alpha$  can upregulate COX-2 in neurovascular endothelial cells, which promotes local prostaglandin synthesis (68) and disrupts BBB permeability (69) (**Figure 2**). TNF $\alpha$ , IL-1 $\beta$ , and IL-6 have all been found in hippocampal tissue in rats (69–71) and in human CSF (42, 72) following surgical trauma, suggestive of a breakdown in the BBB. Cytokine elevation in the CNS has also been associated with memory dysfunction in mice (73) and cognitive dysfunction (measured by different neurocognitive metrics—see **Table 2**) in humans (41, 42). These data suggest that BBB breakdown is associated with cytokine influx and cognitive impairment, however this evidence does not rule out the possibility that the cytokine elevation may be generated locally within the CNS. More convincingly, immunoglobulin G (IgG), which is not present normally in the brain, has also been identified in hippocampal slices in rats following surgery (56, 74). Similarly, CNS-specific proteins such as S100 $\beta$  and neuron-specific enolase (NSE) are found in plasma following cardiac and non-cardiac surgery in patients with POCD (43, 75, 76). TNF $\alpha$  can also upregulate transcription of MMPs, particularly MMP-9; this aberrant MMP expression can degrade extracellular matrix proteins *in vitro*, further breaking down the BBB (66) (**Figure 2**). Unfortunately, there is only limited *in vivo* evidence concerning the role of MMPs in BBB disruption (66). At a functional level however, MMP-9 gene deletion mice exposed to surgical trauma have been shown to exhibit better cognitive performance (in terms of fear conditioning) compared to wild-type mice (77).

Lastly, once the BBB is disrupted, circulating BMDMs in the periphery are able to enter the CNS and augment neuroinflammation *via* cytokine expression and microglial activation (**Figure 2**). While mast cells and microglia exist in the CNS, there are no normally occurring populations of dendritic cells or monocytes (78). In the setting of inflammation and BBB breakdown however, BMDMs are recruited to the CNS (79) *via* interactions between the chemokine monocyte chemo-attractant protein 1 [MCP-1, also called C-C motif ligand 2 (CCL2)] and the BMDM cell surface receptor chemokine receptor type 2 (CCR2) (**Figure 2**). Once the BMDMs are present in the CNS, they continue to secrete pro-inflammatory cytokines *via* upregulation of NF- $\kappa$ B transcription (34), and activate microglia in the CNS, further amplifying the neuroinflammation. In mice it has been shown that preoperative depletion of BMDMs reduced POCD (80), suggesting that BMDM migration plays a pivotal role in POCD. Taken together, once the BBB is disrupted, cytokines can freely enter the CNS, causing trafficking of BMDMs to neural tissues and initiating poorly regulated immune functions.

## Microglial Activation

Microglia are known as the “resident macrophages” of the CNS (81) and have many important contributory functions in the CNS, including synaptic pruning during development (82) and


synaptic scaling in neural plasticity (83). Derived from yolk-sac cells, microglia migrate to the CNS early in development, before the differentiation of many cell types in the CNS (81). As a part of the innate immune system, microglia surveil brain parenchyma (84) and are the first responders to pathogens in the CNS. Although a fully differentiated cell, microglia have the unique ability to self-replenish within the CNS (85).

Normally, microglia are in an inactive state maintained by binding of the CX3CR1 protein to the microglial CX3CR1 receptor (86). However, in the setting of inflammation and BBB breakdown, they can differentiate into one of two activated phenotypes, M1 and M2 (87). The M1 phenotype has high phagocytic properties and is pro-inflammatory (88, 89), while the M2 phenotype is involved in tissue repair and remodeling and is anti-inflammatory (90). Not surprisingly, pro-inflammatory mediators such as TNF $\alpha$  or lipopolysaccharide promote microglial differentiation into the M1 phenotype (91). Moreover, TNF $\alpha$  blockade can suppress microglial activation in mice (35). Conversely, anti-inflammatory cytokines such as IL-4 are known to play a role in promoting the alternative M2 phenotype (88). However, recent evidence is beginning to challenge the dichotomy of the M1/M2 phenotypes, suggesting that there are many overlapping phenotypes with various functions and activation pathways (92). One such new area is the role of mast-cell degeneration in activating microglia: In a recent rat study, Zhang et al. (93) showed that peripheral surgery induced CNS mast cell degranulation and subsequent microglial activation. Further, administration of cromolyn sodium (which inhibits mast cell degranulation) inhibits microglial activation in rats (93, 94), demonstrating a new microglial interaction and a possible new therapeutic target for POCD.

Once microglia are activated, they continue to upregulate expression of pro-inflammatory cytokines, thus amplifying neuroinflammation and contributing to the development of POCD (**Figure 2**). Activated microglia are known to release HMGB1, TNF $\alpha$ , and IL-1 $\beta$  in a variety of rodent models (95–97). Further, astrocytes and microglia both upregulate expression of MCP-1/CCL2 (98), and astrocyte CCL2 can induce further microglial activation *in vitro* (99, 100). These chemokines cause further influx of BMDMs into the CNS: Trafficked BMDMs in turn can activate microglia to the M1 phenotype *via* TNF $\alpha$ /IL-1 expression, and activated microglia recruit more BMDMs into the CNS *via* reciprocal TNF $\alpha$  expression (101). In aged mice, microglial activation is increased in POCD (37, 49, 102). Moreover, in mice, both perioperative microglial depletion (103) and promotion of an M2 phenotype *via* erythropoietin administration (99) improved memory dysfunction as measured by passive avoidance and novel object recognition tests.

## The Role of Oxidative Stress

In addition to the inflammatory pathways described above, surgical trauma can also produce oxidative stress and deplete the body of antioxidants (57); these oxidative processes, when superimposed on the inflammatory pathway, can contribute to the development of POCD. Surgical stimulation in rodents can raise the levels of CNS nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, an enzyme compound that



generates superoxide in response to stress (104). The superoxide radicals in turn generate other reactive oxygen species (ROS), potentially causing direct damage of neural tissues. Additionally, peripheral oxidative stress can also disrupt the BBB (105), representing a convergence of oxidative stress with the neuroinflammatory pathway. Within the CNS, microglia have been shown to release ROS (106) in response to both HMGB1 (107) and S100 $\beta$  (108). Of note, activated microglia are known to release HMGB1 (97), creating the opportunity for yet another neuroinflammatory positive feedback loop.

Recent evidence from animal and human studies suggests that oxidative stress alone can contribute to POCD. Hippocampal neurons are very metabolically active and are some of the most sensitive neurons to oxidative stress (109); it follows that hippocampal injury from oxidative stress can have profound effects on memory formation and spatial navigation. In aged rats, tibial fracture surgery was associated with memory impairments (measured by open field task and novel object recognition task) on postoperative day 1 with corresponding increases in oxidative damage in the hippocampus and prefrontal cortex (109). Oxidative injury from hypoglycemia has also been shown to induce cognitive impairment in rats, and inhibition of NADPH oxidase has been shown to mitigate such impairments (110). In humans, levels of the ROS nitric oxide are correlated with

development of POCD (*via* neurocognitive battery) at 4 days and 3 months following cardiac surgery (111).

## Functional Consequences of CNS Inflammation

Memory formation occurs in the hippocampus and is achieved by a process known as long-term potentiation (LTP). Although the mechanisms of induction and maintenance of LTP at various synapses in the CNS are very complex and somewhat controversial, LTP is thought to be achieved by high frequency glutamatergic activation of hippocampal neurons (112). At rest, presynaptic glutamatergic Schaffer cells signal to postsynaptic CA1 collateral neurons. The CA1 neurons themselves contain three types of glutamate receptors: the metabotropic Glu2 receptor and the ionotropic AMPA and NMDA receptors. During normal, low-frequency stimulation of CA1 neurons, glutamate acts on all receptors, but the NMDA channels are blocked by magnesium. With high frequency stimulation however, postsynaptic depolarization causes an activation of NMDA receptors, which causes an influx of calcium and activation of second messenger systems (112). Downstream, the number and sensitivity of AMPA receptors is increased through phosphorylation, and synaptic strength is increased, resulting in memory formation (113).


The presence of pro-inflammatory cytokines can have detrimental effects on the regulation of neurotransmitter signaling in the hippocampus, ultimately resulting in excitotoxic neuronal damage and resulting cognitive impairment. First, the hippocampus has a large number of cytokine receptors, rendering it susceptible to high concentrations of pro-inflammatory cytokines such as IL-1 and TNF $\alpha$  in neuroinflammatory processes (114, 115). Once these cytokine receptors are activated at high levels, there is a downregulation of metabotropic Glu2 receptors causing enhanced AMPA/NMDA signaling, disrupting the process of LTP (116). Meanwhile, HMGB1 can also potentiate glutamate signaling through NMDA, causing an increased influx of glutamate in hippocampal neurons, which ultimately results in glutamate toxicity (117). Further, TNF $\alpha$  can depress inhibitory neurotransmission *via* downregulation of GABA receptors, disrupting the delicate balance of excitatory and inhibitory neurotransmission and ultimately favoring glutamate toxicity (118). These detrimental effects are compounded by the T-cell mediated release of glutamate from activated microglia *via* a separate glutamate transporter subtype (119). Collectively, the aforementioned mechanisms contribute to glutamate toxicity in the hippocampus, resulting in neuronal death and cognitive impairment.

### Cholinergic Anti-inflammatory Pathway

Although peripheral pro-inflammatory cytokines are the primary initiator of neuroinflammation, they are also involved in regulating the inflammatory response *via* a vagal reflex arc (34) (Figure 3). This serves to help limit the degree of inflammation and protect organ systems from further damage. In this arc, DAMPs released from surgical trauma are sensed by vagal afferents that terminate on the nucleus tractus solitarius (NTS) (120). The efferent arc of this reflex originates from

fibers within the dorsal motor nucleus of the vagus, sending signals to the celiac ganglion. Within the celiac ganglion, vagal efferents regulate postganglionic catecholaminergic fibers *via* functional connections within the splenic nerve (121). The splenic nerve endings are in close anatomical position with T lymphocytes, which express  $\beta_2$  adrenergic receptors (122). When activated, T lymphocytes upregulate transcription of choline acetyltransferase, facilitating synthesis of acetylcholine (ACh) (120); this newly synthesized ACh can then activate circulating macrophages that express alpha-7 nicotinic ACh receptors ( $\alpha_7$  nAChRs). Ultimately, activation of  $\alpha_7$  nAChR-expressing macrophages causes inactivation of NF- $\kappa$ B, which decreases cytokine release (34). In addition, vagal stimulation is known to induce regulatory T-cells and secretion of anti-inflammatory cytokines IL-4 (which promotes microglial differentiation to the M2 phenotype) and IL-10 (123, 124). One experiment in rats treated with the cholinesterase inhibitor physostigmine following laparotomy showed a reduction in hippocampal IL-1 $\beta$  and TNF  $\alpha$  expression and hippocampal damage (125). In humans, anticholinergic drugs are widely known to precipitate POCD (126), although it is unclear whether the cholinergic anti-inflammatory pathway is involved in this process. Thus, it has been proposed that vagal stimulation may mitigate the development of POCD (127), although this remains untested in human literature.

The vagus nerve also regulates pro-resolving lipid mediators known as resolvins, lipoxins, and macrophage mediators in resolving inflammation (maresins), all of which are derived from polyunsaturated fatty acids (4, 128). Resolvins act to block the migration of neutrophils and monocytes, and can reduce the oxidative burst of neutrophils (129). Similarly to  $\alpha_7$  nAChR-expressing cells, maresins can inhibit NF- $\kappa$ B activity in macrophages and help promote microglial differentiation



to the M2 phenotype (130). Together, these lipid mediators represent possible new therapeutic targets for POCD. Lastly, the vagus nerve can also promote the restoration of BBB integrity *via* netrin-1, a protein involved in cell migration and axonal pathfinding during development (34), however netrin-1 has yet to be explored as a therapeutic target for POCD in human studies.

## ETIOLOGY OF POCD

It is difficult to determine the etiology of POCD as surgery and anesthesia are virtually inseparable in modern society. As a result, surgery and anesthesia act as natural confounders of each other, hindering an understanding of a causal relationship and spurring controversy in the literature. Carefully designed animal and human studies have been developed to tease out the contributions of surgery or anesthesia to the development of POCD, however there is great variability in experimental design, limiting the interpretation of these results.

### Evidence From Animal Models

Animal models can provide strong insight into the etiology of POCD by exposing a genetically identical group to different anesthetic or surgical regimens and comparing the rates of POCD across groups. Moreover, animal models have the advantage of assessing neuroinflammation at the level of brain parenchyma in terminal experiments, creating a vital link to the neuroinflammatory hypothesis. Much of the evidence supports the notion that surgery and not anesthesia causes both neuroinflammation and POCD: For example, studies in rodents have shown that hippocampal pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF $\alpha$  (70) and HMGB1 (56) are increased with surgery and isoflurane anesthesia, but not with isoflurane anesthesia alone. Moreover, the same studies have shown a higher incidence of POCD (measured using spatial learning paradigms) with surgery and isoflurane compared to isoflurane alone. Increases in hippocampal IL-1 $\beta$ , and TNF $\alpha$  and impaired spatial learning have also been observed in carotid exploration surgery with propofol anesthesia but not propofol anesthesia alone (63), and no differences have been observed in POCD (measured by fear conditioning and spatial learning) between total intravenous anesthesia (TIVA) and volatile anesthetic (131). More convincingly, a recent study demonstrated that open abdominal surgery under local anesthesia caused increases in hippocampal IL-6, TNF $\alpha$ , and memory impairments (71), suggesting that anesthesia *per se* is not necessary for the production of neuroinflammation and subsequent development of POCD.

However, studies looking solely at the effects of anesthesia yield mixed conclusions, with anesthesia being implicated as either a causal or protective agent. Administration of “balanced anesthesia” (consisting of both intravenous and volatile anesthetic agents) during early postnatal life has been shown to produce neurotoxic effects in rats (132), and repeated exposure to the volatile anesthetic sevoflurane has been shown to affect the cognitive function of young, but not adult, mice (133). Similarly, repeated exposure to 5 h of isoflurane (end-tidal isoflurane = 0.7–1.5 vol %) in infant Rhesus macaque monkeys exposed on

postnatal day (P)6, P9, and P12 resulted in evidence of motor and socioemotional deficits when tested 12 months later; infants that were only exposed once on P5 had no such alterations (134). In older mice, isoflurane alone has been associated with hippocampal inflammation and impairment of spatial memory (135), however in rats, isoflurane alone did not have an effect on spatial memory processes, even with repeat anesthetics (136). In contrast, in the setting of myocardial ischemia-reperfusion injury in rats, sevoflurane seems to exert a protective effect, mitigating impairments in long-term potentiation (LTP) and improving memory function (137, 138). While the discrepancies between these studies may be partially explained by the different experimental paradigms and the different metrics used to evaluate POCD, it may also be possible that anesthesia induces more subtle changes in cognitive function compared to surgery. One study showed that the combination of isoflurane and intraperitoneal ketamine alone decreased spatial memory and learning, but to a lesser degree than with combined anesthesia and surgery (139). Moreover, hippocampal pro-inflammatory cytokines were only increased with the combination of surgery and anesthesia, suggesting that if anesthesia alone can cause POCD, it may do so *via* non-inflammatory mechanisms. A summary of the findings of relevant animal studies can be found in **Table 1**.

### Evidence From Human Studies

Although human studies rely on heterogeneous populations and are limited in scope by ethical considerations, it is possible to tease out the relative contributions of surgery vs. anesthesia to the development of POCD by comparing outcomes in patients undergoing different anesthetic regimens, including general anesthesia, neuraxial anesthesia, and sedation. Indeed, a large ( $n = 636$ ) prospective observational study comparing coronary artery bypass grafting (CABG) under general anesthesia, hip replacement under spinal anesthesia, and percutaneous coronary angiography under sedation showed no difference in POCD rates between groups (7). This result was especially interesting as rates of POCD were long thought to be higher in cardiac surgery due to the inflammation associated with CPB (2, 13). These results have been supported by prospective observational studies showing no difference in POCD between spinal vs. general anesthesia for orthopedic surgery (41, 42). Moreover, a large systematic review was unable to demonstrate a clear connection between general anesthesia and POCD (140), although the majority of studies examined were underpowered and used variable methodologies. As in animal studies, it has even been proposed that volatile anesthesia may be protective in the setting of ischemic organ damage, ultimately mitigating POCD from organ ischemia (141).

Results from randomized controlled trials, while rigorous, are inconsistent and merit further investigation into the causes of POCD. As seen in observational studies, a prospective randomized clinical trial comparing the use of general vs. spinal anesthesia in extracorporeal shock wave lithotripsy showed no significant difference in the incidence of POCD defined by a neurocognitive battery (142), suggesting that surgery and not anesthesia causes POCD. In a separate study of patients undergoing CABG using high-dose vs. low-dose fentanyl

**TABLE 1** | Selected relevant pre-clinical studies on etiology of POCD.

Study	Animal model	Experimental model	Cognitive testing	Cellular/Molecular findings	Neurocognitive findings
Cao et al. (70)	Adult (3–6 month) and aged (20–24 month) old Sprague Dawley rats	Partial hepatectomy under sevoflurane anesthesia vs. sevoflurane alone	Morris water maze	Upregulated expression of IL-1 $\beta$ and IL-6 on postoperative day 1 in all rats, and in aged rats until postoperative day 3	Surgery and anesthesia, but not anesthesia alone, caused impairments in latency and distance in all rats on postoperative day 1, and in aged rats until postoperative day 3
He et al. (56)	22–23 month old Sprague-Dawley rats	Splenectomy under general anesthesia vs. 2 h isoflurane anesthesia vs. naïve control	Reversal learning version of Morris water maze	Upregulation of HMGB1 and RAGE levels in surgical group BBB disruption (by TEM) in surgical group	Surgery and anesthesia, but not anesthesia alone, caused cognitive impairments from surgery to postoperative day 3
Qian et al. (139)	20–22 month old BALB/c mice	Splenectomy with isoflurane vs. isoflurane alone vs. control	Y-maze testing	Splenectomy increased hippocampal expression of IL-1 $\beta$ and TNF $\alpha$	Splenectomy with anesthesia and anesthesia alone both impaired cognitive testing on postoperative days 1 and 3
Tasbihgou et al. (138)	Adult male Wistar rats	Deep vs. light propofol anesthesia, with and without subsequent exposure to hypoxia	Novel object recognition test	Light anesthesia group with hypoxia had lower neurogenesis, but higher BDNF and microglia-ramification	No impairment in cognitive function in either deep or light anesthesia
Walters et al. (136)	Adult Sprague-Dawley rats	Four exposures to isoflurane anesthesia (2, 2, 4, and 6 h) over 7 weeks	Fixed consecutive number, incremental repeated acquisition, progressive ratio tasks <sup>†</sup>	none	No deficits in any cognitive tasks after single or repeat anesthetic exposure
Wang et al. (135)	6–8 month old male C57BL/6 mice; 14 month old male C57BL/6 mice	Isoflurane vs. no anesthetic exposure	Morris water maze	Older but not younger mice had increased hippocampal expression of NLRP3 <sup>‡</sup>	Older but not younger mice had cognitive impairment after isoflurane anesthesia compared to no anesthetic exposure
Xu et al. (71)	9 and 18 month old female C57BL/6J mice	Laparotomy under local anesthesia (no sedation) vs. sham procedure (no incision)	Fear conditioning system	Surgery increased hippocampal levels of IL-6 and TNF $\alpha$ in all mice, with larger increases in older mice	Cognitive deficits with surgery alone in both young and older mice
Zhang et al. (63)	4 month old male Fischer 344 rats	Right carotid exploration with propofol and buprenorphine anesthesia vs. anesthesia alone	Barnes maze Fear conditioning system	Surgery decreased cytoplasmic hippocampal NF- $\kappa$ B, increased IL-1 $\beta$ , IL-6, MMP-9	Surgery and anesthesia, but not anesthesia alone caused impairments in cognitive metrics
Zhang et al. (131)	20 month old male Fischer 344 rats	Right carotid exploration with propofol-buprenorphine anesthesia vs. isoflurane-buprenorphine anesthesia	Barnes maze Fear conditioning system	No difference in hippocampal TNF $\alpha$ and IL-1 $\beta$ expression in propofol vs. isoflurane anesthesia	Surgery caused impairments in cognitive metrics independent of anesthetic type
Zhu et al. (137)	Adult male Wistar rats	Transient coronary artery occlusion with and without sevoflurane preconditioning vs. sham operation	N/A	Coronary occlusion increased hippocampal TNF $\alpha$ and IL-1 $\beta$ mRNA expression 1–3 days postoperatively; cytokine levels attenuated by sevoflurane	Coronary occlusion inhibited LTP compared to sham operation; sevoflurane preconditioning reversed this effect on postoperative days 1 and 3

BBB, blood-brain barrier; BDNF, brain-derived neurotrophic factor; HMGB1, high mobility group box-1 protein; IL-1 $\beta$ , interleukin-1 beta; IL-6, interleukin 6; LTP, long-term potentiation; MMP-9, matrix metalloproteinase 9; N/A, non-applicable; NF- $\kappa$ B, nuclear factor-kappa B; NLRP3, NOD-like receptor protein 3 inflammasome; RAGE, receptor for advanced glycosylation end products; TEM, transmission electron microscopy; TNF $\alpha$ , tumor necrosis factor alpha.

<sup>†</sup>Rats were trained to perform these tasks for at least 15 months prior to anesthetic exposure.

<sup>‡</sup>NLRP3 causes maturation and secretion of cytokines IL-1 $\beta$  and IL-18.



anesthesia, the same group showed no difference in POCD at 3 and 12 months following surgery, although low-dose fentanyl did have higher rates of POCD at 1 week following surgery (143). However, randomized controlled trials comparing propofol and volatile anesthesia in laparoscopic cholecystectomy (144) and esophageal resection (145) have shown a higher incidence of POCD and pro-inflammatory markers with volatile anesthesia. It is important to note that these trials used different neurocognitive assessments to identify POCD, including the mini-mental status exam (MMSE) and the Montreal Cognitive Assessment (MoCA). Other randomized clinical trials attempting to show a dose-response effect with volatile anesthesia have shown that high-dose anesthetic is associated with an increased incidence of POCD (146, 147). However, these trials used the Bispectral Index™ (BIS™) as a proxy for anesthetic depth, which has been shown to be influenced by a variety of non-anesthetic factors (148–150) and is often discordant with brain activity observed under anesthesia (151); thus, BIS™ may not be an accurate representation of anesthetic depth and may limit the interpretation of these studies. A summary of the findings of relevant clinical studies can be found in **Table 2**.

## PROPOSED TREATMENTS FOR POCD

The neuroinflammatory hypothesis provides many varied targets for candidate treatments for POCD. These treatments largely fall into one of three strategies: blocking inflammation by inhibiting inflammatory mediators (anti-inflammatory), preventing the oxidative component of inflammation (anti-oxidative), or protecting neurons during and promoting neuronal health before surgery (pro-neuronal). We present an overview of multiple candidate treatments, with a brief discussion of their hypothesized mechanisms of action and their plausibility established from pre-clinical models where appropriate. We will focus on the existing human data for each treatment, where available, including ongoing human trials from the United States National Library of Medicine (ClinicalTrials.gov), the European Union Clinical Trials Register (clinicaltrialsregister.eu), and the Australian New Zealand Clinical Trials Registry (anzctr.org.au). Please see **Table 3** for a summary of clinical studies for proposed treatments. For an in-depth review of the pre-clinical and human data supporting various treatments for POCD, please see Skvarc et al. (57).

### Anti-inflammatory

#### COX-2 Inhibitors

The cyclooxygenase 2 (COX-2) enzyme is responsible for catalyzing the conversion of arachidonic acid to pro-inflammatory prostaglandins (68) and can increase BBB permeability (69). For these reasons, COX-2 is considered an important mediator of neuroinflammation and a potential target for treatment of POCD. Indeed, rodent models have shown that the COX-2 inhibitor parecoxib is capable of downregulating IL-1  $\beta$  and TNF $\alpha$  expression (168); furthermore, meloxicam, a non-steroidal anti-inflammatory drug (NSAID) with relative selectivity for COX-2 has been shown to decrease short-term

deficits in recognition memory following surgery (169). So far, two human trials have evaluated the efficacy of COX-2 inhibition on POCD, both in geriatric patients undergoing total knee arthroplasty (152, 153). In a trial of 134 elderly patients, parecoxib was shown to decrease pro-inflammatory markers and POCD incidence (as assessed using a neurocognitive battery) compared to placebo at 1 week, but not 3 months following surgery (152), although this negative result was largely due to improved cognitive performance in the placebo group. Similarly, a trial of 178 elderly patients showed that celecoxib reduced pro-inflammatory markers and POCD (determined by reduction in performance of  $\geq 2$  of 5 cognitive tests) at 1 week following surgery compared to placebo (153). There are no ongoing registered clinical trials testing the use of NSAIDs or selective COX-2 inhibitors in POCD.

#### Minocycline

Minocycline is a second-generation tetracycline antibiotic that has anti-inflammatory properties; it has shown to be useful in reducing cognitive deficits in animal models of cerebral ischemia, Alzheimer's disease, and Parkinson's disease (170). Minocycline readily crosses the BBB, and thus may also be useful in inhibiting neuroinflammation. In rats, minocycline has been shown to block IL-1 $\beta$ , with a concomitant reduction in surgery-induced hippocampal-dependent memory impairment (determined by fear conditioning test) (73). In mice undergoing appendectomy, preoperative administration of minocycline has been shown to downregulate production of IL-1 $\beta$ , IL-6, and TNF $\alpha$ , inhibit microglial activation, and impair learning (measured *via* Morris water maze and fear conditioning test) (171). However, it has been recently demonstrated in aged rats undergoing abdominal surgery that minocycline may simply delay microglial activation (172). Thus, while it has been proposed that minocycline may be useful for reducing POCD, it may not prevent development of delayed POCD. Currently, there is a multicenter randomized Phase 3 clinical trial recruiting patients to investigate the efficacy of preoperative minocycline in reducing POCD in patients with colon cancer undergoing colorectal surgery (ClinicalTrials.gov identifier NCT02928692).

#### Dexamethasone

Dexamethasone is a corticosteroid with glucocorticoid actions and powerful (>30 times more potent than cortisol) anti-inflammatory properties. As with other steroid hormones, dexamethasone inhibits the infiltration of leukocytes into the target inflammatory region (173); moreover, it can downregulate the transcription of cytokines and other cell adhesion molecules (174). Although dexamethasone has well-demonstrated anti-inflammatory actions, it is unclear whether it may have an effect on the development of POCD. In a study by Karaman et al. (175), male rats given sevoflurane were shown to develop memory deficits (measured *via* Morris water maze) at 7 and 30 days post anesthesia. Administration of 0.1 mg/kg dexamethasone before anesthetic administration reversed these deficits at both time points, suggesting its utility in

**TABLE 2 |** Relevant clinical studies on etiology of POCD.

Study	Study Type	Cohort	Sample size (n)	Surgical procedure(s)	Anesthetic exposure	Cognitive metrics	Key findings
Evered et al. (7)	Prospective observational	CABG, hip replacement: adults > 55 CA: adults > 50	636	Elective CABG, hip replacement, CA	CABG: general anesthesia Hip replacement: spinal anesthesia CA: sedation	Battery of seven neuropsychological tests	No difference in POCD rates between groups (CABG—16%; hip replacement—16%; CA—21%)
Geng et al. (144)	Prospective randomized	Adults > 60	150	Laparoscopic cholecystectomy	Propofol vs. sevoflurane vs. isoflurane anesthesia	Battery of eight neuropsychological tests	Lower POCD in propofol compared to sevoflurane or isoflurane on postoperative days 1 and 3
Hirsch et al. (42)	Prospective observational	Adults ≥ 55	10	Elective major knee surgery	Spinal anesthesia with propofol sedation and femoral nerve catheter	Word list test Verbal fluency test Digit symbol test	40% POCD on postoperative day 1; 20% POCD on postoperative day 2; 40% POCD on postoperative day 3
Hou et al. (147)	Prospective randomized	Adults ≥ 60; ASA 1-2	66	Elective total knee arthroplasty	Deep vs. light anesthesia <sup>†</sup> with sevoflurane and propofol, femoral and sciatic nerve blocks	MoCA Z-score < 1.96	Higher POCD in deep (20%) compared to light (3%) anesthesia
Ji et al. (41)	Prospective observational	Adults ≥ 65	83	Elective total hip replacement	Spinal anesthesia	Digit symbol substitution test Concentration endurance test Number connection test <sup>‡</sup>	POCD rate 24.6% on postoperative day 7
Qiao et al. (145)	Prospective randomized	Adults 65–75	90	Esophageal resection	Sevoflurane vs. methylprednisone and sevoflurane vs. propofol	MoCA MMSE	Higher POCD in sevoflurane group on postoperative days 1, 3, 7
Shu et al. (146)	Prospective randomized	Females 20–60	192	Gynecologic laparoscopic surgery	Sevoflurane with remifentanyl, titrated to BIS <sup>††</sup>	MMSE Trail-making test	Lower POCD in 40 ≤ BIS ≤ 50 group on postoperative day 1
Silbert et al. (142)	Prospective randomized	Adults > 55 without previous neurologic deficit	100	Extracorporeal shock wave lithotripsy	General vs. spinal anesthesia	Battery of eight neuropsychological tests	No difference in POCD rates between groups
Silbert et al. (143)	Prospective randomized	Adults > 55 without previous neurologic deficit	350	Elective CABG	High-dose vs. low-dose fentanyl anesthesia	Battery of eight neuropsychological tests	Higher POCD in low-dose fentanyl group 1 week following surgery. No difference in POCD at 3 and 12 months following surgery

ASA, American Society of Anesthesiologists Classification Scale; BIS, Bispectral Index; CA, coronary angiography; CABG, coronary artery bypass graft; MMSE, Mini-mental Status Examination; MoCA, Montreal Cognitive Assessment.

<sup>†</sup> Deep vs. light anesthesia determined by BIS values 40–50 vs. 55–65, respectively. <sup>††</sup> BIS values stratified to three groups: 30 ≤ BIS ≤ 40, 40 ≤ BIS ≤ 50, 50 ≤ BIS ≤ 50.

<sup>‡</sup> Neurocognitive tests in this study amended to a Chinese protocol.

mitigating POCD. However, a randomized clinical trial of patients given 1 mg/kg intraoperative dexamethasone during cardiac surgery failed to demonstrate a difference in cognitive performance both at 1 month and at 12 months following surgery (154). There is only one registered clinical trial on dexamethasone and POCD (ClinicalTrials.gov identifier NCT01332812); this Phase 4 study of 300 patients compared administration of 8 mg of dexamethasone following anesthesia

induction vs. no injection and measured POCD *via* a cognitive battery up to 180 days post-surgery. Currently no results are reported.

### Cholinergic Agents

The cholinergic anti-inflammatory pathway provides a variety of potential therapeutic targets for POCD. Both the  $\alpha 7$  nAChR agonist PHA 586487 (176) and physostigmine (125) have been

**TABLE 3** | Clinical studies for proposed treatments for POCD.

Treatment	Published clinical studies and relevant findings	Registered ongoing clinical trials (if applicable)
<b>ANTI-INFLAMMATORY</b>		
COX-2 inhibitors	Zhu et al. (152): intraoperative and postoperative parecoxib vs. placebo in total knee arthroplasty (reduction in POCD at 1 week but not 3 months postoperatively) Zhu et al. (153): 1 week of preoperative celecoxib vs. placebo in total knee arthroplasty (reduction in POCD at postoperative day 7)	none
Minocycline	none	NCT02928692: preoperative minocycline vs. no treatment in colorectal surgery
Dexamethasone	Ottens et al. (154): intraoperative dexamethasone bolus vs. no treatment during cardiac surgery (no difference in cognitive performance at 1 and 12 months postoperatively)	NCT01332812: intraoperative dexamethasone bolus vs. no treatment in general surgery
Cholinergic agents	Doraiswamy et al. (155): 12-week course of donepezil >6 months vs. no treatment after CABG (improved memory recall, no improved cognition)	NCT02419352: sugammadex vs. neostigmine/atropine at end of general anesthesia NCT02927522: donepezil vs. placebo for 7 days following general surgery
Targeted cytokine inhibition	None	None
<b>ANTIOXIDATIVE</b>		
Statins	Das et al. (156): postoperative statin vs. placebo in off-pump CABG (reduced POCD on postoperative day 6)	None
N-acetylcysteine	None	PANACEA trial ACTRN12614000411640: NAC vs. placebo twice daily for 4 days beginning on day of non-cardiac surgery
Edaravone	None	None
<b>PRO-NEURONAL</b>		
Dexmedetomidine	Li et al. (157): dexmedetomidine bolus and intraoperative infusion vs. saline in laparoscopic cholecystectomy (reduced POCD on postoperative day 1) Chen et al. (158): correlation between reduction of pro-inflammatory cytokines and reduced POCD on postoperative day 1 in general surgery	NCT02275182: intraoperative dexmedetomidine vs. placebo in general surgery NEUROPRODEX trial 2013-000823-15: intraoperative dexmedetomidine vs. placebo in cardiac and abdominal surgery NCT03480061: intraoperative dexmedetomidine bolus and postoperative infusion vs. standard sedation in cardiac surgery NCT02923128: postoperative dexmedetomidine vs. sufentanil infusion in elective non-cardiac surgery
Amantadine	None	NCT03527134: five-day postoperative amantadine vs. no treatment in general surgery
Enhancing cognitive reserve	None	NCT02747784: three-month postoperative cognitive training regimen vs. no treatment in breast/urogynecological surgery
<b>VARIOUS TARGETS</b>		
Local anesthetics	Wang et al. (159): lidocaine bolus and intraoperative infusion in CABG surgery (improved working memory, verbal associative learning compared to saline controls) Chen et al. (160): lidocaine bolus and intraoperative infusion in spinal surgery (slight improvement in cognitive function)	NCT00975910: lidocaine bolus and intraoperative infusion vs. placebo in supratentorial craniotomy NCT02848599: bupivacaine vs. morphine PCA for 72 h following general surgery
Ketamine	Hudetz et al. (161): ketamine bolus at anesthetic induction during cardiac surgery (improved memory/executive function) Nagels et al. (162): ketamine bolus and intraoperative infusion following anesthetic induction during cardiac surgery (no change in POCD compared to placebo)	NCT02892916: ketamine bolus following anesthetic induction vs. placebo in elective orthopedic surgery
Lipid mediators	none	None
Cannabinoid receptors	none	none
Melatonin	Hansen et al. (163): melatonin 6 mg/kg daily for 3 months improved sleep but had no effect on POCD in women having breast cancer surgery Fan et al. (164): melatonin 1 mg/kg daily for 6 days improved sleep and improves MMSE scores in patients undergoing hip arthroplasty	none
Turmeric	none	none
Acupuncture	Gao et al. (165): Electroacupuncture preserved MMSE scores 2 and 4 days following general surgery. Lin et al. (166): Electroacupuncture preserved MMSE scores on day 3 following intestinal surgery for cancer. (167): Electroacupuncture preserved MMSE scores in elderly patients on day 3 following colorectal surgery for cancer.	none

CABG, coronary artery bypass graft; MMSE, Mini-Mental State Exam; NAC, N-acetylcysteine; PCA, patient-controlled analgesia.

shown to reduce pro-inflammatory cytokines and neuronal damage in rat hippocampus following surgery. However, neither of these studies evaluated behavioral impairments, limiting their generalizability to POCD. In humans, during anesthetic emergence, patients are often given cholinesterase inhibitors such as neostigmine to reverse neuromuscular blocking agents (which are routinely administered to help provide optimal surgical conditions). However, the cyclic oligosaccharide sugammadex, which rapidly and profoundly reverses neuromuscular blockade by encapsulating nondepolarizing steroidal neuromuscular blocking agents such as rocuronium and vecuronium (177), has significantly reduced the use of cholinesterase inhibitors during surgery and provides a unique way to test the association of cholinesterase inhibitors on POCD. Indeed, one registered clinical trial (ClinicalTrials.gov identifier NCT02419352) has randomized 160 patients to receive either sugammadex or the combination of neostigmine and atropine at the end of surgery and anesthesia; results have not yet been published. Other human studies have focused on the anticholinesterase drug donepezil as a potential therapy. In a pilot randomized clinical trial of 44 patients, Doraiswamy et al. (155) showed that a 12-week course of donepezil given at least 6 months following CABG surgery improved memory recall but not cognition. A new Phase 3 clinical trial (ClinicalTrials.gov identifier NCT02927522) plans to randomize over 500 patients to receive donepezil or placebo for 7 days following surgery, and evaluate for POCD 1 week following surgery (although it is unclear what psychological tests are used to define POCD in this study). Again, although vagal stimulation has been proposed to mitigate the development of POCD (127), there are no current human trials designed to test this hypothesis.

### Targeted Cytokine Inhibition

Although there are currently no human data and no registered clinical trials, drugs that block specific cytokines are already utilized as treatment for chronic inflammatory diseases such as rheumatoid arthritis (RA) and may be a potential target for POCD therapies. The IL-1 receptor antagonist anakinra represents one such target: It has been shown that IL-1 knockout mice have lower levels of IL-6 following peripheral surgery, and less memory impairment (73). Similarly, intracisternal administration of an IL-1 receptor antagonist immediately preceding abdominal surgery in aged rats prevented a decrease in memory consolidation on postoperative day 4 (178). The anti-TNF $\alpha$  antibody Etanercept (also used in RA) may be another target for POCD, as preoperative administration of anti-TNF $\alpha$  antibody inhibited IL-1 $\beta$  production in mice and mitigated memory impairments in mice (35). Further, the IL-6 receptor antibody tocilizumab has been shown to reduce memory impairments in mice following surgery (179).

### Antioxidative Statins

Statins are reversible competitive inhibitors of the enzyme 3-hydroxy-3-methylglutaryl-CoA reductase (HMG-CoA reductase). This enzyme catalyzes the conversion of HMG-CoA to mevalonate, and is the rate-limiting step of cholesterol synthesis from fatty acids (180). As part of this enzymatic

process, NADPH is produced; by inhibiting HMG-CoA reductase, NADPH production is lowered, which can reduce the levels of oxidative species (181). Statins have been widely proposed to be beneficial for neurological disorders including dementia (182) and postoperative delirium (183). In POCD, a small randomized controlled trial comparing postoperative statin vs. placebo administration in patients undergoing off-pump CABG showed a significant reduction in memory dysfunction (measured by postgraduate institute memory scale) on postoperative day 6 (156). Unfortunately, no other prospective clinical trials are currently underway to investigate the otherwise promising effects of a widely utilized drug.

### N-Acetylcysteine

N-acetylcysteine (NAC) has antioxidant properties which are related to its role as a precursor for glutathione synthesis (184). Additionally, in pre-clinical studies, NAC has been shown to downregulate pro-inflammatory cytokine synthesis including HGMB-1 (185), upregulate anti-inflammatory cytokine synthesis (186), and reduce microglial activation (187). A systematic review of the human literature has suggested that NAC supplementation can have beneficial cognitive effects for patients with a wide variety of neurological and psychiatric disorders, including Alzheimer's disease, traumatic brain injury, Parkinson's disease, and addictive behavior (184), thus raising the possibility of NAC as a potential treatment for POCD. Only one randomized controlled trial, The Post-Anesthesia N-acetylcysteine Cognitive Evaluation (PANACEA) trial (Australian New Zealand Clinical Trials Registry identifier ACTRN12614000411640) is currently being conducted to investigate the utility of NAC in POCD. This single center trial has randomized patients recovering from non-cardiac surgery to receive 1,200 mg of NAC or placebo twice daily beginning on the day of surgery and continuing for four consecutive days. POCD will be assessed *via* a neurocognitive battery at 1 week, 3, and 12 months following surgery (188). The study is ongoing and no results have been reported at this time.

### Edaravone

Edaravone is a free radical scavenger that is used as an adjunct therapy for acute ischemic stroke in Japan, and as therapy for amyotrophic lateral sclerosis (ALS) in Japan and the United States. These uses are based on small randomized clinical trials that have shown modest efficacy in stroke (189) and early-stage ALS (190). Edaravone readily crosses the BBB, and has been shown to mitigate or ameliorate impairments in spatial and working memory in rats at 3 and 7 days following left nephrectomy and lipopolysaccharide administration (38). Moreover, the same group showed an increase in hippocampal levels of the antioxidant superoxide dismutase and a decrease in microglial activation on postoperative day 3. Taken together, this evidence suggests that edaravone has antioxidative and anti-inflammatory properties and may be a potential treatment for POCD in humans, however there are no published human studies or registered clinical trials.

## Pro-neuronal Dexmedetomidine

Dexmedetomidine is a centrally-acting presynaptic  $\alpha_2$  adrenergic receptor antagonist used for sedation in the operating room and intensive care unit; its mechanism of action is inhibition of norepinephrine release from adrenergic neurons projecting from the locus coeruleus to the basal forebrain, anterior cortex, intralaminar nucleus of the thalamus, and the preoptic area of the hypothalamus (191). Dexmedetomidine's sedative properties are largely believed to be due to norepinephrine inhibition in the preoptic area of the hypothalamus, an important nucleus in regulating arousal and sleep pathways. Dexmedetomidine is also hypothesized to have actions in the spinal cord, and is used as an adjunct for intraoperative analgesia (192) and the prolongation of regional nerve blockade (193). Recently, dexmedetomidine has been shown to enhance HMGB1 resolution in mice, likely *via* a vagotonic mechanism (194), suggesting that it also has downstream effects on reducing inflammation. Human studies have shown that dexmedetomidine bolus followed by infusion throughout laparoscopic cholecystectomy reduces serum pro-inflammatory cytokines and POCD (as measured *via* MMSE scores) compared to saline on postoperative day 1 (157). Moreover, Chen et al. (158) showed a correlation between the level of reduction of pro-inflammatory cytokines and POCD on postoperative day 1 (measured *via* MMSE), providing a much-needed link between cytokine levels and the severity of cognitive dysfunction. There are several registered ongoing Phase 4 clinical trials examining the efficacy of dexmedetomidine on POCD, comparing intraoperative dexmedetomidine to placebo (ClinicalTrials.gov identifier NCT02275182, NEUROPRODEX trial–EudraCT number 2013-000823-15), looking at late (12 months following surgery) POCD (ClinicalTrials.gov identifier NCT03480061), and comparing postoperative dexmedetomidine vs. sufentanil infusion (ClinicalTrials.gov identifier NCT02923128). So far, no data have been reported from these clinical trials.

## Amantadine

Amantadine was initially marketed as an antiviral agent but was found to have dopaminergic actions which led to its use in Parkinson's disease (74). *In vivo*, amantadine has also been demonstrated to promote the production of glial cell line-derived neurotrophic factor (GDNF), an important pro-neuronal agent that promotes glial growth, protects glia, and inhibits microglial activation (195). In a rat surgical model, animals treated with intraperitoneal amantadine or intracerebroventricular GDNF showed a reduction of memory impairment compared to controls 1 day following surgery (74). Further, amantadine inhibited surgery induced neuroinflammation on postoperative day 1. In humans, there is only one randomized clinical trial in the recruitment phase investigating the use of a 5-day course of amantadine (beginning with one dose preoperatively) on POCD (ClinicalTrials.gov identifier NCT03527134).

## Enhancing Cognitive Reserve

Poor cognitive function preoperatively is a potential risk factor for development of POCD, and pro-cognitive activities

such as sleep, exercise, and education level seem to have a protective effect on POCD (34). As a result, it has been proposed that preoperative cognitive training may have a beneficial effect on reducing the incidence and severity of POCD. In rats, a cognitively stimulating environment has shown to attenuate surgery induced cognitive memory impairments (measured *via* novel object recognition test) and hippocampal cytokine increases (196). There is one registered clinical trial (REACT trial, ClinicalTrials.gov identifier NCT02747784) currently recruiting female patients with breast or urogynecological surgery for a 3-month postoperative cognitive training regimen compared to no treatment. Patients will be measured for POCD *via* a neurocognitive battery at 3 months following surgery; data from this trial are not available at this time.

## Candidate Treatments With Various Targets Local Anesthetics

Local anesthetics such as lidocaine and bupivacaine work by stabilizing the open, inactive state of voltage-gated sodium channels; when injected peri-neuronally, the preferential local diffusion of local anesthetics to pain fibers produces its analgesic actions (197, 198). Because pain is a trigger for inflammatory pathways, it has been proposed that local anesthetics may reduce peripheral inflammation (and thus neuroinflammation and POCD). Despite the plausibility of this hypothesis, human data has not been convincing. Patients undergoing CABG surgery given lidocaine bolus 1.5 mg/kg and infusion of 4 mg/kg/h throughout surgery showed improvements in working memory and verbal associative learning compared to saline controls on postoperative day 9, however both groups had deficits in short-term memory, processing speed, and executive function (159). In spinal surgery, lidocaine bolus of 1 mg/kg followed by 1.5 mg/kg/h infusion showed a slight improvement in MMSE scores (160). Currently, there are two registered randomized controlled trials investigating the use of local anesthetics in preventing POCD. One Phase 2 trial of 100 patients with supratentorial craniotomy tested the efficacy of lidocaine bolus 1.5 mg/kg and infusion 2 mg/kg/h after induction of surgery until anesthetic emergence on POCD (ClinicalTrials.gov identifier NCT00975910), although no results have been published. Similarly, a small Phase 2 trial of 70 patients (currently under recruitment) is testing the use of postoperative bupivacaine vs. morphine patient-controlled analgesia for 72 h following surgery on POCD (ClinicalTrials.gov identifier NCT02848599), although the primary cognitive endpoint is MMSE scores at postoperative day 5.

## Ketamine

Ketamine is an NMDA receptor antagonist with sedative, hypnotic, and analgesic properties; it is used as an anesthetic agent as well as an adjunct for neuropathic pain (191, 199). By virtue of its NMDA receptor antagonism, ketamine reduces glutamate transmission in the brain; coupled with its analgesic properties, ketamine has been proposed to reduce neuroinflammation (200). In pre-clinical models, ketamine

seems to have differential effects on the levels of pro-inflammatory cytokines (201, 202), however ketamine has been shown to attenuate cognitive impairment in rodents (202, 203). Human data are equally unclear: one small clinical trial ( $n = 60$ ) using a bolus of 0.5 mg/kg ketamine at the induction of cardiac surgery showed improved metrics of memory and executive function compared to control 1 week following surgery (161), however in a similar (but smaller) population, a 2.5 mg/kg ketamine bolus followed by 0.125 mg/kg infusion throughout the intraoperative period showed no change in POCD (measured by neurocognitive battery) compared to placebo at 1 or 10 weeks following surgery (162). It is unclear whether the discrepancies observed may be due to different dosing regimens, different cognitive assessments, or small sample size. There is currently a large ( $n = 900$ ) randomized Phase 3 clinical trial (ClinicalTrials.gov identifier NCT02892916) recruiting patients undergoing elective orthopedic surgeries to receive a 0.5 mg/kg ketamine bolus following anesthetic induction with POCD assessment as a secondary outcome (determined by MoCA score) at 1 week and 3 months following surgery. Results are not available at this time.

### Lipid Mediators (Resolvins, Lipoxins, Maresins)

As opposed to preventing the production of pro-inflammatory cytokines or oxidative species, lipid mediators such as resolvins, lipoxins, and maresins have begun to receive attention as possible resolvers of neuroinflammation (4, 129). In a rat model of CPB with deep hypothermic circulatory arrest, the resolution agonist annexin A1 was shown to (1) reduce systemic and neural pro-inflammatory cytokines due to inhibition of NF- $\kappa$ B, (2) inhibit microglial activation, and mitigate declines in Morris water maze performance at postoperative day 3 (204). Currently, no human trials exist on the role of these lipid mediators in POCD, although these agents may become more promising as more animal data become available.

### Cannabinoid Receptors

Cannabinoids are a variety of substances that can modulate neurotransmitter release *via* cannabinoid receptors and regulate a variety of physiological processes including appetite, mood, and pain (205). The most widely known cannabinoid is tetrahydrocannabinol (THC), the psychoactive ingredient in plants of the genus *Cannabis*. Cannabinoids are known to suppress TLR-mediated inflammatory responses, and immune cells themselves can produce endogenous cannabinoids, possibly representing homeostatic mechanisms (206). In mice, the activation of cannabinoid receptor 2 (CR2) was shown to attenuate hippocampal memory impairment (*via* fear conditioning test) and decrease pro-inflammatory cytokines in the hippocampus and prefrontal cortex at 1, 3, and 7 days following tibial fracture surgery (207). Due to the controlled nature of exogenous and synthetic cannabinoids, there are no human data on the effects of cannabinoids on POCD, although this may represent a new area of study as cannabinoids are beginning to be used as therapy for a range of disorders including depression, anorexia, epilepsy, and multiple sclerosis (208, 209).

### Melatonin

Melatonin is an endogenous hormone synthesized from L-tryptophan and secreted from the pineal gland. Its production is inhibited by 460–480 nm light in the blue portion of the electromagnetic spectrum and functions in maintaining circadian rhythms (210). Melatonin is also known to modulate production of pro- and anti-inflammatory cytokines and reduce cell adhesion molecules, and scavenge free radicals (211). In rodents, exogenous melatonin attenuates volatile (isoflurane)-induced memory impairment in adult and aged animals (212–214); this effect appears to result from improvements in the sleep-wake cycle (213, 214). Results from two published trials in human subjects offer no insight as to the efficacy of melatonin for the prevention of POCD. In the first instance, 54 women aged 30–75 years undergoing surgery for breast cancer were given 6 mg/day melatonin vs. placebo for 3 months beginning preoperatively again improved sleep-efficiency but without a discernable effect on POCD as measured using the ISPOCD test battery (163). In a more age-appropriate cohort of patients scheduled for hip arthroplasty (age > 65 years;  $n = 139$ ), melatonin (1 mg/day taken orally beginning the day before surgery and continued for 5 days consecutively postoperatively) again improved sleep quality and appeared to preserve basic aspects of cognition as measured by the MMSE in the immediate (within 7 days) postoperative period (164); however, a lack of more appropriate neurocognitive assessments over a more extended time frame, preclude supporting melatonin as prophylaxis against POCD. There are no registered clinical trials currently investigating the use of melatonin in POCD.

### Turmeric

Turmeric is a plant of the ginger family whose roots are boiled and ground for coloring and flavoring in many Eastern cultures. As such, it is comprised of many biological compounds with varying concentrations depending on the manufacturing method. One compound, curcumin, has been shown to have antioxidant and anti-inflammatory properties, possibly by inhibition of NF- $\kappa$ B (215). In “aged” male ICR mice (age 12 months) who underwent midline laparotomy, curcumin attenuated surgery-induced impairment in novel object recognition as well as spatial learning and memory (216); here, the anesthetic consisted solely of a neuroleptic anesthetic using fentanyl plus droperidol, so the relevance to current clinical practice is unclear. There are no published or open registered clinical trials currently investigating the use of turmeric or curcumin in POCD.

### Acupuncture

Acupuncture is a well-known therapy in alternative medicine, having been developed more than 3,000 years ago in China. It is gaining popularity in the Western world and is being tested as a treatment for a variety of inflammatory disorders including asthma, carpal tunnel syndrome, and fibromyalgia (217). While little is known about acupuncture and POCD, recent evidence suggests that electroacupuncture increases hippocampal expression of  $\alpha 7$  nAChRs, downregulates TNF $\alpha$  and IL-1 $\beta$  expression in hippocampal neurons, and can improve

spatial memory at 1, 3, and 7 days following partial hepatectomy in rats (218). While there are no animal studies on acupuncture and POCD, three human studies (published in Chinese) were identified (165–167); sample sizes were 120, 124, and 83 subjects, respectively. Although subjects were randomized, the reported methods in each report raise enough concern as to render the validity of the data uncertain, thereby precluding a clear assessment as to the efficacy of the technique. There are no registered clinical trials currently investigating the use of acupuncture in POCD.

## CONCLUSION

POCD is a widespread phenomenon following the surgical experience and can have detrimental effects on an individual's functional status and quality of life. People with preexisting neurocognitive impairments seem to be exceptionally prone to developing POCD, and POCD may unmask such impairments even in the absence of clinical detection. A large and growing body of evidence from pre-clinical and clinical studies has implicated the roles of neuroinflammation in the pathogenesis of POCD, from peripheral injury to neuronal death and functional manifestations. However, the data are not entirely conclusive because of heterogeneities in animal models and human populations studied, as well as variability in pre-clinical and clinical assessments of POCD. While both animal and human studies demonstrate a variety of neuroinflammatory mechanisms at play in the perioperative period, the root causes of that inflammation, whether surgery, anesthesia, or even prior inflammation from sources such as infection are unknown. Data from randomized clinical trials seem to more strongly favor surgery as the main inciting factor of POCD, but again these data are not wholly consistent across populations, surgeries, and time scales. An alternative hypothesis is that the combination of surgery and anesthesia contributes to the pathogenesis of POCD: anesthesia may weaken the BBB by modulating tight junction protein expression (219) in a dose-dependent manner (220), while surgery provides the peripheral nidus for inflammation that is ultimately amplified in the CNS. Whatever the cause,

neuroinflammation has been shown to be a common feature underlying many chronic and neurodegenerative diseases; a better understanding of such mechanisms may aid in improved diagnosis and treatment of a family of neurocognitive disorders.

The neuroinflammatory hypothesis has already generated a variety of potential candidates for treatment of POCD. The utility of many of these proposed treatment options have shown promising results in animal studies, however when applied to human populations, the treatment options yield more modest results. At this time, the lack of a formal definition of POCD is a critical barrier to future research; without a formal definition, the results of any one study may not be applicable to any other population than the one tested. Moreover, without a formal definition our understanding of the pathogenesis of POCD lacks generalizability to other neurodegenerative disorders that share common cellular mechanisms and clinical features. Only by standardizing our metrics and timepoints of POCD assessment will we be able to better understand the true incidence of POCD, compare the contributions of potential risk factors, and evaluate treatments across a large patient cohort (49, 221). Nevertheless, the sheer number of proposed treatments is suggestive of a growing interest in understanding POCD, and will hopefully benefit patients *via* a diverse array of therapies.

## AUTHOR CONTRIBUTIONS

SAS drafted the manuscript. SAS and PAG contributed to the literature review, manuscript revision, and read and approved the submitted manuscript.

## FUNDING

This work was supported by the Department of Anesthesiology, Weill Cornell Medical College.

## ACKNOWLEDGMENTS

We gratefully acknowledge Mo' Safavynia for all figure preparation associated with this manuscript.

## REFERENCES

- Edwards ML, Bause GS. From dental to mental institutions: an american dentist and a british psychiatrist highlight insanity following nitrous-oxide administration. *J Anesth Hist.* (2018) 4:133–4. doi: 10.1016/j.janh.2018.02.002
- Bedford PD. Adverse cerebral effects of anaesthesia on old people. *Lancet* (1955) 269:259–63.
- Evered L, Silbert B, Knopman D, Scott D, DeKosky S, Rasmussen L, et al. Recommendations for the nomenclature of cognitive change associated with anaesthesia and surgery. *Br J Anaesth.* (2018) 121:1005–12. doi: 10.1016/j.bja.2017.11.087
- Vacas S, Degos V, Feng X, Maze M. The neuroinflammatory response of postoperative cognitive decline. *Br Med Bull.* (2013) 106:161–78. doi: 10.1093/bmb/ldt006
- Berger M, Nadler JW, Browndyke J, Terrando N, Ponnusamy V, Cohen HJ, et al. Postoperative cognitive dysfunction: minding the gaps in our knowledge of a common postoperative complication in the elderly. *Anesthesiol Clin.* (2015) 33:517–50. doi: 10.1016/j.anclin.2015.05.008
- Androsova G, Krause R, Winterer G, Schneider R. Biomarkers of postoperative delirium and cognitive dysfunction. *Front Aging Neurosci.* (2015) 7:1347–16. doi: 10.3389/fnagi.2015.00112
- Evered L, Scott DA, Silbert B, Maruff P. Postoperative cognitive dysfunction is independent of type of surgery and anesthetic. *Anesth Analg.* (2011) 112:1179–85. doi: 10.1213/ANE.0b013e318215217e
- Moller JT, Cluitmans P, Rasmussen LS, Houx P, Rasmussen H, Canet J, et al. Long-term postoperative cognitive dysfunction in the elderly: ISPOCD1 study. *Lancet* (1998) 351:857–61. doi: 10.1016/S0140-6736(97)07382-0
- Silbert B, Evered L, Scott DA, McMahon S, Choong P, Ames D, et al. Preexisting cognitive impairment is associated with postoperative cognitive dysfunction after hip joint replacement surgery. *Anesthesiology* (2015) 122:1224–34. doi: 10.1097/ALN.0000000000000671
- Newman MF, Kirchner JL, Phillips-Bute B, Gaver V, Grocott H, Jones RH, et al. Longitudinal assessment of neurocognitive function after

- coronary-artery bypass surgery. *N Engl J Med.* (2001) 344:395–402. doi: 10.1056/NEJM200102083440601
11. Steinmetz J, Christensen KB, Lund T, Lohse N, Rasmussen LS, Group I. Long-term consequences of postoperative cognitive dysfunction. *Anesthesiology* (2009) 110:548–55. doi: 10.1097/ALN.0b013e318195b569
  12. Sprung J, Roberts RO, Knopman DS, Olive DM, Gappa JL, Sifuentes VL, et al. Association of mild cognitive impairment with exposure to general anesthesia for surgical and nonsurgical procedures: a population-based study. *Mayo Clin Proc.* (2016) 91:208–17. doi: 10.1016/j.mayocp.2015.10.023
  13. Nussmeier NA, Arlund C, Slogoff S. Neuropsychiatric complications after cardiopulmonary bypass: cerebral protection by a barbiturate. *Anesthesiology* (1986) 64:165–70.
  14. Monk TG, Weldon BC, Garvan CW, Dede DE, van der Aa MT, Heilman KM, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology* (2008) 108:18–30. doi: 10.1097/01.anes.0000296071.19434.1e
  15. Robinson TN, Walston JD, Brummel NE, Deiner S, Brown CH, Kennedy M, et al. Frailty for surgeons: review of a national institute on aging conference on frailty for specialists. *J Am Coll Surg.* (2015) 221:1083–92. doi: 10.1016/j.jamcollsurg.2015.08.428
  16. Hall DE, Arya S, Schmid KK, Blaser C, Carlson MA, Bailey TL, et al. Development and initial validation of the risk analysis index for measuring frailty in surgical populations. *JAMA Surg.* (2017) 152:175–82. doi: 10.1001/jamasurg.2016.4202
  17. Hall DE, Arya S, Schmid KK, Carlson MA, Lavedan P, Bailey TL, et al. Association of a frailty screening initiative with postoperative survival at 30:180, and 365 Days. *JAMA Surg.* (2017) 152:233–40. doi: 10.1001/jamasurg.2016.4219
  18. Esses G, Andreopoulos E, Lin HM, Arya S, Deiner S. A comparison of three frailty indices in predicting morbidity and mortality after on-pump aortic valve replacement. *Anesth Analg.* (2018) 126:39–45. doi: 10.1213/ANE.0000000000002411
  19. Moskven E, Bourassa-Moreau Md MFE, Charest-Morin R, Flexman A, and Street J. The impact of frailty and sarcopenia on postoperative outcomes in adult spine surgery. A systematic review of the literature. *Spine J.* (2018) 18:2354–69. doi: 10.1016/j.spinee.2018.07.008
  20. Rothrock RJ, Steinberger JM, Badgery H, Hecht AC, Cho SK, Caridi JM, et al. Frailty status as a predictor of 3-month cognitive and functional recovery following spinal surgery: a prospective pilot study. *Spine J.* (2019) 19:104–112. doi: 10.1016/j.spinee.2018.05.026
  21. Robinson TN, Wu DS, Pointer LF, Dunn CL, and Moss M. Preoperative cognitive dysfunction is related to adverse postoperative outcomes in the elderly. *J Am Coll Surg.* (2012) 215:12–7; discussion 17–18. doi: 10.1016/j.jamcollsurg.2012.02.007
  22. Culley DJ, Flaherty D, Fahey MC, Rudolph JL, Javedan H, Huang CC, et al. Poor performance on a preoperative cognitive screening test predicts postoperative complications in older orthopedic surgical patients. *Anesthesiology* (2017) 127:765–74. doi: 10.1097/ALN.0000000000001859
  23. Gasparini M, Vanacore N, Schiaffini C, Brusa L, Panella M, Talarico G, et al. A case-control study on Alzheimer's disease and exposure to anesthesia. *Neurol Sci.* (2002) 23:11–4. doi: 10.1007/s100720200017
  24. Horsburgh K, McCarron MO, White F, Nicoll JA. The role of apolipoprotein E in Alzheimer's disease, acute brain injury and cerebrovascular disease: evidence of common mechanisms and utility of animal models. *Neurobiol Aging* (2000) 21:245–55. doi: 10.1016/S0197-4580(00)00097-X
  25. Heyer EJ, Wilson DA, Sahlein DH, Mocco J, Williams SC, Sciacca R, et al. APOE-epsilon4 predisposes to cognitive dysfunction following uncomplicated carotid endarterectomy. *Neurology* (2005) 65:1759–63. doi: 10.1212/01.wnl.0000184579.23624.6b
  26. Avidan MS, Searleman AC, Storandt M, Barnett K, Vannucci A, Saager L, et al. Long-term cognitive decline in older subjects was not attributable to noncardiac surgery or major illness. *Anesthesiology* (2009) 111:964–70. doi: 10.1097/ALN.0b013e3181bc9719
  27. Berger M, Nadler JW, Friedman A, McDonagh DL, Bennett ER, Cooter M, et al. The effect of propofol versus isoflurane anesthesia on human cerebrospinal fluid markers of alzheimer's disease: results of a randomized trial. *J Alzheimers Dis.* (2016) 52:1299–310. doi: 10.3233/JAD-151190
  28. Whittington RA, Bretteville A, Dickler MF, Planel E. Anesthesia and tau pathology. *Prog Neuropsychopharmacol Biol Psychiatry* (2013) 47:147–55. doi: 10.1016/j.pnpbp.2013.03.004
  29. Whittington RA, Bretteville A, Virag L, Emala CW, Maurin TO, Marcouiller F, et al. Anesthesia-induced hypothermia mediates decreased ARC gene and protein expression through ERK/MAPK inactivation. *Sci Rep.* (2013) 3:1388. doi: 10.1038/srep01388
  30. Whittington RA, Virag L, Gratuze M, Petry FR, Noel A, Poitras I, et al. Dexmedetomidine increases tau phosphorylation under normothermic conditions *in vivo* and *in vitro*. *Neurobiol Aging* (2015) 36:2414–28. doi: 10.1016/j.neurobiolaging.2015.05.002
  31. Hudetz JA, Iqbal Z, Gandhi SD, Patterson KM, Byrne AJ, Pagel PS. Postoperative delirium and short-term cognitive dysfunction occur more frequently in patients undergoing valve surgery with or without coronary artery bypass graft surgery compared with coronary artery bypass graft surgery alone: results of a pilot study. *J Cardiothorac Vasc Anesth.* (2011) 25:811–6. doi: 10.1053/j.jvca.2010.05.003
  32. Hudetz JA, Patterson KM, Iqbal Z, Gandhi SD, Pagel PS. Metabolic syndrome exacerbates short-term postoperative cognitive dysfunction in patients undergoing cardiac surgery: results of a pilot study. *J Cardiothorac Vasc Anesth.* (2011) 25:282–7. doi: 10.1053/j.jvca.2010.06.008
  33. Dallmeier D, Larson MG, Vasani RS, Keaney JF, Fontes JD, Meigs JB, et al. Metabolic syndrome and inflammatory biomarkers: a community-based cross-sectional study at the Framingham Heart Study. *Diabetol Metabol Syndr.* (2012) 4:28. doi: 10.1186/1758-5996-4-28
  34. Saxena S, Maze M. Impact on the brain of the inflammatory response to surgery. *La Presse Med.* (2018) 47:e73–e81. doi: 10.1016/j.lpm.2018.03.011
  35. Terrando N, Monaco C, Ma D, Foxwell BMJ, Feldmann M, Maze M. Tumor necrosis factor-alpha triggers a cytokine cascade yielding postoperative cognitive decline. *Proc Natl Acad Sci USA.* (2010) 107:20518–22. doi: 10.1073/pnas.1014557107
  36. Fidalgo AR, Cibelli M, White JPM, Nagy I, Maze M, Ma D. Systemic inflammation enhances surgery-induced cognitive dysfunction in mice. *Neurosci. Lett.* (2011) 498:63–6. doi: 10.1016/j.neulet.2011.04.063
  37. Hovens IB, van Leeuwen BL, Nyakas C, Heineman E, van der Zee EA, Schoemaker RG. Prior infection exacerbates postoperative cognitive dysfunction in aged rats. *Am J Physiol. Regul Integr. Compar Physiol.* (2015) 309:R148–59. doi: 10.1152/ajpregu.00002.2015
  38. Wang P, Cao J, Liu N, Ma L, Zhou X, Zhang H, et al. Protective effects of edaravone in adult rats with surgery and lipopolysaccharide administration-induced cognitive function impairment. *PLoS ONE* (2016) 11:e0153708. doi: 10.1371/journal.pone.0153708
  39. Buvaendran A, Kroin JS, Berger RA, Hallab NJ, Saha C, Negrescu C, et al. Upregulation of prostaglandin E2 and interleukins in the central nervous system and peripheral tissue during and after surgery in humans. *Anesthesiology* (2006) 104:403–10.
  40. van Harten AE, Scheeren TWL, Absalom AR. A review of postoperative cognitive dysfunction and neuroinflammation associated with cardiac surgery and anaesthesia. *Anaesthesia* (2012) 67:280–93. doi: 10.1111/j.1365-2044.2011.07008.x
  41. Ji M-H, Yuan H-M, Zhang G-F, Li X-M, Dong L, Li W-Y, et al. Changes in plasma and cerebrospinal fluid biomarkers in aged patients with early postoperative cognitive dysfunction following total hip-replacement surgery. *J Anesth.* (2013) 27:236–42. doi: 10.1007/s00540-012-1506-3
  42. Hirsch J, Vacas S, Terrando N, Yuan M, Sands LP, Kramer J, et al. Perioperative cerebrospinal fluid and plasma inflammatory markers after orthopedic surgery. *J Neuroinflammat.* (2016) 13:211. doi: 10.1186/s12974-016-0681-9
  43. Peng L, Xu L, Ouyang W. Role of peripheral inflammatory markers in postoperative cognitive dysfunction (POCD): a meta-analysis. *PLoS ONE* (2013) 8:e79624–e79610. doi: 10.1371/journal.pone.0079624
  44. Liu X, Yu Y, Zhu S. Inflammatory markers in postoperative delirium (POD) and cognitive dysfunction (POCD): a meta-analysis of observational studies. *PLoS One* (2018) 13:e0195659. doi: 10.1371/journal.pone.0195659
  45. Goto T, Maekawa K. Cerebral dysfunction after coronary artery bypass surgery. *J Anesth.* (2014) 28:242–8. doi: 10.1007/s00540-013-1699-0
  46. Öztürk S, and Saçar M, Baltalarlı A, and Öztürk I. (2016). Effect of the type of cardiopulmonary bypass pump flow on postoperative cognitive function



- in patients undergoing isolated coronary artery surgery. *Anatol J Cardiol*. 16:875–80. doi: 10.14744/AnatolJCardiol.2015.6572
47. Rasmussen LS, Larsen K, Houx P, Skovgaard LT, Hanning CD, Moller JT, et al. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand*. (2001) 45:275–89. doi: 10.1034/j.1399-6576.2001.045003275.x
  48. Evered L, Scott DA, Silbert B. Cognitive decline associated with anesthesia and surgery in the elderly. *Curr Opin Psychiatry* (2017) 30:220–6. doi: 10.1097/YCO.0000000000000321
  49. Hovens IB, Schoemaker RG, van der Zee EA, Heineman E, Izaks GJ, van Leeuwen BL. Thinking through postoperative cognitive dysfunction: how to bridge the gap between clinical and pre-clinical perspectives. *Brain Behav Immunity* (2012) 26:1169–79. doi: 10.1016/j.bbi.2012.06.004
  50. APA. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*, *American Psychiatric Pub*. Arlington, VA: American Psychiatric Association, (2013).
  51. Berger M, Terrando N, Smith SK, Browndyke JN, Newman MF, Mathew JP. Neurocognitive function after cardiac surgery: from phenotypes to mechanisms. *Anesthesiology* (2018) 129:829–51. doi: 10.1097/ALN.0000000000002194
  52. Evered LA, Silbert BS. Postoperative cognitive dysfunction and noncardiac surgery. *Anesth Analg*. (2018) 127:496–505. doi: 10.1213/ANE.0000000000003514
  53. Maldonado JR. Neuropathogenesis of delirium: review of current etiologic theories and common pathways. *Am J Geriatr Psychiatry* (2013) 21:1190–222. doi: 10.1016/j.jagp.2013.09.005
  54. Zhang Q, Raoof M, Chen Y, Sumi Y, Sursal T, Junger W, et al. Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* (2010) 464:104–7. doi: 10.1038/nature08780
  55. Lotze MT, Tracey KJ. High-mobility group box 1 protein (HMGB1): nuclear weapon in the immune arsenal. *Nat Rev Immunol*. (2005) 5:331–42. doi: 10.1038/nri1594
  56. He H-J, Wang Y, Le Y, Duan K-M, Yan X-B, Liao Q, et al. Surgery upregulates high mobility group box-1 and disrupts the blood-brain barrier causing cognitive dysfunction in aged rats. *CNS Neurosci Therapeut*. (2012) 18:994–1002. doi: 10.1111/cns.12018
  57. Skvarc DR, Berk M, Byrne LK, Dean OM, Dodd S, Lewis M, et al. Post-operative cognitive dysfunction\_ an exploration of the inflammatory hypothesis and novel therapies. *Neurosci Biobehav Rev*. (2018) 84:116–33. doi: 10.1016/j.neubiorev.2017.11.011
  58. Li R-L, Zhang Z-Z, Peng M, Wu Y, Zhang J-J, Wang C-Y, et al. Postoperative impairment of cognitive function in old mice: a possible role for neuroinflammation mediated by HMGB1, S100B, and RAGE. *J Surg Res*. (2013) 185:815–24. doi: 10.1016/j.jss.2013.06.043
  59. Terrando N, Yang T, Wang X, Fang J, Cao M, Andersson U, et al. Systemic HMGB1 neutralization prevents postoperative neurocognitive dysfunction in aged rats. *Front Immunol*. (2016) 7:441. doi: 10.3389/fimmu.2016.00441
  60. Lin G-X, Wang T, Chen M-H, Hu Z-H, Ouyang W. Serum high-mobility group box 1 protein correlates with cognitive decline after gastrointestinal surgery. *Acta Anaesthesiol. Scand*. (2014) 58:668–74. doi: 10.1111/aas.12320
  61. Hem S, Albite R, Loresi M, Rasmussen J, Ajler P, Yampolsky C, et al. Pathological changes of the hippocampus and cognitive dysfunction following frontal lobe surgery in a rat model. *Acta Neurochirurg*. (2016) 158:2163–71. doi: 10.1007/s00701-016-2938-6
  62. Li Z, Ni C, Xia C, Jaw J, Wang Y, Cao Y, et al. Calcineurin/nuclear factor- $\kappa$ B signaling mediates isoflurane-induced hippocampal neuroinflammation and subsequent cognitive impairment in aged rats. *Mol Med Rep*. (2017) 15:201–9. doi: 10.3892/mmr.2016.5967
  63. Zhang J, Jiang W, Zuo Z. Pyrrolidine dithiocarbamate attenuates surgery-induced neuroinflammation and cognitive dysfunction possibly via inhibition of nuclear factor  $\kappa$ B. *Neuroscience* (2014) 261:1–10. doi: 10.1016/j.neuroscience.2013.12.034
  64. Ma Y, Cheng Q, Wang E, Li L, Zhang X. Inhibiting tumor necrosis factor- $\alpha$  signaling attenuates postoperative cognitive dysfunction in aged rats. *Mol Med Rep*. (2015) 12:3095–100. doi: 10.3892/mmr.2015.3744
  65. Cheon SY, Kim JM, Kam EH, Ho C-C, Kim EJ, Chung S, et al. Cell-penetrating interactive inhibition of nuclear factor- $\kappa$ B in a mouse model of postoperative cognitive dysfunction. *Sci Rep* (2017) 7:13482. doi: 10.1038/s41598-017-14027-2
  66. Rempe RG, Hartz AM, Bauer B. Matrix metalloproteinases in the brain and blood-brain barrier: versatile breakers and makers. *J Cereb Blood Flow Metabol*. (2016) 36:1481–507. doi: 10.1177/0271678X16655551
  67. Zlokovic BV. The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron* (2008) 57:178–201. doi: 10.1016/j.neuron.2008.01.003
  68. Engblom D, Ek M, Saha S, Ericsson-Dahlstrand A, Jakobsson P-J, Blomqvist A. Prostaglandins as inflammatory messengers across the blood-brain barrier. *J Mol Med*. (2002) 80:5–15. doi: 10.1007/s00109-001-0289-z
  69. Terrando N, Eriksson LI, Kyu Ryu J, Yang T, Monaco C, Feldmann M, et al. Resolving postoperative neuroinflammation and cognitive decline. *Ann Neurol*. (2011) 70:986–95. doi: 10.1002/ana.22664
  70. Cao X-Z, Ma H, Wang J-K, Liu F, Wu B-Y, Tian A-Y, et al. Postoperative cognitive deficits and neuroinflammation in the hippocampus triggered by surgical trauma are exacerbated in aged rats. *Progr Neuropsychopharmacol Biol Psychiatry* (2010) 34:1426–32. doi: 10.1016/j.pnpbp.2010.07.027
  71. Xu Z, Dong Y, Wang H, Culley DJ, Marcantonio ER, Crosby G, et al. Peripheral surgical wounding and age-dependent neuroinflammation in mice. *PLoS ONE* (2014) 9:e96752. doi: 10.1371/journal.pone.0096752
  72. Reinsfelt B, Ricksten S-E, Zetterberg H, Blennow K, Fredén-Lindqvist J, Westerlind A. Cerebrospinal fluid markers of brain injury, inflammation, and blood-brain barrier dysfunction in cardiac surgery. *Ann Thorac Surg*. (2012) 94:549–55. doi: 10.1016/j.athoracsur.2012.04.004
  73. Cibelli M, Fidalgo AR, Terrando N, Ma D, Monaco C, Feldmann M, et al. Role of interleukin-1beta in postoperative cognitive dysfunction. *Ann Neurol*. (2010) 68:360–8. doi: 10.1002/ana.22082
  74. Zhang J, Tan H, Jiang W, Zuo Z. Amantadine alleviates postoperative cognitive dysfunction possibly by increasing glial cell line-derived neurotrophic factor in rats. *Anesthesiology* (2014) 121:773–85. doi: 10.1097/ALN.0000000000000352
  75. Bayram H, Hidiröglu M, Cetin L, Kucuker A, Iriz E, Uguz E, et al. Comparing S-100 beta protein levels and neurocognitive functions between patients undergoing on-pump and off-pump coronary artery bypass grafting. *J Surg Res*. (2013) 182:198–202. doi: 10.1016/j.jss.2012.10.047
  76. Silva FP, Schmidt AP, Valentin LS, Pinto KO, Zeferino SP, Oses JP, et al. S100B protein and neuron-specific enolase as predictors of cognitive dysfunction after coronary artery bypass graft surgery. *Eur J Anaesthesiol*. (2016) 33:681–9. doi: 10.1097/EJA.0000000000000450
  77. Bi J, Shan W, Luo A, Zuo Z. Critical role of matrix metalloproteinase 9 in postoperative cognitive dysfunction and age-dependent cognitive decline. *Oncotarget* (2017) 8:51817–29. doi: 10.18632/oncotarget.15545
  78. Dong H, Zhang X, Qian Y. Mast cells and neuroinflammation. *Med Sci Monit Basic Res*. (2014) 20:200–6. doi: 10.12659/MSMBR.893093
  79. Wohleb ES, McKim DB, Sheridan JF, Godbout JP. Monocyte trafficking to the brain with stress and inflammation: a novel axis of immune-to-brain communication that influences mood and behavior. *Front Neurosci*. (2014) 8:447. doi: 10.3389/fnins.2014.00447
  80. Degos V, Vacas S, Han Z, van Rooijen N, Gressens P, Su H, et al. Depletion of bone marrow-derived macrophages perturbs the innate immune response to surgery and reduces postoperative memory dysfunction. *Anesthesiology* (2013) 118:527–36. doi: 10.1097/ALN.0b013e3182834d94
  81. Frost JL, Schafer DP. Microglia: architects of the developing nervous system. *Trends Cell Biol*. (2016) 26:587–97. doi: 10.1016/j.tcb.2016.02.006
  82. Cunningham CL, Martínez-Cerdeño V, Noctor SC. Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *J Neurosci*. (2013) 33:4216–33. doi: 10.1523/JNEUROSCI.3441-12.2013
  83. Li Q, Barres BA. Microglia and macrophages in brain homeostasis and disease. *Nat Rev Immunol*. (2018) 18:225–42. doi: 10.1038/nri.2017.125
  84. Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma *in vivo*. *Science* (2005) 308:1314–8. doi: 10.1126/science.1110647
  85. Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, et al. Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity* (2013) 38:792–804. doi: 10.1016/j.immuni.2013.04.004
  86. Biber K, Neumann H, Inoue K, Boddeke HWGM. Neuronal 'On' and 'Off' signals control microglia. *Trends Neurosci*. (2007) 30:596–602. doi: 10.1016/j.tins.2007.08.007

87. Szalay G, Martinez B, Lénárt N, Környei Z, Orsolits B, Judák L, et al. Microglia protect against brain injury and their selective elimination dysregulates neuronal network activity after stroke. *Nat Commun.* (2016) 7:11499. doi: 10.1038/ncomms11499
88. Gordon S. Alternative activation of macrophages. *Nat Rev Immunol.* (2003) 3:23–35. doi: 10.1038/nri978
89. Thériault P, ElAli A, Rivest S. The dynamics of monocytes and microglia in Alzheimer's disease. *Alzheimer Res Ther.* (2015) 7:41. doi: 10.1186/s13195-015-0125-2
90. Martinez FO, Helming L, Gordon S. Alternative activation of macrophages: an immunologic functional perspective. *Ann Rev Immunol.* (2009) 27:451–83. doi: 10.1146/annurev.immunol.021908.132532
91. Machado-Pereira M, Santos T, Ferreira L, Bernardino L, Ferreira R. Anti-inflammatory strategy for M2 microglial polarization using retinoic acid-loaded nanoparticles. *Med Inflamm.* (2017) 2017:6742427–11. doi: 10.1155/2017/6742427
92. Tay TL, Savage JC, Hui CW, Bisht K, Tremblay M-È. Microglia across the lifespan: from origin to function in brain development, plasticity and cognition. *J Physiol.* (2017) 595:1929–45. doi: 10.1113/JP272134
93. Zhang X, Dong H, Li N, Zhang S, Sun J, Zhang S, et al. Activated brain mast cells contribute to postoperative cognitive dysfunction by evoking microglia activation and neuronal apoptosis. *J Neuroinflamm.* (2016) 13:127. doi: 10.1186/s12974-016-0592-9
94. Zhang S, Dong H, Zhang X, Li N, Sun J, and Qian Y. Cerebral mast cells contribute to postoperative cognitive dysfunction by promoting blood brain barrier disruption. *Behav Brain Res.* (2016) 298:158–66. doi: 10.1016/j.bbr.2015.11.003
95. Clausen BH, Lambertsens KL, Babcock AA, Holm TH, Dagnaes-Hansen F, Finsen B. Interleukin-1beta and tumor necrosis factor-alpha are expressed by different subsets of microglia and macrophages after ischemic stroke in mice. *J Neuroinflamm.* (2008) 5:46. doi: 10.1186/1742-2094-5-46
96. Henkel JS, Beers DR, Zhao W, Appel SH. Microglia in ALS: the good, the bad, and the resting. *J Neuroimmune Pharmacol.* (2009) 4:389–98. doi: 10.1007/s11481-009-9171-5
97. Terrando N, Rei Fidalgo A, Vizcaychipi M, Cibelli M, Ma D, Monaco C, et al. The impact of IL-1 modulation on the development of lipopolysaccharide-induced cognitive dysfunction. *Crit Care* (2010) 14:R88. doi: 10.1186/cc9019
98. Prinz M, Priller J. Tickets to the brain: role of CCR2 and CX3CR1 in myeloid cell entry in the CNS. *J Neuroimmunol.* (2010) 224:80–4. doi: 10.1016/j.jneuroim.2010.05.015
99. Lee JH, Kam EH, Kim SY, Cheon SY, Kim EJ, Chung S, et al. Erythropoietin attenuates postoperative cognitive dysfunction by shifting macrophage activation toward the M2 phenotype. *Front Pharmacol.* (2017) 8:839. doi: 10.3389/fphar.2017.00839
100. Xu J, Dong H, Qian Q, Zhang X, Wang Y, Jin W, et al. Astrocyte-derived CCL2 participates in surgery-induced cognitive dysfunction and neuroinflammation via evoking microglia activation. *Behav Brain Res.* (2017) 332:145–53. doi: 10.1016/j.bbr.2017.05.066
101. D'Mello C, Le T, Swain MG. Cerebral microglia recruit monocytes into the brain in response to tumor necrosis factoralpha signaling during peripheral organ inflammation. *J Neurosci.* (2009) 29:2089–102. doi: 10.1523/JNEUROSCI.3567-08.2009
102. Wang H-L, Ma R-H, Fang H, Xue Z-G, Liao Q-W. Impaired spatial learning memory after isoflurane anesthesia or appendectomy in aged mice is associated with microglia activation. *J Cell Death* (2015) 8:9–19. doi: 10.4137/JCD.S30596
103. Feng X, Valdearcos M, Uchida Y, Lutrin D, Maze M, Koliwad SK. Microglia mediate postoperative hippocampal inflammation and cognitive decline in mice. *JCI Insight* (2017) 2:e91229. doi: 10.1172/jci.insight.91229
104. Zhang T, Leng Z, Liu W, Wang X, Yan X, Yu L. Suppressed expression of mitogen-activated protein kinases in hyperthermia induced defective neural tube. *Neurosci Lett.* (2015) 594:6–11. doi: 10.1016/j.neulet.2015.03.046
105. Abdul-Muneer PM, Chandra N, Haorah J. Interactions of oxidative stress and neurovascular inflammation in the pathogenesis of traumatic brain injury. *Mol Neurobiol.* (2015) 51:966–79. doi: 10.1007/s12035-014-8752-3
106. Heneka MT, Kummer MP, Latz E. Innate immune activation in neurodegenerative disease. *Nat Rev Immunol.* (2014) 14:463–77. doi: 10.1038/nri3705
107. Gao H-M, Zhou H, Zhang F, Wilson BC, Kam W, Hong J-S. HMGB1 acts on microglia Mac1 to mediate chronic neuroinflammation that drives progressive neurodegeneration. *J Neurosci.* (2011) 31:1081–92. doi: 10.1523/JNEUROSCI.3732-10.2011
108. Adami C, Bianchi R, Pula G, Donato R. S100B-stimulated NO production by BV-2 microglia is independent of RAGE transducing activity but dependent on RAGE extracellular domain. *Biochim Biophys Acta* (2004) 1742:169–77. doi: 10.1016/j.bbamcr.2004.09.008
109. Netto MB, de Oliveira Junior AN, Goldim M, Mathias K, Fileti ME, da Rosa N, et al. Oxidative stress and mitochondrial dysfunction contributes to postoperative cognitive dysfunction in elderly rats. *Brain Behav Immunity* (2018) 73:661–9. doi: 10.1016/j.bbi.2018.07.016
110. Won SJ, Yoo BH, Kauppinen TM, Choi BY, Kim JH, Jang BG, et al. Recurrent/moderate hypoglycemia induces hippocampal dendritic injury, microglial activation, and cognitive impairment in diabetic rats. *J Neuroinflamm.* (2012) 9:182. doi: 10.1186/1742-2094-9-182
111. Harmon D, Eustace N, Ghori K, Butler M, O'Callaghan S, O'Donnell A, et al. Plasma concentrations of nitric oxide products and cognitive dysfunction following coronary artery bypass surgery. *Eur J Anaesthesiol.* (2005) 22:269–76.
112. Nicoll RA. A brief history of long-term potentiation. *Neuron* (2017) 93:281–90. doi: 10.1016/j.neuron.2016.12.015
113. Segal M. Dendritic spines and long-term plasticity. *Nat Rev Neurosci.* (2005) 6:277–84. doi: 10.1038/nrn1649
114. Rothwell NJ, Luheshi G, Toulmond S. Cytokines and their receptors in the central nervous system: physiology, pharmacology, and pathology. *Pharmacol Ther.* (1996) 69:85–95.
115. Rachal Pugh C, Fleschner M, Watkins LR, Maier SF, Rudy JW. The immune system and memory consolidation: a role for the cytokine IL-1beta. *Neurosci Biobehav Rev.* (2001) 25:29–41.
116. Riazi K, Galic MA, Kentner AC, Reid AY, Sharkey KA, Pittman QJ. Microglia-dependent alteration of glutamatergic synaptic transmission and plasticity in the hippocampus during peripheral inflammation. *J Neurosci.* (2015) 35:4942–52. doi: 10.1523/JNEUROSCI.4485-14.2015
117. Pedrazzi M, Averna M, Sparatore B, Patrone M, Salamino F, Marcoli M, et al. Potentiation of NMDA receptor-dependent cell responses by extracellular high mobility group box 1 protein. *PLoS ONE* (2012) 7:e44518. doi: 10.1371/journal.pone.0044518
118. Pribrag H, Stellwagen D. TNF-alpha downregulates inhibitory neurotransmission through protein phosphatase 1-dependent trafficking of GABA(A) receptors. *J Neurosci.* (2013) 33:15879–93. doi: 10.1523/JNEUROSCI.0530-13.2013
119. Evonuk KS, Baker BJ, Doyle RE, Moseley CE, Sestero CM, Johnston BP, et al. Inhibition of system Xc(-) transporter attenuates autoimmune inflammatory demyelination. *J Immunol.* (2015) 195:450–63. doi: 10.4049/jimmunol.1401108
120. Wu H, Li L, Su X. Vagus nerve through alpha7 nAChR modulates lung infection and inflammation: models, cells, and signals. *BioMed. Res. Int.* (2014) 2014:283525. doi: 10.1155/2014/283525
121. Rosas-Ballina M, Ochani M, Parrish WR, Ochani K, Harris YT, Huston JM, et al. Splenic nerve is required for cholinergic antiinflammatory pathway control of TNF in endotoxemia. *Proc Natl Acad Sci USA.* (2008) 105:11008–13. doi: 10.1073/pnas.0803237105
122. Sanders VM. The beta2-adrenergic receptor on T and B lymphocytes: do we understand it yet? *Brain Behav Immun.* (2012) 26:195–200. doi: 10.1016/j.bbi.2011.08.001
123. Ghia J-E, Blennerhassett P, El-Sharkawy RT, Collins SM. The protective effect of the vagus nerve in a murine model of chronic relapsing colitis. *Am J Physiol Gastrointest Liver Physiol.* (2007) 293:G711–718. doi: 10.1152/ajpgi.00240.2007
124. O'Mahony C, van der Kleij H, Bienenstock J, Shanahan F, O'Mahony L. Loss of vagal anti-inflammatory effect: *in vivo* visualization and adoptive transfer. *Am J Physiol Regul Integr Compar Physiol.* (2009) 297, R1118–26. doi: 10.1152/ajpregu.90904.2008
125. Kalb A, von Haefen C, Sifringer M, Tegethoff A, Paeschke N, Kostova M, et al. Acetylcholinesterase inhibitors reduce neuroinflammation and -degeneration in the cortex and hippocampus of a surgery stress rat model. *PLoS ONE* (2013) 8:e62679. doi: 10.1371/journal.pone.0062679

126. Kashyap M, Belleville S, Mulsant BH, Hilmer SN, Paquette A, Tu LM, et al. Methodological challenges in determining longitudinal associations between anticholinergic drug use and incident cognitive decline. *J Am Geriatr Soc.* (2014) 62:336–41. doi: 10.1111/jgs.12632
127. Xiong J, Xue FS, Liu JH, Xu YC, Liao X, Zhang YM, et al. Transcutaneous vagus nerve stimulation may attenuate postoperative cognitive dysfunction in elderly patients. *Med Hypotheses* (2009) 73:938–41. doi: 10.1016/j.mehy.2009.06.033
128. Serhan CN. Treating inflammation and infection in the 21st century: new hints from decoding resolution mediators and mechanisms. *FASEB J.* (2017) 31:1273–88. doi: 10.1096/fj.201601222R
129. Terrando N, Gómez-Galán M, Yang T, Carlström M, Gustavsson D, Harding RE, et al. Aspirin-triggered resolvin D1 prevents surgery-induced cognitive decline. *FASEB J.* (2013) 27:3564–71. doi: 10.1096/fj.13-230276
130. Serhan CN, Yang R, Martinod K, Kasuga K, Pillai PS, Porter TF, et al. Maresins: novel macrophage mediators with potent antiinflammatory and proresolving actions. *J Exp Med.* (2009) 206:15–23. doi: 10.1084/jem.20081880
131. Zhang J, Tan H, Jiang W, Zuo Z. The choice of general anesthetics may not affect neuroinflammation and impairment of learning and memory after surgery in elderly rats. *J Neuroimm Pharmacol.* (2015) 10:179–89. doi: 10.1007/s11481-014-9580-y
132. Jevtic-Todorovic V, Hartman RE, Izumi Y, Benschoff ND, Dikranian K, Zorumski CF, et al. Early exposure to common anesthetic agents causes widespread neurodegeneration in the developing rat brain and persistent learning deficits. *J Neurosci.* (2003) 23:876–82. doi: 10.1523/JNEUROSCI.23-03-00876.2003
133. Shen X, Dong Y, Xu Z, Wang H, Miao C, Soriano SG, et al. Selective anesthesia-induced neuroinflammation in developing mouse brain and cognitive impairment. *Anesthesiology* (2013) 118:502–15. doi: 10.1097/ALN.0b013e3182834d77
134. Coleman K, Robertson ND, Dissen GA, Neuringer MD, Martin LD, Cuzon Carlson VC, et al. Isoflurane anesthesia has long-term consequences on motor and behavioral development in infant rhesus macaques. *Anesthesiology* (2017) 126:74–84. doi: 10.1097/ALN.0000000000001383
135. Wang Z, Meng S, Cao L, Chen Y, Zuo Z, Peng S. Critical role of NLRP3-caspase-1 pathway in age-dependent isoflurane-induced microglial inflammatory response and cognitive impairment. *J Neuroinflammat.* (2018) 15:109. doi: 10.1186/s12974-018-1137-1
136. Walters JL, Chelonis JJ, Fogle CM, Orser BA, Paule MG. Single and repeated exposures to the volatile anesthetic isoflurane do not impair operant performance in aged rats. *Neurotoxicology* (2016) 56:159–69. doi: 10.1016/j.neuro.2016.07.012
137. Zhu J, Jiang X, Shi E, Ma H, Wang J. Sevoflurane preconditioning reverses impairment of hippocampal long-term potentiation induced by myocardial ischaemia-reperfusion injury. *Eur J Anaesthesiol.* (2009) 26:961–8. doi: 10.1097/EJA.0b013e328330e968
138. Tasbihgou SR, Netkova M, Kalar AF, Doorduyn J, Struys MMRF, Schoemaker RG, et al. Brain changes due to hypoxia during light anaesthesia can be prevented by deepening anaesthesia; a study in rats. *PLoS ONE* (2018) 13:e0193062. doi: 10.1371/journal.pone.0193062
139. Qian X-L, Zhang W, Liu M-Z, Zhou Y-B, Zhang J-M, Han L, et al. Dexmedetomidine improves early postoperative cognitive dysfunction in aged mice. *Eur J Pharmacol.* (2015) 746:206–12. doi: 10.1016/j.ejphar.2014.11.017
140. Newman S, Stygall J, Hirani S, Shaefi S, Maze M. Postoperative cognitive dysfunction after noncardiac surgery: a systematic review. *Anesthesiology* (2007) 106:572–90.
141. Sanders RD, Maze M. Neuroinflammation and postoperative cognitive dysfunction: can anaesthesia be therapeutic? *Eur J Anaesthesiol.* (2010) 27:3–5. doi: 10.1097/EJA.0b013e3283318ef9
142. Silbert BS, Evered LA, Scott DA. Incidence of postoperative cognitive dysfunction after general or spinal anaesthesia for extracorporeal shock wave lithotripsy. *Br J Anaesth.* (2014) 113:784–91. doi: 10.1093/bja/aeu163
143. Silbert BS, Scott DA, Evered LA, Lewis MS, Kalpokas M, Maruff P, et al. A comparison of the effect of high- and low-dose fentanyl on the incidence of postoperative cognitive dysfunction after coronary artery bypass surgery in the elderly. *Anesthesiology* (2006) 104:1137–45.
144. Geng Y-J, Wu Q-H, Zhang R-Q. Effect of propofol, sevoflurane, and isoflurane on postoperative cognitive dysfunction following laparoscopic cholecystectomy in elderly patients: a randomized controlled trial. *J Clin Anesth.* (2017) 38:165–71. doi: 10.1016/j.jclinane.2017.02.007
145. Qiao Y, Feng H, Zhao T, Yan H, Zhang H, Zhao X. Postoperative cognitive dysfunction after inhalational anesthesia in elderly patients undergoing major surgery: the influence of anesthetic technique, cerebral injury and systemic inflammation. *BMC Anesthesiol.* (2015) 15:154. doi: 10.1186/s12871-015-0130-9
146. Shu A-H, Wang Q, Chen X-B. Effect of different depths of anesthesia on postoperative cognitive function in laparoscopic patients: a randomized clinical trial. *Curr Med Res Opin.* (2015) 31:1883–7. doi: 10.1185/03007995.2015.1075968
147. Hou R, Wang H, Chen L, Qiu Y, Li S. POCD in patients receiving total knee replacement under deep vs. light anesthesia: a randomized controlled trial. *Brain Behav.* (2018) 8:e00910. doi: 10.1002/brb3.910
148. Erdogan MA, Demirbilek S, Erdil F, Aydogan MS, Ozturk E, Tugal T, et al. The effects of cognitive impairment on anaesthetic requirement in the elderly. *Eur J Anaesthesiol.* (2012) 29:326–31. doi: 10.1097/EJA.0b013e32835475c6
149. Babiloni C, Carducci F, Lizio R, Vecchio F, Baglieri A, Bernardini S, et al. Resting state cortical electroencephalographic rhythms are related to gray matter volume in subjects with mild cognitive impairment and Alzheimer's disease. *Hum Brain Mapp.* (2013) 34:1427–46. doi: 10.1002/hbm.22005
150. Schuller PJ, Newell S, Strickland PA, Barry JJ. Response of bispectral index to neuromuscular block in awake volunteers. *Br J Anaesth.* (2015) 115 Suppl 1:i95–103. doi: 10.1093/bja/aev072
151. Avidan MS, Graetz TJ. Monitoring the brain strikes a discordant note for anesthesiologists. *Can J Anaesth.* (2018) 65:501–6. doi: 10.1007/s12630-018-1086-2
152. Zhu Y-Z, Yao R, Zhang Z, Xu H, Wang L-W. Parecoxib prevents early postoperative cognitive dysfunction in elderly patients undergoing total knee arthroplasty: a double-blind, randomized clinical consortium study. *Medicine* (2016) 95:e4082. doi: 10.1097/MD.0000000000004082
153. Zhu Y, Yao R, Li Y, Wu C, Heng L, Zhou M, et al. Protective effect of celecoxib on early postoperative cognitive dysfunction in geriatric patients. *Front Neurol.* (2018) 9:633. doi: 10.3389/fneur.2018.00633
154. Ottens TH, Dieleman JM, Sauer AM, Peelen LM, Nierich AP, de Groot WJ, et al. Effects of dexamethasone on cognitive decline after cardiac surgery: a randomized clinical trial. *Anesthesiology* (2014) 121:492–500. doi: 10.1097/ALN.0000000000000336
155. Doraiswamy PM, Babyak MA, Hennig T, Trivedi R, White WD, Mathew JB, et al. Donepezil for cognitive decline following coronary artery bypass surgery: a pilot randomized controlled trial. *Psychopharmacol Bull.* (2007) 40:54–62.
156. Das S, Nanda SK, Bisoi AK, Wadhawan AN. Effect of preoperative statin therapy on early postoperative memory impairment after off-pump coronary artery bypass surgery. *Ann Card Anaesth.* (2016) 19:38–44. doi: 10.4103/0971-9784.173018
157. Li Y, He R, Chen S, Qu Y. Effect of dexmedetomidine on early postoperative cognitive dysfunction and peri-operative inflammation in elderly patients undergoing laparoscopic cholecystectomy. *Exp Therapeut Med.* (2015) 10:1635–42. doi: 10.3892/etm.2015.2726
158. Chen W, Liu B, Zhang F, Xue P, Cui R, Lei W. The effects of dexmedetomidine on post-operative cognitive dysfunction and inflammatory factors in senile patients. *Int J Clin Exp Med.* (2015) 8:4601–5.
159. Wang HL, Yan HD, Liu YY, Sun BZ, Huang R, Wang XS, et al. Intraoperative intravenous lidocaine exerts a protective effect on cell-mediated immunity in patients undergoing radical hysterectomy. *Mol Med Rep.* (2015) 12:7039–44. doi: 10.3892/mmr.2015.4235
160. Chen K, Wei P, Zheng Q, Zhou J, Li J. Neuroprotective effects of intravenous lidocaine on early postoperative cognitive dysfunction in elderly patients following spine surgery. *Med Sci Monit.* (2015) 21:1402–7. doi: 10.12659/MSM.894384
161. Hudetz JA, Iqbal Z, Gandhi SD, Patterson KM, Byrne AJ, Hudetz AG, et al. Ketamine attenuates post-operative cognitive dysfunction

- after cardiac surgery. *Acta Anaesthesiol Scand.* (2009) 53:864–72. doi: 10.1111/j.1399-6576.2009.01978.x
162. Nagels W, Demeyere R, Van Hemelrijck J, Vandenbussche E, Gijbels K, Vandermeersch E. Evaluation of the neuroprotective effects of S(+)-ketamine during open-heart surgery. *Anesth Analg.* (2004) 98:1595–603, table of contents. doi: 10.1213/01.ANE.0000117227.00820.0C
  163. Hansen MV, Madsen MT, Andersen LT, Hageman I, Rasmussen LS, Bokmand S, et al. Effect of melatonin on cognitive function and sleep in relation to breast cancer surgery: a randomized, double-blind, placebo-controlled trial. *Int J Breast Cancer* (2014) 2014:416531. doi: 10.1155/2014/416531
  164. Fan Y, Yuan L, Ji M, Yang J, Gao D. The effect of melatonin on early postoperative cognitive decline in elderly patients undergoing hip arthroplasty: a randomized controlled trial. *J Clin Anesth.* (2017) 39:77–81. doi: 10.1016/j.jclinane.2017.03.023
  165. Gao XQ, Zhang ZY, Ma WH. Effects of electroacupuncture assistant general anesthesia on postoperative cognitive dysfunction of aged patients. *Zhongguo Zhong Xi Yi Jie He Za Zhi* (2012) 32:591–3.
  166. Lin SY, Gao J, Yin ZL, Zhou LJ, Chen X. [Impacts of the different frequencies of electroacupuncture on cognitive function in patients after abdominal operation under compound anesthesia of acupuncture and drugs]. *Zhongguo Zhen Jiu* (2013) 33:1109–12.
  167. Lin SY, Yin ZL, Gao J, Zhou LJ, Chen X. [Effect of acupuncture-anesthetic composite anesthesia on the incidence of POC and TNF- $\alpha$ , IL-1 $\beta$ , IL-6 in elderly patients]. *Zhongguo Zhong Xi Yi Jie He Za Zhi* (2014) 34:795–9.
  168. Peng M, Wang Y-L, Wang F-F, Chen C, Wang C-Y. The cyclooxygenase-2 inhibitor parecoxib inhibits surgery-induced proinflammatory cytokine expression in the hippocampus in aged rats. *J Surg Res.* (2012) 178:e1–8. doi: 10.1016/j.jss.2012.08.030
  169. Kamer AR, Galoyan SM, Haile M, Kline R, Boutajangout A, Li Y-S, et al. Meloxicam improves object recognition memory and modulates glial activation after splenectomy in mice. *Eur J Anaesthesiol.* (2012) 29:332–7. doi: 10.1097/EJA.0b013e3283534f56
  170. Fan L, Wang T-L, Xu YC, Ma YH, Ye WG. Minocycline may be useful to prevent/treat postoperative cognitive decline in elderly patients. *Med Hypotheses* (2011) 76:733–6. doi: 10.1016/j.mehy.2011.02.010
  171. Wang H-L, Liu H, Xue Z-G, Liao Q-W, Fang H. Minocycline attenuates post-operative cognitive impairment in aged mice by inhibiting microglia activation. *J Cell Mol Med.* (2016) 20:1632–9. doi: 10.1111/jcmm.12854
  172. Li W, Chai Q, Zhang H, Ma J, Xu C, Dong J, et al. High doses of minocycline may induce delayed activation of microglia in aged rats and thus cannot prevent postoperative cognitive dysfunction. *J Int Med Res.* (2018) 46:1404–13. doi: 10.1177/0300060517754032
  173. Tsurufuji S, Kurihara A, Kiso S, Suzuki Y, Ohuchi K. Dexamethasone inhibits generation in inflammatory sites of the chemotactic activity attributable to leukotriene B<sub>4</sub>. *Biochem Biophys Res Commun.* (1984) 119:884–90.
  174. Barnes PJ. Anti-inflammatory actions of glucocorticoids: molecular mechanisms. *Clin Sci.* (1998) 94:557–72.
  175. Karaman T, Karaman S, Dogru S, Tapar H, Sahin A, Süren M. Short-term and long-term effects of dexamethasone on cognitive dysfunction induced by sevoflurane in adult rats. *Turk J Anaesthesiol Reanimat.* (2017) 45:158–63. doi: 10.5152/TJAR.2017.98624
  176. Han Z, Li L, Wang L, Degos V, Maze M, Su H. Alpha-7 nicotinic acetylcholine receptor agonist treatment reduces neuroinflammation, oxidative stress, and brain injury in mice with ischemic stroke and bone fracture. *J Neurochem.* (2014) 131:498–508. doi: 10.1111/jnc.12817
  177. Puhlinger FK, Rex C, Sielenkammer AW, Claudius C, Larsen PB, Prins ME, et al. Reversal of profound, high-dose rocuronium-induced neuromuscular blockade by sugammadex at two different time points: an international, multicenter, randomized, dose-finding, safety assessor-blinded, phase II trial. *Anesthesiology* (2008) 109:188–97. doi: 10.1097/ALN.0b013e31817f5bc7
  178. Barrientos RM, Hein AM, Frank MG, Watkins LR, Maier SF. Intracisternal interleukin-1 receptor antagonist prevents postoperative cognitive decline and neuroinflammatory response in aged rats. *J Neurosci.* (2012) 32:14641–8. doi: 10.1523/JNEUROSCI.2173-12.2012
  179. Hu J, Feng X, Valdearcos M, Lutrin D, Uchida Y, Koliwad SK, et al. Interleukin-6 is both necessary and sufficient to produce perioperative neurocognitive disorder in mice. *Br J Anaesth.* (2018) 120:537–45. doi: 10.1016/j.bja.2017.11.096
  180. Cerqueira NM, Oliveira EF, Gesto DS, Santos-Martins D, Moreira C, Moorthy HN, et al. Cholesterol biosynthesis: a mechanistic overview. *Biochemistry* (2016) 55:5483–506. doi: 10.1021/acs.biochem.6b00342
  181. Margaritis M, Channon KM, Antoniadis C. Statins as regulators of redox state in the vascular endothelium: beyond lipid lowering. *Antioxid Redox Signal.* (2014) 20:1198–215. doi: 10.1089/ars.2013.5430
  182. Swiger KJ, Manalac RJ, Blumenthal RS, Blaha MJ, Martin SS. Statins and cognition: a systematic review and meta-analysis of short- and long-term cognitive effects. *Mayo Clin Proc.* (2013) 88:1213–21. doi: 10.1016/j.mayocp.2013.07.013
  183. Katznelson R, Djaiani GN, Borger MA, Friedman Z, Abbey SE, Fedorko L, et al. Preoperative use of statins is associated with reduced early delirium rates after cardiac surgery. *Anesthesiology* (2009) 110:67–73. doi: 10.1097/ALN.0b013e318190b4d9
  184. Skvarc DR, Dean OM, Byrne LK, Gray L, Lane S, Lewis M, et al. The effect of N-acetylcysteine (NAC) on human cognition—A systematic review. *Neurosci Biobehav Rev.* (2017) 78:44–56. doi: 10.1016/j.neubiorev.2017.04.013
  185. Gabryel B, Bielecka A, Bernacki J, Labuzek K, Herman ZS. Immunosuppressant cytoprotection correlates with HMGB1 suppression in primary astrocyte cultures exposed to combined oxygen-glucose deprivation. *Pharmacol Rep.* (2011) 63:392–402. doi: 10.1016/S1734-1140(11)70505-9
  186. Santiago FM, Bueno P, Olmedo C, Muffak-Granero K, Comino A, Serradilla M, et al. Effect of N-acetylcysteine administration on intraoperative plasma levels of interleukin-4 and interleukin-10 in liver transplant recipients. *Transplant Proc.* (2008) 40:2978–80. doi: 10.1016/j.transproceed.2008.08.103
  187. Berman AE, Chan WY, Brennan AM, Reyes RC, Adler BL, Suh SW, et al. N-acetylcysteine prevents loss of dopaminergic neurons in the EAAC1<sup>-/-</sup> mouse. *Ann Neurol.* (2011) 69:509–20. doi: 10.1002/ana.22162
  188. Skvarc DR, Dean OM, Byrne LK, Gray LJ, Ives K, Lane SE, et al. The Post-Anaesthesia N-acetylcysteine cognitive evaluation (PANACEA) trial: study protocol for a randomised controlled trial. *Trials* (2016) 17:395. doi: 10.1186/s13063-016-1529-4
  189. Edaravone Acute Infarction Study G. Effect of a novel free radical scavenger, edaravone (MCI-186), on acute brain infarction. Randomized, placebo-controlled, double-blind study at multicenters. *Cerebrovasc Dis.* (2003) 15:222–9. doi: 10.1159/000069318
  190. Abe K, Itoyama Y, Sobue G, Tsuji S, Aoki M, Doyu M, et al. Confirmatory double-blind, parallel-group, placebo-controlled study of efficacy and safety of edaravone (MCI-186) in amyotrophic lateral sclerosis patients. *Amyotroph Lateral Scler Frontotemporal Degener.* (2014) 15:610–7. doi: 10.3109/21678421.2014.959024
  191. Purdon PL, Sampson A, Pavone KJ, Brown EN. Clinical Electroencephalography for Anesthesiologists: Part I: background and basic signatures. *Anesthesiology* (2015) 123:937–60. doi: 10.1097/ALN.0000000000000841
  192. Wang X, Liu N, Chen J, Xu Z, Wang F, Ding C. Effect of intravenous dexmedetomidine during general anesthesia on acute postoperative pain in adults: a systematic review and meta-analysis of randomized controlled trials. *Clin J Pain* (2018). doi: 10.1097/AJP.0000000000000630
  193. Schnabel A, Reichl SU, Weibel S, Kranke P, Zahn PK, Pogatzki-Zahn EM, et al. Efficacy and safety of dexmedetomidine in peripheral nerve blocks: a meta-analysis and trial sequential analysis. *Eur J Anaesthesiol.* (2018) 35:745–58. doi: 10.1097/EJA.0000000000000870
  194. Hu J, Vacas S, Feng X, Lutrin D, Uchida Y, Lai IK, et al. Dexmedetomidine prevents cognitive decline by enhancing resolution of high mobility group box 1 protein-induced inflammation through a vagomimetic action in mice. *Anesthesiology* (2018) 128:921–31. doi: 10.1097/ALN.0000000000002038
  195. Rocha SM, Cristovao AC, Campos FL, Fonseca CP, Baltazar G. Astrocyte-derived GDNF is a potent inhibitor of microglial activation. *Neurobiol Dis.* (2012) 47:407–15. doi: 10.1016/j.nbd.2012.04.014
  196. Kawano T, Eguchi S, Iwata H, Tamura T, Kumagai N, Yokoyama M. Impact of preoperative environmental enrichment on prevention of development of cognitive impairment following abdominal surgery in a rat model. *Anesthesiology* (2015) 123:160–70. doi: 10.1097/ALN.0000000000000697

197. Sheets MF, Hanck DA. Molecular action of lidocaine on the voltage sensors of sodium channels. *J Gen Physiol.* (2003) 121:163–75. doi: 10.1085/jgp.20028651
198. Amir R, Argoff CE, Bennett GJ, Cummins TR, Durieux ME, Gerner P, et al. The role of sodium channels in chronic inflammatory and neuropathic pain. *J Pain* (2006) 7(5 Suppl. 3):S1–29. doi: 10.1016/j.jpain.2006.01.444
199. Maher DP, Chen L, Mao J. Intravenous ketamine infusions for neuropathic pain management: a promising therapy in need of optimization. *Anesth Analg.* (2017) 124:661–74. doi: 10.1213/ANE.0000000000001787
200. Bell JD. In Vogue: ketamine for neuroprotection in acute neurologic injury. *Anesth Analg.* (2017) 124:1237–43. doi: 10.1213/ANE.0000000000001856
201. De Kock M, Loix S, and Lavand'homme P. (2013). Ketamine and peripheral inflammation. *CNS Neurosci Ther.* 19:403–410. doi: 10.1111/cns.12104
202. Li Y, Shen R, Wen G, Ding R, Du A, Zhou J, et al. Effects of ketamine on levels of inflammatory cytokines IL-6, IL-1beta, and TNF-alpha in the hippocampus of mice following acute or chronic administration. *Front Pharmacol.* (2017) 8:139. doi: 10.3389/fphar.2017.00139
203. Della Giustina A, Goldim MP, Danielski LG, Florentino D, Mathias K, Garbossa L, et al. Alpha-lipoic acid attenuates acute neuroinflammation and long-term cognitive impairment after polymicrobial sepsis. *Neurochem Int.* (2017) 108:436–47. doi: 10.1016/j.neuint.2017.06.003
204. Zhang Z, Ma Q, Shah B, Mackensen GB, Lo DC, Mathew JP, et al. Neuroprotective effects of annexin A1 tripeptide after deep hypothermic circulatory arrest in rats. *Front Immunol.* (2017) 8:1050. doi: 10.3389/fimmu.2017.01050
205. Aizpuru-Olaizola O, Elezgarai I, Rico-Barrío I, Zarandona I, Etxebarria N, Usobiaga A. Targeting the endocannabinoid system: future therapeutic strategies. *Drug Discov Today* (2017) 22:105–10. doi: 10.1016/j.drudis.2016.08.005
206. McCoy KL. Interaction between cannabinoid system and toll-like receptors controls inflammation. *Med Inflamm.* (2016) 2016:5831315. doi: 10.1155/2016/5831315
207. Sun L, Dong R, Xu X, Yang X, Peng M. Activation of cannabinoid receptor type 2 attenuates surgery-induced cognitive impairment in mice through anti-inflammatory activity. *J Neuroinflammat.* (2017) 14:138. doi: 10.1186/s12974-017-0913-7
208. Papaseit E, Perez-Mana C, Perez-Acevedo AP, Hladun O, Torres-Moreno MC, Muga R, et al. Cannabinoids: from pot to lab. *Int J Med Sci.* (2018) 15:1286–95. doi: 10.7150/ijms.27087
209. Poleszak E, Wosko S, Slawinska K, Szopa A, Wrobel A, Serefko A. Cannabinoids in depressive disorders. *Life Sci.* (2018) 213:18–24. doi: 10.1016/j.lfs.2018.09.058
210. Cipolla-Neto J, and do Amaral FG. (2018). Melatonin as a hormone: new physiological and clinical insights. *Endocr Rev.* 39:990–1028. doi: 10.1210/er.2018-00084
211. Esposito E, Cuzzocrea S. Antiinflammatory activity of melatonin in central nervous system. *Curr Neuropharmacol.* (2010) 8:228–42. doi: 10.2174/157015910792246155
212. Liu Y, Ni C, Tang Y, Tian X, Zhou Y, Qian M, et al. Melatonin attenuates isoflurane-induced acute memory impairments in aged rats. *Basic Clin Pharmacol Toxicol.* (2013) 113:215–20. doi: 10.1111/bcpt.12079
213. Xia T, Cui Y, Chu S, Song J, Qian Y, Ma Z, et al. Melatonin pretreatment prevents isoflurane-induced cognitive dysfunction by modulating sleep-wake rhythm in mice. *Brain Res.* (2016) 1634:12–20. doi: 10.1016/j.brainres.2015.10.036
214. Song J, Chu S, Cui Y, Qian Y, Li X, Xu F, et al. Circadian rhythm resynchronization improved isoflurane-induced cognitive dysfunction in aged mice. *Exp Neurol.* (2018) 306:45–54. doi: 10.1016/j.expneurol.2018.04.009
215. Kim JH, Gupta SC, Park B, Yadav VR, Aggarwal BB. Turmeric (Curcuma longa) inhibits inflammatory nuclear factor (NF)-kappaB and NF-kappaB-regulated gene products and induces death receptors leading to suppressed proliferation, induced chemosensitization, and suppressed osteoclastogenesis. *Mol Nutr Food Res.* (2012) 56:454–65. doi: 10.1002/mnfr.201100270
216. Wu X, Chen H, Huang C, Gu X, Wang J, Xu D, et al. Curcumin attenuates surgery-induced cognitive dysfunction in aged mice. *Metab Brain Dis.* (2017) 32:789–98. doi: 10.1007/s11011-017-9970-y
217. Sierpina VS, Frenkel MA. Acupuncture: a clinical review. *South Med J.* (2005) 98:330–7. doi: 10.1097/01.SMJ.0000140834.30654.0F
218. Liu P-R, Zhou Y, Zhang Y, Diao S. Electroacupuncture alleviates surgery-induced cognitive dysfunction by increasing  $\alpha 7$ -nAChR expression and inhibiting inflammatory pathway in aged rats. *Neurosci Lett.* (2017) 659:1–6. doi: 10.1016/j.neulet.2017.08.043
219. Thal SC, Luh C, Schaible E-V, Timaru-Kast R, Hedrich J, Luhmann HJ, et al. Volatile Anesthetics influence blood-brain barrier integrity by modulation of tight junction protein expression in traumatic brain injury. *PLoS ONE* (2012) 7:e50752–e50752. doi: 10.1371/journal.pone.0050752
220. Tetrault S, Chever O, Sik A, Amzica F. Opening of the blood-brain barrier during isoflurane anaesthesia. *Eur J Neurosci.* (2008) 28:1330–41. doi: 10.1111/j.1460-9568.2008.06443.x
221. Royall DR, Palmer R, Chiodo LK, Polk MJ. Declining executive control in normal aging predicts change in functional status: the Freedom House Study. *J Am Geriatr Soc.* (2004) 52:346–52. doi: 10.1111/j.1532-5415.2004.52104.x

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Safavynia and Goldstein. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.