



SARS-CoV-2 Infection in Healthcare Professionals and General Population During “First Wave” of COVID-19 Pandemic: A Cross-Sectional Study Conducted in Sicily, Italy

Claudio Costantino^{1,2*}, Emanuele Cannizzaro¹, Maria Gabriella Verso¹, Fabio Tramuto^{1,2}, Carmelo Massimo Maida^{1,2}, Guido Lacca¹, Davide Alba¹, Livia Cimino¹, Arianna Conforto¹, Luigi Cirrincione¹, Giorgio Graziano², Sara Palmeri¹, Stefano Pizzo¹, Vincenzo Restivo^{1,2}, Alessandra Casuccio¹, Francesco Vitale^{1,2} and Walter Mazzucco^{1,2}

OPEN ACCESS

Edited by:

Nicola Mucci,
University of Florence, Italy

Reviewed by:

Massimiliano Barattucci,
University of eCampus, Italy
Georgios Merekoullas,
University of Patras, Greece

*Correspondence:

Claudio Costantino
claudio.costantino01@unipa.it

Specialty section:

This article was submitted to
Occupational Health and Safety,
a section of the journal
Frontiers in Public Health

Received: 19 December 2020

Accepted: 08 March 2021

Published: 13 May 2021

Citation:

Costantino C, Cannizzaro E, Verso MG, Tramuto F, Maida CM, Lacca G, Alba D, Cimino L, Conforto A, Cirrincione L, Graziano G, Palmeri S, Pizzo S, Restivo V, Casuccio A, Vitale F and Mazzucco W (2021) SARS-CoV-2 Infection in Healthcare Professionals and General Population During “First Wave” of COVID-19 Pandemic: A Cross-Sectional Study Conducted in Sicily, Italy.
Front. Public Health 9:644008.
doi: 10.3389/fpubh.2021.644008

¹ Department of Health Promotion, Maternal and Infant Care, Internal Medicine and Medical Specialties (PROMISE) “G. D’Alessandro,” University of Palermo, Palermo, Italy, ² COVID-19 Surveillance Western Sicily Reference Laboratory, Palermo University Hospital, Palermo, Italy

On December 31, 2019, an outbreak of lower respiratory infections was documented in Wuhan caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Since the beginning, SARS-CoV-2 has caused many infections among healthcare workers (HCWs) worldwide. Aims of this study were: a. to compare the distribution among the HCWs and the general population of SARS-CoV-2 infections in Western Sicily and Italy; b. to describe the characteristics of HCWs infected with SARS-CoV-2 in the western Sicilian healthcare context during the first wave of the epidemic diffusion in Italy. Incidence and mean age of HCWs infected with SARS-CoV-2 were comparable in Western Sicily and in the whole Italian country. The 97.6% of infections occurred in HCWs operating in non-coronavirus disease 2019 (COVID-19) working environments, while an equal distribution of cases between hospital and primary care services context was documented. Nurses and healthcare assistants, followed by physicians, were the categories more frequently infected by SARS-CoV-2. The present study suggests that healthcare workers are easily infected compared to the general population but that often infection could equally occur in hospital and non-hospital settings. Safety of HCWs in counteracting the COVID-19 pandemic must be strengthened in hospital [adequate provision of personal protective equipment (PPE), optimization of human resources, implementation of closed and independent groups of HCWs, creation of traffic control building and dedicated areas in every healthcare context] and non-hospital settings (influenza vaccination, adequate psychophysical support, including refreshments during working shifts, adequate rest, and family support).

Keywords: SARS-CoV-2 infection, COVID-19, healthcare workers, personal protective equipment, traffic control building, influenza vaccination, psycho-physical support

INTRODUCTION

On December 31, 2019, an outbreak of unexplained lower respiratory infections was documented in Wuhan, the largest metropolitan area in China's Hubei Province, and reported for the first time to the WHO National Office in China (1). The etiology of this disease was attributed to a new virus belonging to the family of coronaviruses (CoVs), renamed severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (2).

Previous outbreaks of CoV have been documented in the past, such as severe acute respiratory syndrome renamed coronavirus (SARS-CoV-1) in 2002/2003 and Middle East respiratory syndrome coronavirus (MERS-CoV) in 2012 (3).

In early March 2020, the new virus demonstrated high contagiousness, and its worldwide spread has reached the epidemiological criteria necessary to be declared a pandemic (4). To date, about 73 million cases of SARS-CoV-2 infection and more than 1.6 million deaths due to coronavirus disease 2019 (COVID-19) have been observed worldwide (4).

In the last months, in Europe, there has been an epidemiological transition of the SARS-CoV-2 epidemic, with a sharp drop to 43 years in the average age of the population contracting the infection (5, 6).

Virus circulation occurs more frequently in younger age groups due to a greater reopening of commercial activities (including meeting places) and greater mobility, and this variation in transmission dynamics (with the onset of cases and outbreaks linked to recreational activities) was recently confirmed by epidemiological data supporting the evidence of the occurrence of a possible second wave of the pandemic (6).

SARS-CoV-2 has also caused many infections among healthcare workers (HCWs) worldwide (7). The Chinese National Health Commission stated that more than 3,300 health workers nationwide had been infected, and many of them died during the first epidemic wave that occurred in China (8).

HCWs are particularly vulnerable compared to the general population, as demonstrated by the continuous increase in cases, because the coronavirus is highly contagious and members of medical staff are increasingly exposed to viral particles, while often not being provided with adequate protective equipment (9). In addition, a combination of stress, long working hours, and night work could make their immune system more vulnerable than normal (10–13).

In addition, the Commission Directive (EU) 2020/739 included SARS-CoV-2 in the list of biological agents known to infect humans and amending Commission Directive (EU) 2019/1833 (14).

In Italy, to date, more than 80,000 HCWs have been infected with SARS-CoV-2 and hundreds have died due to COVID-19 (6).

Aims of this observational study were to compare the distribution among the HCWs and the general population of SARS-CoV-2 infection cases that occurred in Western Sicily and Italy and to describe the main characteristics of HCWs infected with SARS-CoV-2 in the western Sicilian healthcare context during the first wave of the SARS-CoV-2 epidemic diffusion in Italy. This analysis could contribute to investigate any differences of SARS-CoV-2 infection

incidence according to health care setting, professional role, or working environment.

MATERIALS AND METHODS

A cross-sectional study was conducted in Western Sicily, to date, a COVID-19 low-incidence area accounting for about 47.5% of the total Sicilian resident population (4,968,410 inhabitants), to highlight the characteristics of HCWs infected with SARS-CoV-2 during the period between February 26, 2020 (first laboratory-confirmed case of SARS-CoV-2 infection diagnosed in Sicily among the general population), and May 3, 2020 (the last days of the “lockdown” that began in Italy on March 8, 2020).

Data were obtained from the COVID-19 surveillance systems in place at the local health facilities (LHFs) of four western Sicilian provinces: Agrigento (429,611 inhabitants), Caltanissetta (260,779 inhabitants), Palermo (1,243,328 inhabitants), and Trapani (428,337 inhabitants).

To this end, a dedicated electronic format has been structured and then shared with officials from the prevention departments of recalled LHFs, identified to collect data on HCWs who resulted positive to SARS-CoV-2.

The variables collected were gender, age (mean, standard deviation), period of SARS-CoV-2 infection diagnosis (I subperiod = from February 26, 2020, to March 31, 2020; II subperiod = from April 1, 2020, to May 3, 2020), province (Agrigento, Caltanissetta, Palermo, and Trapani), healthcare context (hospital, local health structure, or primary care service), professional role (physician, nurse, healthcare assistant, laboratory technician), working environment (COVID-19 or not COVID-19 hospital/ward/primary care service).

The diagnosis of SARS-CoV-2 infection, in accordance with WHO guidelines, was considered for nasopharyngeal swab specimens confirmed at real-time polymerase chain reaction (RT-PCR) by one of the diagnostic laboratories authorized by the Sicilian regional health department or by the Italian Ministry of Health (4, 6).

Lastly, data obtained from the LHFs were compared with data on the general population available from the open-access database provided by the Italian national surveillance system for COVID-19 epidemic and with data on HCWs provided by the Italian National Health Institute (6).

The study was approved by the Ethical Committee Palermo 1 of the University Hospital of Palermo (session n. 7 of July 13, 2020).

Statistical Analysis

Quantitative variables were normally distributed and summarized as means with their standard deviations (\pm SD), while absolute and relative frequencies were calculated for qualitative variables. Chi-square test was performed to compare the distribution of SARS-CoV-2-positive cases. Statistical significance was set with a $p < 0.05$.

All data were entered into an electronic database created by Excel 16.0 software. Descriptive statistics were performed using EpiInfo[®] ver. 3.5.1 software.

TABLE 1 | Distribution of laboratory-confirmed diagnosis of SARS-CoV-2 infections among HCWs and the general population.

Location	SARS-CoV-2 laboratory-confirmed infection			p-value
	Overall	HCWs	General population	
	<i>n</i> (%)			
Western sicily*	944 (100)	85 (9.0)	859 (91.0)	0.18
Remaining part of Italy*	209,773 (100)	21,795 (11.6)	187,978 (88.4)	
Italy overall	210,717 (100)	21,880 (10.4)	188,837 (89.6)	

*Comparison between Western Sicily and Italy (February 26, 2020, to May 3, 2020). HCWs, healthcare workers; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

RESULTS

All SARS-CoV-2 infection cases among HCWs in Western Sicily registered by the Sicilian Health Department were contacted and reported in the analysis.

In **Table 1**, the comparison between Western Sicily and Italy of the distribution of SARS-CoV-2 infection cases, confirmed by RT-PCR molecular test, among HCWs and the general population is represented, which occurred during the first wave of the COVID-19 pandemic.

In Western Sicily, during the period in this study, 944 SARS-CoV-2-positive subjects were documented overall. Of these, 85 (9%) were HCWs and the remaining 859 (91%) occurred in the general population.

The total number of SARS-CoV-2-positive subjects reported in the remaining part of Italy in the same period was of 209,773, with 21,795 (11.6%) cases occurring in HCWs and 187,978 (88.4%) documented in the general population.

However, no statistical difference was reported when comparing the distribution of SARS-CoV-2-positive cases between Western Sicily and the remaining part of Italy (*p*-value 0.18).

Table 2 summarizes the characteristics of the 85 HCWs working in the Western Sicily healthcare context with a laboratory-confirmed diagnosis of SARS-CoV-2 infection reported during the first pandemic. Among them, 52 (61.2%) were males and 33 (38.8%) were females, with an average age of 46.4 (SD ± 10.8) years. The majority of HCWs (61; 71.8%) were infected in the first subperiod of the study (from February 26, 2020, to March 31, 2020), while the remaining part (24; 28.2%) during the second subperiod surveyed (from April 1, 2020, to May 3, 2020).

The cases were distributed in the four LHF as follows: 21 (24.7%) Agrigento, nine (10.6%) Caltanissetta, 35 (41.2%) Palermo, 20 (23.5%) Trapani.

Among the laboratory-confirmed diagnosis of SARS-CoV-2 infection occurring in HCWs from the western Sicilian healthcare context, 43 (50.6%) worked in a hospital and 42 (49.4%) in the primary care services or in other structures belonging to the LHF.

With regard to the working environment, two (2.4%) HCWs operated in a COVID-19 hospital/ward or primary care service

TABLE 2 | Characteristics of 85 HCWs with a laboratory-confirmed diagnosis of SARS-CoV-2 infection working in the western Sicilian healthcare context (February 26, 2020, to May 3, 2020).

Characteristics		<i>n</i>	(%)
Gender	Male	52	61.2
	Female	33	38.8
Period of diagnosis	I subperiod in study*	61	71.8
	II subperiod in study**	24	28.2
Age	Mean (± SD)	46.4	± 10.8
Province	Agrigento	21	24.7
	Caltanissetta	9	10.6
	Palermo	35	41.2
	Trapani	20	23.5
	Healthcare context	Hospital	43
	Local health structure or primary care service	42	49.4
Professional role	Physician	22	26.8
	Nurse	29	34.1
	Healthcare assistant	29	34.1
	Laboratory technician	5	4.9
Working environment (hospital/ward/primary care service)	COVID-19	2	2.4
	Non-COVID-19	83	97.6

COVID-19, coronavirus disease 2019; HCWs, healthcare workers; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; SD, standard deviation.

*From February 26, 2020, to March 31, 2020; **from April 1, 2020, to May 3, 2020.

considered at high risk of exposure to SARS-CoV-2, while 83 (97.6%) of the SARS-CoV-2-infected HCWs were from a non-COVID-19 working context.

Lastly, of the 85 SARS-CoV-2-positive HCWs, 22 (26.8%) were physicians, 29 (34.1%) were nurses, five (4.9%) were laboratory technicians, and 29 (34.1%) were healthcare assistants.

DISCUSSION

Since the end of February 2020, when the first evidence of local transmission of SARS-CoV-2 was observed in Italy, many HCWs have also been infected by the pandemic virus (6).

We conducted a cross-sectional study to compare the distribution among the HCWs and the general population of SARS-CoV-2 infection cases occurring in Western Sicily and Italy and to further document the main characteristics of the SARS-CoV-2-infected HCWs operating in the western Sicilian healthcare system, laying in an Italian COVID-19 low-incidence context, during the first wave of the SARS-CoV-2 pandemic. To this end, we accessed data from the COVID-19 surveillance systems in place at the Sicilian LHF and the open-access database provided by the Italian national surveillance system for COVID-19 epidemic.

During the first wave of the pandemic, the distribution of SARS-CoV-2-positive cases among HCWs and the general population was comparable in Western Sicily and in the whole Italian country.

This finding related to the proportion of SARS-CoV-2-infected HCWs on the overall number of SARS-CoV-2-positive

cases should be further analyzed according to the different incidences of COVID-19 cases decreasing gradient from north to south reported in Italy (6). The greatest burden of the disease documented for northern Italian regions could be attributable to several factors such as the delay in applying the lockdown, the import-export with China, the influenza vaccination coverage rates, the mean seasonal temperatures, and the different impacts of air pollution among different Italian regions, and the role of these factors should be considered when investigating the impact of COVID-19 on HCWs as well (6, 15).

Of interest, the risk of contracting COVID-19 among HCWs is one of the highest because regardless of the specific exposure related to healthcare settings, outside working hours, they are also exposed to the same risk and pathways of contagion reported in the community (16–18).

More in depth, HCWs resulted at very high risk of contracting SARS-CoV-2 infection because of several factors: the healthcare setting (not necessarily an at-risk ward that required tight preventive measures), longer working hours, night shifts that adversely affect the normal waking-sleep rhythm, and poor hand hygiene following contact with patients potentially affected by COVID-19 or the inadequacy of the personal protective equipment (PPE) and of the training in its correct use, provided by the healthcare organization during a pandemic (17, 18).

Our study documented that HCWs at higher risk of SARS-CoV-2 infection during the first pandemic wave were males and had an average age of 46.6 years. These findings are in line with the ones reported in Italy overall during the same time frame (6).

Of relevance, the majority of COVID-19 cases occurring in western Sicilian HCWs were reported during the first month of the first epidemic wave, probably reflecting an improvement over time of the prevention and control measures implemented in the healthcare setting. During the COVID-19 pandemic, HCWs represent the most valuable resource in each country worldwide (19, 20). However, knowledge about the effectiveness of PPE on HCWs caring for patients infected with the novel coronavirus is continuously increasing (21).

Practical Implication

A recent clinical case described the clinical outcome of 41 HCWs who were exposed to aerosol-generating procedures for at least 10 min at a distance of <2 m from a patient with severe pneumonia before the laboratory-confirmed diagnosis of COVID-19 was known (22). Despite the fact that 85% of HCWs were exposed during a procedure involving aerosol generation by the patient while wearing a surgical mask or N95 mask, none of them developed symptoms and all molecular tests resulted negative (22).

These pieces of evidence suggest that surgical masks, hand hygiene, and other standard procedures are effective in protecting HCWs from the SARS-CoV-2 infection. A single case report is not sufficient to determine the best way to protect HCWs from COVID-19, but it is sufficient to suggest that further studies are needed (23).

Moreover, with the present study, we tried to assess the possible increased vulnerability of HCWs to develop the SARS-CoV-2 infection according to their healthcare context and

professional role. In particular, the number of HCWs with a confirmed laboratory diagnosis of SARS-CoV-2 working in hospitals was comparable to the one documented for HCWs operating in the primary care services or in other structures belonging to the LHF, suggesting a similar level of risk for contracting the disease (23, 24). A possible interpretation of these data can be related to a contagion in the LHF, especially among HCWs working in long-term care facilities for elderly and disabled persons (25).

Therefore, HCWs operating at the territorial services or in a non-COVID-19 setting, on one side, and the ones working at the hospital wards (not dedicated to the treatment of patients affected by COVID-19), could not have greater protection from the virus than their colleagues working in COVID-19 units (26).

In accordance with this hypothesis, 97.6% of the HCWs infected by SARS-CoV-2 in western Sicilian healthcare settings worked in environments irresolute to COVID-19 patterns of care. In this perspective, the experiences conducted in low-resource countries in the management of epidemics within outpatient settings while having a reduced number of available health personnel could be of interest (27–29).

However, one of the most important reasons for early infection among general ward medical staff is the frequent misconduct of admitting patients to the ward without implementing preventive and protective measures (30).

On the contrary, infection rates in the most protected intensive care and emergency wards were lower in unannounced cases of illness (31).

Recent research published in the *Journal of the American Medical Association* found that of the 138 patients studied in a hospital in Wuhan, 29% were HCWs, of whom 31 (77.5%) were working in general wards, seven (17.5%) in emergency rooms, and two (5%) in intensive care units (32).

Of interest, it was assumed that more than 10 HCWs in this ward were infected by a patient with abdominal symptoms and admitted to the surgical ward (32).

One of the reasons for the difficulties in diagnosing COVID-19 and the consequent placement of the patient in wards without adequate measures to contain the virus was found in the atypical presentation symptoms in the initial phase in some patients infected with SARS-CoV-2, such as the gastrointestinal symptoms of the patient under examination, which probably contributed to the rapid spread of the infection among HCWs (32).

Of interest, Traffic Control Bundling (TCB) tool was proven to be effective in drastically reducing infection rates among HCWs in Taiwan during the SARS epidemic (33). Furthermore, the greatest susceptibility to respiratory infectious diseases, such as SARS, was documented for HCWs (24); moreover, data from seven hospitals in China showed an incidence of up to 13.53% of SARS-infected HCWs in intensive care (intensive care units) (26).

More in depth, TCB approach requires the identification of (1) a pre-triage outside hospital (in tents or other shelters), ensuring that patients are triaged in outdoor screening stations so that sick patients are directed to a contamination zone; and (2) risk zones, clearly delimiting separate zones, including a contamination,

transition, and clean zone, each separated by control points. Therefore, it has been suggested to implement the TCB model to face manage the COVID-19 outbreak (33).

HCWs should be adequately trained on TCB protocols including good dressing and undressing practices, proper use of all appropriate personal safety equipment (e.g., respirators and eye protection), and how to move safely between areas (34).

Last but not least, some differences between the working categories were observed in the present study. Nurses and healthcare assistants were the categories most affected by the pandemic virus as compared to physicians probably because their role involves greater physical proximity to the patient (dressing, cleaning, feeding) and, therefore, a greater chance of viral transmission, differently from laboratory technicians who manipulate the swabs in a controlled laboratory (35).

Generally, HCWs could also be easily infected outside work environment in the familiar context. The lack or incorrect use of preventive measures in order to prevent the transmission of SARS-CoV-2 infection is usually more frequent among family members or friends (36).

The strong recommendation of influenza vaccination should be of paramount importance during the 2020/2021 influenza season for certain categories such as HCWs, medical and healthcare residents/students/trainees, police officers, fire workers, and other workers of public utility (37).

In that direction, three Italian regions have tried to introduce local policies or laws that mandate influenza vaccination among HCWs, considering them at high risk for contracting and spreading influenza viruses that could contribute to the burden and overcrowding of health care systems together with SARS-CoV-2 during the next cold season (38, 39).

Could be also useful is the possibility for some HCWs, directly involved in the care of COVID-19-infected patients, to separate temporarily from their families in dedicated COVID-19 hotels for HCWs in order to protect their parents from the risk of secondary infection (40, 41).

Also the mental health and the COVID-19-related mental health effects in the workplace of frontline health and social care professionals should be taken into account in the preventive measures to be considered in the future (42, 43). For instance, HCWs infected by SARS-CoV-2 expected to experience higher levels of stigma among colleagues and in the working setting, reporting increased psychological distress (44–46).

Limitations and Strengths

The main limitation of the present study is linked to its purely descriptive design and to the small number of cases collected that could not be representative of the national context where a different epidemiologic impact was observed, with a COVID-19 incidence decreasing from north to south.

Moreover, further analytical studies, such as ones with a case-control design, should be conducted in the near future to evaluate the different risk factors associated with the infection among HCWs who tested positive for SARS-CoV-2.

Also for future research on SARS-CoV-2 infection of HCWs, an in-depth analysis of psychological variables should be considered.

Finally, the present study contributes to the research conducted on SARS-CoV-2 infections among HCWs in other countries. Specifically, the main elements of originality of the present research can be summarized as follows:

1. HCWs, independently if working in COVID-19 or non-COVID-19 environment, are easily infected in geographical areas with a higher incidence of SARS-CoV-2 infection (Northern vs. Southern Italy during the “first-wave” interesting Italy) (6).
2. There are no significant differences in SARS-CoV-2 infection incidence in accordance to working environment or healthcare context, suggesting a higher risk of transmission among HCWs in non-healthcare settings, where preventive measures are usually neglected.

CONCLUSIONS

In conclusion, it is essential to highlight how the safety of HCWs must be strengthened at a global level through certain essential preventive measures (23).

During the next phases of the COVID-19 pandemic, some preventive measures in order to reduce the SARS-CoV-2 infections among HCWs should be strongly encouraged.

Adequately providing suitable PPE, optimizing human resources in the different healthcare settings, implementing closed and independent groups of HCWs that can be easily isolated, creating traffic control building and dedicated areas for suspected and confirmed cases of SARS-CoV-2 infection, offering influenza vaccination to all HCWs, and providing everything that can guarantee adequate psychophysical support (including refreshments during working shifts, guaranteeing adequate rest, and family support) are among the most important strategies to be considered.

Finally, it should be considered that HCWs are easily infected compared to the general population, but often that infection could occur in hospital and non-hospital settings.

Preventive measures in order to contrast SARS-CoV-2 infection should be continuously applied by the general population until the end of this pandemic, also contributing to a reduction of HCW infections (36).

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Ethical Committee Palermo 1 of the University Hospital of Palermo (session n.7 of July 13, 2020). Written informed consent for participation was not required for this study in accordance with the national legislation and the institutional requirements.

AUTHOR CONTRIBUTIONS

CC and WM were responsible for the study conception, methodology, and writing the original draft of the manuscript. EC and FV were responsible for reviewing and editing the manuscript and supervision of the project. ACa, VR, GL, and

MV were responsible for the statistical analysis. CM and FT were responsible for data curation. LCir, GG, SPa, and SPI were responsible for data collection. DA, ACo, and LCim were responsible for the investigation in the western Sicilian local health facilities. All authors contributed to the article and approved the submitted version.

REFERENCES

- Hongzhou L, Stratton CW, Tang, YW. Outbreak of pneumonia of unknown etiology in Wuhan, China: the mystery and the miracle. *J Med Virol.* (2020) 92:401–2. doi: 10.1002/jmv.25678
- Li H, Liu SM, Yu XH, Tang SL, Tang CK. Coronavirus disease 2019 (COVID-19): current status and future perspective. *Int J Antimicrob Agents.* (2020) 55:105951. doi: 10.1016/j.ijantimicag.2020.105951
- Al-Tawfiq JA. Asymptomatic coronavirus infection: MERS-CoV and SARS-CoV-2 (COVID-19). *Travel Med Infect Dis.* (2020) 35:101608. doi: 10.1016/j.tmaid.2020.101608
- World Health organization. *WHO Coronavirus Disease (COVID-19) Dashboard.* (2020). Available online at: https://covid19.who.int/?gclid=Cj0KCQiA2uH-BrCCARIsAEef3m3GsDyJAW0WSvWcOjP_HhPib3C44pLp3TuuNKDCXwTR0r4Zd-ALj4aAqXhEALw_wcB (accessed December 15, 2020).
- Costantino C, Cannizzaro E, Alba D, Conforto A, Cimino L, Mazzucco W. Sars-Cov-2 pandemic in the mediterranean area: epidemiology and perspectives. *EuroMediterranean Biomed J.* (2020) 15:102–106. doi: 10.3269/1970-5492.2020.15.25
- Istituto Superiore di Sanità. *Epicentro Dati Della Sorveglianza Integrata COVID-19 in Italia.* (2020). Available online at: <https://www.epicentro.iss.it/coronavirus/sars-cov-2-dashboard> (accessed December 15, 2020).
- Li R, Xuyu C, Ying W, Wenwen W, Ling Z, Xiaodong T. Risk factors of healthcare workers with coronavirus disease 2019: a retrospective cohort study in a designated hospital of Wuhan in China. *Clin Infect Dis.* (2020) 17:2218–21. doi: 10.1093/cid/ciaa287
- Lai X, Wang M, Qin C, Tan L, Ran L, Chen D, et al. Coronavirus disease 2019 (COVID-2019) infection among health care workers and implications for prevention measures in a tertiary hospital in Wuhan, China. *JAMA Netw Open.* (2020) 3:e209666. doi: 10.1001/jamanetworkopen.2020.9666
- Sommerstein R, Fux CA, Vuichard-Gysin D, Abbas M, Marschall J, Balmelli C, et al. Risk of SARS-CoV-2 transmission by aerosols, the rationale use of masks, and protection of healthcare workers from COVID-19. *Antimicrob Resist Infect Control.* (2020) 9:100. doi: 10.1186/s13756-020-00763-0
- Torales J, O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int J Soc Psychiatry.* (2020) 66:317–20. doi: 10.1177/0020764020915212
- Cannizzaro E, Ramaci T, Cirrincione L, Plescia F. Work-Related stress, physio-pathological mechanisms, and the influence of environmental genetic factors. *Int J Environ Res Public Health.* (2019) 16:4031. doi: 10.3390/ijerph16204031
- Lahner E, Dilaghi E, Prestigiacomo C, Alessio G, Marcellini L, Simmaco M, et al. Prevalence of sars-Cov-2 infection in health workers (HWs) and diagnostic test performance: the experience of a teaching hospital in central Italy. *Int J Environ Res Public Health.* (2020) 17:4417. doi: 10.3390/ijerph17124417
- Ramaci T, Barattucci M, Vella F, Senia P, Cannizzaro E, Scorciapino A, et al. Straining at work and its relationship with personality profiles and individual consequences in healthcare workers (HCWs). *Int J Environ Res Public Health.* (2020) 17:610. doi: 10.3390/ijerph17020610
- European Union Law. *Commission Directive (EU) 2020/739 of 3 June 2020 Amending Annex III to Directive 2000/54/EC of the European Parliament and of the Council as Regards the Inclusion of SARS-CoV-2 in the List of Biological Agents Known to Infect Humans and Amending Commission Directive (EU) 2019/1833.* (2020). Available online at <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32020L0739> (accessed December 11, 2020).
- Amato M, Werba JP, Frigerio B, Coggi D, Sansaro D, Ravani A, et al. Relationship between influenza vaccination coverage rate and COVID-19 outbreak: an Italian ecological study. *Vaccines.* (2020) 8:535. doi: 10.3390/vaccines8030535
- Costantino C, Fiacchini D. Rationale of the WHO document on risk communication and community engagement (RCCE) readiness and response to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and of the Italian decalogue for prevention departments. *J Prev Med Hyg.* (2020) 61:E1–2. doi: 10.15167/2421-4248/jpmh2020.61.1.1502
- World Health Organization. *Coronavirus Disease (COVID-19) Outbreak: Rights, Roles and Responsibilities of Health Workers, Including Key Considerations for Occupational Safety and Health.* (2020). Available online at: [https://www.who.int/publications-detail/coronavirus-disease-\(covid-19\)-outbreak-rights-roles-and-responsibilities-of-health-workers-including-key-considerations-for-occupational-safety-and-health](https://www.who.int/publications-detail/coronavirus-disease-(covid-19)-outbreak-rights-roles-and-responsibilities-of-health-workers-including-key-considerations-for-occupational-safety-and-health) (accessed December 11, 2020).
- Shah ASV, Wood R, Gribben C, Caldwell D, Bishop J, Weir A, et al. Risk of hospital admission with coronavirus disease 2019 in healthcare workers and their households: nationwide linkage cohort study. *BMJ.* (2020) 371:m3582. doi: 10.1136/bmj.m3582
- The Lancet. COVID-19: protecting health-care workers. *Lancet.* (2020) 395:922. doi: 10.1016/S0140-6736(20)30644-9
- Cirrincione L, Plescia F, Ledda C, Rapisarda V, Martorana D, Moldovan RE, et al. COVID-19 pandemic: prevention and protection measures to be adopted at the workplace. *Sustainability.* (2020) 12:3603. doi: 10.3390/su12093603
- Bahl P, Doolan C, de Silva C, Chughtai AA, Bourouiba L, MacIntyre CR, et al. Airborne or droplet precautions for health workers treating COVID-19? *J Infect Dis.* (2020) 16:jiaa189. doi: 10.1093/infdis/jiaa189
- Ng, K, Poon BH, Kiat Puar TH, Li Shan Quah J, Loh WJ, et al. COVID-19 and the risk to health care workers: a case report. *Ann Intern Med.* (2020) 172:766–67. doi: 10.7326/L20-0175
- World Health Organization. *Preventing and Managing COVID-19 Across Long-Term Care Services: Policy Brief.* (2020). Available online at: https://www.who.int/publications/i/item/WHO-2019-nCoV-Policy_Brief-Long-term_Care-2020.1 (accessed December 11, 2020).
- Schwartz J, King CC, Yen MY. Protecting healthcare workers during the coronavirus disease 2019 (COVID-19) outbreak: lessons from taiwan's severe acute respiratory syndrome response. *Clin Infect Dis.* (2020) 28:71:858–60. doi: 10.1093/cid/ciaa255
- Martin C, Montesinos I, Dauby N, Gilles C, Dahma H, Van Den Wijngaert S, et al. Dynamics of SARS-CoV-2 RT-PCR positivity and seroprevalence among high-risk healthcare workers and hospital staff. *J Hosp Infect.* (2020) 106:102–6. doi: 10.1016/j.jhin.2020.06.028
- Huh S. How to train health personnel to protect themselves from SARS-CoV-2 (novel coronavirus) infection when caring for a patient or suspected case. *J Educ Eval Health Prof.* (2020) 17:10. doi: 10.3352/jeehp.2020.17.10
- Quaglio G, Tognon F, Finos L, Bome D, Sesay S, Kebbie A, et al. Impact of Ebola outbreak on reproductive health services in a rural district of sierra leone: a prospective observational study. *BMJ Open.* (2020) 9:e029093. doi: 10.1136/bmjopen-2019-029093
- Pizzol D, Veronese N, Marotta C, Di Gennaro F, Moiane J, Chhaganlal K, et al. Predictors of therapy failure in newly diagnosed pulmonary tuberculosis cases in Beira, Mozambique. *BMC Res Notes.* (2018) 11:99. doi: 10.1186/s13104-018-3209-9
- Marotta C, Giaquinto C, Di Gennaro F, Chhaganlal KD, Saracino A, Moiane J, et al. Pathways of care for HIV infected children in Beira, Mozambique: pre-post intervention study to assess impact of task shifting. *BMC Public Health.* (2018) 18:703. doi: 10.1186/s12889-018-5646-8

30. Ong S, Tan Y, Sutjipto S, Chia PY, Young BE, Gum M, et al. Absence of contamination of personal protective equipment (PPE) by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Infect Control Hosp Epidemiol.* (2020) 41:614–6. doi: 10.1017/ice.2020.91
31. Hu X, Zhang Z, Li N, Liu D, Zhang L, He W, et al. Self-Reported Use of personal protective equipment among Chinese critical care clinicians during 2009 h1n1 influenza pandemic. *PLoS ONE.* (2012) 7:e44723. doi: 10.1371/journal.pone.0044723
32. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *JAMA.* (2020) 323:1061–9. doi: 10.1001/jama.2020.1585
33. Yen MY, Lin YE, Lee CH, Ho MS, Huang FY, Chang SC, et al. Taiwan's traffic control bundle and the elimination of nosocomial severe acute respiratory syndrome among healthcare workers. *J Hosp Infect.* (2011) 77:332–7. doi: 10.1016/j.jhin.2010.12.002
34. Bartoszko JJ, Farooqi MAM, Alhazzani W, Loeb M. Medical masks vs N95 respirators for preventing COVID-19 in healthcare workers: a systematic review and meta-analysis of randomized trials. *Influenza Other Respir. Viruses.* (2020) 14:365–73. doi: 10.1111/irv.12745
35. Tostmann A, Bradley J, Bousema T, Yiek WK. Strong associations and moderate predictive value of early symptoms for SARS-CoV-2 test positivity among healthcare workers, the Netherlands. *Euro Surveill.* (2020) 25:2000508. doi: 10.2807/1560-7917.ES.2020.25.16.2000508
36. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ. COVID-19 systematic urgent review group effort (SURGE) study authors. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet.* (2020) 395:1973–87. doi: 10.1016/j.lancet.2020.07.040
37. Costantino C, Restivo V, Gaglio V, Lanza GLM, Marotta C, Maida CM, et al. Effectiveness of an educational intervention on seasonal influenza vaccination campaign adherence among healthcare workers of the Palermo University Hospital, Italy. *Ann Ig.* (2019) 31:35–44. doi: 10.7416/ai.2019.2256
38. Costantino C, Restivo V, Tramuto F, Casuccio A, Vitale F. Influenza vaccination of healthcare workers in Italy: could mandatory vaccination be a solution to protect patients? *Future Microbiol.* (2020) 14:45–9. doi: 10.2217/fmb-2018-0238
39. Costantino C, Casuccio A, Caracci F, Bono S, Calamusa G, Ventura G, et al. Impact of communicative and informative strategies on influenza vaccination adherence and absenteeism from work of health care professionals working at the University hospital of Palermo, Italy: a quasi-experimental field trial on twelve influenza seasons. *Vaccines.* (2020) 8:E5. doi: 10.3390/vaccines8010005
40. Vimercati L, Tafuri S, Chironna M, Loconsole D, Fucilli FIM, Migliore G, et al. The COVID-19 hotel for healthcare workers: AN Italian best practice. *J Hosp Infect.* (2020) 105:387–8. doi: 10.1016/j.jhin.2020.05.018
41. Vimercati L, Dell'Erba A, Migliore G, De Maria L, Caputi A, Quarato M, et al. Prevention and protection measures of healthcare workers exposed to SARS-CoV-2 in a University hospital in Bari, Apulia, southern Italy. *J Hosp Infect.* (2020) 105:454–8. doi: 10.1016/j.jhin.2020.05.024
42. Pollock A, Campbell P, Cheyne J, Cowie J, Davis B, McCallum J, et al. Interventions to support the resilience and mental health of frontline health and social care professionals during and after a disease outbreak, epidemic or pandemic: a mixed methods systematic review. *Cochrane Database Syst Rev.* (2020) 11:CD013779. doi: 10.1002/14651858.CD013779
43. Giorgi G, Lecca LI, Alessio F, Finstad GL, Bondanini G, Lulli LG, et al. COVID-19-Related mental health effects in the workplace: a narrative review. *Int J Environ Res Public Health.* (2020) 17:7857. doi: 10.3390/ijerph17217857
44. Ramaci T, Barattucci M, Ledda C, Rapisarda V. Social stigma during COVID-19 and its impact on HCWs outcomes. *Sustainability.* (2020) 12:3834. doi: 10.3390/su12093834
45. Barattucci M, Chirico A, Kuvačić G, De Giorgio A. Rethinking the role of affect in risk judgment: what we have learned from COVID-19 during the first week of quarantine in Italy. *Front Psychol.* (2020) 11:554561. doi: 10.3389/fpsyg.2020.554561
46. Cannizzaro E, Cannizzaro C, Martorana D, Moscadini S, Lo Coco D. Effects of shift work on cardiovascular activity, serum cortisol and white blood cells count in a group of Italian fishermen. *EuroMediterranean Biomed J.* (2012) 7:109–13. doi: 10.3269/1970-5492.2012.7.23

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Costantino, Cannizzaro, Verso, Tramuto, Maida, Lacca, Alba, Cimino, Conforto, Cirrincione, Graziano, Palmeri, Pizzo, Restivo, Casuccio, Vitale and Mazucco. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.