



# New Insights on the Use of Polyphenols as Natural Preservatives and Their Emerging Safety Concerns

Fred Kwame Ofofu<sup>1</sup>, Eric Banan-Mwine Daliri<sup>1</sup>, Fazle Elahi<sup>1</sup>, Ramachandran Chelliah<sup>1</sup>, Byong-Hoon Lee<sup>2</sup> and Deog-Hwan Oh<sup>1\*</sup>

<sup>1</sup> Department of Food Science and Biotechnology, College of Agriculture and Life Sciences, Kangwon National University, Chuncheon, South Korea, <sup>2</sup> Department of Microbiology & Immunology, McGill University, Montreal, QC, Canada

## OPEN ACCESS

### Edited by:

Juana Fernández-López,  
Miguel Hernández University of  
Elche, Spain

### Reviewed by:

Guadalupe Virginia Nevárez-Moonillón,  
Autonomous University of  
Chihuahua, Mexico  
Vânia Regina Nicoletti,  
São Paulo State University, Brazil  
Ana C. Pinheiro,  
University of Minho, Portugal

### \*Correspondence:

Deog-Hwan Oh  
deoghwa@kangwon.ac.kr

### Specialty section:

This article was submitted to  
Sustainable Food Processing,  
a section of the journal  
Frontiers in Sustainable Food Systems

**Received:** 10 January 2020

**Accepted:** 16 October 2020

**Published:** 10 November 2020

### Citation:

Ofofu FK, Daliri EB-M, Elahi F, Chelliah R, Lee B-H and Oh D-H (2020) New Insights on the Use of Polyphenols as Natural Preservatives and Their Emerging Safety Concerns. *Front. Sustain. Food Syst.* 4:525810. doi: 10.3389/fsufs.2020.525810

Polyphenols are natural high-valued secondary metabolites of plant that demonstrate strong antimicrobial potential as natural preservatives besides their well-established health promoting benefits. This review highlights the challenges, novel strategies to achieve industrial production of polyphenols and their safe use as potential alternative natural preservative. Since plant extraction presents considerable limitations of high cost due to excessive energy and solvent requirements, climate and long growth cycles, microbial biosynthesis using highly advanced omics techniques and metabolic engineering tools could provide superior alternative for commercial production of environmentally sustainable and cost-effective high-value metabolites in a short time. In spite of the many beneficial effects, some plant metabolites and polyphenol compounds at high dosage are found to be pro-oxidant or mutagenic with toxicity. Due to the controversial findings from sub-chronic and oral toxicity studies, more detailed safety and efficacy studies are needed to substantiate findings. Furthermore, extensive research efforts are required to ascertain optimal dosage for safe use in various foods to avoid potential harmful side effects.

**Keywords:** polyphenols, natural preservatives, microbial biosynthesis, high dose, safety

## INTRODUCTION

Microbial contamination not only causes food loss and public health concerns but also affects the sensory characteristics and overall quality of food products. Spoilage and foodborne pathogenic microorganisms are one of the major causes of enormous economic losses despite several advances in food production technologies, distribution, hygiene standards, and consumer education. Food preservatives are defined as compounds that retard spoilage caused by microorganisms and other means such as oxidation. Several traditional food preservation technologies such as freezing, modified atmosphere packaging, thermal and non-thermal physical treatments (high hydrostatic pressure and pulsed electric fields) have been employed to control spoilage microorganisms and extend shelf life of products. More recently, synthetic chemical preservatives such as, nitrate, nitrite, citric acid, propionate, tartaric acid, sorbate, benzoate, sulfites, to mention a few, approved by regulatory agencies are widely used to control microbial growth (Russell, 1991; Manas and Pagán, 2005; Kuorwel et al., 2011; Silva and Lidon, 2016).

However, prolonged use of synthetic preservatives have been shown to pose severe health problems such as liver and kidney damage, gastrointestinal disorders, asthma, certain cancers, and many allergies (Varraso and Camargo, 2014; Etemadi et al., 2017).

Consumers are becoming increasingly conscious of their health and safety and the negative health impacts of chemically synthesized antimicrobials. Due to the increasing demand for minimally processed and healthier foods, the use of natural antimicrobials rather than synthetic preservatives for food preservation is highly sought after by both consumers and food manufacturers (Rico et al., 2007; Bouarab-Chibane et al., 2019a). Phenolic compounds are well documented for their antibacterial activities, besides their antioxidant, antidiabetic, anti-inflammatory, antihypertensive, anticancer, immune enhancing, and cognitive function that are beneficial for optimal health (Milner and Goldberg, 1994; Sun et al., 2017; Araya-Cloutier et al., 2018). A number of reviews on specific biological activities of polyphenols have been reported elsewhere and can be referred to for more details (Duthie and Brown, 1994; Pasinetti and Ho, 2010; Salehi et al., 2018).

Extracts of plant origin are rich sources of polyphenols. Numerous studies have elucidated the antimicrobial activities of various plant extracts, hence, their potential as target alternatives for synthetic preservatives. Owing to this growing interest, several governments and companies are investing heavily for the development of natural food preservatives (Carocho et al., 2014). Currently, commercial production of polyphenols is largely dependent on plant extraction and chemical synthesis (Figure 1). However, plant extraction has considerable limitations of high cost due to excessive energy and solvent requirements, climate and long growth cycles. Microbial biosynthesis using highly advanced omics techniques and system metabolic engineering tools could provide superior alternative for large scale production of environmentally sustainable and cost-effective high-value metabolites in a short time frame to address the increasing demand of the food and nutraceutical market (Schempp et al., 2017; Ong et al., 2018; Zhao and Li, 2018). In spite of the beneficial effects, some plant metabolites and polyphenol compounds at high dosage are found to be pro-oxidant or mutagenic with toxicity (Cory et al., 2018). Due to the controversial results from sub-chronic and oral toxicity studies, more detailed safety and efficacy studies are needed to substantiate findings. Moreover, extensive research efforts are required to ascertain optimal dosage for safe and beneficial use in various foods to avoid potential harmful side effects. This review highlights the current challenges, novel design and strategies to achieve commercial production of polyphenols using advanced systems metabolic engineering tools and their use as alternative natural food preservatives. Furthermore, the safety and potential health risks associated with high dosage polyphenol consumption in food applications are discussed.

## CHEMISTRY AND CLASSIFICATION OF POLYPHENOLS

Plant-derived antimicrobials are promising natural preservatives considered safe, healthy and possess additional properties from their bioactivity and nutritional value. Traditionally, plant extracts have been consumed by mankind for centuries. Polyphenols are secondary plant metabolites derived from


shikimate and or the polyketide pathway. They vary widely within different plants and play key roles in pigment formation, resisting environmental stresses including defense mechanisms against pathogens or ultraviolet radiation, and acting as chemical messengers. They consist of one or more aromatic rings with hydroxyl groups attached as shown in Figure 2 (Fantini et al., 2015; Lyu et al., 2019).

Polyphenols are highly diverse, numerous and widely distributed in plants. Their structure and content are greatly affected by environmental conditions and plant species. Over 10,000 phenolic compounds are presently identified, with flavonoids constituting the largest group. They are largely found in foods such as fruits, vegetables, whole grains, coffee, wine, tea, and chocolate. Polyphenols are generally categorized according to their source of origin, structural differences, and biological activity. Therefore, polyphenols can be grouped into phenolic acids, flavonoids, lignans, stilbenoids, coumarins, and tannin polymers (Table 1). These naturally occurring compounds exist as glycosides with various sugar molecules and their acylated forms attached at different positions in the carbon skeleton (Naczka and Shahidi, 2004; Tsao, 2010; Liu, 2013). Polyphenol classification and their sources are shown in Table 1.

## ANTIMICROBIAL ACTIVITIES OF POLYPHENOLS

Globally, antibiotic resistance has been one of the serious health problems over the past few decades. Harvey et al. (2015) disclosed that more than 70% of pathogenic bacteria have shown resistance to antibiotic treatment. This has untold public health impact and together with other findings has necessitated the urgent search for novel antimicrobial compounds. Since time immemorial, plants have long been used as food and medicine for humans (Duthie and Brown, 1994). Numerous studies have demonstrated antimicrobial activities from several plants and their extracts (Savoia, 2012). Naturally occurring compounds including phenolic secondary metabolites (polyphenols), terpenoids, alkaloids, and peptides have been reported to exhibit antimicrobial properties (Cowan, 1999; Gibbons, 2008; Obied, 2013; Reichling, 2018). These plant metabolites have been found to elicit both direct and indirect inhibitory activities against efflux pump, biofilm formation, and or quorum sensing (Savoia, 2012).


Recently, (Bouarab-Chibane et al., 2019b) designed a quantitative model to establish the structure-activity relationship between polyphenols and pathogenic or food spoilage bacteria. These authors predicted the mechanisms of toxic action of 35 polyphenols ( $1\text{g L}^{-1}$ ) against Gram-positive (*Staphylococcus aureus*, *Bacillus subtilis*, and *Listeria monocytogenes*) and Gram-negative (*Escherichia coli*, *Pseudomonas aeruginosa*, and *Salmonella enteritidis*) bacteria. Their findings revealed a strong interaction between bacterial cell surface and polyphenols in the exertion of antibacterial activity. This was mainly attributed to the lipophilicity, electronic and charge properties of the polyphenols. Furthermore, they also observed a strain and dose-dependent effect of polyphenols.



They found that *L. monocytogenes* was sensitive to polyphenols whereas *P. aeruginosa* was not. Strong antibacterial effect was demonstrated by 5,8-dihydroxy-1,4-naphthoquinone and butyl gallate against five out of the six bacterial strains. However, further investigations on surface properties of bacteria with other techniques besides adhesion of microorganisms to solvents are needed. Generally, extracts have shown stronger antimicrobial potency than individual polyphenols. In addition, the antimicrobial properties of plant extracts also depend on extraction solvent polarity, composition, concentration, and the type of pathogen used. Different research groups have reported the antimicrobial properties of olive leaf polyphenol extracts (Markin et al., 2003; Pereira et al., 2007; Sudjana et al., 2009; Lee and Lee, 2010; Karygianni et al., 2014; Lim et al., 2016). Wine polyphenols were found to exhibit antibacterial effect against pathogenic *E. coli* and *L. monocytogenes* in fish and could find application in the preservation of fish (Rodríguez-Vaquero et al., 2013). Recently, Aziz and Karboune (2018) extensively reviewed the application of natural antimicrobials from plant (phenolic extracts and essential oils), animal (peptides and lactoperoxidase system), and microbial origin (chitosan, bacteriocins, bacteriophages, fermented ingredients) for prolonging the shelf life of fruits, vegetables, and meat products. Flavonoid rich green tea and grape seed extracts containing flavanols such as epicatechin, catechin, epicatechin gallate, procyanidin oligomers, and epigallocatechin were reported to control the growth of pathogenic bacteria in meat products. Cooked beef samples treated with 1% grape seed

extracts (GSE) significantly reduced *E. coli* O157:H7 and *Salmonella typhimurium* counts and controlled the growth of *L. monocytogenes* and *Aeromonas hydrophila* (Ahn et al., 2007). With minimum inhibitory concentrations (MICs) below 10 µg phenol/mL and 15%, cranberry extract with proanthocyanidins as its main antimicrobial compound showed strong inhibitions against several pathogenic bacteria including *E. coli* O157:H7 EDL 933, *E. coli* ATCC 25922, *L. monocytogenes* HPB 2812, *P. aeruginosa* ATCC 15442, *Klebsiella pneumoniae*, *S. Typhimurium* SL1344, *Enterococcus faecium*, *Enterobacter aerogenes*, and *S. aureus* ATCC 29213 (Sagdica et al., 2006; Côté et al., 2011).

Akhtar et al. (2019) investigated the antimicrobial properties of phenolic extracts of different spices and herbs in minced beef during refrigeration storage (4°C). The methanolic extracts of ginger and coriander showed the highest total phenolic content (TPC) of 70.8 and 69.8 mg GAE/100 g, respectively. They found that ginger and coriander extracts at a concentration of 6% significantly inhibited the growth of *S. aureus* (0.24; 0.32 log<sub>10</sub> CFU/g, respectively) and *E. coli* (0.23; 0.36 log<sub>10</sub> CFU/g, respectively) compared to the antibiotic, gentamycin (0.60; 0.77 log<sub>10</sub> CFU/g, respectively) in minced beef stored at 4°C for 9 days and could be developed as natural preservatives. Ghosia et al. (2019) illustrated the antioxidant and antibacterial properties of the leaves, roots and bark of *daphne mucronata* plant extracts. The antibacterial activity of methanolic extracts of roots, bark and leaves was stronger compared to other solvents. The methanolic, hexane, chloroform and ethyl acetate extracts of the roots against *Acinetobacter baumannii* was 86.95, 78.26, 60.86, and 69.56%,



**FIGURE 2** | Schematic representation of biosynthetic pathway of polyphenols. Green, pink, yellow, purple, and blue areas depicts phenolic acids, lignans, coumarins, stilbenes, and flavonoids, respectively [Adopted with permission from Lyu et al. (2019)]. Copyright 2019 American Chemical Society. Intermediates: PEP, phosphoenolpyruvate; E4P, erythrose4-phosphate; DAHP, 3-deoxy-D-arabino-heptulosonate-7-phosphate; DHQ, 3-dehydroquininate. Enzymes: ARO4/ARO3, DAHP synthase; ARO1, pentafunctional arom protein; ARO2, bifunctional chorismate synthase and flavin reductase; ARO7, chorismate mutase; TRP2, anthranilate synthase; TRP3, indole-3-glycerol-phosphate synthase; ARO10, phenylpyruvate decarboxylase; PDC5, minor isoform of pyruvate decarboxylase; TAL, tyrosine ammonia lyase; PAL, phenylalanine ammonia lyase; C4H, cinnamate 4-hydroxylase; 4CL, p-coumaric acid; CoA ligase; CHS, chalcone synthase. Key enzymes are marked in red.

respectively. The hexane and chloroform root extracts did not show any activity against *E. coli*. However, methanolic and ethyl acetate root extracts showed 85.18 and 62.96% activity, respectively, against *E. coli*. Moreover, *S. aureus* was strongly inhibited by methanolic (84.61%), chloroform (61.53%) and ethyl acetate root extracts (69.23%). Good activity of methanolic leave extracts was exhibited against *A. burmanni* (78.26%), *E. coli* (77.78%) and *S. aureus* (73.07%). They also found moderate activity of methanolic bark extracts against *A. burmanni* (65.21%) and *E. coli* (62.96%). Furthermore, a good antioxidant activity with EC<sub>50</sub> values ranging from 157.82 to 361.61 µg/ml was obtained for the extracts. It is worthy to note the antimicrobial activities of *Azadirachta indica* (*A. indica*) phenolic extracts evaluated by Ouerfelli et al. (2019) on shelf-life stability of raw beef patties during storage at 4°C for 11 days. *A. indica* leaf extracts was found to inhibit microbial growth and retard lipid oxidation of chilled beef patties. *A. indica* extracts showed no activity on *Listeria* and *B. cereus* strains. However, strong antimicrobial activity was observed against *E. coli* and *S. aureus* with an inhibition zone of 21 and 19 mm, respectively. Moderate activity was obtained against *Micrococcus luteus* and *Salmonella paratyphi* with inhibition zones of 12 and 10 mm, respectively. Beef samples treated with 0.7% (w/w) fresh powdered *A. indica* leaves retarded the growth of mesophilic bacteria (<10<sup>4</sup> CFU/g) compared to the control without treatment (4.2 × 10<sup>4</sup> CFU/g) after 11 days refrigeration storage (4°C). Findings from these studies can be used to assess the safety of these extracts in different food applications to ascertain their beneficial use and consumer safety.

## POLYPHENOL EXTRACTION FROM PLANT SOURCES

As mentioned earlier, plants represent an abundant resource for polyphenol-rich metabolites synthesized during their growth and under stressful environmental conditions. Literature is replete with numerous studies on the extraction of polyphenols from diverse plant materials. Generally, these phytochemicals are extracted from either fresh or dried plant sources prior to their utilization as food ingredients or in nutraceutical, pharmaceutical, and cosmetic products. Because of the diverse composition and structural differences of polyphenols in various plant materials and the varying nature of plant matrix, the establishment of a standard protocol for extracting all target metabolites from plant sources remains a huge challenge. The type of solvent, plant to solvent ratio, extraction time, temperature and agitation rate are among several parameters or conditions that affect the efficiency of the extraction process (Jovanović et al., 2017; Asofiei et al., 2019). Recent years has seen the development of more efficient extraction techniques, owing to the low efficiency of conventional methods such as maceration, Soxhlet extraction and percolation with long extraction times, large solvent volume, high temperatures and polyphenol degradation. These extraction methods include ultrasound and microwave assisted, simultaneous ultrasound-microwave assisted, supercritical and pressurized fluid, microwave and enzyme co-assisted, and pulsed electric-field. Compared to








the conventional methods, these novel methods have the advantages of a shorter extraction time, reduced solvent usage, reduced thermolabile polyphenolic compound degradation, less laborious, environmentally friendly, and reduced energy and operational cost (Ameer et al., 2017; Jovanović et al., 2017; Asofiei et al., 2019). Asofiei et al. (2019) developed an efficient semi-continuous microwave assisted extraction of polyphenols from sea buckthorn leaves using a modified standard reactor. The total phenolic content (TPC) increased from 144.66 to 176.66 mg GAE/g using microwave assisted extraction at a solvent flow rate of 6 ml/min. TPC was found to increase with increasing stirring rate using larger magnetic stirrer. In addition, pre-heating of solvent (50% ethanol) at 60°C increased polyphenol extraction efficiency by 80%. Comparatively, the antioxidant capacity of sea buckthorn leaves from microwave assisted extracts (405.88 mg TE/g) was higher than the conventional extracts (354.56 mg TE/g). Furthermore, the composition of phytochemicals was not affected by the optimized semi-continuous microwave assisted process. Compared with the conventional extraction, the developed semi-continuous microwave assisted extraction resulted in a significantly lower energy consumption. However, the optimization of extraction kinetic parameters or conditions, taking into consideration plant matrices and the composition and structure of compounds of interest are some bottlenecks for industrial scale up. In addition to the aforementioned, the impact of climate change and long growth cycles poses significant limitations to polyphenol extraction from plants for commercialization.

## BIOSYNTHESIS AND METABOLIC ENGINEERING FOR PRODUCTION OF POLYPHENOLS

Literature on the biosynthesis of polyphenols has been thoroughly discussed by other authors (Knaggs, 2001; Tsao, 2010; Lyu et al., 2019) and because of that only a summary will be given in this work. In plants and microbes, the shikimate and phenylpropanoid metabolic pathways represent the main routes for polyphenol biosynthesis (Figure 2). The central carbon metabolism is linked to aromatic amino acids biosynthesis via the shikimate pathway. Shikimate intermediates such as chorismate are key precursors for the biosynthesis of various aromatic compounds including phenylalanine, tryptophan, tyrosine and polyphenolic compounds (Herrmann, 1995). Intermediates from the phenylpropanoid pathway such as 4-coumaroyl CoA serves as good precursors for the biosynthesis of complex polyphenols such as flavonoids and stilbenes (Lyu et al., 2019). Hence, gaining a complete understanding of polyphenol biosynthesis pathways and developing efficient strategies for their production is extremely important for the *de novo* synthesis of diverse high value metabolites for industrial applications.






Metabolic engineering is a technique used to manipulate and optimize genetic and regulatory processes in living organisms to increase the production of target metabolites. Metabolically engineered microbes have gained increasing attention in pharmaceutical, nutraceutical and food applications. In recent years, the development of high-performance strains

**TABLE 1** | Classification of polyphenols and their sources [adapted and modified from Papuc et al. (2017)].

Class name and backbone structure	Examples	Sources	References
<b>1. Phenolic acids</b>			
 Benzoic acid	Gallic acid Syringic acid Vanillic acid Protocatechuic acid	Fruits, vegetables and cereals	Rice-Evans et al., 1996; Manach et al., 2004
 Cinnamic acid	<i>p</i> -coumaric acid Caffeic acid Ferulic acid Chlorogenic acid Sinapic acid	Fruits, vegetables, cereals, coffee	Manach et al., 2004; Ofosu et al., 2020
<b>2. Flavonoids</b>			
 Anthocyanin	Cyanidin Pelargonidin Petunidin Malvidin Delphinidin Peonidin	Fruits, vegetables, cereals, red and blue flower petals	Mazza and Francis, 1995; Bravo, 1998; Pietta, 2000
 Flavone	Luteolin Apigenin Chrysin Baicalein	Alfalfa Indian trumpet flower	Kawaii et al., 1999; Chen et al., 2003; El-Shafey and Abdelgawad, 2012; Mishima et al., 2015; Kim et al., 2016
 Flavanone	Hesperetin Naringenin	Citrus and grapefruit peels	Lien et al., 2008; Mullen et al., 2008; Tomas-Navarro et al., 2014
 Flavonol	Quercetin Kaempferol Galangin Fisetin Myricetin Morin	Propolis, honey, fruits, vegetables, cereals	Chan et al., 2007; Bertelli et al., 2012; Kurtagić et al., 2013
 Flavan-3-ol	(+)-Catechin (-)-Epicatechin (-)-Epigallocatechin (-)-Epicatechin-3-gallate (-)-Epigallocatechin-3-gallate	Green tea, black tea, cocoa, cereals	Zaveri, 2006; Subhashini et al., 2010; Gadkari and Balaraman, 2015

(Continued)

TABLE 1 | Continued

Class name and backbone structure	Examples	Sources	References
 <p>Isoflavone</p>	Genistein Genistin Daidzein Daidzin Biochanin A Formononetin	Legumes, red clovers	Wang and Murphy, 1994; Mazur et al., 1998; Dixon and Ferreira, 2002; Tsao et al., 2003; Kim et al., 2012; Kuligowski et al., 2017
 <p>Chalcone</p>	Isoliquiritigenin Flavokawain A Flavokawain B Flavokawain C Gymnogrammene	Apples, flowers, hop, beer	Tsao et al., 2003; Tsao and McCallum, 2009; Aksöz and Ertan, 2012; Suwito et al., 2014
<b>3. Stilbenoids</b> 	<i>trans</i> -Resveratrol <i>trans</i> -Piceatannol <i>trans</i> -Piceid <i>trans</i> -Pterostilbene Cajanotone Cajanamide	Red grapes, wine, blueberries, peanuts, dark chocolate, <i>Cajanus cajan</i> , sorghum	Sanders et al., 2000; Burns et al., 2002; Lyons et al., 2003; Counet et al., 2006; Bröhan et al., 2011; Zhang et al., 2012
<b>4. Lignan</b> 	Sesamin Matairesinol Pinoresinol Medioresinol	Flax seed, sesame seed, fruits, vegetables, legumes and many grains	Mazur et al., 1998; Milder et al., 2005; Peñalvo et al., 2005; Smeds et al., 2007, 2012
<b>5. Coumarins</b> 	Ostruthin Ammoresinol Anthogenol Felamidin Agasyllin	Fruits, flowers, seeds	Basile et al., 2009; Rosselli et al., 2009; Chakthong et al., 2012

has seen tremendous acceleration with the emergence of systems metabolic engineering. This interdisciplinary field employs new tools and strategies from systems biology, synthetic biology and evolutionary engineering with conventional metabolic engineering (Choi et al., 2019). Recent reviews have focused on sustainable biosynthesis of polyphenols in bioengineered yeast; advances in tools and strategies for systems metabolic engineering with significant potential for industrial applications (Choi et al., 2019; Lyu et al., 2019).

Several microorganisms have been engineered to synthesize various polyphenols. The advantages of using bioengineered microorganisms over enzymatic and chemical syntheses for production of natural products include but are not limited to inexpensive cofactor requirement, and specific regioselectivity

and stereoselectivity. *E. coli*, *Saccharomyces cerevisiae* and *Corynebacterium glutamicum* with GRAS (generally regarded as safe) status have been widely used as microbial cell factories to produce diverse high-value compounds. Nevertheless, industrial application of polyphenols produced using metabolic engineered microbes still faces quite a number of challenges. The limitation of low productivity or yield, high toxicity to cell growth or low host tolerance to target compounds and expensive feedstock supply poses significant difficulty for their commercial production (Lyu et al., 2019). The development of highly advanced omics techniques and systems metabolic engineering tools is expected to unveil and provide a complete understanding of polyphenol biosynthesis pathways needed to circumvent microbial biosynthesis productivity challenges. Choi et al.

(2019) provides a thorough description of the selection and improvement of high performance strains using novel design and strategies in genetic and genomic engineering, *in silico* metabolic simulation and high-throughput screening to reconstruct metabolic pathways, increase host tolerance and optimize metabolic fluxes for enhanced target compound production. Chung et al. (2017) used plant aromatic aldehyde synthase (AAS) and uridine diphosphate-dependent glycosyltransferases (UGTs) to synthesize polyphenols in *E. coli*. Using engineered *E. coli* harboring AAS and 12 UGTs, tyrosol, hydroxytyrosol and salidroside were synthesized at 531, 208, and 288 mg/L, respectively. When glucose was used as carbon source, naringenin was produced in *E. coli* at a titer of 29 mg/L (Santos et al., 2011). Other studies have reported the production of flavonoids; kaempferol, quercetin and fisetin using *p*-coumaric acid and L-tyrosine as precursor in *E. coli*, respectively (Leonard et al., 2006; Stahlhut et al., 2015). Rodriguez et al. (2017) reported the synthesis of the flavonoids; naringenin, liquiritigenin, kaempferol, resokaempferol, quercetin, and fisetin in an engineered *S. cerevisiae* directly from glucose. Generally, extracts have shown stronger antimicrobial potency than individual polyphenols. Compared to pure flavonoids and phenolic acids, phenolic compounds (naringenin, phloretic acid, phenylacetaldehyde, and homogentisic acid) from engineered *S. cerevisiae* strain N2 was found to exert more potent antibacterial activity against pathogenic *E. coli* ATCC 25922 and *S. aureus* ATCC 29213 (Ng et al., 2019). N2 extracts of concentrations up to 400 µg showed inhibition zones of 4 and 8.33 mm against *E. coli* ATCC 25922 and *S. aureus* ATCC 29213, respectively. On the contrary, pure naringenin did not show any inhibition against either strains, while 6-prenylnaringenin and 8-prenylnaringenin demonstrated inhibition zones of 4 and 7.67 mm against *S. aureus*, respectively. It is therefore imperative to show the safety of these extracts in human trials to prove their viability as food preservatives. Using engineered *C. glutamicum*, Kallscheuer et al. (2016) produced resveratrol at titers of 5 mg/L independent from endogenous aromatic amino acid pathway. Using a different strategy which avoided ammonia lyase activity, a rate limiting step in microbial synthesis, the authors reversed the β-oxidative phenylpropanoid degradation pathway and could produce polyphenols from cheap benzoic acids. A more recent review by Kogure and Inui (2018) summarizes the potential of metabolically engineered *C. glutamicum* for commercial production of several valuable aromatic and natural products including plant polyphenols at high titer from renewable feedstocks.

## POLYPHENOL SAFETY EVALUATION AND FUTURE PERSPECTIVES

It is a common belief that compounds from natural sources are safe, even at high doses. Thus, the limited evidence in literature regarding toxicity of natural plant extracts. Moreover, the safety aspects on the long term consumption of high doses of polyphenols either as dietary supplements or food additives is not clearly known in humans and needs to be thoroughly

investigated. Although numerous animal studies have confirmed the beneficial role of polyphenols, findings from several sub-chronic and oral toxicity studies, still remain controversial (Table 2). Therefore, assessing the safe levels of polyphenols and other natural compounds for use as food preservatives is crucial for determining potential cytotoxicity. Findings from Yamakoshi et al. (2002) showed a lack acute and subchronic toxicity of proanthocyanin-rich grape seed extracts administered orally in rats at high doses of 2 and 4 g/kg. They found the lethal dose to be higher than 4 g/kg in the acute study on the 14th day of clinical observation. However, the no-observed-adverse effect level (NOAEL) during 90 days subchronic toxicity study was found to be 2 g/kg of GSE (about 240 times the estimated daily proanthocyanidin intake by humans). Grape seed polyphenolic extract (GSPE) has shown high tolerability and found to be safe in animal models fed with chow and extract in a 90 days subchronic toxicity study. No detectable adverse effects were observed after treatment with doses from 200 to 2150 mg/kg/day (Bentivegna and Whitney, 2002; Wang et al., 2008). Moreover, treatment of pre-hypertensive people with GSPE 300 mg/day showed a significant reduction in blood pressure with no observed adverse effects (Sivaprakasapillai et al., 2009). This evidence suggests the safety and tolerability of GSPE for treatment in humans. In addition, Charradi et al. (2018) have reported the safety of high repeated dosing of GSPE (8 and 16 g/kg bw) fed to healthy rats for 2 months. Their findings showed no toxicity and support the potential applications in different foods for biotic or abiotic stress-induced multi-organ dysfunction. European Food Safety Authority (EFSA) Panel on Additives and Products or Substances used in Animal Feed (FEEDAP) has established the safety and efficacy of dry grape extract used as feed flavoring for all animals and categories except dogs at a maximum level of 100 mg/kg complete feed. The panel emphasized no safety concerns for consumers at this level in feeds (EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2016). Despite these results, more detailed safety and efficacy human studies are needed to substantiate findings for other plant extracts.

On the contrary, certain studies have delineated the pro-oxidant and adverse effect of plant metabolites and polyphenol compounds (Galati et al., 2002; Shao et al., 2003). Thus, it is imperative to mention that natural compounds may not be safe as previously thought at high concentrations and cause deleterious effects to humans (Li et al., 2008). Higher doses of GSPE (100 or 500 µg/mL) showed pro-oxidant effects and toxicity to cells (Shao et al., 2003). According to a recent systematic review on toxicological studies on green tea extracts (rich in catechin phenolic profiles) by Oketch-Rabah et al. (2020), adverse effects of hepatotoxicity was associated with (–)-epigallocatechin-3-gallate (EGCG) consumption from 140 mg to 1,000 mg/day. The authors also concluded that individual genetic susceptibility and underlying liver health could contribute to these effects and that product label should include caution of usage.

Therefore, extensive research efforts in humans are required to ascertain optimal dosage for safe use to avoid potential harmful side effects. Likewise, recent advances in omics tools and strategies, together with more carefully designed animal



**TABLE 2 |** *In-vivo* and *in-vitro* studies on the safety and toxicity potential of polyphenol consumption.

Plant source	Major polyphenol	Duration of intervention/Cell lines	Purpose of the study	Clinical outcomes	References
Grape seed extracts (GSE)	Proanthocyanidin	90 days	Examined GSE for acute and subchronic oral toxicity in Fischer 344 rats and their mutagenicity potential.	No evidence of mutagenicity and acute oral toxicity at doses of 2 and 4 g/kg. 2% (w/w) GSE was the NOAEL in subchronic toxicity study. Lack of toxicity of GSE.	Yamakoshi et al., 2002
Unripe Apple polyphenol extract	Procyanidins, epicatechin, catechin, chlorogenic acid, phloridzin	90 days	Evaluated the mutagenicity, genotoxicity, the acute oral and subchronic toxicity of Applephenon in Sprague-Dawley rats.	Lethal dose of >2 g/kg in acute oral toxicity in both male and female rats. No abnormal hematological, clinical, histopathological or urinary effects at a dose of 2 g/kg in subchronic oral toxicity study.	Shoji et al., 2004
<i>Capparis spinosa</i> L. leave extracts	Gallic acid, caffeic acid, coumaric acid, ferulic acid, chlorogenic acid, rutin, quercetin	3 days	Investigated the oral acute toxicity of polyphenolic extract of <i>Capparis spinosa</i> L. leaves in female rats and <i>in vitro</i> antibacterial effect.	No adverse effect or mortality at doses of 100 mg/kg. The extract showed antibacterial activity against <i>S. aureus</i> (12 mm), <i>B. subtilis</i> (11 mm), <i>E. coli</i> (12 mm), <i>P. aeruginosa</i> (12 mm).	Oudah et al., 2019
Seventeen freshly prepared plant extracts, two commercial plant extracts, naringin, kaempferol and resveratrol.	Not provided	HepG2, Caco-2, A549, HMEC-1, and 3T3 cell lines.	Screened the toxicity potential of 22 polyphenol-rich compounds on mitochondrial membrane potential, cell membrane integrity and nuclear size using high content analysis.	Buckthorn bark, walnut husk and hollyhock extracts showed high cytotoxicity to mitochondrion and cell membrane but not the nucleus. On the contrary, spent hop and kale leaf extracts showed low cytotoxicity to mitochondrial membrane, cell membrane integrity and nuclear area. Kaempferol exerted strong toxicity on mitochondrial and nuclear compartments while naringin showed least cytotoxicity.	Boncler et al., 2017
Green tea	Epigallocatechin gallate (EGCG)	30 days	Examined the safety and pharmacokinetics of chronic green tea polyphenol or polyphenon E consumption in healthy men and women with Fitzpatrick skin type II or III.	Mild effects of excess gas, stomach upsets, heartburn, abdominal pain, headache, dizziness and muscle pain at doses of 800 mg EGCG once/day or 400 mg EGCG twice/day in treated group. Daily dose of 800 mg EGCG was safe and well-tolerated and resulted in >60% increase in systemic EGCG but did not provide protection against UV-induced erythema.	Chow et al., 2003
Green tea	Epigallocatechin-3- gallate (EGCG), propyl gallate, epicatechin-3-gallate, epigallocatechin, epicatechin, gallic acid	1 day/rats hepatocytes	Examined the cytotoxicities of major tea phenolics on isolated rat hepatocytes and its hepatotoxicity in mice administered intraperitoneally.	Tea phenolics showed cytotoxicity on hepatocytes. Epigallocatechin-3- gallate exerted strong toxicity on mitochondrial membrane collapse and inducing ROS formation. Tea phenolics at doses of 50–800 mg/kg caused liver injury in mice after 24 h.	Galati et al., 2006
Green tea extract	Epigallocatechin gallate	13 weeks	Evaluated the dermal, acute and short-term safety studies of EGCG in rats and dogs.	Lethal dose of 2 g/kg, no oral toxicity at a dose of 0.2 g/kg in rats. The NOAEL was 0.5 g/kg/day for subchronic toxicity study.	Isbrucker et al., 2006
Herbal mixture	Gallic acid, Cinnamic acid, Coumarin, Benzoic acid, Paoniflorin, Albiflorin, Amygdalin, Oxypaeoniflorin	13 weeks	Assessed the safety of oral administration of traditional herbal formula of <i>Gyejibokryeong-hwan</i> in male and female Sprague-Dawley (SD) rats.	At a dose of 5 g/kg/day, ulcers and dilated stomach glands were observed in males, while squamous cell hyperplasia and epithelial atrophy in stomach were observed in both male and female rats. Hematological examinations showed a significant increase in white blood cells, neutrophils and fibrinogen at 5 g/kg/day. The NOAEL was 2 g/kg/day.	Jin et al., 2018
Grape seed and skin extracts	Proanthocyanidin	3 months	Investigated the subchronic oral toxicity of grape seed extracts (GSE) and grape skin extracts (GSKE) in SD rats.	No significant histological changes were observed in all tissues. The NOAEL for male rats was 1.78 g/kg/day GSE or GSKE and 2.15 g/kg/day GSE or GSKE in female rats.	Bentivegna and Whitney, 2002

(Continued)

TABLE 2 | Continued

Plant source	Major polyphenol	Duration of intervention/Cell lines	Purpose of the study	Clinical outcomes	References
Grape seed powder (GSP)	Procyanidins, catechin, epicatechin	2 months	Evaluated the antioxidant, anti-inflammatory effects and the subchronic toxicity of GSP administered orally in Wistar rats.	No adverse effects at the high dose of 16 g/kg. GSP reduced lipoperoxidation, increased antioxidant enzyme activities (CAT, GPx and SOD), reduced plasma IL17A and CRP, increased IL10 and adiponectin, improved heart and renal microcirculation.	Charradi et al., 2018
<i>Morus nigra</i> L. leaves	Quercetin, caffeic acid	28 days	Examined the phytochemical composition of ethanolic extracts of <i>Morus nigra</i> L. leaves, their oral acute and subacute toxicities in male and female Wistar rats.	Lethal dose of 2 g/kg was observed in renal and hepatic organs in acute studies. The NOAEL in male rats was 0.75 g and 1 g/kg and 1 g/kg in female rats.	Figueredo et al., 2018
Grape seed extract	Proanthocyanidin	Chick cardiomyo-cytes.	Investigated the effects of grape seed proanthocyanidin extract (GSPE) on ROS generation, cell survival, LDH and caspase-3 activity using chick cardiomyocytes.	Higher doses (100 or 500 $\mu$ g/mL) of GSPE increased ROS generation, LDH release and caused cell death. At 500 $\mu$ g/mL, caspase-3 activity was significantly increased.	Shao et al., 2003
<i>Vernonia mespilifolia</i> Less.	Not provided	28 days	Evaluated the acute and subacute toxicity of aqueous extract of <i>Vernonia mespilifolia</i> Less in male and female Wistar rats.	Lethality dose was > 5 g/kg in acute toxicity (single dose) study. No evidence of heart, hepatic and renal toxicity at doses up to 0.6 g/kg in subacute study.	Unuofin et al., 2018
<i>Olea europaea</i> L. (olive tree)	Oleuropein, hydroxytyrosol, luteolin-7-glucoside, apigenin-7-glucoside, verbascoside	90 days	Investigated the genotoxicity and repeated-dose oral toxicity of water-soluble extract of olive tree leaves in male and female Wistar rats.	No evidence of mutagenicity and genotoxicity in mice micronucleus up to doses of 2 g/kg/day. No toxic effects or mortality in both male and female rats at high dose of 1 g/kg/day in subchronic study.	Clewell et al., 2016
<i>Olea europaea</i> L.	Oleuropein, Verbascoside, hydroxytyrosol, Rutin, Oleanolic acid	28 days	Investigated the acute and subacute oral toxicities of olive leaves ethanolic extracts in male and female Wistar rats.	No adverse effects or mortality was observed at a single dose of 2 g/kg. Subacute repeated doses up to 0.4 g/kg showed no toxicities in both male and female rats.	Guex et al., 2018
<i>Olea europaea</i> L.	Oleuropein, Verbascoside, hydroxytyrosol, Rutin, Oleanolic acid	6 weeks	Evaluated the effect of olive leaf extracts (OLE) on liver and kidney toxicity in Wistar albino rats.	Dosage of 0.9% OLE showed hepatocellular and renal abnormalities.	Omer et al., 2012
<i>Olea europaea</i> L.	Oleuropein, Verbascoside, hydroxytyrosol, Rutin, Ligstroside	5 weeks	Assessed the effect of OLE administered orally on busulfan (BU) induced damages in rat testes and the safety profile.	OLE at doses of 0.25, 0.5 and 0.75 g/kg repaired defects in rats testes. 0.25 and 0.5 g/kg OLE significantly reduced apoptotic spermatogonia cells. However, OLE at a dose of 0.75 g/kg increased markers of liver damages (ALP and AST).	Hakemi et al., 2019
<i>Campomanesia guazumifolia</i> (Cambess.) O. Berg.	quercetin pentose, quercetin deoxyhexoside, myricetin deoxy-hexoside, quinic acid	28 days	Evaluated the toxicity and anti-inflammatory activities of leave extracts of <i>C. guazumifolia</i> in female mice.	Lethal dose of > 5 g/kg in acute oral toxicity. Doses of up to 1 g/kg did not show any histological and hematological toxicity in subacute study. Doses of 0.3 g and 0.7 g/kg showed anti-inflammatory activity by reducing mechanical hyperalgesia, leukocyte migration and extravasation protein in pleural cavity.	Catelan et al., 2018

LDH, Lactate dehydrogenase; SOD, superoxide dismutase; GPx, glutathione peroxidase; CAT, catalase; NOAEL, No Observed Adverse Effect Level; IL17A, interleukin-17A; IL10, Interleukin-10; ROS, Reactive oxygen species; AST, Aspartate aminotransferase; ALP, alkaline phosphatase.

studies will contribute immensely to provide clearer insight into emerging safety concerns. Further works on understanding structure-activity relationship of natural antimicrobials and their mechanisms of action may help facilitate their use in different food systems, thus mitigating antibiotic resistance. Finally, in food applications regulatory authorities should assess and ensure acceptable levels for long term use to guarantee consumer safety.

## CONCLUSION

Polyphenols are natural high-valued secondary metabolites with well-established benefits in human health. This had led to an increased intake of dietary polyphenol supplements. Moreover, due to the surge in demand for minimally processed and healthier foods, polyphenols have gained tremendous interest as promising alternative natural antimicrobial preservatives for inactivating spoilage and pathogenic microorganisms and to enhance microbial safety. Therefore, gaining a complete understanding of polyphenol biosynthesis pathways using highly advanced omics techniques and systems metabolic engineering tools could help circumvent microbial biosynthesis productivity challenges and provide superior alternative to plant

extraction and chemical synthesis for large scale production of environmentally sustainable and cost-effective high-value metabolites in a short time frame to address the increasing demand of the food and nutraceutical market. Thus, phenolic compounds producing GRAS microbes could be promising source as natural food preservatives. Hence, direct or indirect incorporation of these antimicrobial extracts into foods or their packaging materials should be done with caution to ensure consumer safety since safe levels are not fully known. However, the controversial findings from sub-chronic and oral toxicity animal studies require that more detailed safety and efficacy human studies be conducted. Furthermore, extensive research efforts are required to ascertain optimal dosage for safe use in various foods that are beneficial to avoid potential harmful side effects.

## AUTHOR CONTRIBUTIONS

This review manuscript was conceived and written by FO, ED, and FE. FO, ED, and RC revised the manuscript. This work was revised for its intellectual content by BH-L and D-HO. All authors read and approved the final manuscript.

## REFERENCES

- Ahn, J., Grun, I. U., and Mustapha, A. (2007). Effects of plant extracts on microbial growth, color change, and lipid oxidation in cooked beef. *Food Microbiol.* 24, 7–14. doi: 10.1016/j.fm.2006.04.006
- Akhtar, S., Waseem, M., Ahmad, N., Ismail, T., Ahmad, Z., Manzoor, M. F., et al. (2019). Polyphenol-rich extracts of traditional culinary spices and herbs and their antibacterial activity in minced beef. *J. Food Qual.* 2019:1702086. doi: 10.1155/2019/1702086
- Aksöz, B. E., and Ertan, R. (2012). Spectral properties of chalcones II. *FABAD J. Pharm. Sci.* 37, 205–216.
- Ameer, K., Shahbaz, H. M., and Kwon, J. H. (2017). Green extraction methods for polyphenols from plant matrices and their byproducts: a review. *Compr. Rev. Food Sci. Food Saf.* 16, 295–315. doi: 10.1111/1541-4337.12253
- Araya-Cloutier, C., Vincken, J.-P., Van Ederen, R., Den Besten, H. M., and Gruppen, H. (2018). Rapid membrane permeabilization of *Listeria monocytogenes* and *Escherichia coli* induced by antibacterial prenylated phenolic compounds from legumes. *Food Chem.* 240, 147–155. doi: 10.1016/j.foodchem.2017.07.074
- Asofei, I., Calinescu, I., Trifan, A., and Gavrila, A. I. (2019). A Semi-continuous process for polyphenols extraction from Sea Buckthorn Leaves. *Sci. Rep.* 9, 1–7. doi: 10.1038/s41598-019-48610-6
- Aziz, M., and Karboune, S. (2018). Natural antimicrobial/antioxidant agents in meat and poultry products as well as fruits and vegetables: a review. *Crit. Rev. Food Sci. Nutr.* 58, 486–511. doi: 10.1080/10408398.2016.1194256
- Basile, A., Sorbo, S., Spadaro, V., Bruno, M., Maggio, A., Faraone, N., et al. (2009). Antimicrobial and antioxidant activities of coumarins from the roots of *Ferulago campestris* (Apiaceae). *Molecules* 14, 939–952. doi: 10.3390/molecules14030939
- Bentivegna, S., and Whitney, K. (2002). Subchronic 3-month oral toxicity study of grape seed and grape skin extracts. *Food Chem. Toxicol.* 40, 1731–1743. doi: 10.1016/S0278-6915(02)00155-2
- Bertelli, D., Papotti, G., Bortolotti, L., Marazzan, G. L., and Plessi, M. (2012). 1H-NMR simultaneous identification of health-relevant compounds in propolis extracts. *Phytochem. Anal.* 23, 260–266. doi: 10.1002/pca.1352
- Boncler, M., Golanski, J., Lukasiak, M., Redzynia, M., Dastych, J., and Watala, C. (2017). A new approach for the assessment of the toxicity of polyphenolic compounds with the use of high content screening analysis. *PLoS ONE* 12:e0180022. doi: 10.1371/journal.pone.0180022
- Bouarab-Chibane, L., Degraeve, P., Ferhout, H., Bouajila, J., and Oulahal, N. (2019a). Plant antimicrobial polyphenols as potential natural food preservatives. *J. Sci. Food Agric.* 99, 1457–1474. doi: 10.1002/jsfa.9357
- Bouarab-Chibane, L., Forquet, V., Lantéri, P., Clément, Y., Léonard-Akkari, L., Oulahal, N., et al. (2019b). Antibacterial properties of polyphenols: characterization and qsar (quantitative structure-activity relationship) models. *Front. Microbiol.* 10:829. doi: 10.3389/fmicb.2019.00829
- Bravo, L. (1998). Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr. Rev.* 56, 317–333. doi: 10.1111/j.1753-4887.1998.tb01670.x
- Bröhan, M., Jerkovic, V., and Collin, S. (2011). Potentiality of red sorghum for producing stilbenoid-enriched beers with high antioxidant activity. *J. Agric. Food Chem.* 59, 4088–4094. doi: 10.1021/jf1047755
- Burns, J., Yokota, T., Ashihara, H., Lean, M. E., and Crozier, A. (2002). Plant foods and herbal sources of resveratrol. *J. Agric. Food Chem.* 50, 3337–3340. doi: 10.1021/jf0112973
- Côté, J., Caillet, S., Doyon, G., Dussault, D., Sylvain, J. F., and Lacroix, M. (2011). Antimicrobial effect of cranberry juice and extracts. *Food Control* 22, 1413–1418. doi: 10.1016/j.foodcont.2011.02.024
- Carocho, M., Barreiro, M. F., Morales, P., and Ferreira, I. C. (2014). Adding molecules to food, pros and cons: a review on synthetic and natural food additives. *Compr. Rev. Food Sci. Food Saf.* 13, 377–399. doi: 10.1111/1541-4337.12065
- Catelan, T. B. S., Radai, J. A. S., Leitão, M. M., Branquinho, L. S., de Paula Vasconcelos, P. C., Heredia-Vieira, S. C., et al. (2018). Evaluation of the toxicity and anti-inflammatory activities of the infusion of leaves of *Campomanesia guazumifolia* (Cambess.) O. Berg. *J. Ethnopharmacol.* 226, 132–142. doi: 10.1016/j.jep.2018.08.015
- Chakthong, S., Wearyee, P., Puangphet, P., Mahabusarakam, W., Plodpai, P., Voravuthikunchai, S. P., et al. (2012). Alkaloid and coumarins from the green fruits of *Aegle marmelos*. *Phytochemistry* 75, 108–113. doi: 10.1016/j.phytochem.2011.11.018

- Chan, P. C., Xia, Q., and Fu, P. P. (2007). Ginkgo biloba leaf extract: biological, medicinal, and toxicological effects. *J. Environ. Sci. Health C* 25, 211–244. doi: 10.1080/10590500701569414
- Charradi, K., Mahmoudi, M., Bedhafi, T., Jebari, K., El May, M. V., Limam, F., et al. (2018). Safety evaluation, anti-oxidative and anti-inflammatory effects of subchronically dietary supplemented high dosing grape seed powder (GSP) to healthy rat. *Biomed. Pharmacother.* 107, 534–546. doi: 10.1016/j.biopha.2018.08.031
- Chen, L. J., Games, D. E., and Jones, J. (2003). Isolation and identification of four flavonoid constituents from the seeds of *Oroxylum indicum* by high-speed counter-current chromatography. *J. Chromatogr. A* 988, 95–105. doi: 10.1016/S0021-9673(02)01954-4
- Choi, K. R., Jang, W. D., Yang, D., Cho, J. S., Park, D., and Lee, S. Y. (2019). Systems metabolic engineering strategies: integrating systems and synthetic biology with metabolic engineering. *Trends Biotechnol.* 37, 817–837. doi: 10.1016/j.tibtech.2019.01.003
- Chow, H. S., Cai, Y., Hakim, I. A., Crowell, J. A., Shahi, F., Brooks, C. A., et al. (2003). Pharmacokinetics and safety of green tea polyphenols after multiple-dose administration of epigallocatechin gallate and polyphenon E in healthy individuals. *Clin. Cancer Res.* 9, 3312–3319.
- Chung, D., Kim, S. Y. and Ahn, J. (2017). Production of three phenylethanoids, tyrosol, hydroxytyrosol, and salidroside, using plant genes expressing in *Escherichia coli*. *Sci. Rep.* 7, 1–8. doi: 10.1038/s41598-017-02042-2
- Clewell, A. E., Béres, E., Vértési, A., Glávits, R., Hirka, G., Endres, J. R., et al. (2016). A comprehensive toxicological safety assessment of an extract of *Olea europaea* L. leaves (bonolive™). *Int. J. Toxicol.* 35, 208–221. doi: 10.1177/1091581815619764
- Cory, H., Passarelli, S., Szeto, J., Tamez, M., and Mattei, J. (2018). The role of polyphenols in human health and food systems: a mini-review. *Front. Nutr.* 5:87. doi: 10.3389/fnut.2018.00087
- Counet, C., Callemien, D., and Collin, S. (2006). Chocolate and cocoa: new sources of trans-resveratrol and trans-piceid. *Food Chem.* 98, 649–657. doi: 10.1016/j.foodchem.2005.06.030
- Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* 12, 564–582. doi: 10.1128/CMR.12.4.564
- Dixon, R. A., and Ferreira, D. (2002). Genistein. *Phytochemistry* 60, 205–211. doi: 10.1016/S0031-9422(02)00116-4
- Duthie, G. G., and Brown, K. M. (1994). “Reducing the risk of cardiovascular disease.” in *Functional Foods: Designer Foods, Pharmafoods, Nutraceuticals*, ed I. Goldberg (New York, NY: Chapman and Hall), 19–38. doi: 10.1007/978-1-4615-2073-3\_2
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed) (2016). Scientific opinion on the safety and efficacy of dry grape extract when used as a feed flavouring for all animal species and categories. *EFSA J.* 14:4476. doi: 10.2903/j.efsa.2016.4476
- El-Shafey, N. M., and Abdelgawad, H. (2012). Luteolin, a bioactive flavone compound extracted from *Cichorium endivia* L. subsp. *divaricatum* alleviates the harmful effect of salinity on maize. *Acta Physiol. Plant.* 34, 2165–2177. doi: 10.1007/s11738-012-1017-8
- Etemadi, A., Sinha, R., Ward, M. H., Graubard, B. I., Inoue-Choi, M., Dawsey, S. M., et al. (2017). Mortality from different causes associated with meat, heme iron, nitrates, and nitrites in the NIH-AARP Diet and Health Study: population based cohort study. *BMJ* 357:j1957. doi: 10.1136/bmj.j1957
- Fantini, M., Benvenuto, M., Masuelli, L., Frajese, G. V., Tresoldi, I., Modesti, A., et al. (2015). *In vitro* and *in vivo* antitumoral effects of combinations of polyphenols, or polyphenols and anticancer drugs: perspectives on cancer treatment. *Int. J. Mol. Sci.* 16, 9236–9282. doi: 10.3390/ijms16059236
- Figueredo, K. C., Guex, C. G., Reginato, F. Z., da Silva, A. R. H., Cassanego, G. B., Lhamas, C. L., et al. (2018). Safety assessment of *Morus nigra* L. leaves: acute and subacute oral toxicity studies in Wistar rats. *J. Ethnopharmacol.* 224, 290–296. doi: 10.1016/j.jep.2018.05.013
- Gadkari, P. V., and Balaraman, M. (2015). Catechins: sources, extraction and encapsulation: a review. *Food Bioprod. Process.* 93, 122–138. doi: 10.1016/j.fbp.2013.12.004
- Galati, G., Lin, A., Sultan, A. M., and O'Brien, P. J. (2006). Cellular and *in vivo* hepatotoxicity caused by green tea phenolic acids and catechins. *Free Rad. Biol. Med.* 40, 570–580. doi: 10.1016/j.freeradbiomed.2005.09.014
- Galati, G., Sabzevari, O., Wilson, J. X., and O'Brien, P. J. (2002). Prooxidant activity and cellular effects of the phenoxyl radicals of dietary flavonoids and other polyphenolics. *Toxicology* 177, 91–104. doi: 10.1016/S0300-483X(02)00198-1
- Ghosia, L., Asma, S., Kafel, A., and Jamila, H. (2019). Phytochemical screening, antioxidant and antibacterial properties of *daphne mucronata*. *J. Trad. Chin. Med.* 39, 764–771.
- Gibbons, S. (2008). Phytochemicals for bacterial resistance—strengths, weaknesses and opportunities. *Planta Med.* 74, 594–602. doi: 10.1055/s-2008-1074518
- Guex, C. G., Reginato, F. Z., Figueredo, K. C., da da Silva, A. R. H., Pires, F. B., da Silva Jesus, R., et al. (2018). Safety assessment of ethanolic extract of *Olea europaea* L. leaves after acute and subacute administration to Wistar rats. *Regul. Toxicol. Pharmacol.* 95, 395–399. doi: 10.1016/j.yrtph.2018.04.013
- Hakemi, S. G., Shariffar, F., Haghpanah, T., Babae, A., and Eftekhari-Vaghefi, S. H. (2019). The effects of olive leaf extract on the testis, sperm quality and testicular germ cell apoptosis in male rats exposed to busulfan. *Int. J. Fertil. Steril.* 13:57–65. doi: 10.22074/ijfs.2019.5520
- Harvey, A. L., Edrada-Ebel, R., and Quinn, R. J. (2015). The re-emergence of natural products for drug discovery in the genomics era. *Nat. Rev. Drug Discov.* 14, 111–129. doi: 10.1038/nrd4510
- Herrmann, K. M. (1995). The shikimate pathway: early steps in the biosynthesis of aromatic compounds. *Plant Cell* 7:907. doi: 10.2307/3870046
- Isbrucker, R., Edwards, J., Wolz, E., Davidovich, A., and Bausch, J. (2006). Safety studies on epigallocatechin gallate (EGCG) preparations. Part 2: dermal, acute and short-term toxicity studies. *Food Chem. Toxicol.* 44, 636–650. doi: 10.1016/j.fct.2005.11.003
- Jin, S. E., Seo, C. S., Lee, M. Y., Shin, H.-K., Yang, M.-J., and Ha, H. (2018). Sub-chronic toxicity of Gyejibokryeong-hwan in Sprague-Dawley rats. *J. Ethnopharmacol.* 224, 441–450. doi: 10.1016/j.jep.2018.06.017
- Jovanović, A., Petrović, P., Đorđević, V., Zdunić, G., Šavikin, K., and Bugarski, B. (2017). Polyphenols extraction from plant sources. *Lekovite sirovine* 37, 45–49. doi: 10.5937/lekir1737045J
- Kallscheuer, N., Vogt, M., and Marienhagen, J. (2016). A novel synthetic pathway enables microbial production of polyphenols independent from the endogenous aromatic amino acid metabolism. *ACS Synth. Biol.* 6, 410–415. doi: 10.1021/acssynbio.6b00291
- Karygianni, L., Cecere, M., Skaltsounis, A. L., Argyropoulou, A., Hellwig, E., Aliagiannis, N., et al. (2014). High-level antimicrobial efficacy of representative Mediterranean natural plant extracts against oral microorganisms. *Biomed Res. Int.* 2014:839019. doi: 10.1155/2014/839019
- Kawai, S., Tomono, Y., Katase, E., Ogawa, K., and Yano, M. (1999). Quantitation of flavonoid constituents in citrus fruits. *J. Agric. Food Chem.* 47, 3565–3571. doi: 10.1021/jf990153
- Kim, E.-H., Kim, S.-L., Kim, S.-H., and Chung, I.-M. (2012). Comparison of isoflavones and anthocyanins in soybean [*Glycine max* (L.) Merrill] seeds of different planting dates. *J. Agric. Food Chem.* 60, 10196–10202. doi: 10.1021/jf3031259
- Kim, J.-H., Ahn, D., Eun, J., and Moon, S. (2016). Antioxidant effect of extracts from the coffee residue in raw and cooked meat. *Antioxidants* 5:21. doi: 10.3390/antiox5030021
- Knaggs, A. R. (2001). The biosynthesis of shikimate metabolites. *Nat. Prod. Rep.* 18, 334–355. doi: 10.1039/b001717p
- Kogure, T., and Inui, M. (2018). Recent advances in metabolic engineering of *Corynebacterium glutamicum* for bioproduction of value-added aromatic chemicals and natural products. *Appl. Microbiol. Biotechnol.* 102, 8685–8705. doi: 10.1007/s00253-018-9289-6
- Kuligowski, M., Pawłowska, K., Jasińska-Kuligowska, I., and Nowak, J. (2017). Isoflavone composition, polyphenols content and antioxidative activity of soybean seeds during tempeh fermentation. *CyTA J. Food* 15, 27–33. doi: 10.1080/19476337.2016.1197316
- Kuorwel, K. K., Cran, M. J., Sonneveld, K., Miltz, J., and Bigger, S. W. (2011). Essential oils and their principal constituents as antimicrobial agents for synthetic packaging films. *J. Food Sci.* 76, 164–177. doi: 10.1111/j.1750-3841.2011.02384.x
- Kurtagić, H., Redžić, S., Memić, M., and Sulejmanović, J. (2013). Identification and quantification of Quercetin, Naringenin and Hesperetin by RP LC-DAD in honey samples from BandH. *Bull. Chem. Technol. Bosnia Herzegovina* 40, 25–30.

- Lee, O.-H., and Lee, B.-Y. (2010). Antioxidant and antimicrobial activities of individual and combined phenolics in *Olea europaea* leaf extract. *Bioresour. Technol.* 101, 3751–3754. doi: 10.1016/j.biortech.2009.12.052
- Leonard, E., Yan, Y., and Koffas, M. A. (2006). Functional expression of a P450 flavonoid hydroxylase for the biosynthesis of plant-specific hydroxylated flavonols in *Escherichia coli*. *Metab. Eng.* 8, 172–181. doi: 10.1016/j.ymben.2005.11.001
- Li, N., Liu, J.-H., Zhang, J., and Yu, B.-Y. (2008). Comparative evaluation of cytotoxicity and antioxidative activity of 20 flavonoids. *J. Agric. Food Chem.* 56, 3876–3883. doi: 10.1021/jf073520n
- Lien, T. F., Yeh, H. S., and Su, W. T. (2008). Effect of adding extracted hesperetin, naringenin and pectin on egg cholesterol, serum traits and antioxidant activity in laying hens. *Arch. Anim. Nutr.* 62, 33–43. doi: 10.1080/17450390701780318
- Lim, A., Subhan, N., Jazayeri, J. A., John, G., Vanniasinkam, T., and Obied, H. K. (2016). Plant phenols as antibiotic boosters: *in vitro* interaction of olive leaf phenols with ampicillin. *Phytother. Res.* 30, 503–509. doi: 10.1002/ptr.5562
- Liu, R. H. (2013). Health-promoting components of fruits and vegetables in the diet. *Adv. Nutr.* 4, 384–392. doi: 10.3945/an.112.003517
- Lyons, M. M., Yu, C., Toma, R., Cho, S. Y., Reiboldt, W., Lee, J., et al. (2003). Resveratrol in raw and baked blueberries and bilberries. *J. Agric. Food Chem.* 51, 5867–5870. doi: 10.1021/jf034150f
- Lyu, X., Lee, J., and Chen, W. N. (2019). Potential natural food preservatives and their sustainable production in yeast: terpenoids and polyphenols. *J. Agric. Food Chem.* 67, 4397–4417. doi: 10.1021/acs.jafc.8b07141
- Manach, C., Scalbert, A., Morand, C., Rémésy, C., and Jiménez, L. (2004). Polyphenols: food sources and bioavailability. *Am. J. Clin. Nutr.* 79, 727–747. doi: 10.1093/ajcn/79.5.727
- Manas, P., and Pagán, R. (2005). Microbial inactivation by new technologies of food preservation. *J. Appl. Microbiol.* 98, 1387–1399. doi: 10.1111/j.1365-2672.2005.02561.x
- Markin, D., Duek, L., and Berdicevsky, I. (2003). *In vitro* antimicrobial activity of olive leaves. Antimikrobielle Wirksamkeit von Olivenblättern *in vitro*. *Mycoses* 46, 132–136. doi: 10.1046/j.1439-0507.2003.00859.x
- Mazur, W. M., Duke, J. A., Wähälä, K., Rasku, S., and Adlercreutz, H. (1998). Isoflavonoids and lignans in legumes: nutritional and health aspects in humans. *J. Nutr. Biochem.* 9, 193–200. doi: 10.1016/S0955-2863(97)00184-8
- Mazza, G., and Francis, F. (1995). Anthocyanins in grapes and grape products. *Crit. Rev. Food Sci. Nutr.* 35, 341–371. doi: 10.1080/10408399509527704
- Milder, I. E., Arts, I. C., Van De Putte, B., Venema, D. P., and Hollman, P. C. (2005). Lignan contents of Dutch plant foods: a database including larciresinol, pinoresinol, secoisolaricresinol and matairesinol. *Br. J. Nutr.* 93, 393–402. doi: 10.1079/BJN20051371
- Milner, J., and Goldberg, I. (1994). *Functional Foods*. London: Chapman and Hall.
- Mishima, K., Kawamura, H., Kawakami, R., Ito, S., Inoue, Y., Hirota, T., et al. (2015). Extraction of apigenin and luteolin from leaves of alfalfa (*Medicago sativa* L.) using mixtures of liquid carbon dioxide and ethanol. *Solvent Extr. Res. Dev.* 22, 177–186. doi: 10.15261/serdj.22.177
- Mullen, W., Archeveque, M.-A., Edwards, C. A., Matsumoto, H., and Crozier, A. (2008). Bioavailability and metabolism of orange juice flavanones in humans: impact of a full-fat yogurt. *J. Agric. Food Chem.* 56, 11157–11164. doi: 10.1021/jf801974v
- Naczka, M., and Shahidi, F. (2004). Extraction and analysis of phenolics in food. *J. Chromatogr. A* 1054, 95–111. doi: 10.1016/S0021-9673(04)01409-8
- Ng, K. R., Lyu, X., Mark, R., and Chen, W. N. (2019). Antimicrobial and antioxidant activities of phenolic metabolites from flavonoid-producing yeast: potential as natural food preservatives. *Food Chem.* 270, 123–129. doi: 10.1016/j.foodchem.2018.07.077
- Obied, H. K. (2013). Biography of biophenols: past, present and future. *Funct. Foods Health Dis.* 3, 230–241. doi: 10.31989/ffhd.v3i6.51
- Ofosu, F. K., Elahi, F., Daliri, E. B.-M., Tyagi, A., Chen, X. Q., Chelliah, R., et al. (2020). UHPLC-ESI-QTOF-MS/MS characterization, antioxidant and antidiabetic properties of sorghum grains. *Food Chem.* 337:127788. doi: 10.1016/j.foodchem.2020.127788
- Oketch-Rabah, H. A., Roe, A. L., Rider, C. V., Bonkovsky, H. L., Giancaspro, G. I., Navarro, V., et al. (2020). United States Pharmacopeia (USP) comprehensive review of the hepatotoxicity of green tea extracts. *Toxicol. Rep.* 7, 386–402. doi: 10.1016/j.toxrep.2020.02.008
- Omer, S. A., Eloheid, M., Elamin, M., Hassan, Z., Virk, P., Daghestani, M., et al. (2012). Toxicity of olive leaves (*Olea europaea* L.) in Wistar albino rats. *Asian J. Anim. Vet. Adv.* 7, 1175–1182. doi: 10.3923/ajava.2012.1175.1182
- Ong, K. L., Kaur, G., Pensupa, N., Uisan, K., and Lin, C. S. K. (2018). Trends in food waste valorization for the production of chemicals, materials and fuels: case study South and Southeast Asia. *Bioresour. Technol.* 248, 100–112. doi: 10.1016/j.biortech.2017.06.076
- Oudah, S. K., Al-Salih, R. M., Gusar, S. H., and Roomi, A. B. (2019). Study of the role of polyphenolic extract of *capparis spinosa* L. leaves as acute toxicity and antibacterial agent. *Plant Arch.* 19, 3821–3829.
- Ouerfelli, M., Villasante, J., Kaab, L. B. B., and Almajano, M. P. (2019). Effect of neem (*Azadirachta indica* L.) on lipid oxidation in raw chilled beef patties. *Antioxidants* 8:305. doi: 10.3390/antiox8080305
- Papuc, C., Goran, G. V., Predescu, C. N., Nicorescu, V., and Stefan, G. (2017). Plant polyphenols as antioxidant and antibacterial agents for shelf-life extension of meat and meat products: classification, structures, sources, and action mechanisms. *Compr. Rev. Food Sci. Food Saf.* 16, 1243–1268. doi: 10.1111/1541-4337.12298
- Pasinetti, G. M., and Ho, L. (2010). Role of grape seed polyphenols in Alzheimer's disease neuropathology. *Nutr. Diet.* 2010, 97–103. doi: 10.2147/NDS.S6898
- Peñalvo, J. L., Haajanen, K. M., Botting, N., and Adlercreutz, H. (2005). Quantification of lignans in food using isotope dilution gas chromatography/mass spectrometry. *J. Agric. Food Chem.* 53, 9342–9347. doi: 10.1021/jf051488w
- Pereira, A., Ferreira, I., Marcelino, F., Valentão, P., Andrade, P., Seabra, R., et al. (2007). Phenolic compounds and antimicrobial activity of olive (*Olea europaea* L. Cv. *Cobrançosa*) leaves. *Molecules* 12, 1153–1162. doi: 10.3390/12051153
- Pietta, P.-G. (2000). Flavonoids as antioxidants. *J. Nat. Prod.* 63, 1035–1042. doi: 10.1021/np9904509
- Reichling, J. (2018). Plant–microbe interactions and secondary metabolites with antibacterial, antifungal and antiviral properties. *Ann. Plant Rev.* 39, 214–347. doi: 10.1002/9781119312994.apr0420
- Rice-Evans, C. A., Miller, N. J., and Paganga, G. (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Rad. Biol. Med.* 20, 933–956. doi: 10.1016/0891-5849(95)02227-9
- Rico, D., Martin-Diana, A. B., Barat, J., and Barry-Ryan, C. (2007). Extending and measuring the quality of fresh-cut fruit and vegetables: a review. *Trends Food Sci. Technol.* 18, 373–386. doi: 10.1016/j.tifs.2007.03.011
- Rodriguez, A., Strucko, T., Stahlhut, S. G., Kristensen, M., Svenssen, D. K., Forster, J., et al. (2017). Metabolic engineering of yeast for fermentative production of flavonoids. *Bioresour. Technol.* 245, 1645–1654. doi: 10.1016/j.biortech.2017.06.043
- Rodriguez-Vaquero, M. J., Aredes Aredes-Fernández, P., and Manca De Nadra, M. C. (2013). Phenolic compounds from wine as natural preservatives of fish meat. *Food Technol. Biotechnol.* 51, 376–382.
- Rosselli, S., Maggio, A. M., Faraone, N., Spadaro, V., Morris-Natschke, S. L., Bastow, K. F., et al. (2009). The cytotoxic properties of natural coumarins isolated from roots of *Ferulago campestris* (Apiaceae) and of synthetic ester derivatives of aegelinol. *Nat. Prod. Commun.* 4:1934578X0900401219. doi: 10.1177/1934578X0900401219
- Russell, A. (1991). Mechanisms of bacterial resistance to non-antibiotics: food additives and food and pharmaceutical preservatives. *J. Appl. Bacteriol.* 71, 191–201. doi: 10.1111/j.1365-2672.1991.tb04447.x
- Sagdica, O., Aksoy, A., and Ozkan, G. (2006). Evaluation of the antibacterial and antioxidant potentials of cranberry (*Gilaburu*, *Viburnum Opulus* L.) fruit extract. *Acta Alimentaria* 35, 487–492. doi: 10.1556/AAlim.35.2006.4.12
- Salehi, B., Mishra, A. P., Nigam, M., Sener, B., Kilic, M., Sharifi-Rad, M., et al. (2018). Resveratrol: A double-edged sword in health benefits. *Biomedicines* 6, 91. doi: 10.3390/biomedicines6030091
- Sanders, T. H., Mcmichael, R. W., and Hendrix, K. W. (2000). Occurrence of resveratrol in edible peanuts. *J. Agric. Food Chem.* 48, 1243–1246. doi: 10.1021/jf990737b
- Santos, C. N. S., Koffas, M., and Stephanopoulos, G. (2011). Optimization of a heterologous pathway for the production of flavonoids from glucose. *Metab. Eng.* 13, 392–400. doi: 10.1016/j.ymben.2011.02.002
- Savoia, D. (2012). Plant-derived antimicrobial compounds: alternatives to antibiotics. *Future Microbiol.* 7, 979–990. doi: 10.2217/fmb.12.68

- Schempp, F. M., Drummond, L., Buchhaupt, M., and Schrader, J. (2017). Microbial cell factories for the production of terpenoid flavor and fragrance compounds. *J. Agric. Food Chem.* 66, 2247–2258. doi: 10.1021/acs.jafc.7b00473
- Shao, Z.-H., Hoek, T. L. V., Xie, J., Wojcik, K., Chan, K. C., Li, C.-Q., et al. (2003). Grape seed proanthocyanidins induce pro-oxidant toxicity in cardiomyocytes. *Cardiovasc. Toxicol.* 3, 331–339. doi: 10.1385/CT:3:4:331
- Shoji, T., Akazome, Y., Kanda, T., and Ikeda, M. (2004). The toxicology and safety of apple polyphenol extract. *Food Chem. Toxicol.* 42, 959–967. doi: 10.1016/j.fct.2004.02.008
- Silva, M. M., and Lidon, F. (2016). Food preservatives—an overview on applications and side effects. *Emir. J. Food Agric.* 26, 366–373. doi: 10.9755/efja.2016-04-351
- Sivaprakasapillai, B., Edirisinghe, I., Randolph, J., Steinberg, F., and Kappagoda, T. (2009). Effect of grape seed extract on blood pressure in subjects with the metabolic syndrome. *Metab. Clin. Exp.* 58, 1743–1746. doi: 10.1016/j.metabol.2009.05.030
- Smeds, A. I., Eklund, P. C., Sjöholm, R. E., Willför, S. M., Nishibe, S., Deyama, T., et al. (2007). Quantification of a broad spectrum of lignans in cereals, oilseeds, and nuts. *J. Agric. Food Chem.* 55, 1337–1346. doi: 10.1021/jf0629134
- Smeds, A. I., Eklund, P. C., and Willför, S. M. (2012). Content, composition, and stereochemical characterisation of lignans in berries and seeds. *Food Chem.* 134, 1991–1998. doi: 10.1016/j.foodchem.2012.03.133
- Stahlhut, S. G., Siedler, S., Malla, S., Harrison, S. J., Maury, J., Neves, A. R., et al. (2015). Assembly of a novel biosynthetic pathway for production of the plant flavonoid fisetin in *Escherichia coli*. *Metab. Eng.* 31, 84–93. doi: 10.1016/j.ymben.2015.07.002
- Subhashini, R., Rao, U. M., Sumathi, P., and Gunalan, G. (2010). A comparative phytochemical analysis of cocoa and green tea. *Indian J. Sci. Technol.* 3, 188–192. doi: 10.17485/ijst/2010/v3i2.3
- Sudjana, A. N., D'orazio, C., Ryan, V., Rasool, N., Ng, J., Islam, N., et al. (2009). Antimicrobial activity of commercial *Olea europaea* (olive) leaf extract. *Int. J. Antimicrob. Agents* 33, 461–463. doi: 10.1016/j.ijantimicag.2008.10.026
- Sun, H., Wang, D., Song, X., Zhang, Y., Ding, W., Peng, X., et al. (2017). Natural prenylchalconaringenins and prenylnaringenins as antidiabetic agents:  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibition and *in vivo* antihyperglycemic and antihyperlipidemic effects. *J. Agric. Food Chem.* 65, 1574–1581. doi: 10.1021/acs.jafc.6b05445
- Suwito, H., Jumina, M., Kristanti, A. N., and Puspaningsih, N. N. T. (2014). Chalcones: synthesis, structure diversity and pharmacological aspects. *J. Chem. Pharm. Res.* 6, 1078–1088.
- Tomas-Navarro, M., Vallejo, F., and Tomas-Barberan, F. A. (2014). “Chapter 40: Bioavailability and metabolism of citrus fruit beverage flavanones in humans,” in *Polyphenols in Human Health and Disease*, eds R. R. Watson, V. R. Preedy, and S. Zibadi (San Diego, CA: Academic Press), 537–51. doi: 10.1016/B978-0-12-398456-2.00040-2
- Tsao, R. (2010). Chemistry and biochemistry of dietary polyphenols. *Nutrients* 2, 1231–1246. doi: 10.3390/nu2121231
- Tsao, R., and McCallum, J. (2009). “Chemistry of flavonoids – chapter 5,” in *Fruit and Vegetable Phytochemicals: Chemistry, Nutritional Value and Stability*, eds L. A. de la Rosa, E. Alvarez-Parrilla, and G. Gonzalez-Aguilar (Ames, IA: Blackwell Publishing), 131–53.
- Tsao, R., Yang, R., and Young, J. C. (2003). Antioxidant isoflavones in osage orange, *Maclura pomifera* (Raf.) Schneid. *J. Agric. Food Chem.* 51, 6445–6451. doi: 10.1021/jf0342369
- Unuofin, J. O., Otunola, G. A., and Afolayan, A. J. (2018). Evaluation of acute and subacute toxicity of whole-plant aqueous extract of *Vernonia mespilifolia* Less. in Wistar rats. *J. Integr. Med.* 16, 335–341. doi: 10.1016/j.joim.2018.07.003
- Varraso, R., and Camargo, C. A. (2014). Processed meat consumption and lung health: more evidence for harm. *Eur. Respir. Soc.* 43, 943–946. doi: 10.1183/09031936.00228213
- Wang, H.-J., and Murphy, P. A. (1994). Isoflavone content in commercial soybean foods. *J. Agric. Food Chem.* 42, 1666–1673. doi: 10.1021/jf00044a016
- Wang, J., Ho, L., Zhao, W., Ono, K., Rosensweig, C., Chen, L., et al. (2008). Grape-derived polyphenolics prevent A $\beta$  oligomerization and attenuate cognitive deterioration in a mouse model of Alzheimer's disease. *J. Neurosci.* 28, 6388–6392. doi: 10.1523/JNEUROSCI.0364-08.2008
- Yamakoshi, J., Saito, M., Kataoka, S., and Kikuchi, M. (2002). Safety evaluation of proanthocyanidin-rich extract from grape seeds. *Food Chem. Toxicol.* 40, 599–607. doi: 10.1016/S0278-6915(02)00006-6
- Zaveri, N. T. (2006). Green tea and its polyphenolic catechins: medicinal uses in cancer and noncancer applications. *Life Sci.* 78, 2073–2080. doi: 10.1016/j.lfs.2005.12.006
- Zhang, N. L., Zhu, Y. H., Huang, R. M., Fu, M. Q., Su, Z. W., Cai, J. Z., et al. (2012). Two new stilbenoids from *Cajanus cajan*. *Zeitschrift für Naturforschung B* 67, 1314–1318. doi: 10.5560/znb.2012-0184
- Zhao, Y.-J., and Li, C. (2018). Biosynthesis of plant triterpenoid saponins in microbial cell factories. *J. Agric. Food Chem.* 66, 12155–12165. doi: 10.1021/acs.jafc.8b04657

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Ofosu, Daliri, Elahi, Chelliah, Lee and Oh. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.