



Regulation of the Intestinal Barrier Function by Host Defense Peptides

Kelsy Robinson¹, Zhuo Deng¹, Yongqing Hou² and Guolong Zhang^{1,2*}

¹ Department of Animal Science, Oklahoma State University, Stillwater, OK, USA, ² Hubei Key Laboratory of Animal Nutrition and Feed Science, Wuhan Polytechnic University, Wuhan, China

OPEN ACCESS

Edited by:

Michael Kogut,
USDA – Agricultural Research
Service, USA

Reviewed by:

Kenneth James Genovese,
USDA – Agricultural Research
Service, USA
Ryan Arsenault,
University of Delaware, USA

Matthew Jon Sylte,
USDA – Agricultural Research
Service, USA

*Correspondence:

Guolong Zhang
zguolon@okstate.edu

Specialty section:

This article was submitted to
Veterinary Infectious Diseases,
a section of the journal
Frontiers in Veterinary Science

Received: 25 August 2015

Accepted: 02 November 2015

Published: 23 November 2015

Citation:

Robinson K, Deng Z, Hou Y and
Zhang G (2015) Regulation of the
Intestinal Barrier Function by Host
Defense Peptides.
Front. Vet. Sci. 2:57.
doi: 10.3389/fvets.2015.00057

Intestinal barrier function is achieved primarily through regulating the synthesis of mucins and tight junction (TJ) proteins, which are critical for maintaining optimal gut health and animal performance. An aberrant expression of TJ proteins results in increased paracellular permeability, leading to intestinal and systemic disorders. As an essential component of innate immunity, host defense peptides (HDPs) play a critical role in mucosal defense. Besides broad-spectrum antimicrobial activities, HDPs promotes inflammation resolution, endotoxin neutralization, wound healing, and the development of adaptive immune response. Accumulating evidence has also indicated an emerging role of HDPs in barrier function and intestinal homeostasis. HDP deficiency in the intestinal tract is associated with barrier dysfunction and dysbiosis. Several HDPs were recently shown to enhance mucosal barrier function by directly inducing the expression of multiple mucins and TJ proteins. Consistently, dietary supplementation of HDPs often leads to an improvement in intestinal morphology, production performance, and feed efficiency in livestock animals. This review summarizes current advances on the regulation of epithelial integrity and homeostasis by HDPs. Major signaling pathways mediating HDP-induced mucin and TJ protein synthesis are also discussed. As an alternative strategy to antibiotics, supplementation of exogenous HDPs or modulation of endogenous HDP synthesis may have potential to improve intestinal barrier function and animal health and productivity.

Keywords: host defense peptides, barrier function, tight junction, gut health, innate immunity

INTRODUCTION

The gastrointestinal (GI) tract is lined by a single layer of epithelial cells that serve to facilitate digestion and absorption of nutrients and also act as a barrier to invading microorganisms, toxins, and dietary antigens. Intestinal barrier function is achieved through coating of the epithelial cells with a

Abbreviations: aPKC, atypical protein kinase C; CCR2, CC chemokine receptor 2; CD, Crohn's disease; DSS, dextran sulfate sodium; EGF, epidermal growth factor; EGFR, EGF receptor; ERK, extracellular signal-regulated kinase; ETEC, enterotoxigenic *E. coli*; GI, gastrointestinal; GSK-3, glycogen synthase kinase 3; GUK, guanylate kinase; HBD, human β -defensin; HDPs, host defense peptides; JAM, junction adhesion molecule; LPS, lipopolysaccharides; MAMPs, microbe-associated molecular patterns; MAPK, mitogen-activated protein kinase; MBD, mouse β -defensin; MLCK, myosin light chain kinase; MUC, mucin; P2X7, P2X purinergic receptor 7; PAMPs, pathogen-associated molecular patterns; PBD, porcine β -defensin; PDZ, post-synaptic density 95, disk-large, and zonula occludens; PI3K, phosphatidylinositol-3-kinase; PKA, protein kinase A; PKC, protein kinase C; PRRs, pattern recognition receptors; SEA, sea urchin sperm protein, enterokinase and agrin; TACE, TNF- α -converting enzyme; TEER, transepithelial electrical resistance; TGF, transforming growth factor; TJ, tight junction; TR, tandem repeat; UC, ulcerative colitis; VWF, von Willebrand factor domain; ZO, zonula occludens.

mucus layer and the formation of a selectively permeable barrier across and between epithelial cells (1). The mucus layer consists primarily of mucin glycoproteins that are secreted by goblet cells, functioning as a physical barrier between the luminal contents and the host and also to facilitate nutrient digestion and absorption (2). However, the primary barrier function of the GI tract resides with epithelial cells, which transport water, ions, and macromolecules through either of two routes, i.e., the transcellular and paracellular pathways (1, 3, 4). The transcellular pathway refers to the movement of small molecules through epithelial cells either by active or passive transport, whereas the paracellular pathway refers to the diffusion of water, macromolecules, and immune cells between epithelial cells. In the presence of intact epithelial cells, the paracellular pathway dictates the intestinal permeability and is regulated by inter-epithelial connections known as tight junctions (TJs) (1, 3, 4).

Maintenance of mucin and TJ assembly ensures proper absorption and transport of nutrients, water, and electrolytes, while shielding the host from pathogens, toxins, intestinal microbiota, and dietary antigens. Disruption of the mucus layer and TJ complex, on the other hand, results in an increase in intestinal permeability, followed by heightened bacterial translocation, inflammation, and possibly the onset of various enteric and systemic disorders (1, 3, 4). In livestock production, impaired intestinal barrier function leads to reduced animal health and growth performance (5, 6). Therefore, it is critically important to understand how the intestinal barrier function is maintained and regulated in order to achieve optimal animal health and productivity.

Host defense peptides (HDPs), also known as antimicrobial peptides, are an important component of the animal innate immune system, and a majority of HDPs are expressed on mucosal surfaces, including the GI tract (7, 8). With potent antimicrobial and immunomodulatory activities, HDPs exert a pleiotropic effect on innate adaptive immune responses (9–11). Recent research has further shed light on the direct involvement of epithelial HDPs in regulating intestinal mucin and TJ protein expression and microbiota composition. The focus of this review is to summarize the latest advances regarding the emerging role of HDPs in maintaining intestinal barrier and homeostasis with a goal of exploring HDP-based therapies to improve gut health and performance of food-producing animals.

HOST DEFENSE PEPTIDES: A CRITICAL COMPONENT OF INNATE IMMUNITY

A variety of HDPs with direct antimicrobial activities are produced by host cells in response to infections. Among them are several major families found in vertebrate species such as defensins, cathelicidins, the S100 family, the RNase A superfamily, regenerating islet-derived III (REGIII) C-type lectins, and peptidoglycan-recognition proteins (12, 13). Defensins are primarily identified by three conserved disulfide bridges that form several antiparallel β -sheets due to the presence of multiple cysteine residues (14). Based on the spacing pattern of six cysteines, vertebrate defensins are further categorized into three

subfamilies, including α -, β -, and θ -defensins. While β -defensins are present in all vertebrate animals, α -defensins are found in most but not all mammals, and θ -defensins only exist in primates (14). Cathelicidins are structurally recognized by the highly conserved cathelin domain found in the precursor that is cleaved off to release the biologically active peptides adopting a variety of structures such as α -helix (15).

The S100 family proteins are 9–14 kDa in mass containing two highly conserved Ca^{2+} -binding EF-hand domains that are separated by four α -helical domains with a variable C-terminal region (16, 17). The RNase A superfamily are characterized by the presence of 6–8 conserved cysteines forming distinct disulfide bridges, together with two invariantly spaced histidines and a lysine (18, 19). The REGIII family proteins are a group of soluble C-type lectins with a conserved carbohydrate-recognition domain that binds to sugars in a Ca^{2+} -dependent manner (20). Peptidoglycan-recognition proteins constitute a family of phylogenetically conserved host defense molecules with a PGRP domain that binds to bacterial peptidoglycans through specific interactions with the muramyl-tripeptide fragments (21).

Expression of HDPs

Six α -defensins (22) and a minimum of 39 β -defensins (23) have been reported in humans. The genomes of cattle and pigs encode at least 57 and 29 β -defensin genes, respectively (24, 25), while the chicken genome harbors a total of 14 β -defensin genes (26, 27), with no α -defensins being found in cattle, pigs, or chickens. All β -defensin genes are located in tandem in a single genomic region in the chicken (26) and are expanded to 4–5 different clusters in humans, cattle, and pigs (23–25). Interestingly, all human α -defensin genes form a single cluster within a β -defensin gene cluster (22), suggesting that α -defensins likely diverged from β -defensins. While four human α -defensins (HNP1–4) are abundantly present in neutrophil granules, the other two α -defensins (HD5 and HD6) are specific to Paneth cells in the crypts of the human small intestinal tract (14). On the other hand, a majority of β -defensins are expressed in a wide range of cell types, particularly the epithelial cells lining the skin, GI, respiratory, and urogenital tracts of all livestock species as well as humans (14).

A single cathelicidin known as LL-37, CAMP, or hCAP-18 is present in humans (28) and four cathelicidins are reported in chickens (29, 30). In cattle and pigs, 10 and 11 cathelicidins have been identified, respectively (31, 32). All cathelicidin genes are located in a syntenic chromosomal region in vertebrate species. Expressions of cathelicidins are widespread with abundant presence in neutrophil granules as well as various epithelial mucosal surfaces of cattle, pigs, and humans. Four chicken cathelicidins are similarly expressed in a broad range of tissues as well as in heterophils (29, 30, 33, 34), which are equivalent to neutrophils in mammals. Additionally, chicken cathelicidin-B1 is highly expressed in M cells of the bursa of Fabricius (30), a type of specialized epithelial cells involved primarily in antigen transportation from the intestinal lumen to submucosal immune cells (35).

The S100 family members have been found in all vertebrates (17). A total of 21 S100 proteins are present in humans, with 17 members clustered in the same 2-Mb region on chromosome

1q21 (17). The tissue expression pattern of S100 proteins is unique and isoform specific (16). For example, S100A7 (also known as psoriasin) isolated initially from the skin of psoriatic patients is mainly expressed in the skin and breast tissues, whereas the heterodimer S100A8/S100A9, or calgranulin A/B, is expressed in keratinocytes, neutrophils, monocytes, and macrophages.

Biological Functions of HDPs

Host defense peptides are an integral part of the innate immune system. Historically, HDPs are known for their ability to function as natural antibiotics with broad-spectrum antimicrobial activities against Gram-negative and Gram-positive bacteria, fungi, viruses, protozoa, and even cancerous cells (7, 36). Human LL-37 and α - and β -defensins are all capable of killing a broad spectrum of pathogens (7, 36). All four chicken cathelicidins have been demonstrated to be active at low micromolar concentrations against both Gram-positive and Gram-negative bacteria, including antibiotic-resistant strains (29, 30, 37–39). Several chicken β -defensins are also potent against a range of human and zoonotic pathogens (40–42). Similarly, β -defensins and cathelicidins in the cattle and pigs are broadly active against multiple pathogens as well (32).

Because of the cationic and amphipathic properties associated with a majority of HDPs, they kill bacteria primarily through disruption of cell membranes and/or interaction with intracellular macromolecules (43). A net positive charge allows HDPs to bind to negatively charged phospholipid groups on the bacterial membrane through electrostatic interactions. The amphipathic nature of HDPs facilitates their insertion into target cellular membranes allowing them to disrupt its integrity. Multiple models of membrane disruption, such as “barrel-stave,” “carpet,” or “toroidal-pore” models, have been proposed (43). Intracellularly, certain HDPs are also capable of inhibiting protein, DNA and RNA synthesis, or binding to specific targets (43). Because of their primary membrane-lytic activities, HDPs are generally equally active among drug-resistant and -susceptible pathogens. It is conceivably more difficult for pathogens to develop resistance to HDPs, although certain bacteria have developed mechanisms to resist their action in order to infect and colonize the hosts (44). It appears that commensal bacteria are generally resistant to the action of the constitutively expressed HDPs, but sensitive to certain inducibly expressed HDPs in the human intestinal tract (45). However, the mechanism by which commensal and probiotic bacteria show a reduced sensitivity to HDPs remains elusive.

Besides their antimicrobial activity, HDPs are involved in the modulation of innate and adaptive immune responses (10, 11) (**Figure 1**). Many human HDPs have been shown to promote the recruitment of neutrophils or monocytes and suppress proinflammatory response. Human HDPs also induce the differentiation and activation of macrophages and dendritic cells. Additionally, human cathelicidin LL-37 facilitates the resolution of inflammation by promoting re-epithelialization and wound healing as well as autophagy and apoptosis (10, 11). Three chicken cathelicidins known as fowlicidin 1–3 bind to bacterial lipopolysaccharides (LPS) directly with a strong capacity to neutralize LPS-induced production of inflammatory cytokines in macrophage cells (37–39). Furthermore, chicken

fowlicidin-1 is chemotactic to neutrophils, but not monocytes or lymphocytes (46). Fowlicidin-1 also activates macrophages by inducing modest synthesis of inflammatory cytokines and chemokines and further potentiates the antibody response if co-administered with a model antigen (ovalbumin) in mice (46). Importantly, a single application of fowlicidin-1 is not only able to protect animals from an established infection (47) but also to prevent the disease beyond a 2- to 4-day window (46) in a murine model of methicillin-resistant *Staphylococcus aureus* (MRSA) infection (46).

Among 14 bovine β -defensins examined, three (BNBD3, BNBD9, and EBD) are chemotactic to immature monocyte-derived dendritic cells (48). Porcine cathelicidin PR-39 is also capable of inhibiting phagocyte NADPH oxidase activity and attenuating myocardial ischemia–reperfusion injury (49) by blocking the assembly of the enzyme complex through binding to p47phox, a cytosolic component of the NADPH oxidase (50). PR-39 accelerates wound repair by inducing syndecans (51). Furthermore, PR-39 facilitates angiogenesis and formation of functional blood vessels by inhibiting the ubiquitin–proteasome-dependent degradation of hypoxia-inducible factor (HIF)-1 α (52). Several porcine cathelicidins also help with the update of bacterial DNA and subsequent activation of dendritic cells (53). HDPs with potent antimicrobial activity and the ability to modulate innate and adaptive immunity are, therefore, being actively exploited as novel antibiotics.

Additionally, recent emerging evidence has highlighted the beneficial effect of HDPs on mucosal barrier permeability by directly regulating mucin and TJ protein expression and shaping microbiota composition. This emerging role of HDPs in intestinal barrier function and homeostasis will be the focus of this review.

MUCUS LAYER: A LAYER OF INTIMATE PROTECTION FOR MUCOSAL SURFACE

An intact mucus layer that is composed primarily of secreted mucins plays a critical role in maintaining the intestinal barrier function (54, 55). Mucins are large, highly glycosylated proteins ranging from 0.5 to 20 MDa. Synthesized and released by goblet cells, mucins function to coat the mucosal surface to facilitate the passage of substances, maintain proper cell hydration, act as a permeable barrier for the exchange of gas and nutrients, and also protect the epithelial cells from invading pathogens and toxins (54, 55). Structurally, a hallmark of all mucin protein backbones is the presence of 1–5 tandem repeat (TR) domains, which consist of an excessive number of identical or nearly identical TR sequences rich in serine, threonine, and proline residues (56) (**Figure 2**). The TR domain is heavily glycosylated because of attachment of oligosaccharides to serine and threonine through O-linked glycosylation, giving rise to 50–80% glycans in mass. Saturated sugar coating is beneficial to increase the water-holding capacity and the resistance of mucins to proteolytic cleavage.

In humans, the mucin family consists of up to 20 members, including both secreted and membrane-bound forms. Secreted mucins form homo-oligomeric, gel-like structures constituting the mucosal layer, while the membrane-bound mucins are part




FIGURE 1 | Multifunctional roles of host defense peptides (HDPs). Besides direct antimicrobial activities, HDPs actively participate in systemic and mucosal epithelial defense by modulating a range of host innate and adaptive immune responses as indicated. Recent accumulating evidence has highlighted a direct involvement of HDPs in improving intestinal and epidermal barrier function.

of the epithelial glycocalyx that are involved in cell signaling and interactions with the environment without forming gel or oligomerization (57). Secreted human mucins include MUC2, -5AC, -5B, -6, -7, and -19, and the transmembrane mucins consist of MUC1, -3, -4, -12, -13, -15, -16, -17, and -20 (54, 55). Structurally, most secreted mucins are unique in the presence of multiple von Willebrand factor (VWF) domains and a C-terminal cysteine-knot domain, while a majority of membrane-bound mucins specifically consist of a transmembrane domain, a cytoplasmic tail, 1–2 epidermal growth factor (EGF)-like domains, and a sea urchin sperm protein, enterokinase, and agrin (SEA) domain (56, 58) (**Figure 2**). The VWF domains and cysteine-knot domain of secreted mucins are responsible for formation of higher-order structures through oligomerization, while the EGF-like and SEA domains of membrane-bound mucins mediate signaling transduction and cleavage of the extracellular portion of mucins, respectively (58). Among all secreted mucins, MUC2 is the most abundant in the human small intestine and colon, and MUC5AC is predominant in the stomach. Structurally, similar mucins have been found in most other vertebrate species, including cattle, pigs, and chickens (59–62).

The mucus layer formed by secreted mucins varies in composition along the GI tract. The stomach and large intestine consist

of two distinct mucus layers: a “loose” outer layer and a “thick” inner layer (2). The inner layer closest to epithelial cells is densely packed and holds firmly to the cells. The inner mucus layer is largely free of bacteria, providing a sterile protective environment for the epithelium. The outer mucus layer is much more soluble due to proteolytic cleavages that allow the mucus layer to expand without disrupting mucin polymers. This outer layer provides a habitat for commensal bacteria to bind via specific adhesins and to thrive via breaking down the mucin glycans as a food source. Specificity of bacteria for different glycans is speculated to be important for developing species-specific microbiota (2).

Altered expression or glycosylation of mucins is often associated with intestinal barrier dysfunction (57). For example, Muc2 deficiency in mice causes increased permeability, gross bleeding, spontaneous development of inflammation in the GI tract, as well as severe growth retardation (63, 64). Muc1- or Muc2-deficient mice become more prone to infections with *Campylobacter jejuni*, *Helicobacter pylori*, *Salmonella enterica* serovar Typhimurium, and *Citrobacter rodentium* (64–67). Moreover, mice lacking the enzyme, β 1,3-*N*-acetylglucosaminyltransferase that synthesizes O-glycans on mucins, exhibit a thinner mucus layer showing an enhanced susceptibility to enteric bacterial infections (67) and dextran sodium sulfate (DSS)-induced colitis (68). Additionally,




FIGURE 2 | Schematic diagrams of glycosylated mucin monomers. Representative membrane-bound and secreted mucins are exemplified by mucin 1 (MUC1) and MUC2, respectively. The protein backbone of mucins is characterized by the presence of an excessive number of tandem repeat (TR) sequences. Mucins are heavily O-glycosylated through enriched threonine and serine residues in the TR domain. A transmembrane domain, a cytoplasmic tail, and a sea urchin sperm protein, enterokinase, and agrin (SEA) domain are unique to most membrane-bound mucins, whereas the presence of several von Willebrand factor (VWF) domains and a C-terminal cysteine-knot domain is specific to a majority of secreted mucins. The diagrams were modified primarily from reference (56).

significantly reduced expressions of multiple mucins such as MUC1, MUC3, MUC4, and MUC5B are observed in ileal mucosa of Crohn's disease (CD) patients (69), although the expression changes of mucins are less clear in ulcerative colitis (UC) patients (70).

TIGHT JUNCTIONS: GATE GUARDS FOR BORDER PROTECTION

The intestinal epithelium is made up of several different cell types organized into crypts and villi. These include intestinal epithelial stem cells, enterocytes, and secretory cells, such as Paneth, goblet, and enteroendocrine cells (71). Intestinal stem cells give rise to all other epithelial cell types, while enterocytes primarily function in nutrient absorption with the ability to synthesize and release HDPs and mucins. Paneth and goblet cells are major producers of HDPs and mucins, respectively, while enteroendocrine cells have a primary role of secreting numerous hormones that act as regulators of digestive function (71). All intestinal epithelial cells are linked at lateral membranes through formation of three major types of junctional complexes, i.e., TJs, adherens junctions, and desmosomes (1, 3, 72). Collectively, they form a virtually impermeable seal to the paracellular space. Besides the barrier function, these junctional complexes maintain cell polarity by separating the apical from basolateral membranes. TJs are multi-protein complexes located at the most apical end of the lateral membrane. The TJ assembly is composed of both transmembrane and cytoplasmic plaque proteins that interact directly with the cytoskeleton (Figure 3). Among all three major junctional complexes, only TJs have the ability to control the

selective paracellular permeability for ions, water, and other small molecules (1, 3, 72). Therefore, TJs are the major determinant of mucosal epithelial permeability.

Tight Junction Structures

Among the proteins involved in TJ assembly, claudins, occludin, junctional adhesion molecules (JAM), and tricellulin are the major transmembrane proteins that constitute a selective paracellular barrier, whereas zonula occludens (ZO) and cingulin are the main cytoplasmic plaque proteins located at the peripheral membrane (1, 3, 72). All TJ proteins are highly conserved in vertebrate species. Claudins are a large family of small proteins of 21–34 kDa that make up the backbone of the TJ structure, with at least 26 members reported in humans (73). Remarkably, each claudin shows a unique tissue expression pattern with varied levels of expression in different segments of the GI tract. Occludin (65 kDa) was the first transmembrane TJ protein identified (74), with no homologs being found (75). On the other hand, the JAM family is comprised of three classical members (JAM-1, -2, and -3) and four related molecules (JAM-4, JAM-L, CAR, and ESAM) at ~40 kDa each (76). Tricellulin is a 64-kDa protein located preferentially at tricellular junctions, although it is also present in bicellular junctions (77). Tricellulin shares 32% identity in the amino acid sequence with the C-terminal tail of occludin. ZO proteins belong to the family of membrane-associated guanylate kinase (GUK) homologs that include three members, i.e., ZO-1 (~220 kDa), ZO-2 (~160 kDa), and ZO-3 (~130 kDa) (78), whereas cingulin is an ~140-kDa protein that links ZO proteins to the actin cytoskeleton (79).

Claudins, occludin, and tricellulin are all membrane proteins with four transmembrane domains, one intracellular and two





FIGURE 3 | Schematic drawing of tight junction structures at the apicolateral membranes of the paracellular space. Tight junctions are comprised of transmembrane proteins such as claudins, occludin, junctional adhesion molecules (JAM), and tricellulin as well as cytoplasmic plaque proteins such as three zonula occludens (ZO) proteins and cingulin. Claudins, occludin, and tricellulin consist of four transmembrane domains, while JAMs have a single transmembrane domain. Through largely homophilic interactions, neighboring cells are sealed at the apicolateral region. Cytoplasmic proteins link the tight junction assembly to the cytoskeleton to control the opening of paracellular pores.

extracellular loops, and two cytoplasmic tails, whereas the JAM proteins are single-pass transmembrane proteins consisting of two extracellular immunoglobulin (Ig)-like domains, a single transmembrane domain, and a short intercellular C-terminal tail (72) (**Figure 4**). While the N-terminal tail is generally short, the longer C-terminal tail of claudins consists of a post-synaptic density 95, disk-large, and zonula occludens (PDZ)-binding motif that interacts with the first PDZ domain of ZO-1 (73). The crystal structure of claudins as exemplified by mouse claudin-15 indicates that the four transmembrane segments form a tight four-helix bundle with parts of the two extracellular loops forming a “palm-shaped” structure (80). A model for the architecture of claudin-formed TJ strands in the membrane has been proposed based on the results of crosslinking experiments and electron microscopy (81). In this model, claudins show an antiparallel, double-layer arrangement (**Figures 5A,B**). The association of claudin double layers in neighboring lateral membranes results in the formation of multiple extracellular β -barrel-like pores parallel with the membrane plane to allow the passage of ions through the paracellular space (**Figure 5C**).

As for occludin, its two extracellular loops mediate homophilic interactions and permeability to macromolecules. The occludin-ELL domain in the C-terminal tail of occludin is responsible for interacting with the GUK domain of ZO proteins, while

the N-terminal tail lacks a defined function (75) (**Figure 4**). In comparison with claudins and occludin, tricellulin consists of two long tails, with the C-terminal tail containing an occludin-ELL domain that is likely to interact with the GUK domain of ZO proteins (82). JAM proteins also contain a cytoplasmic tail with a PDZ-binding motif that interacts with the third PDZ domain of ZO-1 (83) (**Figure 4**). The extracellular Ig-like domains of JAM proteins are responsible for homophilic and heterophilic interactions.

The cytoplasmic ZO proteins contain three PDZ domains, a Src homology-3 (SH3) domain, and a GUK domain (78). As stated above, ZO-1 directly interacts with claudins, ZO-2, and JAM-1 through the first, second, and third PDZ domain, respectively. The GUK domain of ZO proteins is known to associate with occludin, while the C-terminal actin-binding region is responsible for bridging with actin, raising the possibility of forming a large protein complex through simultaneous interactions of many TJ proteins with ZO-1 (78). Cingulin is another intracellular plaque protein that is predicted to form a homodimer with globular head and tail at both ends connected by a coiled-coil “rod” domain in the central region. The head of cingulin is known to bind to ZO proteins and the coiled-coil domain interacts with myosin (79). The interactions between the TJ protein complex and cytoskeleton are critical in maintaining and regulating the TJ



structure and function, as the mucosal permeability is regulated heavily by the phosphorylation status of myosin light chain, which can be modified by the kinases such as myosin light chain kinase (MLCK) (72, 73) (Figure 4).

Tight Junctions' Function in Selective Permeability


Tight junctions are distributed at the apical surface of epithelial and endothelial cells throughout the body in vertebrate animals, including the skin, GI, respiratory, and urogenital tracts as well as the blood vessels (72). TJs are the major determinant of mucosal barrier permeability. Ions, water, and macromolecules pass TJs through either of the two major types of pores. The non-restrictive pathway, also known as the “leak” pathway, allows the transport of macromolecules through large pores with no charge selectivity, while the restrictive or “pore” pathway is only permeable to small ions through pores of ~ 4 Å in radius with charge selectivity (1, 3, 4). Claudins are mainly responsible for the “pore” pathway, and the two extracellular loops work as an “electrostatic selective filter” to select the size and charge for small pores (Figure 5). The charge selectivity of individual claudins is determined by the net charge of the amino acid residues in the first extracellular loop. For example, claudin-1 is selective for anions, while claudin-2 prefers cations. In the intestinal tract, claudin-1, -3, -4, -5, -8, -9, -11, and -14 decrease paracellular permeability and are regarded

as barrier-forming claudins, while claudin-2, -7, -12, and -15 are to increase permeability and generally referred as pore-forming claudins (72, 73).

The non-restrictive “leak” pathway is primarily dependent on occludin as evidenced by an increase in paracellular flux of macromolecules without a noticeable effect on transepithelial electrical resistance (TEER) both *in vitro* and *in vivo* after occludin knock-down (84). JAM proteins also enhance the TJ function by decreasing permeability and facilitating the assembly of occludin to the TJ complex (83). In addition, JAMs can regulate the paracellular barrier for the transmigration of leukocytes from the blood vessel to inflamed sites in response to inflammation (76). It is noteworthy that paracellular water permeability is mainly dictated by claudin-2.

Modulation of Paracellular Permeability

A number of agents such as cytokines, growth factors, pathogens, probiotics, nutrients, and phytochemicals have been found to impact TJ permeability and mucosal barrier functions through transcriptional regulation and posttranslational modification of TJ proteins (72, 73). Increased expressions of barrier-forming claudins, occludin, and JAM proteins are commonly associated with reduced paracellular permeability and improved barrier function, whereas an elevation in the expression of pore-forming claudins often leads to barrier dysfunction. For example,



transforming growth factor (TGF)- β enhances barrier integrity of intestinal epithelial cells by augmenting claudin-1 and -4 expression, while IL-1 β , IL-6, and TNF- α increase intestinal cell permeability by increasing claudin-2 expression and/or reducing occludin and ZO-1 expression (72).

Tight junction barrier integrity is also affected by post-translational modifications of transmembrane and cytoplasmic proteins and associated regulatory proteins. Phosphorylation, glycosylation, and/or ubiquitination of the TJ proteins have a profound impact on barrier permeability. For example, claudin-1 is phosphorylated by atypical protein kinase C (aPKC), protein

kinase A (PKA), and mitogen-activated protein kinases (MAPK) and dephosphorylated by protein phosphatase 2A (72, 73). Phosphorylation of claudins generally promotes their assembly into the TJ, whereas dephosphorylation often results in the dissociation of claudins from the TJ (72, 73). Similarly, phosphorylation of occludin enhances the barrier function, while dephosphorylation delays the TJ assembly resulting in barrier dysfunction (72, 75).

Besides those proteins involved directly in the TJ assembly, paracellular permeability is also heavily influenced by actin-myosin filaments that are linked to the TJ proteins. It is well known that up-regulation of MLCK is linked to an increase in the TJ permeability by catalyzing the phosphorylation of myosin light chain, which in turn induces the contraction of actin-myosin filaments and opening of the TJ barrier (85). Both IL-1 β and TNF- α are strong inducers of the MLCK gene transcription and activation, resulting increased myosin light chain phosphorylation and TJ barrier permeability (86, 87).

The expression and posttranslational modifications of TJ proteins are influenced by a complicated network of signaling pathways that intertwine with each other. Activation of nuclear factor (NF)- κ B signaling by pathogens, pathogen-associated molecular patterns (PAMPs), and proinflammatory cytokines often causes an increase in intestinal epithelial permeability through induction of pore-forming claudin-2 and suppression of barrier-forming claudins such as claudin-1, -3, -4, -5, -7, and -8. On the other hand, activation of TGF- β /SMAD and PPAR- α / γ signaling are generally barrier protective by enhancing claudin-1 and -4 expressions while downregulating claudin-2 (73).

Implication of Tight Junction Dysfunction in Disease Pathogenesis

The intestinal barrier helps to maintain homeostasis between gut microbiota and the immune system. TJ dysfunction is associated with many enteric disorders such as CD, UC, and celiac disease (3, 72). In CD patients, the expressions of barrier-forming claudin-3, -5, -8, occludin, and JAM-1 are decreased, while pore-forming claudin-2 is significantly increased; and in UC patients, a down-regulation of claudin-1, -4, and JAM-1, and up-regulation of claudin-2, is observed (3, 72). Moreover, increased MLCK expression and activity are evident in both CD and UC patients (85). These factors collectively exacerbate the intestinal paracellular permeability leading to a “leaky gut syndrome.” However, many of these clinical conditions are also accompanied with increase synthesis of proinflammatory cytokines, which are known to cause barrier dysfunction. Thus, it is difficult to determine whether barrier dysfunction is a cause or effect of many of these diseases.

Several enteric pathogens such as *Vibrio cholera*, enteropathic *Escherichia coli*, *Clostridium perfringens* are known to cause diarrhea mainly through disruption of the intestinal barrier function by secretion of exotoxins (72). For example, claudin-3 and -4 are receptors for *C. perfringens* enterotoxin, and the binding of enterotoxin to the extracellular loops of claudins causes internalization of claudins and disintegration of the TJ assembly (88). Early weaning (<3 weeks of age) is known to impair the development of intestinal barrier functions of pigs leading to

more pronounced diarrhea (89), and was recently found to lead to reduced expressions of occludin, claudin-1, and ZO-1 in the jejunum (90).

REGULATION OF TIGHT JUNCTION AND MUCIN PRODUCTION BY HOST DEFENSE PEPTIDES

Association of HDP Expression with Barrier Dysfunction

Along with decreased TJ protein expression, aberrant HDP expression is common in CD and UC patients (91). The expression of Paneth cell α -defensins (HD5 and HD6) is significantly reduced in ileal CD patients, but unaffected in colonic CD patients Z (92). Instead, a reduced expression of HBD-1 and HBD-2 is observed in colonic CD (93). Moreover, induction of cathelicidin LL-37 and HBD-2, -3, and -4 is also reduced in colonic CD relative to healthy subjects (94–97). This lack of HDP induction in CD patients is thought to play a key role in CD pathogenesis as it indicates a lack of intestinal immune response. A deficiency in HD5 and HD6 synthesis is even more pronounced in patients carrying a mutation in the intracellular NOD2 receptor (98), which is expressed by Paneth cells. Consistently, NOD2-knockout mice show a diminished expression of Paneth cell α -defensins known as cryptdins in mice (99). In contrast to CD patients, UC patients display unchanged HD5 and HD6 expressions (92), while LL-37, HBD-2, -3, and -4 are upregulated (94–97). A thin or even absent mucus layer is evident in UC intestinal segments, which causes intestinal inflammation due to direct adhesion and invasion of bacteria to mucosal epithelial cells. Although UC patients produce HDPs, these peptides are not retained in the intestinal tract. Perhaps the most convincing evidence linking the positive role of HDPs in barrier function comes from the studies with cathelicidin (CRAMP)-deficient mice. These mice show delayed recovery of barrier permeability in response to acute disruption of epidermal barrier, albeit with subtle barrier abnormalities in the epidermis (100). Collectively, these lines of evidence suggest a direct impact of HDPs on intestinal barrier function and homeostasis.

Transcriptional Regulation of Mucins and TJ Proteins by HDPs

Accumulating pieces of evidence suggest a direct involvement of HDPs in regulating the synthesis of mucins and TJ proteins in the intestinal tract. HBD-2 upregulates MUC2, MUC3, but not MUC1 or MUC5AC in human HT-29 colonic epithelial cells (101). MUC2 expression is also enhanced in human Caco-2 colonic epithelial cells in response to HBD-2 (101), and upregulated MUC2 in turn promotes HBD-2 expression (102), suggestive of a positive feedback mechanism between MUC2 and HBD-2. LL-37 also enhances MUC1, MUC2, and MUC3 expressions in HT-29 cells (103, 104) and MUC3 expression only in Caco-2 cells (104). Buforin II, a 21-amino acid HDP isolated from the stomach of an Asian toad (*Bufo bufo garagrizans*), improves intestinal barrier function in weaned piglets challenged with three enterotoxigenic *E. coli* (ETEC) strains (105). Oral administration (twice daily) of buforin II leads to an increase in claudin-1, occludin, and ZO-1

expression in the jejunal segments of *E. coli*-challenged piglets (105). Importantly, buforin II also improves intestinal morphology and growth performance and reduced bacterial shedding in fecal swabs (105). Additionally, administration of a banded krait HDP known as cathelicidin-BF induces ZO-1 expression in the jejunum of healthy mice and also restores LPS-mediated impairment of ZO-1 and intestinal barrier function (106). Furthermore, porcine β -defensin-2 (PBD-2) is capable of restoring the expression of MUC1, MUC2, claudin-1, ZO-1, and ZO-2 as well as the barrier integrity of the colon of DSS-treated mice (107).


Besides direct regulation of the intestinal paracellular permeability, several HDPs also positively influence the barrier effect of the respiratory tract and the skin. LL-37 induces MUC5AC in human NCI-H292 airway epithelial cells (108). LL-37, HBD-3, and S100A7/psoriasin are all able to augment the expression of TJ proteins in human skin keratinocytes (109–111). HBD-3 induces the expressions of all 14 claudins examined, but not occludin or ZO 1–3 in human keratinocytes (110). Similarly, LL-37 dose-dependently enhances the expressions of 11 claudins and occludin in skin keratinocytes, but not JAM 1–3 or ZO 1–3 (109). S100A7 also promotes the expressions of multiple claudins and occludin, but not JAM 1–3 or ZO 1–3 in human keratinocytes (111). Multiple signaling pathways are involved in HDP-induced barrier protein synthesis as detailed below.

MOLECULAR MECHANISMS OF HDP REGULATION OF BARRIER FUNCTIONS

A number of extracellular and intracellular receptors have been reported to be responsible for a range of physiological functions of cathelicidins and defensins in humans and mice. LL-37 and the mouse ortholog (CRAMP) are ligands for P2X purinergic receptor 7 (P2X7), formyl peptide receptor-like (FPRL) 1/2, glyceraldehyde 3-phosphate dehydrogenase, and sequestosome-1/p62, whereas several human and mouse β -defensins bind to CC chemokine receptor 2 (CCR2), CCR6, CXC chemokine receptor 2 (CXCR2), and toll-like receptor 1/2/4 (112, 113). The receptors and signaling pathways by which HDPs induce the expression of mucins and TJ proteins have been studied, but remain elusive in most cases. It is worth noting that, although most published mechanistic studies were based on skin keratinocytes, the following overall conclusions are believed to be applicable to intestinal epithelia as well: (1) HDPs vary greatly in their ability to modulate barrier function, albeit with structural similarities, (2) mucins and TJ proteins are differentially regulated by HDPs, (3) multiple signaling pathways are employed by the same HDP, and (4) receptors appear to be differentially engaged in mediating the induction of mucins and TJ proteins by different HDPs. The current findings are summarized below.

Signaling Mechanisms of HDP-Mediated Mucin Induction

LL-37-induced MUC5AC expression in lung epithelial cells appears to be mediated mainly through transactivation of the EGF receptor (EGFR) (108), although EGFR is not a direct receptor for LL-37 (Figure 6A). Initially, LL-37



triggers the activation of TNF- α -converting enzyme, which in turn cleaves the membrane-bound form of TGF- α , but not heparin binding-EGF. Released TGF- α subsequently interacts with and phosphorylates its receptor, EGFR, which induces MUC5AC gene expression through activation of multiple signaling pathways (108). For LL-37 to induce MUC2 and MUC3 expression in human intestinal epithelial cells, both EGFR and P2X7, but not G-protein-coupled receptors, are involved (104). HBD-2-induced mucin expression in human intestinal epithelial cells is shown to be partially mediated through CCR6 (101). The p38 MAPK, but not extracellular signal-regulated kinase (ERK) or PI3K, is involved in mediating P2X7- and EGFR-activation of MUC2 production in human Caco-2 cells (104).

LL-37-Mediated TJ Protein Induction

Rac1, aPKC ζ/λ , glycogen synthase kinase (GSK)-3 α/β , and PI3K are all phosphorylated and activated in human skin keratinocytes in response to LL-37, and blockage of any enzyme with a specific inhibitor results in a substantial reduction in the TEER and a significant increase in the permeability to FITC-dextran (109) (Figure 6B). Rac1 is a small GTPase that functions upstream of aPKC ζ/λ as part of the Par3/Par6/aPKC ζ/λ polarity complex, which in turn phosphorylates the C-terminal domain of occludin (114) or JAM-1 (115), promoting its assembly into the TJ complex and enhancing the barrier function (116). GSK-3 α/β is involved in a number of signaling pathways (117) and is required for induction of occludin and claudin-1 in intestinal and kidney epithelial cells (118). Consistently, LL-37 triggers phosphorylation

and activation of GSK-3 α/β at Tyr 216 and Tyr 279 in human keratinocytes at 1–2 h after exposure, leading to the improvement of epidermal barrier function (109).

PI3K functions upstream of GSK-3, and the PI3K signaling cascade has been implicated in both the degradation and stimulation of TJ barrier function depending on the stimulating agent (119). PI3K is quickly phosphorylated within 30 min in human keratinocytes upon stimulation with LL-37 (109). In intestinal cells, PI3K plays a key role in directing proper occludin localization and subsequent tightening of epithelial barrier function in response to prostaglandins (120). Inhibition of PI3K in porcine ischemia-injured ileal mucosa attenuates the ability of prostaglandin to recover proper barrier function. In rat Con8 mammary epithelial cells, glucocorticoid recruits Ras and the p85 subunit of PI3K to the TJ complex and increases barrier function (121). However, the specific cellular receptor(s) mediating LL-37-induced TJ protein expression remain unknown and warrant further investigation.

HBD-3-Mediated TJ Protein Induction

Although HBD-2 is capable of inducing mucin expression (101), only HBD-3 triggers the synthesis of multiple TJ proteins (110). CCR6 has been shown to be primarily responsible for HBD-3-induced enhancement of barrier integrity in epidermis (110). Similar to LL-37, HBD-3 is also capable of phosphorylating and activating Rac1, aPKC ζ/λ , PI3K, and GSK-3 α/β in similar kinetics in human skin keratinocytes (110) (Figure 6C). Of note, toll-like receptors, PKA, and MAPK pathways are not involved in mediating HBD-3-induced barrier function improvement (110). Although HBD-1, HBD-2, and HBD-4 fail to alter the epidermal permeability, they also have weak activities in activating Rac1, aPKC, GSK-3, and PI3K (110), suggesting those pathways may not be solely devoted to the TJ functions. It is important to note that, although similar in the tertiary structure, only HBD-3, but not HBD-1, -2 or -4, has the ability to trigger the induction of TJ proteins (115).

S100A7-Mediated TJ Protein Induction

The role of GSK-3 α/β and MAPK in human epidermal barrier function mediated by S100A7 has been studied (111) (Figure 6D). GSK-3 α/β is phosphorylated and activated at Tyr 216 and Tyr 279 within 30 min following exposure of human keratinocytes to S100A7/psoriasis. Specific inhibition of GSK-3 activation abolishes induction of claudins and epidermal TEER by S100A7 (111). Because β -catenin is regulated directly by GSK-3 (117), S100A7 is revealed to phosphorylate and activate β -catenin, which is vital to the assembly of adherens junctions. E-cadherin, another essential component of the adherens junction complex, is also phosphorylated by S100A7 (111), suggesting that, besides the TJs, S100A7 also improves the assembly of adherens junctions.

The MAPK pathway includes three canonical signaling cascades that consist of ERK, c-Jun N-terminal kinase (JNK), and p38 (122). Collectively, they are critical to many important physiological processes ranging from cell division and differentiation to stress and immune responses. Unlike HBD-3, S100A7 is capable of activating all three canonical MAPK cascades (111). ERK is

quickly phosphorylated in 2 min in human skin keratinocytes following exposure to S100A7, while JNK and p38 MAPK are also phosphorylated in 30 min. Inhibition of individual MAPK signaling cascades leads to a substantial reduction in claudin induction and epidermal TEER (111), implying that all three major MAPK pathways are required. However, the involvement of any specific receptors or other signaling pathways remains to be studied.

ROLE OF HDPs IN INTESTINAL MUCOSAL HOMEOSTASIS, IMMUNE DEFENSE, AND GROWTH PERFORMANCE

One of the major functions of the intestinal epithelium is to act as a barrier against the invasion of microorganisms. This task is especially difficult considering that the intestinal mucosa is colonized by over 10¹³ microorganisms (123), with the majority being commensal bacteria that are beneficial to the host through their ability to improve digestion, absorption, and vitamin synthesis while also limiting pathogen growth (124). The two most dominant bacterial phyla present in the intestinal tract of humans and mice are Gram-negative Bacteroidetes and Gram-positive Firmicutes, which together comprise about 70–80% of the total bacteria present (125). Commensal bacteria are vital to the development of normal intestinal morphology and immune system (126, 127). While commensal bacteria are beneficial to the host under homeostatic conditions, a state of dysbiosis, or imbalance of the microbial community, leads to inflammation and disturbed epithelial homeostasis. This is particularly seen in the CD patients in which the host immune system displays increased activation against commensal microbiota.

The intestinal epithelium continuously monitors resident microbes through interactions between pattern recognition receptors (PRRs) and microbe-associated molecular patterns (MAMPs). Activation of PRRs stimulates the synthesis and release of HDPs and mucins from intestinal cells (126, 127). A large amount of HDPs secreted from Paneth cells and enterocytes are retained in the mucus layer to create a strong barrier against bacterial invasion (128). Studies with HDP-knockout and -transgenic mice have illuminated the role of HDPs in intestinal homeostasis and immune defense. Knockout of the mouse cathelicidin CRAMP gene causes exaggerated colitis in the colon of mice, and the disease symptoms are further exacerbated following DSS treatment (129). Adoptive transfer of bone marrow cells from the wild-type mice to CRAMP-knockout mice alleviates DSS-induced colitis (129). Mice carrying the transgene for HD5 show an augmented ability to fight off orally challenged *S. enterica* serovar Typhimurium (130). Conversely, matrix metalloproteinase 7 (MMP7)-knockout mice with a deficiency in producing biologically active enteric defensins display reduced capacity to clear enteric pathogens (131). Furthermore, a comparison between those two complementary mouse models has revealed dramatic defensin-dependent reciprocal shifts in the intestinal bacterial composition. In comparison with wild-type mice, Firmicutes are reduced and Bacteroidetes are increased in small intestine of HD5-transgenic mice, while the opposite is true with MMP7-deficient mice (125, 132). Moreover, overexpression

of HD5 in mice causes a significant loss of segmented filamentous bacteria in the distal small intestine and a reduced presence of Th17 cells in the lamina propria (132), suggesting clearly that enteric HDPs represent a critical factor in shaping the microbiota composition and inflammatory status of the GI tract.

Multiple studies have highlighted the beneficial effects of direct feeding of HDPs on growth, intestinal morphology, and immune status in pigs (133, 134). Dietary supplementation of an *E. coli*-producing bacteriocin, colicin E1, significantly improved weight gain and feed efficiency of ETEC-challenged weanling pigs in a 4-day trial, relative to the control pigs (135). Colicin E1 inclusion also reduced the *E. coli* titers in both the fecal and ileal samples as well as the incidence and severity of diarrhea (135). Moreover, the expression levels of proinflammatory cytokines (IL-1 β and TNF- α) were reduced in the ileum of pigs in response to colicin E1 feeding (135). Similarly, feeding a recombinant silkworm HDP, cecropin A/D, improved growth and feed efficiency and reduced diarrhea incidence in ETEC-challenged weanling pigs, without an obvious impact on intestinal morphology or nitrogen/energy utilization over a period of 6 days (136). Dietary inclusion of a recombinant fusion HDP derived from bovine lactoferrin also enhanced growth performance and decreased the incidence of diarrhea in ETEC-challenged piglets over a 21-day period (137). Across five different commercial farms, feeding a mixture of four recombinant HDPs, including lactoferrin, cecropin, defensin, and plectasin resulted in an enhancement of growth and feed efficiency and a reduction in diarrhea incidence in normally reared weanling pigs (138). Parallel to the studies above, supplementation of a synthetic HDP (AMP-A3 or P5) improved nutrient digestibility, intestinal morphology, and growth performance of normally reared weanling pigs in a 4-week trial, without affecting serum concentrations of IgA, IgG, or IgM (139, 140). Additionally, AMP-A3 and P5 appeared to reduce the titers of potentially harmful *Clostridium* spp. and coliforms in the ileum, cecum, and feces (140). Feeding a combination of two HDPs and a probiotic yeast led to an improvement in intestinal morphology and feed efficiency of piglets challenged with deoxynivalenol, a mycotoxin commonly found in grains (141, 142). In most trials above, HDPs performed indistinguishably with in-feed antibiotics in promoting growth, feed efficiency, and intestinal morphology (133).

The beneficial effects of direct feeding of HDPs are not limited to pigs. Supplementation of AMP-A3 to broiler chickens resulted in an increase in weight gain and feed efficiency over control birds, which was comparable to the birds fed avilamycin, an in-feed antibiotic (143). Intestinal morphology was also improved in broilers as measured by increased villus heights and villus height:crypt depth ratios in the small intestine (143). Similar to the results in pigs, broilers also displayed an improvement in nutrient utilization and a reduction in *Clostridium* spp. and coliforms in the intestinal tract (143). Supplementation of a yeast broth containing recombinant cecropin A/D improved intestinal morphology and nutrient utilization, with a tendency to enhance growth performance of broiler chickens in a 4-week trial (144). Cecropin A/D inclusion also reduced the total aerobic bacterial counts in both the jejunal and cecal contents of 42-day-old chickens (144). Collectively, these animal results have suggested

the beneficial effects of HDP feeding, justifying dietary supplementation of HDPs as an antibiotic-alternative strategy in growth promotion and disease control.

However, because of HDP's proneness to enzymatic degradation and high production costs with either the synthetic or recombinant form, it may not be biologically efficient and economically effective for direct supplementation of HDPs in animal diets. Recently, several classes of small-molecule compounds, such as butyrate, have been found to induce HDP synthesis and enhance bacterial clearance in humans, chickens, pigs, and cattle without triggering inflammatory response (145–151). Dietary supplementation of these simple HDP-inducing compounds or their combinations may prove to be an alternative, cost-effective approach to antibiotics for livestock applications (152). However, the efficacy of these HDP-inducing compounds in promoting growth, intestinal health, and microbiota balance is yet to be demonstrated in animal trials.

CONCLUDING REMARKS

A comprehensive understanding of intestinal barrier function and its regulation is paramount to ensuring the sustainability of the food animal industry because disruption of the barrier results in disease states and decreased production efficiency (6). With potent antimicrobial and immunomodulatory properties, HDPs are further revealed to hold a new capacity to directly regulate barrier function. Aberrant synthesis of epithelial HDPs often leads to barrier dysfunction, and the diseases with impaired barrier integrity are commonly associated with reduced HDP synthesis, raising the possibility of treating barrier dysfunction with HDPs. However, a number of questions remain before HDP-based therapies can be devised for augmentation of intestinal barrier function, animal health, and production efficiency.

On the one hand, several structurally diverse HDPs (e.g., cathelicidins, defensins, and S100A7) have a similar ability to induce the expression of mucins and TJ proteins in both epidermal and intestinal epithelial cells. On the other hand, certain structurally similar HDPs (e.g., HBD-3 vs. HBD-1, -2, and -4) behave quite differently in their capacity to induce TJ proteins. Only HBD-3, but not other HBDs, has the capacity to enhance the barrier effect (110). Structure–activity relationship studies of HDPs may reveal whether there is an optimal physicochemical or structural feature for maximal induction of mucins and TJ proteins.

A number of different extracellular and intracellular receptors have been identified for human HDPs to mediate different physiological functions. However, the identities of the receptors utilized by different classes of HDPs to regulate paracellular permeability remain largely unclear in most cases. There are a number of questions on the signaling mechanisms of HDP-mediated barrier function that need to be answered. For example, what are the major receptors involved in HDP-induced synthesis of mucins and TJ proteins in humans? Do the same set of receptors that are utilized by human HDPs work similarly in the livestock species? Are there any new, unidentified receptors specific for regulation of barrier function? Besides the Rac1-aPKC, PI3K-GSK-3, and MAPK pathways, what are other major signaling pathways that

mediate the barrier effect? How do these pathways cross-talk with each other? What are those major transcription factors that are required for induction of different mucin and TJ proteins? How and whether are mucins and TJ proteins differentially regulated? Do epithelial cells on epidermis, GI, respiratory, and genitourinary tracts engage in different receptors and signaling pathways?

Besides the abundance of TJ proteins, both posttranslational modifications of TJ proteins and the status of associated actomyosin ring have a strong impact on barrier permeability. Many agents are shown to alter the barrier function through phosphorylation of certain TJ proteins or through activation of MLCK, which in turn phosphorylates myosin light chain and causes contraction of the perijunctional actomyosin ring and opening of the paracellular pores. It will be important to examine whether and how HDPs influence posttranslational modifications of TJ proteins as well as the transcription and activity of MLCK.

Nevertheless, it is exciting to reveal a direct involvement of HDPs in barrier function and the potential of HDPs in enhancing

gut health and animal performance. Additional studies along this line may someday turn the HDP-based therapies into reality. Although administration of synthetic peptides may be feasible in human medicine, it is cost-prohibitive in the livestock industry. Supplementation of exogenous recombinant HDPs or dietary compounds with the capacity to induce the synthesis of HDPs in the GI tract has emerged as a cost-effective strategy in antimicrobial therapy and may have potential to replace antibiotics in food animal production.

ACKNOWLEDGMENTS

This work was supported in part by Oklahoma Center for the Advancement of Science and Technology grants (AR12.2-077, HR12-051, and AR15.049), Oklahoma State University Cowboy Technologies Business Development Program grant, and Oklahoma Agricultural Experiment Station project H-2811. KR was supported by a USDA-NIFA National Needs Fellowship grant (2013-38420-20500).

REFERENCES

- Turner JR. Intestinal mucosal barrier function in health and disease. *Nat Rev Immunol* (2009) **9**:799–809. doi:10.1038/nri2653
- Johansson ME, Sjovall H, Hansson GC. The gastrointestinal mucus system in health and disease. *Nat Rev Gastroenterol Hepatol* (2013) **10**:352–61. doi:10.1038/nrgastro.2013.35
- Groschwitz KR, Hogan SP. Intestinal barrier function: molecular regulation and disease pathogenesis. *J Allergy Clin Immunol* (2009) **124**:3–20; quiz 1–2. doi:10.1016/j.jaci.2009.05.038
- Rescigno M. The intestinal epithelial barrier in the control of homeostasis and immunity. *Trends Immunol* (2011) **32**:256–64. doi:10.1016/j.it.2011.04.003
- Liu D, Guo Y, Wang Z, Yuan J. Exogenous lysozyme influences *Clostridium perfringens* colonization and intestinal barrier function in broiler chickens. *Avian Pathol* (2010) **39**:17–24. doi:10.1080/03079450903447404
- Mani V, Weber TE, Baumgard LH, Gabler NK. Growth and development symposium: endotoxin, inflammation, and intestinal function in livestock. *J Anim Sci* (2012) **90**:1452–65. doi:10.2527/jas.2011-4627
- Zaslouff M. Antimicrobial peptides of multicellular organisms. *Nature* (2002) **415**:389–95. doi:10.1038/415389a
- Brogden KA, Ackermann M, McCray PB Jr, Tack BF. Antimicrobial peptides in animals and their role in host defences. *Int J Antimicrob Agents* (2003) **22**:465–78. doi:10.1016/S0924-8579(03)00180-8
- Hancock RE, Nijnik A, Philpott DJ. Modulating immunity as a therapy for bacterial infections. *Nat Rev Microbiol* (2012) **10**:243–54. doi:10.1038/nrmicro2745
- Hilchie AL, Wuertth K, Hancock RE. Immune modulation by multifaceted cationic host defense (antimicrobial) peptides. *Nat Chem Biol* (2013) **9**:761–8. doi:10.1038/nchembio.1393
- Mansour SC, Pena OM, Hancock RE. Host defense peptides: front-line immunomodulators. *Trends Immunol* (2014) **35**:443–50. doi:10.1016/j.it.2014.07.004
- Kolls JK, McCray PB Jr, Chan YR. Cytokine-mediated regulation of antimicrobial proteins. *Nat Rev Immunol* (2008) **8**:829–35. doi:10.1038/nri2433
- Gallo RL, Hooper LV. Epithelial antimicrobial defence of the skin and intestine. *Nat Rev Immunol* (2012) **12**:503–16. doi:10.1038/nri3228
- Selsted ME, Ouellette AJ. Mammalian defensins in the antimicrobial immune response. *Nat Immunol* (2005) **6**:551–7. doi:10.1038/ni1206
- Zanetti M. Cathelicidins, multifunctional peptides of the innate immunity. *J Leukoc Biol* (2004) **75**:39–48. doi:10.1189/jlb.0403147
- Donato R, Cannon BR, Sorci G, Riuzzi F, Hsu K, Weber DJ, et al. Functions of s100 proteins. *Curr Mol Med* (2013) **13**:24–57. doi:10.2174/156652401307010024
- Bresnick AR, Weber DJ, Zimmer DB. S100 proteins in cancer. *Nat Rev Cancer* (2015) **15**:96–109. doi:10.1038/nrc3893
- Rosenberg HF. Rnase a ribonucleases and host defense: an evolving story. *J Leukoc Biol* (2008) **83**:1079–87. doi:10.1189/jlb.1107725
- Simanski M, Kolen B, Schroder JM, Glaser R, Harder J. Antimicrobial RNases in cutaneous defense. *J Innate Immun* (2012) **4**:241–7. doi:10.1159/00035029
- Santaolalla R, Abreu MT. Innate immunity in the small intestine. *Curr Opin Gastroenterol* (2012) **28**:124–9. doi:10.1097/MOG.0b013e3283506559
- Royet J, Gupta D, Dziarski R. Peptidoglycan recognition proteins: modulators of the microbiome and inflammation. *Nat Rev Immunol* (2011) **11**:837–51. doi:10.1038/nri3089
- Patil A, Hughes AL, Zhang G. Rapid evolution and diversification of mammalian alpha-defensins as revealed by comparative analysis of rodent and primate genes. *Physiol Genomics* (2004) **20**:1–11. doi:10.1152/physiolgenomics.00150.2004
- Patil AA, Cai Y, Sang Y, Blecha F, Zhang G. Cross-species analysis of the mammalian beta-defensin gene family: presence of syntenic gene clusters and preferential expression in the male reproductive tract. *Physiol Genomics* (2005) **23**:5–17. doi:10.1152/physiolgenomics.00104.2005
- Meade KG, Cormican P, Narciandi F, Lloyd A, O'Farrelly C. Bovine beta-defensin gene family: opportunities to improve animal health? *Physiol Genomics* (2014) **46**:17–28. doi:10.1152/physiolgenomics.00085.2013
- Choi MK, Le MT, Nguyen DT, Choi H, Kim W, Kim JH, et al. Genome-level identification, gene expression, and comparative analysis of porcine ss-defensin genes. *BMC Genet* (2012) **13**:98. doi:10.1186/1471-2156-13-98
- Xiao Y, Hughes AL, Ando J, Matsuda Y, Cheng JF, Skinner-Noble D, et al. A genome-wide screen identifies a single beta-defensin gene cluster in the chicken: implications for the origin and evolution of mammalian defensins. *BMC Genomics* (2004) **5**:56. doi:10.1186/1471-2164-5-56
- Lynn DJ, Higgs R, Lloyd AT, O'Farrelly C, Herve-Grepinet V, Nys Y, et al. Avian beta-defensin nomenclature: a community proposed update. *Immunol Lett* (2007) **110**:86–9. doi:10.1016/j.imlet.2007.03.007
- Gudmundsson GH, Agerberth B, Odeberg J, Bergman T, Olsson B, Salcedo R. The human gene fall39 and processing of the cathelin precursor to the antibacterial peptide ll-37 in granulocytes. *Eur J Biochem* (1996) **238**:325–32. doi:10.1111/j.1432-1033.1996.03252.x
- Xiao Y, Cai Y, Bommineni YR, Fernando SC, Prakash O, Gilliland SE, et al. Identification and functional characterization of three chicken cathelicidins with potent antimicrobial activity. *J Biol Chem* (2006) **281**:2858–67. doi:10.1074/jbc.M507180200
- Goitsuka R, Chen CL, Benyon L, Asano Y, Kitamura D, Cooper MD. Chicken cathelicidin-b1, an antimicrobial guardian at the mucosal m cell gateway. *Proc Natl Acad Sci U S A* (2007) **104**:15063–8. doi:10.1073/pnas.0707037104

31. Whelehan CJ, Barry-Reidy A, Meade KG, Eckersall PD, Chapwanya A, Nanciandi F, et al. Characterisation and expression profile of the bovine cathelicidin gene repertoire in mammary tissue. *BMC Genomics* (2014) **15**:128. doi:10.1186/1471-2164-15-128
32. Sang Y, Blecha F. Porcine host defense peptides: expanding repertoire and functions. *Dev Comp Immunol* (2009) **33**:334–43. doi:10.1016/j.dci.2008.05.006
33. Lynn DJ, Higgs R, Gaines S, Tierney J, James T, Lloyd AT, et al. Bioinformatic discovery and initial characterisation of nine novel antimicrobial peptide genes in the chicken. *Immunogenetics* (2004) **56**:170–7. doi:10.1007/s00251-004-0675-0
34. Van Dijk A, Veldhuizen EJ, Van Asten AJ, Haagsman HP. Cmap27, a novel chicken cathelicidin-like antimicrobial protein. *Vet Immunol Immunopathol* (2005) **106**:321–7. doi:10.1016/j.vetimm.2005.03.003
35. Mabbott NA, Donaldson DS, Ohno H, Williams IR, Mahajan A. Microfold (m) cells: important immunosurveillance posts in the intestinal epithelium. *Mucosal Immunol* (2013) **6**:666–77. doi:10.1038/mi.2013.30
36. Wang G. Human antimicrobial peptides and proteins. *Pharmaceuticals (Basel)* (2014) **7**:545–94. doi:10.3390/ph7050545
37. Xiao Y, Dai H, Bommineni YR, Soulaiges JL, Gong YX, Prakash O, et al. Structure-activity relationships of fowlicidin-1, a cathelicidin antimicrobial peptide in chicken. *FEBS J* (2006) **273**:2581–93. doi:10.1111/j.1742-4658.2006.05261.x
38. Bommineni YR, Dai H, Gong YX, Soulaiges JL, Fernando SC, Desilva U, et al. Fowlicidin-3 is an alpha-helical cationic host defense peptide with potent antibacterial and lipopolysaccharide-neutralizing activities. *FEBS J* (2007) **274**:418–28. doi:10.1111/j.1742-4658.2006.05589.x
39. Xiao Y, Herrera AI, Bommineni YR, Soulaiges JL, Prakash O, Zhang G. The central kink region of fowlicidin-2, an alpha-helical host defense peptide, is critically involved in bacterial killing and endotoxin neutralization. *J Innate Immun* (2009) **1**:268–80. doi:10.1159/000174822
40. Van Dijk A, Veldhuizen EJ, Haagsman HP. Avian defensins. *Vet Immunol Immunopathol* (2008) **124**:1–18. doi:10.1016/j.vetimm.2007.12.006
41. Cuperus T, Coorens M, Van Dijk A, Haagsman HP. Avian host defense peptides. *Dev Comp Immunol* (2013) **41**:352–69. doi:10.1016/j.dci.2013.04.019
42. Zhang G, Sunkara LT. Avian antimicrobial host defense peptides: from biology to therapeutic applications. *Pharmaceuticals (Basel)* (2014) **7**:220–47. doi:10.3390/ph7030220
43. Brogden KA. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat Rev Microbiol* (2005) **3**:238–50. doi:10.1038/nrmicro1098
44. Peschel A, Sahl HG. The co-evolution of host cationic antimicrobial peptides and microbial resistance. *Nat Rev Microbiol* (2006) **4**:529–36. doi:10.1038/nrmicro1441
45. Nuding S, Zabel LT, Enders C, Porter E, Fellermann K, Wehkamp J, et al. Antibacterial activity of human defensins on anaerobic intestinal bacterial species: a major role of hbd-3. *Microbes Infect* (2009) **11**:384–93. doi:10.1016/j.micinf.2009.01.001
46. Bommineni YR, Pham GH, Sunkara LT, Achanta M, Zhang G. Immune regulatory activities of fowlicidin-1, a cathelicidin host defense peptide. *Mol Immunol* (2014) **59**:55–63. doi:10.1016/j.molimm.2014.01.004
47. Bommineni YR, Achanta M, Alexander J, Sunkara LT, Ritchey JW, Zhang G. A fowlicidin-1 analog protects mice from lethal infections induced by methicillin-resistant *Staphylococcus aureus*. *Peptides* (2010) **31**:1225–30. doi:10.1016/j.peptides.2010.03.037
48. Mackenzie-Dyck S, Attah-Poku S, Juillard V, Babiuk LA, Van Drunen Littel-Van Den Hurk S. The synthetic peptides bovine enteric beta-defensin (ebd), bovine neutrophil beta-defensin (bnbd) 9 and bnbd 3 are chemotactic for immature bovine dendritic cells. *Vet Immunol Immunopathol* (2011) **143**:87–107. doi:10.1016/j.vetimm.2011.06.028
49. Ikeda Y, Young LH, Scalia R, Ross CR, Lefer AM. Pr-39, a proline/arginine-rich antimicrobial peptide, exerts cardioprotective effects in myocardial ischemia-reperfusion. *Cardiovasc Res* (2001) **49**:69–77. doi:10.1016/S0008-6363(00)00226-1
50. Shi J, Ross CR, Leto TL, Blecha F. Pr-39, a proline-rich antibacterial peptide that inhibits phagocyte nadph oxidase activity by binding to src homology 3 domains of p47 phox. *Proc Natl Acad Sci U S A* (1996) **93**:6014–8. doi:10.1073/pnas.93.12.6014
51. Gallo RL, Ono M, Povsic T, Page C, Eriksson E, Klagsbrun M, et al. Syndecans, cell surface heparan sulfate proteoglycans, are induced by a proline-rich antimicrobial peptide from wounds. *Proc Natl Acad Sci U S A* (1994) **91**:11035–9. doi:10.1073/pnas.91.23.11035
52. Li J, Post M, Volk R, Gao Y, Li M, Metais C, et al. Pr39, a peptide regulator of angiogenesis. *Nat Med* (2000) **6**:49–55. doi:10.1038/71527
53. Baumann A, Demoulines T, Python S, Summerfield A. Porcine cathelicidins efficiently complex and deliver nucleic acids to plasmacytoid dendritic cells and can thereby mediate bacteria-induced ifn-alpha responses. *J Immunol* (2014) **193**:364–71. doi:10.4049/jimmunol.1303219
54. Corfield AP. Mucins: a biologically relevant glycan barrier in mucosal protection. *Biochim Biophys Acta* (2015) **1850**:236–52. doi:10.1016/j.bbagen.2014.05.003
55. Linden SK, Sutton P, Karlsson NG, Korolik V, McGuckin MA. Mucins in the mucosal barrier to infection. *Mucosal Immunol* (2008) **1**:183–97. doi:10.1038/mi.2008.5
56. Rose MC, Voynow JA. Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol Rev* (2006) **86**:245–78. doi:10.1152/physrev.00010.2005
57. Etzold S, Juge N. Structural insights into bacterial recognition of intestinal mucins. *Curr Opin Struct Biol* (2014) **28**:23–31. doi:10.1016/j.sbi.2014.07.002
58. Hollingsworth MA, Swanson BJ. Mucins in cancer: protection and control of the cell surface. *Nat Rev Cancer* (2004) **4**:45–60. doi:10.1038/nrc1251
59. Hoorens PR, Rinaldi M, Li RW, Goddeeris B, Claerebout E, Vercruyse J, et al. Genome wide analysis of the bovine mucin genes and their gastrointestinal transcription profile. *BMC Genomics* (2011) **12**:140. doi:10.1186/1471-2164-12-140
60. Kim CH, Oh Y, Ha Y, Ahn Q, Kim SH, Cho KD, et al. Expression of mucins in the mucosal surface of small intestines in 1 week-old pigs. *J Vet Med Sci* (2010) **72**:245–7. doi:10.1292/jvms.09-0308
61. Kim CH, Lee K, Han K, Oh Y, Kim D, Seo HW, et al. Expression of mucins on the mucosal surface of lungs of 4-week-old pigs. *Vet Res Commun* (2011) **35**:255–60. doi:10.1007/s11259-011-9465-y
62. Lang T, Hansson GC, Samuelsson T. An inventory of mucin genes in the chicken genome shows that the mucin domain of Muc13 is encoded by multiple exons and that ovomucin is part of a locus of related gel-forming mucins. *BMC Genomics* (2006) **7**:197. doi:10.1186/1471-2164-7-197
63. Van Der Sluis M, De Koning BA, De Bruijn AC, Velcich A, Meijerink JP, Van Goudoever JB, et al. Muc2-deficient mice spontaneously develop colitis, indicating that Muc2 is critical for colonic protection. *Gastroenterology* (2006) **131**:117–29. doi:10.1053/j.gastro.2006.04.020
64. Bergstrom KS, Kisson-Singh V, Gibson DL, Ma C, Montero M, Sham HP, et al. Muc2 protects against lethal infectious colitis by disassociating pathogenic and commensal bacteria from the colonic mucosa. *PLoS Pathog* (2010) **6**:e1000902. doi:10.1371/journal.ppat.1000902
65. McAuley JL, Linden SK, Png CW, King RM, Pennington HL, Gendler SJ, et al. Muc1 cell surface mucin is a critical element of the mucosal barrier to infection. *J Clin Invest* (2007) **117**:2313–24. doi:10.1172/JCI26705
66. McGuckin MA, Every AL, Skene CD, Linden SK, Chionh YT, Swierczak A, et al. Muc1 mucin limits both *Helicobacter pylori* colonization of the murine gastric mucosa and associated gastritis. *Gastroenterology* (2007) **133**:1210–8. doi:10.1053/j.gastro.2007.07.003
67. Zarepour M, Bhullar K, Montero M, Ma C, Huang T, Velcich A, et al. The mucin muc2 limits pathogen burdens and epithelial barrier dysfunction during *Salmonella enterica* serovar typhimurium colitis. *Infect Immun* (2013) **81**:3672–83. doi:10.1128/IAI.00854-13
68. An G, Wei B, Xia B, McDaniel JM, Ju T, Cummings RD, et al. Increased susceptibility to colitis and colorectal tumors in mice lacking core 3-derived O-glycans. *J Exp Med* (2007) **204**:1417–29. doi:10.1084/jem.20061929
69. Buisine MP, Desreumaux P, Debailleul V, Gambiez L, Geboes K, Ectors N, et al. Abnormalities in mucin gene expression in Crohn's disease. *Inflamm Bowel Dis* (1999) **5**:24–32. doi:10.1002/ibd.3780050105
70. Corfield AP, Carroll D, Myerscough N, Probert CS. Mucins in the gastrointestinal tract in health and disease. *Front Biosci* (2001) **6**:D1321–57. doi:10.2741/Corfield
71. Peterson LW, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol* (2014) **14**:141–53. doi:10.1038/nri3608
72. Suzuki T. Regulation of intestinal epithelial permeability by tight junctions. *Cell Mol Life Sci* (2013) **70**:631–59. doi:10.1007/s00018-012-1070-x

73. Gunzel D, Yu AS. Claudins and the modulation of tight junction permeability. *Physiol Rev* (2013) **93**:525–69. doi:10.1152/physrev.00019.2012
74. Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S, et al. Occludin: a novel integral membrane protein localizing at tight junctions. *J Cell Biol* (1993) **123**:1777–88. doi:10.1083/jcb.123.6.1777
75. Cummins PM. Occludin: one protein, many forms. *Mol Cell Biol* (2012) **32**:242–50. doi:10.1128/MCB.06029-11
76. Luissint AC, Nusrat A, Parkos CA. JAM-related proteins in mucosal homeostasis and inflammation. *Semin Immunopathol* (2014) **36**:211–26. doi:10.1007/s00281-014-0421-0
77. Kojima T, Ninomiya T, Konno T, Kohno T, Taniguchi M, Sawada N. Expression of tricellulin in epithelial cells and non-epithelial cells. *Histol Histopathol* (2013) **28**:1383–92. doi:10.14670/HH-28.1383
78. Herve JC, Derangeon M, Sarrouihe D, Bourmeyster N. Influence of the scaffolding protein zonula occludens (ZO) on membrane channels. *Biochim Biophys Acta* (2014) **1838**:595–604. doi:10.1016/j.bbame.2013.07.006
79. Citi S, Pulimeno P, Paschoud S. Cingulin, paracingulin, and plekha7: signaling and cytoskeletal adaptors at the apical junctional complex. *Ann N Y Acad Sci* (2012) **1257**:125–32. doi:10.1111/j.1749-6632.2012.06506.x
80. Suzuki H, Nishizawa T, Tani K, Yamazaki Y, Tamura A, Ishitani R, et al. Crystal structure of a claudin provides insight into the architecture of tight junctions. *Science* (2014) **344**:304–7. doi:10.1126/science.1248571
81. Suzuki H, Tani K, Tamura A, Tsukita S, Fujiyoshi Y. Model for the architecture of claudin-based paracellular ion channels through tight junctions. *J Mol Biol* (2015) **427**:291–7. doi:10.1016/j.jmb.2014.10.020
82. Mariano C, Sasaki H, Brites D, Brito MA. A look at tricellulin and its role in tight junction formation and maintenance. *Eur J Cell Biol* (2011) **90**:787–96. doi:10.1016/j.ejcb.2011.06.005
83. Ebnet K, Suzuki A, Ohno S, Vestweber D. Junctional adhesion molecules (JAMs): more molecules with dual functions? *J Cell Sci* (2004) **117**:19–29. doi:10.1242/jcs.00930
84. Al-Sadi R, Khatib K, Guo S, Ye D, Youssef M, Ma T. Occludin regulates macromolecule flux across the intestinal epithelial tight junction barrier. *Am J Physiol Gastrointest Liver Physiol* (2011) **300**:G1054–64. doi:10.1152/ajpgi.00055.2011
85. Cunningham KE, Turner JR. Myosin light chain kinase: pulling the strings of epithelial tight junction function. *Ann N Y Acad Sci* (2012) **1258**:34–42. doi:10.1111/j.1749-6632.2012.06526.x
86. Ye D, Ma I, Ma TY. Molecular mechanism of tumor necrosis factor- α modulation of intestinal epithelial tight junction barrier. *Am J Physiol Gastrointest Liver Physiol* (2006) **290**:G496–504. doi:10.1152/ajpgi.00318.2005
87. Al-Sadi R, Ye D, Dokladny K, Ma TY. Mechanism of IL-1 β -induced increase in intestinal epithelial tight junction permeability. *J Immunol* (2008) **180**:5653–61. doi:10.4049/jimmunol.180.8.5653
88. Sonoda N, Furuse M, Sasaki H, Yonemura S, Katahira J, Horiguchi Y, et al. *Clostridium perfringens* enterotoxin fragment removes specific claudins from tight junction strands: evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol* (1999) **147**:195–204. doi:10.1083/jcb.147.1.195
89. Smith F, Clark JE, Overman BL, Tozel CC, Huang JH, Rivier JE, et al. Early weaning stress impairs development of mucosal barrier function in the porcine intestine. *Am J Physiol Gastrointest Liver Physiol* (2010) **298**:G352–63. doi:10.1152/ajpgi.00081.2009
90. Hu CH, Xiao K, Luan ZS, Song J. Early weaning increases intestinal permeability, alters expression of cytokine and tight junction proteins, and activates mitogen-activated protein kinases in pigs. *J Anim Sci* (2013) **91**:1094–101. doi:10.2527/jas.2012-5796
91. Antoni L, Nuding S, Wehkamp J, Stange EF. Intestinal barrier in inflammatory bowel disease. *World J Gastroenterol* (2014) **20**:1165–79. doi:10.3748/wjg.v20.i5.1165
92. Wehkamp J, Salzman NH, Porter E, Nuding S, Weichenthal M, Petras RE, et al. Reduced paneth cell alpha-defensins in ileal crohn's disease. *Proc Natl Acad Sci U S A* (2005) **102**:18129–34. doi:10.1073/pnas.0505256102
93. Peyrin-Biroulet L, Beisner J, Wang G, Nuding S, Oommen ST, Kelly D, et al. Peroxisome proliferator-activated receptor gamma activation is required for maintenance of innate antimicrobial immunity in the colon. *Proc Natl Acad Sci U S A* (2010) **107**:8772–7. doi:10.1073/pnas.0905745107
94. Wehkamp J, Fellermann K, Herrlinger KR, Baxmann S, Schmidt K, Schwind B, et al. Human beta-defensin 2 but not beta-defensin 1 is expressed preferentially in colonic mucosa of inflammatory bowel disease. *Eur J Gastroenterol Hepatol* (2002) **14**:745–52. doi:10.1097/00042737-200207000-00006
95. Wehkamp J, Harder J, Weichenthal M, Mueller O, Herrlinger KR, Fellermann K, et al. Inducible and constitutive beta-defensins are differentially expressed in Crohn's disease and ulcerative colitis. *Inflamm Bowel Dis* (2003) **9**:215–23. doi:10.1097/00054725-200307000-00001
96. Fahlgren A, Hammarstrom S, Danielsson A, Hammarstrom ML. Beta-defensin-3 and -4 in intestinal epithelial cells display increased mRNA expression in ulcerative colitis. *Clin Exp Immunol* (2004) **137**:379–85. doi:10.1111/j.1365-2249.2004.02543.x
97. Schaubert J, Rieger D, Weiler F, Wehkamp J, Eck M, Fellermann K, et al. Heterogeneous expression of human cathelicidin hCAP18/LL-37 in inflammatory bowel diseases. *Eur J Gastroenterol Hepatol* (2006) **18**:615–21. doi:10.1097/00042737-200606000-00007
98. Wehkamp J, Harder J, Weichenthal M, Schwab M, Schaffeler E, Schlee M, et al. Nod2 (card15) mutations in Crohn's disease are associated with diminished mucosal alpha-defensin expression. *Gut* (2004) **53**:1658–64. doi:10.1136/gut.2003.032805
99. Kobayashi KS, Chamallard M, Ogura Y, Henegariu O, Inohara N, Nunez G, et al. Nod2-dependent regulation of innate and adaptive immunity in the intestinal tract. *Science* (2005) **307**:731–4. doi:10.1126/science.1104911
100. Aberg KM, Man MQ, Gallo RL, Ganz T, Crumrine D, Brown BE, et al. Co-regulation and interdependence of the mammalian epidermal permeability and antimicrobial barriers. *J Invest Dermatol* (2008) **128**:917–25. doi:10.1038/sj.jid.5701099
101. Otte JM, Werner I, Brand S, Chromik AM, Schmitz F, Kleine M, et al. Human beta defensin 2 promotes intestinal wound healing in vitro. *J Cell Biochem* (2008) **104**:2286–97. doi:10.1002/jcb.21787
102. Cobo ER, Kissoon-Singh V, Moreau F, Chadee K. Colonic muc2 mucin regulates the expression and antimicrobial activity of beta-defensin 2. *Mucosal Immunol* (2015) **8**(6):1360–72. doi:10.1038/mi.2015.27
103. Tai EK, Wong HP, Lam EK, Wu WK, Yu L, Koo MW, et al. Cathelicidin stimulates colonic mucus synthesis by up-regulating Muc1 and Muc2 expression through a mitogen-activated protein kinase pathway. *J Cell Biochem* (2008) **104**:251–8. doi:10.1002/jcb.21615
104. Otte JM, Zdebek AE, Brand S, Chromik AM, Strauss S, Schmitz F, et al. Effects of the cathelicidin LL-37 on intestinal epithelial barrier integrity. *Regul Pept* (2009) **156**:104–17. doi:10.1016/j.regpep.2009.03.009
105. Tang ZR, Deng H, Zhang XL, Zen Y, Xiao DF, Sun WZ, et al. Effects of orally administering the antimicrobial peptide buforin II on small intestinal mucosal membrane integrity, the expression of tight junction proteins and protective factors in weaned piglets challenged by enterotoxigenic *Escherichia coli*. *Anim Feed Sci Technol* (2013) **186**:177–85. doi:10.1016/j.anifeedsci.2013.10.012
106. Song D, Zong X, Zhang H, Wang T, Yi H, Luan C, et al. Antimicrobial peptide cathelicidin-BF prevents intestinal barrier dysfunction in a mouse model of endotoxemia. *Int Immunopharmacol* (2015) **25**:141–7. doi:10.1016/j.intimp.2015.01.017
107. Han F, Zhang H, Xia X, Xiong H, Song D, Zong X, et al. Porcine beta-defensin 2 attenuates inflammation and mucosal lesions in dextran sodium sulfate-induced colitis. *J Immunol* (2015) **194**:1882–93. doi:10.4049/jimmunol.1402300
108. Zhang Y, Zhu M, Yang Z, Pan X, Jiang Y, Sun C, et al. The human cathelicidin LL-37 induces muc5ac mucin production by airway epithelial cells via TACE-TGF- α -EGFR pathway. *Exp Lung Res* (2014) **40**:333–42. doi:10.3109/01902148.2014.926434
109. Akiyama T, Niyonsaba F, Kiatsurayanon C, Nguyen TT, Ushio H, Fujimura T, et al. The human cathelicidin LL-37 host defense peptide upregulates tight junction-related proteins and increases human epidermal keratinocyte barrier function. *J Innate Immun* (2014) **6**:739–53. doi:10.1159/000362789
110. Kiatsurayanon C, Niyonsaba F, Smithrithee R, Akiyama T, Ushio H, Hara M, et al. Host defense (antimicrobial) peptide, human beta-defensin-3, improves the function of the epithelial tight-junction barrier in human keratinocytes. *J Invest Dermatol* (2014) **134**:2163–73. doi:10.1038/jid.2014.143
111. Hattori F, Kiatsurayanon C, Okumura K, Ogawa H, Ikeda S, Okamoto K, et al. The antimicrobial protein S100a7/psoriasin enhances the expression of keratinocyte differentiation markers and strengthens the skin's tight junction barrier. *Br J Dermatol* (2014) **171**:742–53. doi:10.1111/bjd.13125

112. Choi KY, Chow LN, Mookherjee N. Cationic host defence peptides: multifaceted role in immune modulation and inflammation. *J Innate Immun* (2012) **4**:361–70. doi:10.1159/000336630
113. Erridge C. Endogenous ligands of TLR2 and TLR4: agonists or assistants? *J Leukoc Biol* (2010) **87**:989–99. doi:10.1189/jlb.1209775
114. Iitaka D, Moodley S, Shimizu H, Bai XH, Liu M. PKCdelta-iPLA2-PGE2-PPARgamma signaling cascade mediates TNF-alpha induced claudin 1 expression in human lung carcinoma cells. *Cell Signal* (2015) **27**:568–77. doi:10.1016/j.cellsig.2014.12.015
115. Iden S, Misselwitz S, Peddibhotla SS, Tuncay H, Rehder D, Gerke V, et al. aPKC phosphorylates JAM-A at Ser285 to promote cell contact maturation and tight junction formation. *J Cell Biol* (2012) **196**:623–39. doi:10.1083/jcb.201104143
116. Suzuki A, Ishiyama C, Hashiba K, Shimizu M, Ebnet K, Ohno S. aPKC kinase activity is required for the asymmetric differentiation of the premature junctional complex during epithelial cell polarization. *J Cell Sci* (2002) **115**:3565–73. doi:10.1242/jcs.00032
117. Beurel E, Michalek SM, Joep RS. Innate and adaptive immune responses regulated by glycogen synthase kinase-3 (GSK3). *Trends Immunol* (2010) **31**:24–31. doi:10.1016/j.it.2009.09.007
118. Severson EA, Kwon M, Hilgarth RS, Parkos CA, Nusrat A. Glycogen synthase kinase 3 (GSK-3) influences epithelial barrier function by regulating occludin, claudin-1 and e-cadherin expression. *Biochem Biophys Res Commun* (2010) **397**:592–7. doi:10.1016/j.bbrc.2010.05.164
119. González-Mariscal L, Tapia R, Chamorro D. Crosstalk of tight junction components with signaling pathways. *Biochim Biophys Acta* (2008) **1778**:729–56. doi:10.1016/j.bbame.2007.08.018
120. Little D, Dean RA, Young KM, McKane SA, Martin LD, Jones SL, et al. PI3K signaling is required for prostaglandin-induced mucosal recovery in ischemia-injured porcine ileum. *Am J Physiol Gastrointest Liver Physiol* (2003) **284**:G46–56. doi:10.1152/ajpgi.00121.2002
121. Woo PL, Ching D, Guan Y, Firestone GL. Requirement for ras and phosphatidylinositol 3-kinase signaling uncouples the glucocorticoid-induced junctional organization and transepithelial electrical resistance in mammary tumor cells. *J Biol Chem* (1999) **274**:32818–28. doi:10.1074/jbc.274.46.32818
122. Su B, Karin M. Mitogen-activated protein kinase cascades and regulation of gene expression. *Curr Opin Immunol* (1996) **8**:402–11. doi:10.1016/S0952-7915(96)80131-2
123. Lozupone CA, Stombaugh JJ, Gordon JI, Jansson JK, Knight R. Diversity, stability and resilience of the human gut microbiota. *Nature* (2012) **489**:220–30. doi:10.1038/nature11550
124. Saulnier DM, Kolida S, Gibson GR. Microbiology of the human intestinal tract and approaches for its dietary modulation. *Curr Pharm Des* (2009) **15**:1403–14. doi:10.2174/138161209788168128
125. Salzman NH. Paneth cell defensins and the regulation of the microbiome: detente at mucosal surfaces. *Gut Microbes* (2010) **1**:401–6. doi:10.4161/gmic.1.6.14076
126. McDermott AJ, Huffnagle GB. The microbiome and regulation of mucosal immunity. *Immunology* (2014) **142**:24–31. doi:10.1111/imm.12231
127. Kabat AM, Srinivasan N, Maloy KJ. Modulation of immune development and function by intestinal microbiota. *Trends Immunol* (2014) **35**:507–17. doi:10.1016/j.it.2014.07.010
128. Ostaff MJ, Stange EF, Wehkamp J. Antimicrobial peptides and gut microbiota in homeostasis and pathology. *EMBO Mol Med* (2013) **5**:1465–83. doi:10.1002/emmm.201201773
129. Koon HW, Shih DQ, Chen J, Bakirtzi K, Hing TC, Law I, et al. Cathelicidin signaling via the toll-like receptor protects against colitis in mice. *Gastroenterology* (2011) **141**(1852–63):e1–3. doi:10.1053/j.gastro.2011.06.079
130. Salzman NH, Ghosh D, Huttner KM, Paterson Y, Bevins CL. Protection against enteric salmonellosis in transgenic mice expressing a human intestinal defensin. *Nature* (2003) **422**:522–6. doi:10.1038/nature01520
131. Wilson CL, Ouellette AJ, Satchell DP, Ayabe T, Lopez-Boado YS, Stratman JL, et al. Regulation of intestinal alpha-defensin activation by the metalloproteinase matrilysin in innate host defense. *Science* (1999) **286**:113–7. doi:10.1126/science.286.5437.113
132. Salzman NH, Hung K, Haribhai D, Chu H, Karlsson-Sjoberg J, Amir E, et al. Enteric defensins are essential regulators of intestinal microbial ecology. *Nat Immunol* (2010) **11**:76–83. doi:10.1038/ni.1825
133. Xiao H, Shao F, Wu M, Ren W, Xiong X, Tan B, et al. The application of antimicrobial peptides as growth and health promoters for swine. *J Anim Sci Biotechnol* (2015) **6**:19. doi:10.1186/s40104-015-0018-z
134. Wang S, Thacker PA, Watford M, Qiao S. Functions of antimicrobial peptides in gut homeostasis. *Curr Protein Pept Sci* (2015) **16**:582–91. doi:10.2174/1389203716666150630135847
135. Cutler SA, Lonergan SM, Cornick N, Johnson AK, Stahl CH. Dietary inclusion of colicin E1 is effective in preventing postweaning diarrhea caused by F18-positive *Escherichia coli* in pigs. *Antimicrob Agents Chemother* (2007) **51**:3830–5. doi:10.1128/AAC.00360-07
136. Wu S, Zhang F, Huang Z, Liu H, Xie C, Zhang J, et al. Effects of the antimicrobial peptide cecropin ad on performance and intestinal health in weaned piglets challenged with *Escherichia coli*. *Peptides* (2012) **35**:225–30. doi:10.1016/j.peptides.2012.03.030
137. Tang XS, Fatufe AA, Yin YL, Tang ZR, Wang SP, Liu ZQ, et al. Dietary supplementation with recombinant lactoferrin-lactoferricin improves growth performance and affects serum parameters in piglets. *J Anim Vet Adv* (2012) **11**:2548–55. doi:10.3923/javaa.2012.2548.2555
138. Xiong X, Yang HS, Li L, Wang YF, Huang RL, Li FN, et al. Effects of antimicrobial peptides in nursery diets on growth performance of pigs reared on five different farms. *Livest Sci* (2014) **167**:206–10. doi:10.1016/j.livsci.2014.04.024
139. Yoon JH, Ingale SL, Kim JS, Kim KH, Lohakare J, Park YK, et al. Effects of dietary supplementation with antimicrobial peptide-P5 on growth performance, apparent total tract digestibility, faecal and intestinal microflora and intestinal morphology of weanling pigs. *J Sci Food Agric* (2013) **93**:587–92. doi:10.1002/jsfa.5840
140. Yoon JH, Ingale SL, Kim JS, Kim KH, Lee SH, Park YK, et al. Effects of dietary supplementation of synthetic antimicrobial peptide-A3 and P5 on growth performance, apparent total tract digestibility of nutrients, fecal and intestinal microflora and intestinal morphology in weanling pigs. *Livest Sci* (2014) **159**:53–60. doi:10.1016/j.livsci.2013.10.025
141. Xiao H, Wu MM, Tan BE, Yin YL, Li TJ, Xiao DF, et al. Effects of composite antimicrobial peptides in weanling piglets challenged with deoxynivalenol: I. Growth performance, immune function, and antioxidation capacity. *J Anim Sci* (2013) **91**:4772–80. doi:10.2527/jas.2013-6426
142. Xiao H, Tan BE, Wu MM, Yin YL, Li TJ, Yuan DX, et al. Effects of composite antimicrobial peptides in weanling piglets challenged with deoxynivalenol: II. Intestinal morphology and function. *J Anim Sci* (2013) **91**:4750–6. doi:10.2527/jas.2013-6427
143. Choi SC, Ingale SL, Kim JS, Park YK, Kwon IK, Chae BJ. An antimicrobial peptide-A3: effects on growth performance, nutrient retention, intestinal and faecal microflora and intestinal morphology of broilers. *Br Poult Sci* (2013) **54**:738–46. doi:10.1080/00071668.2013.838746
144. Wen LF, He JG. Dose-response effects of an antimicrobial peptide, a cecropin hybrid, on growth performance, nutrient utilisation, bacterial counts in the digesta and intestinal morphology in broilers. *Br J Nutr* (2012) **108**:1756–63. doi:10.1017/S0007114511007240
145. Schaubert J, Svanholm C, Termen S, Iffland K, Menzel T, Scheppach W, et al. Expression of the cathelicidin LL-37 is modulated by short chain fatty acids in colonocytes: relevance of signalling pathways. *Gut* (2003) **52**:735–41. doi:10.1136/gut.52.5.735
146. Raqib R, Sarker P, Bergman P, Ara G, Lindh M, Sack DA, et al. Improved outcome in shigellosis associated with butyrate induction of an endogenous peptide antibiotic. *Proc Natl Acad Sci U S A* (2006) **103**:9178–83. doi:10.1073/pnas.0602888103
147. Ochoa-Zarzosa A, Villarreal-Fernandez E, Cano-Camacho H, Lopez-Meza JE. Sodium butyrate inhibits *Staphylococcus aureus* internalization in bovine mammary epithelial cells and induces the expression of antimicrobial peptide genes. *Microb Pathog* (2009) **47**:1–7. doi:10.1016/j.micpath.2009.04.006
148. Sunkara LT, Achanta M, Schreiber NB, Bommineni YR, Dai G, Jiang W, et al. Butyrate enhances disease resistance of chickens by inducing antimicrobial host defense peptide gene expression. *PLoS One* (2011) **6**:e27225. doi:10.1371/journal.pone.0027225
149. Sunkara LT, Jiang W, Zhang G. Modulation of antimicrobial host defense peptide gene expression by free fatty acids. *PLoS One* (2012) **7**:e49558. doi:10.1371/journal.pone.0049558

150. Zeng X, Sunkara LT, Jiang W, Bible M, Carter S, Ma X, et al. Induction of porcine host defense peptide gene expression by short-chain fatty acids and their analogs. *PLoS One* (2013) **8**:e72922. doi:10.1371/journal.pone.0072922
151. Sunkara LT, Zeng X, Curtis AR, Zhang G. Cyclic AMP synergizes with butyrate in promoting beta-defensin 9 expression in chickens. *Mol Immunol* (2014) **57**:171–80. doi:10.1016/j.molimm.2013.09.003
152. Lyu W, Curtis AR, Sunkara LT, Zhang G. Transcriptional regulation of antimicrobial host defense peptides. *Curr Protein Pept Sci* (2015) **16**:672–9. doi:10.2174/1389203716666150630133432

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Robinson, Deng, Hou and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.