



Erratum: Cingulate Cortex Atrophy Is Associated With Hearing Loss in Presbycusis With Cochlear Amplifier Dysfunction

Frontiers Production Office*

Frontiers Media SA, Lausanne, Switzerland

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Frontiers Production Office
production.office@frontiersin.org

Received: 16 May 2019

Accepted: 17 May 2019

Published: 12 June 2019

Citation:

Frontiers Production Office (2019)
Erratum: Cingulate Cortex Atrophy Is
Associated With Hearing Loss in
Presbycusis With Cochlear Amplifier
Dysfunction.
Front. Aging Neurosci. 11:131.
doi: 10.3389/fnagi.2019.00131

Keywords: presbycusis, neurodegeneration, dementia, cognition, cochlear amplifier

An Erratum on

Cingulate Cortex Atrophy Is Associated With Hearing Loss in Presbycusis With Cochlear Amplifier Dysfunction

by Belkhiria, C., Vergara, R. C., San Martín, S., Leiva, A., Marcenaro, B., Martinez, M., et al. (2019).
Front. Aging Neurosci. 11:97. doi: 10.3389/fnagi.2019.00097

Due to a production error, **Table 4** was erroneously typeset, thereby three rows were deleted. The correct Table is provided below. The publisher apologizes for this mistake. The original article has been updated.

Copyright © 2019 Frontiers Production Office. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 4 | Pearson partial correlation corrected by age, sex, and education between cognitive tests and volume and thickness of anterior and posterior cingulate cortex for PCF and CD groups (* $p < 0.05$).

Cognitive tests	ACC volume				ACC thickness				PCC volume				PCC thickness			
	PCF		CD		PCF		CD		PCF		CD		PCF		CD	
	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right	r-Left	r-Right
Cognitive functions (MMSE)	0.14	0.46*	-0.19	0.00	0.22	0.17	0.03	-0.13	0.05	0.40	-0.10	0.00	0.14	0.14	-0.01	-0.19
Executive functions (FAB)	0.29	0.07	-0.01	0.21	0.14	0.26	0.62	0.13	-0.12	-0.13	-0.01	0.02	-0.03	0.04	-0.11	-0.24
Visuospatial capacities (Rey figure)	0.05	0.41*	0.14	0.30	0.01	0.28	0.15	0.49*	0.32	0.27	0.35	0.54*	0.00	0.02	0.35	0.45*
Processing speed (TMT A)	-0.27	0.01	-0.37	-0.04	-0.22	-0.17	-0.34*	-0.26	-0.40*	-0.21	-0.41*	-0.31	0.02	-0.01	-0.20	-0.11
Episodic Memory (FCSRT)	-0.20	0.19	0.44*	0.31	0.01	0.08	0.34	0.10	-0.26	0.05	0.28	0.22	0.05	0.11	0.08	-0.18
Nomination (Boston)	0.21	0.03	0.47*	0.29	0.03	0.07	0.48*	0.41*	0.19	0.11	0.23	0.31	-0.16	0.03	0.28	0.20
Working memory (backward digit n ^o)	0.10	-0.14	0.64*	0.40	-0.06	-0.42	0.18	0.07	0.10	0.05	0.45*	0.18	0.12	0.40	-0.35	0.00
Depression score (GDS-15)	0.32	-0.20	-0.53*	-0.01	-0.17	0.36	0.05	0.00	0.17	0.00	-0.12	0.04	0.27	0.35	0.25	-0.09