



## OPEN ACCESS

## EDITED BY

Yi Min Wan,  
National University Health System  
(Singapore), Singapore, Singapore

## REVIEWED BY

Christian Barbato,  
National Research Council (CNR), Roma, Italy  
Carla Masala,  
University of Cagliari, Cagliari, Italy

## \*CORRESPONDENCE

Ding Ding  
✉ dingding@huashan.org.cn

RECEIVED 02 April 2024

ACCEPTED 24 May 2024

PUBLISHED 10 June 2024

## CITATION

Liang X, Xiao Z, Wu J, Ma X, Zhao Q and  
Ding D (2024) Serum TRPA1 mediates the  
association between olfactory function and  
cognitive function.

*Front. Aging Neurosci.* 16:1411031.

doi: 10.3389/fnagi.2024.1411031

## COPYRIGHT

© 2024 Liang, Xiao, Wu, Ma, Zhao and Ding.  
This is an open-access article distributed  
under the terms of the [Creative Commons  
Attribution License \(CC BY\)](#). The use,  
distribution or reproduction in other forums is  
permitted, provided the original author(s) and  
the copyright owner(s) are credited and that  
the original publication in this journal is cited,  
in accordance with accepted academic  
practice. No use, distribution or reproduction  
is permitted which does not comply with  
these terms.

# Serum TRPA1 mediates the association between olfactory function and cognitive function

Xiaoniu Liang<sup>1,2,3</sup>, Zhenxu Xiao<sup>1,2,3</sup>, Jie Wu<sup>1,2,3</sup>, Xiaoxi Ma<sup>1,2,3</sup>,  
Qianhua Zhao<sup>1,2,3</sup> and Ding Ding<sup>1,2,3\*</sup>

<sup>1</sup>Institute of Neurology, Huashan Hospital, Fudan University, Shanghai, China, <sup>2</sup>National Clinical Research Center for Aging and Medicine, Huashan Hospital, Fudan University, Shanghai, China, <sup>3</sup>National Center for Neurological Disorders, Huashan Hospital, Fudan University, Shanghai, China

**Background:** Olfactory dysfunction was associated with poorer cognition. However, the association between transient receptor potential cation channel subfamily A member 1 (TRPA1) and cognitive function have not been studied. This study aimed to evaluate the mediation effect of TRPA1 on the association between olfactory and cognitive function among Chinese older adults.

**Methods:** We recruited 121 participants with cognitive impairment (CI) and 135 participants with normal cognition (NC) from a memory clinic and the “Shanghai Aging Study.” Olfactory identification of each participant was measured by the Sniffin’ Sticks Screening Test 12 (SSST-12). Serum TRPA1 were quantified using the Enzyme-Linked Immunosorbent Assay. The mediation effects of TRPA1 on the association between olfactory function and cognitive function were explored using mediation analysis.

**Results:** The CI group had a significantly higher proportion of the high level of serum TRPA1 (58.7%) than the NC group (42.2%) ( $p = 0.0086$ ). After adjusted for gender, age, and years of education, mediation analysis verified that TRPA1 partially mediated the association between SSST-12 and Mini Mental State Examination (MMSE). It also verified that TRPA1 partially mediated the association between the identification of peppermint and MMSE.

**Conclusion:** Our study emphasizes the mediation role of TRPA1 in the relationship between olfactory and cognitive function among older adults. Further research is necessary to explore the mechanism of TRPA1 on the relationship between olfactory and cognitive decline.

## KEYWORDS

cognitive function, olfactory identification, TRPA1, mediation analysis, elderly

## Introduction

Olfactory dysfunction affects 24–75% of older adults, with its prevalence substantially increasing by age (Choi et al., 2018). Olfactory dysfunction is an important clinical symptom indicating an early stage of neurodegenerative diseases, including Alzheimer’s disease (AD), frontotemporal lobar degeneration, dementia with Lewy bodies, and Parkinson’s disease (Attems et al., 2014). Olfactory identification is likely to involve the most elaborate network of brain regions associated with language processing, semantic and episodic memory, as well as olfactory perception (Wilson et al., 2006). Several epidemiologic studies demonstrated that

poorer olfactory function was significantly associated with poorer cognition in population-based cohorts (Stanciu et al., 2014; Devanand et al., 2015; Liang et al., 2016; Roberts et al., 2016; Yaffe et al., 2017).

Our previous cross-sectional study showed that participants with mild cognitive impairment (MCI) performed worse olfactory function, especially the ability to identify peppermint among 12 odors in the Sniffin' Sticks Screening Test 12 (SSST-12), than those with normal cognition (NC) (Liang et al., 2016). Our cohort study showed that an inability to smell peppermint was related to a higher risk for incident dementia, and was inversely associated with the annual rate of change in the Mini Mental State Examination (MMSE) score (Liang et al., 2020).

Peppermint was found to enhance memory (Moss et al., 2009) and the sustained visual attention task (Warm et al., 1991). Peppermint was mainly composed of L-menthol, carboxyl esters, menthone, menthyl acetate (Thomson Healthcare, 2007) and other components (Leung, 1980). The main cellular receptor for L-menthol is transient receptor potential cation channel subfamily A member 1 (TRPA1), which is associated with AD (Lee et al., 2016; Bosson et al., 2017). Studies have shown that the protein expression of TRPA1 channel significantly increased in AD transgenic mice, mainly in astrocytes of the hippocampus, and TRPA1 could be involved in the pathogenesis of AD through multiple pathways (Lee et al., 2016; Bosson et al., 2017; Paumier et al., 2022).

In this pilot study, we aimed to evaluate the potential mediation effect of TRPA1 on the association between olfactory and cognitive function in Chinese older adults.

## Materials and methods

### Study participants

Participants with cognitive impairment (CI) were from the memory clinic of the department of neurology, Huashan Hospital, Shanghai, China from December 2020 to May 2021. Participants met the inclusion criteria if they (1) visited the memory clinic due to memory complaints from themselves or the proxy; (2) were diagnosed with AD clinical syndrome or MCI; (3) had the ability to cooperate with neuropsychological tests and physical examinations; (4) completed the olfactory assessment; (5) 60 years or older; and (6) agreed to the blood draw.

The Shanghai Aging Study (SAS) is a community-based cohort in downtown Shanghai, China. SAS aimed to explore the prevalence, incidence, and risk factors for MCI and dementia among older residents. The detailed recruitment procedure and study design have been published elsewhere (Ding et al., 2014). In the SAS, the participants were diagnosed with NC from the third wave of follow-up between June 2020 and November 2021 if they (1) completed the olfactory assessment; (2) had the ability to cooperate with neuropsychological tests and physical examinations; (3) 60 years or older; and (4) agreed to the blood draw.

We excluded participants who (1) had histories of rhinal or paranasal sinuses diseases; (2) underwent maxillofacial surgery; (3) had chronic sinusitis, asthma, chronic obstructive pulmonary disease, or acute upper respiratory tract infection within 7 days before the olfactory assessment; and (4) alcohol or drug abuse.

### Demographics, lifestyles, and medical history

The lifestyle and demographic characteristics were collected from the participants and/or proxy through questionnaires, including gender, age, and years of education. Hypertension and diabetes mellitus were confirmed by the medical records (Ding et al., 2014).

### Olfactory identification test

Olfactory identification was a procedure in which a participant needed to accurately identify olfactory stimuli using alternative choices. Olfactory identification test was assessed by the SSST-12, which includes 12 common odors (orange, cinnamon, leather, banana, peppermint, liquorice, lemon, cloves, coffee, rose, pineapple, and fish) presented on felt-tip sticks (Hummel et al., 2001). The SSST-12 was devised by G. Kobal in Erlangen, Germany, and it is a portable, rapid (approximately 6 min), suited for inexpensive and repetitive screening of olfactory identification. The administrator of SSST-12 was blind for the cognitive diagnosis of each participant. Before the test, participants were reminded to stay away from chewing sweets, cigarettes or gum. Testing was performed in an air-conditioned and quiet room. A brief history was recorded, including questions related to the participant's previous diseases, olfactory experience, occupation, drug intake and smoking habits. When presenting the odors, the administrator was wearing cotton gloves. The opened odor sticks were positioned about 2 cm in front of both nostrils of each participant. Participants were then asked to sniff for no longer than 3–4 s and to choose one of four answers from a list that described the best odor. An interval of 30 s was set between the different sticks. Detailed instructions of SSST-12 were reported elsewhere (Liang et al., 2016).

### Neurological, neuropsychological assessments, and consensus diagnosis

Comprehensive neuropsychological tests were administered by the certified psychometrists in accordance with the education level of each participant. All tests conducted had been translated, adapted, and validated within the local Chinese population.

Each participant from the memory clinic received a battery of neuropsychological tests including (Xiao et al., 2021): (1) Montreal Cognitive Assessment-Basic (MoCA-B); (2) MMSE; (3) Auditory Verbal Learning Test; (4) Boston Naming Test; (5) Rey-Osterrieth Complex Figure test; (6) Symbol Digit Modalities Test; and (7) Trail-making test A&B. For those who were unable or refused to complete the whole battery of tests, only MoCA-B and MMSE were administered. Because the MMSE is less sensitive for MCI detection (Nasreddine et al., 2005), MMSE was used together with MoCA-B to discriminate MCI and dementia.

As for the participants from SAS, a battery of similar neuropsychological tests was administered due to the study design of SAS (Zhang et al., 1990; Ding et al., 2015): (1) MMSE; (2) Auditory Verbal Learning Test; (3) Conflicting Instructions Task (Go/No Go Task); (4) Modified Fuld Object Memory Evaluation; (5) Modified Common Objects Sorting Test; (6) RMB (Chinese currency) test; (7) Stick Test; and (8) Trail-making test A&B.

Two study neurologists, one neuroepidemiologist and one neuropsychologist reviewed the medical, functional, neuropsychological, neurological, and psychiatric data and reached a consensus regarding the absence or presence of dementia using DSM-IV criteria (American Psychiatric Association, 1994). Probable AD was diagnosed using the NINCDS-ADRDA criteria (McKhann et al., 1984). Participants who met the criterion of probable AD were regarded as having AD clinical syndrome. The diagnosis of MCI was based on Petersen's criteria (Petersen, 2004).

## Serum TRPA1 measurement

Blood was collected from the study participants. Serum and plasma samples were centrifuged, aliquoted, and stored at  $-80^{\circ}\text{C}$ . Serum TRPA1 were quantified using the Enzyme-Linked Immunosorbent Assay test on the Enzyme labeling apparatus (352 Labsystems Multiskan MS), and serum samples were diluted at a 1:4 ratio following the manufacturer's instructions. Duplicate measurements were taken for calibrators and quality controls. Sample measurements were conducted in a single run using kits with the identical lot numbers. Operators remained unaware of participants' characteristics. The statistical median value of serum TRPA1 was used to categorize participants into low or high levels of serum TRPA1.

## Statistical analysis

Continuous variables were expressed as the mean (standard deviation) or median (lower quartile[25%], upper quartile[75%]), and categorical variables were expressed as frequencies (%). The Wilcoxon rank-sum test and Pearson chi-square test were used to compare continuous and categorical variables.

Scatter plots were performed to analyze the correlations between SSST-12 and serum TRPA1, between SSST-12 and MMSE, and between serum TRPA1 and MMSE. Their correlations were assessed using the generalized linear model with the adjustment for gender, age, and years of education.

The logistic regression model was used to detect the association between serum TRPA1, the level of serum TRPA1, SSST-12, olfactory dysfunction, or the identification of peppermint and CI after adjusted for gender, age, and years of education. Risk was presented as odds ratio (OR) and 95% confidence interval (95%CI).

Mediation analysis was conducted to decompose the total effect of olfactory function on cognitive function into a natural direct effect and a natural indirect effect through TRPA1 after adjusted for gender, age, and years of education.

Two-tailed tests were used to estimate all  $p$  values and 95% confidence intervals. Statistically significant differences were identified at  $p < 0.05$ . Mediation analysis was conducted using R packages "lavaan" and "mediation." Data analysis was conducted in SAS 9.4 (SAS Institute Inc., Cary, NC, United States) and R Software (version 4.1.2).

## Results

### Characteristics of the participants

We recruited 256 participants (164 females and 92 males), including 135 participants with NC and 121 participants with CI. Table 1 displayed the characteristics of the study participants. Significant differences were observed in the level of serum TRPA1 ( $p = 0.0086$ ), education year ( $p = 0.0002$ ), MMSE ( $p < 0.0001$ ), and SSST-12 ( $p < 0.0001$ ) between the two groups. The CI groups had higher proportion of the high level of serum TRPA1 (58.7%), the lower median education year (median = 11), the lower median MMSE score (median = 27), and the lower median SSST-12 score (median = 7).



### Association analysis

SSST-12 were inversely correlated with serum TRPA1 in total participants ( $\beta = -8.0848$ , 95%CI:  $-12.6423, -3.5272$ ), participants with NC ( $\beta = -7.0532$ , 95%CI:  $-13.9778, -0.1286$ ) and participants with CI ( $\beta = -7.2309$ , 95%CI:  $-13.8897, -0.5721$ ) after adjusted for gender, age, and years of education (Figure 1). SSST-12 were

TABLE 1 Characteristics of study participants.

Clinical features	Total (N = 256)	Normal cognition (N = 135)	Cognitive impairment (N = 121)	p-value
Level of TRPA1, n (%)				
Low	128 (50.0)	78 (57.8)	50 (41.3)	<b>0.0086</b>
High	128 (50.0)	57 (42.2)	71 (58.7)	
TRPA1, pg./mL, mean $\pm$ SD	482.8 $\pm$ 88.0	472.6 $\pm$ 82.7	494.1 $\pm$ 92.5	0.0626
Gender, female, n (%)	164 (64.1)	89 (65.9)	75 (62.0)	0.5116
Age, year, median (Q1, Q3)	74 (71, 80)	73 (71, 78)	76 (71, 82)	0.0553
Education, year, median (Q1, Q3)	12 (9, 13)	12 (10, 15)	11 (9, 12)	<b>0.0002</b>
MMSE, score, median (Q1, Q3)	28 (26, 29)	29 (28, 30)	27 (25, 28)	<b>&lt;0.0001</b>
SSST-12, score, median (Q1, Q3)	8 (6, 9)	8 (7, 9)	7 (5, 9)	<b>&lt;0.0001</b>
Correct identification of peppermint, n (%)	210 (82.0)	115 (85.2)	95 (78.5)	0.2083

MMSE, mini-mental state examination; Q1, lower quartile; Q3, upper quartile; SSST-12, Sniffin' Sticks Screening Test 12; SD, standard deviation; TRPA1, transient receptor potential cation channel subfamily A member 1.





positively correlated with MMSE in total participants ( $\beta = 0.7217$ , 95%CI: 0.5454, 0.8980) and participants with CI ( $\beta = 0.9544$ , 95%CI: 0.6528, 1.2559) after adjusted for gender, age, and years of education (Figure 2). The serum TRPA1 level was inversely correlated with MMSE in total participants ( $\beta = -0.0088$ , 95%CI:  $-0.0138, -0.0037$ ) and participants with CI ( $\beta = -0.0093$ , 95%CI:  $-0.0182, -0.0003$ ) after adjusted for gender, age, and years of education (Figure 3).

As shown in Table 2, the higher serum TRPA1 level was associated with an increased risk for participants with CI (OR = 1.889, 95%CI: 1.125, 3.172) after adjusted for gender, age, and years of education. Lower SSST-12 score (OR = 0.776, 95%CI: 0.685, 0.878) and olfactory dysfunction (OR = 3.116, 95%CI: 1.592, 6.101) were associated with an increased risk for participants with CI after adjusted for gender, age, and years of education. The correct identification of peppermint may be a protector for participants with


CI (OR = 0.858, 95%CI: 0.428, 1.719), although the finding did not reach statistical significance.

## Mediation analysis

Figure 4 showed the total, direct and indirect effects for the mediating role of TRPA1 on the relationship between olfactory function and cognitive function in mediation models. As shown in Figure 4B, after adjusted for gender, age, and years of education, the estimated average causal mediated effect (ACME) (indirect effect estimate  $\beta = 0.0402$ , 95%CI: 0.0023, 0.0906), average direct effect (ADE) (direct effect estimate  $\beta = 0.6816$ , 95%CI: 0.3906, 0.9630), and total effects (total effect estimate  $\beta = 0.7217$ , 95%CI: 0.4193, 1.0015) were all statistically significant, suggesting a partial mediation effect of TRPA1 in the association between olfactory function and MMSE, and TRPA1 had the mediation effect with







a proportion of mediation up to 5.57%. After adjusted for gender, age, and years of education, the estimated ADE (direct effect estimate  $\beta = -0.0513$ , 95%CI:  $-0.0741, -0.0275$ ) and total effects (total effect estimate  $\beta = -0.0543$ , 95%CI:  $-0.0766, -0.0301$ ) were statistically significantly different from zero, but the estimated ACME were not (**Figure 4D**). The results suggested that lower SSST-12 may be associated with higher serum TRPA1, which in turn made participants more likely to have worse cognitive function.

**Figure 5** showed the total, direct and indirect effects for the mediating role of TRPA1 on the relationship between the identification of peppermint and cognitive function in mediation models. As shown in **Figure 5B**, after adjusted for gender, age, and years of education, the estimated ACME (indirect effect estimate  $\beta = 0.2975$ , 95%CI:  $0.0443, 0.6200$ ), ADE (direct effect estimate  $\beta = 2.8716$ , 95%CI:  $0.9727, 4.9200$ ), and total effects (total effect estimate  $\beta = 3.1691$ , 95%CI:  $1.2178, 5.3100$ ) were all statistically significant, suggesting a partial mediation effect of TRPA1 in the association between the identification of peppermint and MMSE, and TRPA1 had the mediation effect with a proportion of mediation up

to 9.39%. However, the estimated ACME, the estimated ADE and total effects were all not statistically significant (**Figure 5D**). The results suggested that the incorrect identification of peppermint may be associated with higher serum TRPA1, which in turn made participants more likely to have worse cognitive function.

## Discussion

This study marks the initial attempt to partition the effects of olfactory function on cognitive function into direct effects and indirect effects (mediated by serum TRPA1) among Chinese older adults. Mediation analysis of olfactory function (both SSST-12 and peppermint) with MMSE showed a partial mediation effect acting through serum TRPA1.

The association between SSST-12 and CI has been reported in a few epidemiologic studies. We previously demonstrated that lower SSST-12 score was related to MCI (OR = 1.19, 95%CI: 1.11, 1.27) in the older adults by using the baseline data of SAS (Liang et al., 2016). Poor

TABLE 2 Odds ratios for serum TRPA1 between participants with normal cognition and cognitive impairment.

	OR (95%CI)	OR (95%CI)*
Serum TRPA1	1.003(1.000, 1.006)	1.003(1.000, 1.006)
Level of TRPA1		
Low	1	1
High	1.943(1.181, 3.196)	1.889(1.125, 3.172)
SSST-12	0.754(0.671, 0.847)	0.776(0.685, 0.878)
Olfactory dysfunction		
No	1	1
Yes	3.537(1.853, 6.751)	3.116(1.592, 6.101)
Peppermint		
Wrong	1	1
Right	0.661(0.346, 1.263)	0.858(0.428, 1.719)

\*The OR (95%CI) was calculated by logistic regression after adjusted for gender, age, and years of education.

95%CI, 95% confidence interval; OR, odds ratio; SSST-12, Sniffin' Sticks Screening Test 12; TRPA1, transient receptor potential cation channel subfamily A member 1.

olfactory dysfunction (assessed by 16-item Sniffin' Sticks identification test [SSST-16]) was significantly associated with an increased risk for non-amnesic MCI, amnesic MCI, and MCI (Dong et al., 2023). However, to our knowledge, the effect of TRPA1 on olfactory dysfunction or CI has not been reported in any population-based studies. TRPA1 is closely associated with inflammation, pruritus, and chronic pain, and TRPA1 is considered to be a promising treatment for them (Hu et al., 2023). Increased TRPA1 mRNA expression in whole blood cells was significantly related to decreased pain symptoms in chronic pain patients (Bell et al., 2014; Sukenaga et al., 2016). Epidemiologic studies have shown that patients with AD reported less pain (Scherder et al., 1999; Mantyselka et al., 2004; Achterberg et al., 2010; Jensen-Dahm et al., 2012), and patients with dementia were less likely to use analgesics (Horgas and Tsai, 1998; Morrison and Siu, 2000; Mantyselka et al., 2004).

Multiple pathways may implicate TRPA1 in the pathogenesis of AD mice. TRPA1 receptors mediate deteriorating effects in the decline of memory (Borbely et al., 2019). Functional ablation of the TRPA1 channel in mice improved hippocampal functions, demonstrating by reduced anxiety-like behavior, improved fear-related or spatial learning and memory, novel location recognition and social interactions (Lee et al., 2017). Astrocytic TRPA1 and GABA coordinately suppress hippocampal circuit function (Cheng et al., 2023). TRPA1 is expressed on astrocytes in the hippocampus, and the production of amyloid- $\beta$  activates this channel, thereby initiated this hyperactivity and subsequently induced the hyperactivity of nearby neurons, which is a pivotal factor in the progression of AD (Bosson et al., 2017). The toxic effect of amyloid- $\beta$  on astrocytes, triggered by TRPA1 channel activation, is crucial to the progression of AD, and TRPA1 blockade prevents irreversible neuronal dysfunction (Paumier et al., 2022). TRPA1-Ca<sup>2+</sup>-PP2B signaling may be crucial to regulate pathogenesis of AD and astrocyte-derived inflammation (Lee et al., 2016). Melatonin may be an effective option in the treatment and prophylaxis of AD by reducing cytosolic Ca<sup>2+</sup> concentration, apoptosis and intracellular ROS through TRPA1 channels (Ozsimsek and Ovey, 2022). These studies showed that blocking TRPA1

could prevent irreversible neuronal dysfunction, and TRPA1 might be a potential therapeutic target for neuroprotection (Hu et al., 2023).

Several advantages existed in our study. Firstly, neuropsychological assessments and the diagnosis of CI were administered by the certified neurologists and neuropsychologists. Secondly, TRPA1 was tested among the patients with CI for the first time. Thirdly, we studied the mediating effect of serum TRPA1 on the causal pathway from olfactory function to cognitive function. Our findings warrant cautious interpretation in light of several limitations. The inherent cross-sectional design precludes establishing causal relationships, while the small sample size may have affected the detection of the mediation role of TRPA1 in the relationship between olfactory function and cognitive impairment. Future studies with larger samples are necessary to validate our findings. Moreover, as our study focused on older adults, further basic research is needed to confirm our results. Additionally, due to the absence of an established cutoff value for serum TRPA1, we utilized the statistical median to categorize participants into low or high levels of serum TRPA1, tailored specifically for this study. Despite the relative arbitrariness of this cutoff, our study still revealed a significant association.

In conclusion, our study explored the mediation role of TRPA1 in the relationship between olfactory and cognitive function among the Chinese older adults. Our findings provide preliminary evidence suggesting a mechanism linking olfactory and cognitive function, highlighting the potential significance of TRPA1 as a biomarker for cognitive impairment.

## Data availability statement

The datasets presented in this article are available from the corresponding author upon reasonable request and with permission of Huashan Hospital. Requests to access the datasets should be directed to DD, [dingding@huashan.org.cn](mailto:dingding@huashan.org.cn).

## Ethics statement

The studies involving humans were approved by Medical Ethics Committee of Huashan Hospital, Fudan University, Shanghai, China. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

## Author contributions

XL: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing. ZX: Conceptualization, Data curation, Investigation, Methodology, Writing – review & editing. JW: Investigation, Methodology, Visualization, Writing – review & editing. XM: Data curation, Methodology, Supervision, Writing – review & editing. QZ: Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Visualization, Writing – review & editing. DD: Conceptualization, Funding acquisition, Investigation,

Methodology, Project administration, Resources, Supervision, Validation, Visualization, Writing – review & editing.

Wenfan Zhu, Zhi Zhou, Xiaoying Liu, Fuqin Gao, Peng Gong, Lin Lu, Meng Wang, Ting Zhang, Yaru Guo, Xiaoli Jin, Shiqi Li, Qiongyi Xu, and Yiping Wang for their efforts to the study, and all the participants for their cooperation.

## Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. This work was supported by the National Natural Science Foundation of China (82173599, 82071200), Shanghai Sailing Program (20YF1404000), Shanghai Municipal Science and Technology Major Project (2018SHZDZX01), and Project of the Ministry of Science and Technology of China (STI2030-Major Projects 2021ZD0200800).

## Acknowledgments

The authors thank Fang Pei, Yan Zhang, Zhaolan Ding, Meihua Jin, Meirong Chen, Zeya Wang, Meizheng Shi, Jingping Ye, Meiping He, Lanfang Yu, Deping Chen, Fusheng Gong, Meili Shi, Wenying Zhou, Shumin Chen, Xiudi Xu, Meiling Huang, Linghua Ding,

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Achterberg, W. P., Gambassi, G., Finne-Soveri, H., Liperoti, R., Noro, A., Frijters, D., et al. (2010). Pain in European long-term care facilities: cross-national study in Finland, Italy and the Netherlands. *Pain* 148, 70–74. doi: 10.1016/j.pain.2009.10.008
- American Psychiatric Association (1994). *DSM-IV: Diagnostic and statistical manual of mental disorders*. Washington, DC: American Psychiatric Press Inc.
- Attems, J., Walker, L., and Jellinger, K. A. (2014). Olfactory bulb involvement in neurodegenerative diseases. *Acta Neuropathol.* 127, 459–475. doi: 10.1007/s00401-014-1261-7
- Bell, J. T., Loomis, A. K., Butcher, L. M., Gao, F., Zhang, B., Hyde, C. L., et al. (2014). Differential methylation of the TRPA1 promoter in pain sensitivity. *Nat. Commun.* 5:2978. doi: 10.1038/ncomms3978
- Borbely, E., Payrits, M., Hunyady, A., Mezo, G., and Pinter, E. (2019). Important regulatory function of transient receptor potential ankyrin 1 receptors in age-related learning and memory alterations of mice. *Geroscience* 41, 643–654. doi: 10.1007/s11357-019-00083-1
- Bosson, A., Paumier, A., Boisseau, S., Jacquier-Sarlin, M., Buisson, A., and Albrieux, M. (2017). TRPA1 channels promote astrocytic ca(2+) hyperactivity and synaptic dysfunction mediated by oligomeric forms of amyloid-beta peptide. *Mol. Neurodegener.* 12:53. doi: 10.1186/s13024-017-0194-8
- Cheng, Y. T., Woo, J., Luna-Figueroa, E., Maleki, E., Harmanci, A. S., and Deneen, B. (2023). Social deprivation induces astrocytic TRPA1-GABA suppression of hippocampal circuits. *Neuron* 111, 1301–1315.e5. doi: 10.1016/j.neuron.2023.01.015
- Choi, J. S., Hur, K., Chow, M., Shen, J., and Wrobel, B. (2018). Olfactory dysfunction and cognition among older adults in the United States. *Int. Forum Allerg. Rhinol.* 8, 648–654. doi: 10.1002/alr.22078
- Devanand, D. P., Lee, S., Manly, J., Andrews, H., Schupf, N., Doty, R. L., et al. (2015). Olfactory deficits predict cognitive decline and Alzheimer dementia in an urban community. *Neurology* 84, 182–189. doi: 10.1212/WNL.0000000000001132
- Ding, D., Zhao, Q., Guo, Q., Meng, H., Wang, B., Luo, J., et al. (2015). Prevalence of mild cognitive impairment in an urban community in China: a cross-sectional analysis of the Shanghai aging study. *Alzheimers Dement.* 11, 300–309.e2. doi: 10.1016/j.jalz.2013.11.002
- Ding, D., Zhao, Q., Guo, Q., Meng, H., Wang, B., Yu, P., et al. (2014). The Shanghai aging study: study design, baseline characteristics, and prevalence of dementia. *Neuroepidemiology* 43, 114–122. doi: 10.1159/000366163
- Dong, Y., Li, Y., Liu, K., Han, X., Liu, R., Ren, Y., et al. (2023). Anosmia, mild cognitive impairment, and biomarkers of brain aging in older adults. *Alzheimers Dement.* 19, 589–601. doi: 10.1002/alz.12777
- Horgas, A. L., and Tsai, P. F. (1998). Analgesic drug prescription and use in cognitively impaired nursing home residents. *Nurs. Res.* 47, 235–242. doi: 10.1097/00006199-199807000-00009
- Hu, Z., Zhang, Y., Yu, W., Li, J., Yao, J., Zhang, J., et al. (2023). Transient receptor potential ankyrin 1 (TRPA1) modulators: recent update and future perspective. *Eur. J. Med. Chem.* 257:115392. doi: 10.1016/j.ejmech.2023.115392
- Hummel, T., Konnerth, C. G., Rosenheim, K., and Kobal, G. (2001). Screening of olfactory function with a four-minute odor identification test: reliability, normative data, and investigations in patients with olfactory loss. *Ann. Otol. Rhinol. Laryngol.* 110, 976–981. doi: 10.1177/000348940111001015
- Jensen-Dahm, C., Vogel, A., Waldorff, F. B., and Waldemar, G. (2012). Discrepancy between self- and proxy-rated pain in Alzheimer's disease: results from the Danish Alzheimer intervention study. *J. Am. Geriatr. Soc.* 60, 1274–1278. doi: 10.1111/j.1532-5415.2012.04036.x
- Lee, K. I., Lee, H. T., Lin, H. C., Tsai, H. J., Tsai, F. C., Shyue, S. K., et al. (2016). Role of transient receptor potential ankyrin 1 channels in Alzheimer's disease. *J. Neuroinflammation* 13:92. doi: 10.1186/s12974-016-0557-z
- Lee, K. I., Lin, H. C., Lee, H. T., Tsai, F. C., and Lee, T. S. (2017). Loss of transient receptor potential Ankyrin 1 channel deregulates emotion, learning and memory, cognition, and social behavior in mice. *Mol. Neurobiol.* 54, 3606–3617. doi: 10.1007/s12035-016-9908-0
- Leung, A. Y. (1980). *Encyclopedia of common natural ingredients used in food, drugs and cosmetics*. New York: John Wiley & Sons.
- Liang, X., Ding, D., Zhao, Q., Guo, Q., Luo, J., Hong, Z., et al. (2016). Association between olfactory identification and cognitive function in community-dwelling elderly: the Shanghai aging study. *BMC Neurol.* 16:199. doi: 10.1186/s12883-016-0725-x
- Liang, X., Ding, D., Zhao, Q., Wu, W., Xiao, Z., Luo, J., et al. (2020). Inability to smell peppermint is related to cognitive decline: a prospective community-based study. *Neuroepidemiology* 54, 258–264. doi: 10.1159/000505485
- Mantyselka, P., Hartikainen, S., Louhivuori-Laako, K., and Sulkava, R. (2004). Effects of dementia on perceived daily pain in home-dwelling elderly people: a population-based study. *Age Ageing* 33, 496–499. doi: 10.1093/ageing/afh165
- McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., and Stadlan, E. M. (1984). Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA work group under the auspices of Department of Health and Human Services Task Force on Alzheimer's disease. *Neurology* 34, 939–944. doi: 10.1212/wnl.34.7.939
- Morrison, R. S., and Siu, A. L. (2000). A comparison of pain and its treatment in advanced dementia and cognitively intact patients with hip fracture. *J. Pain Symptom Manag.* 19, 240–248. doi: 10.1016/s0885-3924(00)00113-5
- Moss, M., Hewitt, S., Moss, L., and Wesnes, K. (2009). Modulation of cognitive performance and mood by aromas of peppermint and ylang-ylang. *Int. J. Neurosci.* 118, 59–77. doi: 10.1080/00207450601042094
- Nasreddine, Z. S., Phillips, N. A., Bedirian, V., Charbonneau, S., Whitehead, V., Collin, I., et al. (2005). The Montreal cognitive assessment, MoCA: a brief screening tool for mild cognitive impairment. *J. Am. Geriatr. Soc.* 53, 695–699. doi: 10.1111/j.1532-5415.2005.53221.x



- Ozsimsek, A., and Ovey, I. S. (2022). Potential effects of melatonin on TRPA1 channels in the prevention and treatment of Alzheimer's disease. *Noro Psikiyatr. Ars.* 59, 188–192. doi: 10.29399/npa.28183
- Paumier, A., Boisseau, S., Jacquier-Sarlin, M., Pernet-Gallay, K., Buisson, A., and Albrieux, M. (2022). Astrocyte-neuron interplay is critical for Alzheimer's disease pathogenesis and is rescued by TRPA1 channel blockade. *Brain* 145, 388–405. doi: 10.1093/brain/awab281
- Petersen, R. C. (2004). Mild cognitive impairment as a diagnostic entity. *J. Intern. Med.* 256, 183–194. doi: 10.1111/j.1365-2796.2004.01388.x
- Roberts, R. O., Christianson, T. J. H., Kremers, W. K., Mielke, M. M., Machulda, M. M., Vassilaki, M., et al. (2016). Association between olfactory dysfunction and amnesic mild cognitive impairment and Alzheimer disease dementia. *JAMA Neurol.* 73, 93–101. doi: 10.1001/jamaneurol.2015.2952
- Scherder, E., Bouma, A., Borkent, M., and Rahman, O. (1999). Alzheimer patients report less pain intensity and pain affect than non-demented elderly. *Psychiatry* 62, 265–272. doi: 10.1080/00332747.1999.11024871
- Stanciu, I., Larsson, M., Nordin, S., Adolfsson, R., Nilsson, L.-G., and Olofsson, J. K. (2014). Olfactory impairment and subjective olfactory complaints independently predict conversion to dementia: a longitudinal, population-based study. *J. Int. Neuropsychol. Soc.* 20, 209–217. doi: 10.1017/s1355617713001409
- Sukenaga, N., Ikeda-Miyagawa, Y., Tanada, D., Tunetoh, T., Nakano, S., et al. (2016). Correlation between DNA methylation of TRPA1 and chronic pain states in human whole blood cells. *Pain Med.* 17, 1906–1910. doi: 10.1093/pm/pnv088
- Thomson Healthcare (2007). *PDR for herbal medicines*. 4th Edn: Physician's Desk Reference. Montvale, NJ: Medical Economics Company, Inc, 580.
- Warm, J. S., Dember, W. N., and Parasuraman, R. (1991). Effects of olfactory stimulation on performance and stress in a visual sustained attention task. *J. Soc. Cosmet. Chem.* 42, 199–210.
- Wilson, R. S., Arnold, S. E., Tang, Y., and Bennett, D. A. (2006). Odor identification and decline in different cognitive domains in old age. *Neuroepidemiology* 26, 61–67. doi: 10.1159/000090250
- Xiao, Z., Wu, X., Wu, W., Yi, J., Liang, X., Ding, S., et al. (2021). Plasma biomarker profiles and the correlation with cognitive function across the clinical spectrum of Alzheimer's disease. *Alzheimers Res. Ther.* 13:123. doi: 10.1186/s13195-021-00864-x
- Yaffe, K., Freimer, D., Chen, H., Asao, K., Rosso, A., Rubin, S., et al. (2017). Olfaction and risk of dementia in a biracial cohort of older adults. *Neurology* 88, 456–462. doi: 10.1212/wnl.0000000000003558
- Zhang, M. Y., Katzman, R., Salmon, D., Jin, H., Cai, G. J., Wang, Z. Y., et al. (1990). The prevalence of dementia and Alzheimer's disease in Shanghai, China: impact of age, gender, and education. *Ann. Neurol.* 27, 428–437. doi: 10.1002/ana.410270412