



An Insight Into Ameliorating Production, Catalytic Efficiency, Thermostability and Starch Saccharification of Acid-Stable α -Amylases From Acidophiles

Deepak Parashar¹ and Tulasi Satyanarayana^{2*}

¹ Functional Genomic Unit, CSIR-Institute of Genomics and Integrative Biology, New Delhi, India, ² Division of Biological Sciences and Engineering, Netaji Subhas Institute of Technology, New Delhi, India

OPEN ACCESS

Edited by:

Caixia Wan,
University of Missouri, United States

Reviewed by:

Fu-Li Li,
Qingdao Institute of Bioenergy and
Bioprocess Technology (CAS), China
Roland Wohlgenuth,
Lodz University of Technology, Poland

*Correspondence:

Tulasi Satyanarayana
tsnarayana@gmail.com

Specialty section:

This article was submitted to
Process and Industrial Biotechnology,
a section of the journal
Frontiers in Bioengineering and
Biotechnology

Received: 30 December 2017

Accepted: 20 August 2018

Published: 28 September 2018

Citation:

Parashar D and Satyanarayana T
(2018) An Insight Into Ameliorating
Production, Catalytic Efficiency,
Thermostability and Starch
Saccharification of Acid-Stable
 α -Amylases From Acidophiles.
Front. Bioeng. Biotechnol. 6:125.
doi: 10.3389/fbioe.2018.00125

Most of the extracellular enzymes of acidophilic bacteria and archaea are stable at acidic pH with a relatively high thermostability. There is, however, a dearth of information on their acid stability. Although several theories have been postulated, the adaptation of acidophilic proteins to low pH has not been explained convincingly. This review highlights recent developments in understanding the structure and biochemical characteristics, and production of acid-stable and calcium-independent α -amylases by acidophilic bacteria with special reference to that of *Bacillus acidicola*.

Keywords: acidophiles, *Bacillus acidicola*, acid-stable α -amylase, thermostability, starch saccharification

INTRODUCTION

Enzyme characteristics such as thermostability, selectivity, solvent tolerance and substrate affinity can be improved through genetic engineering based on the availability of large data on improving these characteristics (Verma and Satyanarayana, 2012; Widersten, 2014; Joshi and Satyanarayana, 2015). Alteration in pH stability is tedious and lacks rational approaches. However, a few reports on enhancing acid stability of α -amylases through protein engineering are available (Liu et al., 2008a, 2012; Yang et al., 2013).

The majority of the enzymes used at commercial scale lack adequate acid stability, thus limiting their applications. For using such enzymes, adjusting pH to their optima is required, which makes the process tedious, expensive and time consuming. In order to overcome the problems, many industries use acid-stable enzymes from fungal sources. Since these lack adequate thermostability, the enzymes get denatured when processes are carried out at elevated process temperatures (Demirjian et al., 2001; Elleuche et al., 2014). In order to overcome these problems, microbes that are capable of tolerating harsh conditions could be exploited for naturally tailored enzymes that are superior to their neutrophilic counterparts for utility under harsh bioprocess conditions (Hough and Danson, 1999; Eichler, 2001; Sharma et al., 2012; Raddadi et al., 2015). It has generally been observed that the enzymes from acidophilic microbes function under their optimal growth conditions (Ferrer et al., 2007), thus find several commercial applications. Furthermore, the study of these enzymes might also enable us to understand the underlying mechanisms to make them functional in extreme acidic conditions (Demirjian et al., 2001). Although several acidophilic microbes have been reported (Table 1), a very few acid-stable amylases have been studied in adequate detail (Matzke et al., 1997; Sharma et al., 2012). In this review, we have explained possible strategies for improving acid-stable α -amylase production to make the process cost-effective.

TABLE 1 | Archaeal and bacterial acidophiles.

Microorganisms	Optimum pH	Optimum temp	References
<i>Bacillus acidicola</i>	4	37	Sharma and Satyanarayana, 2010
<i>Leptospirillum ferroxidans</i>	40	1.6	Zhang et al., 2010
<i>Acidithiobacillus ferrovarans</i>	29	2.1	Hedrich and Johnson, 2013
<i>Acidiphilium organovorum</i>	37	3	Lobos et al., 1986
<i>Acidiphilium symbioticum</i>	37	3–4	Bhattacharyya et al., 1990
<i>Acidiphilium cryptum</i>	35–40	3	Harrison, 1981
MODERATELY THERMOPHILIC BACTERIA			
<i>Alicyclobacillus acidocaldarius</i>	60–65	3–4	Mavromatis et al., 2010
<i>Acidomicrobium ferroxidans</i>	48	2	Clark and Norris, 1996
<i>Sulfobacillus acidophilus</i>	45–48	2	Norris et al., 1996
<i>Sulfobacillus thermosulfidooxidans</i>	45–48	2	Golovacheva and Karavaiko, 1978
<i>Alicyclobacillus acidiphilus</i>	50	3	Matsubara et al., 2002
<i>Leptospirillum thermoferrooxidans</i>	45–50	1.6–1.9	Golovacheva et al., 1992
<i>Hydrogenobacter acidophilus</i>	65	3–4	Shima and Suzuki, 1993
<i>Alicyclobacillus acidoterrestris</i>	35–55	2–5	Orr et al., 2000
Archaea			
<i>Thermogymnomonas acidicola</i>	60	3	Itoh et al., 2007
<i>Sulfurococcus yellowstonii</i>	60–65	2–2.5	Karavaiko et al., 1994
<i>Sulfolobus metallicus</i>	65–68	1–4.5	Huber and Stetter, 1991
<i>Picrophilus torridus</i>	60	0.7	Schleper et al., 1995
<i>Thermoplasma acidophilum</i>	59	1–2	Darland et al., 1970
<i>Thermoplasma volcanium</i>	60	2	Segerer et al., 1988
<i>Picrophilus oshimae</i>	60	0.7	Schleper et al., 1995
HYPERTHERMOPHILIC ARCHAEA			
<i>Acidianus infernus</i>	90	2	Segerer et al., 1986
<i>Desulfurolobus ambivalens</i>	81	2.5	Fuchs et al., 1996
<i>Metallosphaera sedula</i>	75	1.7	Huber et al., 1989
<i>Metallosphaera prunae</i>	75	1–4	Fuchs et al., 1996
<i>Sulfolobus acidocaldarius</i>	80	2–3.5	Brock et al., 1972
<i>Sulfolobus shibatae</i>	80	3	Grogan et al., 1990
<i>Sulfolobus yangmingensis</i>	80	4	Ren-Long et al., 1999
<i>Sulfolobus tengchongensis</i>	85	3.5	Xiang et al., 2003
<i>Sulfurisphaera ohwakuensis</i>	84	2	Kurosawa et al., 1998
<i>Acidilobus aceticus</i>	85	3.8	Prokofeva et al., 2000

NEED FOR ACID-STABLE AMYLASES

Starch is a ubiquitous reserve polysaccharide in plants and one of the most abundant energy sources. The starch hydrolyzing enzymes such as α -amylases, glucoamylases, α -glucosidases, and pullulanases have become increasingly attractive for starch industries because of increasing demand for sugar syrups (Sharma and Satyanarayana, 2013a; Elleuche et al., 2014). Moreover, starch hydrolyzing enzymes have attracted attention in ethanol production and account for 25 % of the global enzyme market today (Sharma et al., 2016).


The conventional industrial conversion of starch to glucose consists of a three-step industrial process: in the first step, 25–30% starch slurry is gelatinized in a jet cooker at 100–105°C

for 5–10 min, in the second step α -amylase and Ca^{2+} (50 ppm) are added with pH adjusted to 6.5 for liquefaction, and in the last step, glucoamylase addition leads to the formation of glucose (**Figure 1**) (Crabb and Mitchinson, 1997; Mehta and Satyanarayana, 2013). A few bottlenecks are associated with the process: firstly, gelatinization at higher temperature (100–105°C) requires high energy input. Secondly, native pH of starch slurry is around 3–4.5, thus, a pH adjustment step is required because α -amylases that are commercially available function best at pH 6–6.5. Thirdly, most of the α -amylases are Ca^{2+} -dependent, and thus, Ca^{2+} is added during the process, which must be removed in the subsequent stages because glucose isomerase used in fructose syrup production is inhibited in the presence of Ca^{2+} . In order to make the process economical and time saving, there has been an emphasis on discovering Ca^{2+} independent, acid-stable and raw starch degrading thermostable α -amylases, which can hydrolyze raw starch at sub-gelatinization temperatures bypassing the energy intensive gelatinization, and avoiding Ca^{2+} addition and pH adjustment steps (Sharma and Satyanarayana, 2010; Mehta and Satyanarayana, 2016; Sharma et al., 2016).

ACIDOPHILES AND THEIR BIOLOGY

Among extreme environments, acidic environments are especially interesting because the low pH of the habitat is a consequence of microbial metabolic activities (e.g., production of acid, deamination of amino acids etc.), and not a condition imposed by the system as in other extreme environments like temperature, radiation and pressure. For example, the extreme conditions of the Tinto River in Southwestern Spain are due to the products of metabolic activity of chemolithotrophic microorganisms residing in its water, but not due to the intensive mining activity carried out in the area as believed earlier (Johnson, 1998; González-Toril et al., 2003). Acidophiles are classified as organisms which can withstand and even thrive in acidic environments having pH values in the range of 1.0 to 5.0. Acidophiles are found in eukaryotes (fungi) as well as prokaryotes (bacteria and archaea) which thrive in a variety of acidic environments, including sulphuric pools and geysers, areas polluted by acid mine drainage, and even our own guts (Baker-Austin and Dopson, 2007; Sharma et al., 2012). Based on the optimum temperature required for growth, acidophiles have been subdivided further into various groups: mesophilic acidophiles (*Acidithiobacillus*, *Ferroplasma*, *Leptospirillum*), moderate thermoacidophiles (*Picrophilus torridus*) and hyper thermoacidophiles (*Acidianus infernus*) (**Table 1**). It has, however, been observed that the most thermotolerant microbes are not the most acid tolerant and vice versa (Auernik et al., 2008). The most thermophilic extreme thermoacidophilic archaeon, *Acidianus infernus* that grows at 65–95°C (T_{opt} 90°C), grows in the pH range between 5.5 and 1.0, with the optimum around 2.0 (Segerer et al., 1986). The members of archaeal Picrophilaceae are the most acidophilic organisms known and are able to grow at pH 0.7 and 60°C (Schleper et al., 1995).

Despite being able to survive in extreme acidic conditions, intracellular pH of acidophiles is similar to that of neutrophiles



because macromolecules such as DNA become unstable at acidic pH. Their pH gradients [$\text{pH gradient (pH)} = \text{pH}_{\text{in}} - \text{pH}_{\text{out}}$], however, remain several orders of magnitude greater than neutrophiles. In order to survive, acidophiles have evolved multiple mechanisms such as highly impermeable cell membranes, small genomes for ease in maintenance, genes for organic acid degradation, DNA and protein repair systems, and a predominance of secondary transporters to remove protons once they have entered cytoplasm (Figure 2) (Johnson, 1998; Baker-Austin and Dopson, 2007; Sharma et al., 2016). Acidophiles are the most widely distributed in the bacterial and archaeal domains (Table 1) and have numerous biotechnological applications (Sharma et al., 2012; Elleuche et al., 2014; Mehta and Satyanarayana, 2016).


α - AMYLASE PRODUCING ACIDOPHILES

Despite extensive research on acidophiles, very few have been exploited for commercial purposes. Matzke et al. (1997) reported α -amylase from acidophilic bacterium *Alicyclobacillus acidocaldarius*, which is thermostable and acidstable with a molecular mass of 140 kDa, with optimum temperature and pH of 75°C and 3.0, respectively. Bai et al. (2012) reported another α -amylase from *Alicyclobacillus* sp. A4 with a molecular mass of 64 kDa and optimal activity at 75°C and pH 4.2. In still another report, an α -amylase from the acidophilic bacterium *Bacillus* sp. DR90, isolated from Dig Rostam hot mineral spring (Iran), was investigated. The enzyme was active in a wide range of pH and

temperature having optimal activity at pH 4.0 and 75°C with a molecular mass of 76 kDa (Asoodeh et al., 2014). Laderman et al. (1993) reported a thermoacidophilic α -amylase from *Pyrococcus furiosus* which was optimally active at $\sim 100^\circ\text{C}$ and pH 5.5–6.0. The enzyme is a homodimer with a subunit molecular mass of 66 kDa. Recently a Ca^{2+} -independent, acid-stable α -amylase (Ba-amy) from the acidophilic bacterium *Bacillus acidicola* TSAS1 has been investigated in detail. This has been found to be a potential candidate for saccharification of starch at its native pH of 3.0–4.5 with $T_{1/2}$ of 25 min at 70°C (Sharma and Satyanarayana, 2010).

PRODUCTION OF ACID-STABLE AMYLASES

At commercial level, amylolytic enzymes are being produced from *Bacillus* spp. (*B. licheniformis*, *B. stearothermophilus* and *B. amyloliquefaciens*) and filamentous fungi such as *Aspergillus oryzae* and *A. niger* (<http://www.novozymes.com/>) (Sharma and Satyanarayana, 2013a; Sharma et al., 2016). They are the preferred sources because of their rapid growth rates, economical production and ability to secrete a large quantity of amylases, which lack either acid stability or thermostability. Amylases from acidophiles have tremendous potential to replace neutrophilic enzymes because they are active in the acidic range. Production on a commercial scale is, however, a major bottleneck because of low titres of extracellular amylases secreted by acidophilic and acid tolerant microbes (Table 4). Schwermann et al. (1994) recorded maximum amylase production (90 U mL⁻¹) in the



presence of maltose as compared to other carbon sources used in the production medium. Bai et al. (2012) reported a 2.3 U mL^{-1} α -amylase production in *Alicyclobacillus* sp. A4 after 48 h in the presence of starch as carbon source. A few researchers have attempted multiple strategies to increase the extracellular acid-stable amylase titres (Table 4). Kanno (1986) improved *A. acidocaldarius* A2 strain by using UV/enrichment method for α -amylase production. A total of 11-fold increase in the enzyme titre was achieved as compared to the wild type strain (220 U mL^{-1}). Brown et al. (1990) attained a low constitutive level of α -amylase from *Pyrococcus furiosus* in the presence of simple sugars, while polysaccharides with α -1,4 linkages stimulated production. The maximum yield of extracellular α -amylase was achieved (100 U mL^{-1}) in *P. furiosus* in the presence of pullulan as a carbon source in submerged fermentation at 98°C .

A detailed investigation was carried out on the production of α -amylase by *B. acidicola* in submerged fermentation. Conventional “one-variable-at-a time” and statistical approaches have been used for optimizing the cultural parameters (Sharma and Satyanarayana, 2011). α -Amylase production by *B. acidicola* was high in the presence of soluble starch (2%) as a carbon source. Among nitrogen sources tested, tryptone (HIMEDIA) [0.5%] in combination with yeast extract (0.5%) supported a high enzyme titre. The α -amylase produced by *B. acidicola* displayed a high activity (8 U/mL) at pH 4.0 and 37°C after 44 h. The enzyme production was high when the cells are in stationary phase as in *Geobacillus thermoleovorans* (Uma Maheswar Rao and Satyanarayana, 2003). Further statistical approaches [Plackett-Burman design and Response Surface Methodology (RSM)] were

employed for optimizing α -amylase production by *B. acidicola* in submerged fermentation. When the effect of 11 variables on α -amylase production was assessed using Plackett-Burman design, four variables (starch, K_2HPO_4 , inoculum size, and temperature) were identified to significantly affect enzyme production. In order to control process parameters like aeration, uniform distribution of nutrients, and heat and oxygen transfer for α -amylase production by *B. acidicola*; fermentation was carried out in a 7 L laboratory fermentor. A reduction in fermentation time for attaining the peak was recorded; this could be due to improvement in mixing of nutrients and the control of dissolved oxygen (Kumar and Satyanarayana, 2007). Further, a 2.9-fold enhancement in enzyme production was attained due to fed-batch fermentation as compared to that in the initial unoptimized medium (3.5 U mL^{-1}) (Sharma and Satyanarayana, 2011).

Bacillus acidicola is amenable to solid state cultivation like some other *Bacillus* spp. (Babu and Satyanarayana, 1995; Chen et al., 2005). Therefore, production of α -amylase by solid state fermentation was also attempted in order to find the prospects of using a wide range of agro-industrial residues as substrates (Babu and Satyanarayana, 1995; Sharma and Satyanarayana, 2012b). A peak in enzyme production was reached in 72 h, when 10 g of wheat bran was used in 250 mL Erlenmeyer flasks as reported for the production of α -amylase by *Bacillus coagulans* (Babu and Satyanarayana, 1995). In contrast, 5 g wheat bran per 250 mL flask was used for α -amylase production by *B. amyloliquefaciens* (Gangadharan et al., 2006). *Bacillus acidicola* secreted high enzyme titre at a_w (water activity) 0.95; below this, α -amylase production declined, with no growth and

enzyme production below a_w 0.85, indicating that the bacterium is desiccation sensitive. Supplementation of wheat bran with ammonium sulfate supported a high enzyme titre. The addition of nitrogen sources to the solid substrates had been shown to enhance the production of various enzymes including α -amylase in solid state fermentation (SSF) (Pedersen and Nielsen, 2000). Moisture (substrate:water, 1:3.5), starch (2.9%) and ammonium chloride (0.38%) supported a high enzyme titre in *B. acidicola*. Statistical optimization of α -amylase production in SSF using response surface methodology led to 5.6-fold [28 ± 2.3 U/g dry bacterial bran (DBB)] increase in the titre as compared to unoptimized conditions (5 ± 1.1 U/g DBB) (Sharma and Satyanarayana, 2012b).

CLONING AND EXPRESSION OF ACID-STABLE α -AMYLASE ENCODING GENES

Despite extensive efforts, attaining higher enzyme titres by wild type microbial strains is a major hurdle for their commercialization. By cloning and expressing acid-stable amylase encoding genes from acidophiles in mesophilic hosts such as *E. coli*, *Pichia pastoris* or *Bacillus subtilis*, the production can be made cost-effective (Table 4). However, improper folding and differences in codon usage often hinder over production of enzymes in heterologous expression systems (Sharma et al., 2012). In order to overcome these bottlenecks, researchers use codon optimized synthetic genes or different host variants (Elleuche et al., 2014; Ranjan and Satyanarayana, 2016). Acid-stable α -amylase gene from *Bacillus* sp. DR90 was successfully cloned in *E. coli* BL21 and expressed as an intracellular active protein. After induction, specific activity was around 600 U/mg (Asoodeh et al., 2010). Matzke et al. (1997) cloned acid-stable α -amylase gene from *Alicyclobacillus acidocaldarius* in *E. coli*. The enzyme was intracellular and the optimum temperature for recombinant acid-stable α -amylase was slightly lower than that of the native enzyme. Extracellular acid-stable α -amylase encoding gene of *P. furiosus* (PFA) was cloned and expressed in *E. coli* (Dong et al., 1997). The recombinant acid-stable α -amylase was mainly expressed in the form of insoluble inclusion bodies. An improved purification method was developed by Wang et al. (2007). The solubilization of the inclusion bodies was achieved by treatment at 90°C for 3 min in Britton–Robinson buffer at pH 10.5. After solubilization, a total of 58,000 U/g wet cells yield was obtained. In another study, Peng et al. (2016) co-expressed PFA with chaperones in *E. coli*. Both chaperonin and a small heat shock protein (sHSP) increased the solubility of PFA to a certain degree, while pre-folding seemed to be the most efficient that increased the enzyme activity to about 60,000 U g⁻¹ wet weight over that of 5,000 U g⁻¹ wet weight without chaperone. Wang et al. (2016) produced soluble PFA by expressing PFA in *B. amyloliquefaciens*. The yield of PFA was 2,000 U mL⁻¹ of supernatant and 2,714 U mL⁻¹ of total culture. Zhu et al. (2017) expressed PFA in *Nicotiana tabacum* and found that plant produced PFA forms functional aggregates with an accumulation level up to 3.4 g kg⁻¹

fresh weight. The aggregates were functional without requiring refolding. As stated above, several attempts have been made to increase acid-stable α -amylase titres. Intracellular accumulation and inclusion body formation make them impractical for industrial applications (Grzybowska et al., 2004; Wang et al., 2007).

A truncated 1,441 bp acid-stable α -amylase gene encoding 479 amino acid α -amylase (Ba-amy) of *B. acidicola* was successfully cloned and expressed in active form in *E. coli*. Various approaches have been developed for efficient secretion of proteins such as increasing the permeability of the outer membrane chemically (adding EDTA, glycine, and Triton X-100) and by enzymatic (lysozyme) treatments (Sharma and Satyanarayana, 2012a; Parashar and Satyanarayana, 2016a). The purified recombinant α -amylase was active at pH 4.0 and 60°C, and retained all characteristics like that of the native α -amylase.

The methylotrophic yeast *Pichia pastoris* has emerged as an important production host for extracellular production of proteins for both basic research and industrial applications (Cregg et al., 2009; Spohner et al., 2015). Codon usage analysis of acid-stable *Ba-amy* revealed the feasibility of its expression in *Pichia pastoris*. In order to increase extracellular production of acid-stable α -amylase, *Ba-amy* was cloned and expressed in *P. pastoris* under dual promoters (*GAP* and *AOX*) and fused with α -factor secretion signal peptide. Mixed fed batch and high cell density cultivation experiments were performed which led to 15- and 7- fold higher extracellular enzyme titres than that of the wild type *B. acidicola* and recombinant *E. coli*, respectively (Parashar and Satyanarayana, 2016b). The recombinant acid-stable *Ba-amy* purified from *P. pastoris* was biochemically characterized, which revealed kinetic properties and thermostability of glycosylated acid-stable *Ba-amy* to be similar to those of the recombinant acid-stable *Ba-amy* expressed in *E. coli*. The engineered *Ba-amy* (*Ba-Gt-amy*) was also cloned and expressed in *P. pastoris* (Parashar and Satyanarayana, 2017a). The combination of multiple transformations and post-transformational vector amplification (PTVA) and high cell density cultivation in fermentor led to a very high production (750 U/mL) of the chimeric *Ba-Gt-amy* (Parashar and Satyanarayana, 2017a).

STRUCTURAL CHARACTERISTICS OF ACID-STABLE α -AMYLASE

Although the adaptation of acidstable enzymes to low pH has not been explored in greater detail, one explanation for pH stability has been offered from the modeling of α -amylases from *B. acidicola* and other acid-stable amylases. A detailed investigation revealed that the acid stability and activity at acidic pH could be attributed to the surface charge density and amino acid composition of these proteins (Table 2). A prominent feature of acidstable α -amylases is the excess of glutamic and aspartic acid (D + E) residues on their surface as compared to their closest relatives. Moreover, the enzyme contains less positively charged amino acid residues (K + R + H) than their neutrophilic counterparts that leads to reduced positive charge

TABLE 2 | Amino acid composition (%) and other characteristics of acidic, basic and neutral amylases.

Characteristics	Basic amino acid (H + R + K)	Acidic amino acid (D + E)	Total amino acid	PI	Molecular weight	Source	Reference
Acidic	6.4	10.5	1276	4.36	137	<i>Alicyclobacillus acidocaldarius</i>	Matzke et al., 1997
Acidic	11.1	10.9	626	5.53	68	<i>Bacillus</i> sp. DR90	Asoodeh et al., 2014
Acidic	11.7	11.9	479	5.41	62	<i>B. acidicola</i>	Sharma and Satyanarayana, 2010
Acidic	10.4	12.9	435	4.82	50	<i>Pyrococcus furiosus</i>	Laderman et al., 1993
Basic	10.5	15.2	923	4.61	103	<i>Bacillus halodurans</i>	Murakami et al., 2007
Basic	9.3	15.5	922	4.44	102	<i>Bacillus</i> sp.	Shirokizawa et al., 1990
Neutral	15.4	12.9	483	6.05	55	<i>Bacillus licheniformis</i>	Joyet et al., 1984
Neutral	15.7	15.7	488	5.62	57	<i>Geobacillus thermoleovorans</i>	Mehta and Satyanarayana, 2014

H, Histidine; R, Arginine; K, lysine; D, aspartic acid; E, Glutamic acid; pi, isoelectric point.

density at the surface of the protein (**Figure 2**) (Reed et al., 2013). This effect was interpreted as follows: if the proteins were to possess a large content of positively charged residues (K + R), the positive charges at the surface will repel each other, leading to unfolding of the protein. On the other hand, protonation of the negatively charged group increases at lower pH that leads to reduction in the negative charge, which aids in stabilizing proteins in acidic conditions. If such proteins were to possess a large excess of negative groups, unfolding might also occur above the isoelectric point of the protein due to disruption of stabilizing structural interactions. In order to be stable and active in a broad pH range, these groups of proteins must have reduced number of D + E residues, which is compensated by an increase in the number of polar residues (Matzke et al., 1997; Schäfer et al., 2004; Reed et al., 2013). These characteristics were also found in other proteins. Huang et al. (2005) reported a high number of acidic residues on the surface of proteins, which causes repulsion due to excess negative charges resulting instability of proteins at high pH. However, a few exceptions have been found to this rule. Schäfer et al. (2004) reported thermo-acid-stable maltose-binding protein from *Alicyclobacillus acidocaldarius*. This protein has higher content of basic residues exposed on its surface, while most acidic residues are buried in the interior. As a consequence, this protein has a highly positive surface charge. This study suggested that there are multiple factors responsible for the acid-stability of proteins.

Another theory, a change in the pH activity of the amylases is depends the pKa of the catalytically important residues which are known to be influenced by the electrostatic field. It has been hypothesized that slight change in the pKa values of the catalytic residues can change the pH activity profile of the enzyme (Nielsen et al., 1999; Nielsen and Borchert, 2000). The pKa of catalytic residues in the active site can be altered by mutating selected residues that can alter the hydrogen bonding network, solvent accessibility or change in the net charge of the molecule (Nielsen and Borchert, 2000). This can be explained as follows: active site residues must be in a catalytically competent protonation state for the enzyme to be active. Thus the proton donor (Glu) is required to be protonated, while the nucleophile (Asp) must be

TABLE 3 | Distinguishable properties of acid-stable and neutral α -amylases.

Characteristics	Acidstable α -amylases	Neutral α -amylases
pH range	3.0–6.0	6.5–8.0
Temperature range (°C)	40–115	37–90
Molecular weight (kDa)	41–160	12.5–70
pi	3.4–4.8	5.0–7.1
Acid-stability	3.5–5.5	Unstable
Thermostability (°C)	60–80	Unstable
Release of CNP from CNP- α -G3	G3 Suppressed by KSCN	Stimulated by KSCN
Cleavage of G5	GGG(α) + GG	GG + GGG(α/β)
Number of subsites	5	7–9


Modified from Sharma and Satyanarayana (2013a).

negatively charged. If an α -amylase is stable over the entire pH range, it is feasible that the pH-activity profile can be changed if the pKa value of either the nucleophile or the proton donor is changed. Typically, charged residues are inserted in the vicinity of the titrable group to change the immediate environment of the active site of enzymes (Wind et al., 1998; Nielsen et al., 1999; Nielsen and Borchert, 2000). In several cases, it has been observed that pH \pm activity profiles shifted in the opposite direction as compared to the shift predicted from electrostatic calculations. This strongly suggests that electrostatic effects cannot be the best method to alter the optimum pH for enzymes. Secondary structural content does not appear to vary greatly in different amylases, which suggests that it is not a contributing factor.

Acid-stable α -amylases appear to degrade starch essentially by the same mechanism as neutrophilic α -amylases despite their distinguishing characteristics (**Table 3**). This contention is based on the deduction that the α -amylase from acid-stable and neutrophilic members conserved the same charge at the catalytic active site (**Figure 3**). Moreover, multiple amino acid alignments and site directed mutagenesis revealed that acid-stable amylases conserved same catalytic residues (Asp189, Glu320, and Asp401) like that of their neutrophilic counterparts (**Figure 4**). Hence, the acid stability of amylases has minor effect on their catalytic sites (Sharma and Satyanarayana, 2013b).

TABLE 4 | Production profile of wild and recombinant acid-stable amylases from various acidophiles.

Source	Optimum pH	Production by the wild strain U mL ⁻¹	Production in the recombinant <i>E. coli</i> (U mg ⁻¹)	Production in the recombinant <i>P. pastoris</i> U mL ⁻¹	Production in <i>Bacillus</i> spp. U mL ⁻¹	Production in plant	References
<i>Alicyclobaillus</i> sp. A4	4.2	2.3	–	–	–	–	Bai et al., 2012
<i>Alicyclobacillus acidocaldarius</i>	–	–	0.33	–	–	–	Schwermann et al., 1994
<i>Alicyclobacillus acidocaldarius</i>	–	0.1	–	–	–	–	Schwermann et al., 1994
<i>B. acidocaldarius</i>	–	2200.0	–	–	–	–	Kanno, 1986
<i>Bacillus</i> sp. DR90	4	–	600.0	–	–	–	Asoodeh et al., 2014
<i>P. furiosus</i>	5.5	–	138.0	–	–	–	Wang et al., 2007
<i>P. furiosus</i>	5.5	100.0	–	–	–	–	Brown et al., 1990
<i>P. furiosus</i>	5.5	–	–	–	2714	–	Wang et al., 2016
<i>P. furiosus</i>	5.5	–	–	–	–	34.0 g Kg ⁻¹ fresh weight	Zhu et al., 2017
<i>P. woesei</i>	5.5	1.0	–	–	–	–	Koch et al., 1991
<i>B. acidicola</i>	4.0	12.0	180.0	750.0	–	–	Sharma and Satyanarayana, 2010, 2012a; Parashar and Satyanarayana, 2017a



PROTEIN ENGINEERING OF α -AMYLASES FOR ACID STABILITY AND STARCH SACCHARIFICATION

Several methods of protein engineering are employed for improving acid stability of thermostable α -amylases to make them suitable for industrial applications, although success achieved so far is negligible. Nielsen et al. (1999) modified thermostable α -amylase from *Bacillus licheniformis* by using site directed mutagenesis, which was predicted to change the

pKa values of the catalytic residues. The observations suggested that pH \pm activity profiles of mutants which change the net charge on the molecule were significantly different from the wild-type pH \pm activity profile. The differences were, however, difficult to correlate with the electrostatic field changes calculated. In another study, two amino acids of α -amylase from *B. licheniformis* were substituted (Leu134 to Arg and Ser320 to Ala) for acid tolerance, and the mutated gene was expressed in *Bacillus subtilis* WB600. The α -amylase variants were found to be more acid tolerant than the native protein. The optimum pH and stable

AAA23810.1	---MRNPTLLQCFHWYYPGGSSWPELAERADGFNDIGINMVLPPAYKGGASGGYVSGYD	57
1UD2_A	--DGLNGTMMQYYEWHLENDGQHWNRLHDDAAALS DAGITAIWI PPAKGN SQ-ADVG YG	57
Ba-amy	----FNGTMMQYFEWYLPDDGTLWTKVANEANLSSLGITALWLPAYKGT SR-SDVG YG	55
1W9X_A	----TNGTMMQYFEWHL PNDGQHWNRLRDDASNLRRNGITAIWI PPAWKGT SQ-NDVG YG	55
1WPC_A	HNNGTNGTMMQYFEWYLPNDGNHWNRLNSDASNLKSKGITAVWI PPAWKGT SQ-NDVG YG	59
	* * : : * : : * : : * : : * : : * : : * : : * : : * : : * : : * : : * : : * : : *	
AAA23810.1	SYDLFDLGEFNDQKGS IPTKYGDKAQLLAAIDALKRNDIAVLLD VVVYHKMGVDEKEAIRV	117
1UD2_A	AYDLYDLGEFNQKGT VRTKYGTKAQLERAIGSLKSN DINVYGDVVMN HKMGADFT EAVQA	117
Ba-amy	VYDLYDLGEFNQKGT VRTKYGTKAQYLQTIQAHAAGM QVYADVVFYHKGGADGTEWVDA	115
1W9X_A	AYDLYDLGEFNQKGT VRTKYGTRS QLESALHALKNNGVQVYGDVVMN HKGGADATENVLA	115
1WPC_A	AYDLYDLGEFNQKGT VRTKYGTRS QLAAVTSLKNNGIQVYGDVVMN HKGGADATEMVRA	119
	*** : ***** : *** : : * * : : * : : * : : * : : * : : * : : * : : *	
AAA23810.1	QRVNADDR TQIDEEIECEGWTRYTFPARAGEYSQFIWDFKCFSGIDHIE NPDE-DGIFK	176
1UD2_A	VQVNPTNRWDI SGAYTIDAWTGFDFSGRNAYSDFKWRWFHFN GVDWDQRYQE-NHIFR	176
Ba-amy	VEVNPDRNQEISGTYQIQAWTKFDFPGRGNTYS SFKWRWYHFDGVDWDESRKL-SRIYK	174
1W9X_A	VEVNPNNRNQEISGDYTI EAWTKFDFPGRGNTYSDFKWRWYHFDGVDWDQSRQFNRIYK	175
1WPC_A	VEVNPNNRNQEVTEYTI EAWTRFDFPGRGNTHS SFKWRWYHFDGVDWDQSRRLNRIYK	179
	. * * : * * : : * * : : * * : : * * : : * * : : * * : : * * : : * * : : *	
AAA23810.1	IVNDYTGEGWNDQVDELGNFYLMGENIDFRNHAVTEEIKYWARWVMEQTQCDGFRILA	236
1UD2_A	F----ANTNWNWRVDEENGYDYL LGSNIDFSHPEVQDELKDWG SWFTDEL DLDG YRLA	232
Ba-amy	FR--GIGKAWDWEVDTEENGYDYL MYADLMDHPEVVTELKNWGK WYVNTNIDGFRILA	232
1W9X_A	FR--GDGKAWDWEVDS ENGYDYL MYADVMDHPEVVNELLRRWGEWY TNLNLDGFRILA	233
1WPC_A	FR--GHGKAWDWEVDTEENGYDYL MYADIDMDHPEVVNELLRNWGVY TNLGLDGFRI LA	237
	: : . * : * * * * : * * : : * * : : * * : : * * : : * * : : * * : : *	
AAA23810.1	VKHI PAWFYKEWIEHVQEVAPKPLFIVAYWSHEVDK LQTYIDQVEGK TMLFDAPLQMKF	296
1UD2_A	IKHIFWYTS DWRHRQNEADQDLFVVG YWKDDVGALEFYLD EMMWEMSLFDVPLNYNF	292
Ba-amy	VKHIFKSSFPDLSYVRSQTGKPLFTVGYWYSYDINKLHNYITKTNGTMSLFDAPLHSHK F	292
1W9X_A	VKHIKYSFTRDWLTHVRNATGKEMFAVAFWKNDLGALENYLNKTNWNH SVFVDFPLHYNL	293
1WPC_A	VKHIKYSFTRDWINHVR SATGKNMFAVAFWKNDLGA IENYLQKTNWNH SVFVDFPLHYNL	297
	: * * * : * * : : * * : : * * : : * * : : * * : : * * : : * * : : *	
AAA23810.1	HEASRMGRDYDMTQIF TGTLVEADPFHAVTLVANHITQPLQALEAPVEPWFKPLAYALIL	356
1UD2_A	YRASQQGGSYDMRNILRGLVEAHPMHA VTFVDNHITQPGESLESWVADWFKPLAYATIL	352
Ba-amy	YTASKSGGAFDMRTLMNTLMK DQPTLAVTFVDNHIT E P VQALQSWVD PWFKPLAYAFIL	352
1W9X_A	YNASNSGGNYDMAKLLNGTVVQKH PMHA VTFVDNHIT SQPGESLESFVQEWFKPLAYALIL	353
1WPC_A	YNASKSGGNYDMRNIFNGTVVQRHPSH AVTFVDNHIT SQPEEALESFVEWFKPLAYALTL	357
	: * * . * : * * : : : : : * * * * * : * : : * * : : * * : : * * : : *	
AAA23810.1	LRENGVPVSFYPDLYGAHYEDVGGDGTYPIDMPI IEQLDELILARQRF AHGVQTLFFFDH	416
1UD2_A	TREGGYPNV FYGDY YGIPNDNIS A-----K KDMI DELLDARQNYAYGTQH DYFDH	402
Ba-amy	TRQEGYPCV FYGDY YGIPQYNI PS-----L KSKIDPLLIARDYAYGTQH DYLDH	402
1W9X_A	TREQGYPSV FYGDY YGIPTHSVPA-----M KAKIDPILEARQNFAYGTQH DYFDH	403
1WPC_A	TREQGYPSV FYGDY YGIPTHGVPA-----M RSKIDPILEARQKYAYGKQNDYLDH	407
	* : * * * * * * * : : . : : : : * : : * * : : * * : : * * : : * * : : *	
AAA23810.1	PNCIAFSRSGTDEFPG--CVVMSNGDDGEKTIHLGENYGNKTWRDFLGNRQERVVTDEN	474
1UD2_A	WDVVGWTRREGSSSRPN SGLATIMSNGPGGSKWMYVGRQ NAGQWTWDLTGNGGASVTINGD	462
Ba-amy	SDIIGWTRREGVTEKPGSGLAALITD GPGGSKWMYVGKQHAGKVFYDLTGNRSDT V TINS D	462
1W9X_A	HNIIGWTRREGNTHPNSGLATIMS D GPGGSKWMYVGRN KAGQVWH DITGNKPGT V T INAD	463
1WPC_A	HNIIGWTRREGNTAHPNSGLATIMS D GAGGSKW M FVGRN KAGQVWSDITGNRTGT V T INAD	467
	: : : : * * * * . : : : : * * * * : : * * : : : : * * : : * * : : * * : : *	
AAA23810.1	GEATFFCNGGSVSVWVIEEVI	495
1UD2_A	GWGEFFTNGGSVSVYVNQ---	480
Ba-amy	GWGEFKVNGGSVSVWVVP----	479
1W9X_A	GWANFVNGGSVSIWVKR---	481
1WPC_A	GWGNFVNGGSVSIWVNK---	485
	* . * * * * * : : *	

FIGURE 4 | Multiple sequence alignment of different α -amylases. Residues responsible for calcium binding sites are highlighted in yellow color. Catalytically important residues are shown in green color. AAA23810.1- α -amylase from *Escherichia coli*; 1UD2_A: Calcium free α -amylase from *Bacillus* Sp. Strain Ksm-K38; Ba-amy: α -amylase from *B. acidicola*; 1W9X_A: *Bacillus halmapalus* α -amylase; 1WPC_A: maltotriose producing α -amylase from *Bacillus* sp. 707. [(*) the residues are identical in all sequences; (:) the conserved substitutions; (.) semi-conserved substitutions].



pH range of the mutein (mutated protein) were 4.5 and 4.0–6.5, as compared to 6.5 and 5.5–7.0 as the optimum pH and pH stability range of the native protein. It has been postulated that mutations changed the net charge on the substituted residues,

which influenced the pKa values of catalytically important amino acid residues (E261 and/or D328) (Liu et al., 2008a, 2012).

Yang et al. (2013) engineered amylase from *Bacillus subtilis* for improving protein stability and catalytic efficiency under acidic

conditions by site-directed mutagenesis. Based on the analysis of three dimensional structure model, four basic histidine (His) residues (His222, His275, His293, and His310) in the catalytic domain were replaced with acidic aspartic acid (Asp) residues.

The acid stability of the enzyme was significantly enhanced after mutation. It has been observed that the hydrogen bonds and salt bridges increased after mutation around the catalytic domain. The higher pKa of Asp was responsible for destabilizing the



protonated form of Glu250, resulting in a decrease of the pKa value of Glu250. These changes around the catalytic domain have been suggested for improvement in protein stability and catalytic efficiency at low pH. Similar efforts have also been made for improving the pH stability of amylases by protein engineering (Shaw et al., 1999; Priyadharshini et al., 2010; Liu et al., 2012).

Despite a few strategies reported in the literature for improving acid stability of proteins (Liu et al., 2008a,b, 2012; Yang et al., 2013), there is no universal strategy that aids in engineering pH-activity profiles. Therefore, further research efforts are needed to find other factors that can contribute to the acid stability of proteins. Investigations have shown that it is possible to increase the thermostability, but not so with acid-stability. Therefore, instead of increasing acid-stability of already existing thermostable enzymes, a better option is to improve the thermostability of acid-stable enzymes. Since α -amylase from *B. acidicola* (Ba-amylase) is stable in acidic conditions with moderate thermostability, an attempt has thus been made to improve thermostability (Parashar and Satyanarayana, 2016c). Several chimeras were constructed with the addition of amino acids at N- and C-terminal ends of acid-stable Ba-amylase from the α -amylase of *Geobacillus thermoleovorans* (Gt-amylase) (Figure 5). All chimeras were successfully expressed in active form in *E. coli*. Among all chimeras, one chimera displayed higher thermostability and specific activity as well as catalytic efficiency without change in its acid stability and pH optimum for activity (Parashar and Satyanarayana, 2016a). Increase in starch binding capacity of the chimeric α -amylase was observed in comparison with that of the wild type. The adsorption of chimeric α -amylase to raw starch suggests that the hydrolysis of raw starch can occur at high temperatures without energy-intensive gelatinization step, which brings down the energy consumption for starch saccharification. Furthermore, the end products of raw starch hydrolysis by the chimera suggested that the addition of residues did not alter the catalytic activity.

In another study, this chimeric α -amylase was fused with acid-stable glucoamylase of *Aspergillus niger* through a linker peptide for saccharifying starch in a single step (Figure 5) (Parashar and Satyanarayana, 2017b). The kinetic properties of the fused enzyme supported its suitability in raw starch saccharification in acidic conditions of native starch that liberates glucose besides maltodextrins as the major starch hydrolysis products. The fused chimeric enzyme can, therefore, be a practical option for the cost effective saccharification of raw starch. Engineering multidomain enzymes that are capable of catalyzing two or more reactions is a potential strategy to reduce enzyme costs in industrial processes because multiple catalytic properties in a single polypeptide simplify production and purification process (Fan et al., 2009; Ribeiro et al., 2011; Parashar and Satyanarayana, 2017b).

CALCIUM BINDING REGION IN α -AMYLASE

Calcium is known to stimulate α -amylases and has also been implicated in enhancing their thermostability (Savchenko

et al., 2002). Acid-stable Ba-amylase was found to be a calcium-independent. A few Ca^{2+} -independent α -amylases have also been reported earlier (Babu and Satyanarayana, 1993; Sajedi et al., 2005; Asoodeh et al., 2010), which were considered to be useful in industrial starch saccharification.

In general, most of the α -amylases possess conserved calcium ion binding sites, which are positioned at the interface between domains A and B, and play a major role in its stability and activity (Figure 6) (Boel et al., 1990; Linden et al., 2003). *Bacillus* α -amylases have been reported to have three Ca^{2+} ions and one Na^{+} ion and a metal triad bridge (calcium-sodium-calcium) (Linden et al., 2003). This metal triad is important for maintaining the compact protein structure and provides thermal stability to the enzyme (Linden et al., 2003). Calcium ion helps in salting out of hydrophobic residues in the protein, resulting in the formation of a compact structure that enhances stability (Linden et al., 2003). Whenever calcium ions are detached, amylases lose their stability, while its restoration recovers the stability (Boel et al., 1990). Sequence analysis revealed that acid-stable Ba-amylase comprises three calcium binding sites despite the fact that Ba-amylase does not show any calcium-dependent activity. Since the calcium binding site is far away from the active site residues, it has been postulated that the role of calcium ions is in maintaining the structure rather than in catalysis. A second theory suggests that calcium is loosely bound, which is replaced by other metal ions such as Na^{+} . These theories get support from the investigations on other calcium-independent enzymes (Nonaka et al., 2003).

COMMERCIAL APPLICATION OF ACIDSTABLE AMYLASES

Extremophiles are potent sources of extremozymes, which display a high stability under extreme bioprocess conditions (Elleuche et al., 2014). Thus biocatalysts from extremophiles have been shown to be useful in industrial bioprocesses. Only a few extremozymes, however, found their way to the market (e.g., thermostable DNA polymerases from *Thermus aquaticus* and *Pyrococcus furiosus* and others). There is a




FIGURE 7 | Applications of acid-stable α -amylases.

tremendous potential for acidstable enzymes from acidophiles to revolutionize existing industrial processes and to make many novel applications possible (Figure 7) (Mehta and Satyanarayana, 2016; Sharma et al., 2016). Moreover, acidstable α -amylases reduce the cost and time required for multistep maltooligosaccharide production from raw starches, which can be used as antistaling agents in baking industry (Parashar and Satyanarayana, 2016c). Other acid-stable enzymes have potential biotechnological and industrial applications (Figure 7).

CONCLUSIONS

α -Amylases produced by acidophilic microorganisms find applications in industrial processes such as starch saccharification and hydrolysis of polysaccharides in plant biomass in bioethanol production. The analysis of structure of these proteins suggests that acid-stable biocatalysts differ in surface charge, amino acid composition, salt bridges, and hydrophobicity. There

are very few successful attempts in improving acid-stability of enzymes through protein engineering. These studies lack adequate rational concept, thus, difficult to apply for other proteins. Extensive as well as intensive efforts are, therefore, called for attaining high titres of acid-stable α -amylases and to understand the mechanisms which make them functional at low pH for ameliorating the existing enzymes for novel industrial applications.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENTS

One of us (TS) is grateful to the University Grants Commission, Govt. of India, New Delhi for granting BSR Faculty Fellowship while writing this review.

REFERENCES

- Asodeh, A., Chamani, J., and Lagzian, M. (2010). A novel thermostable, acidophilic α -amylase from a new thermophilic "Bacillus sp. Ferdowsicus" isolated from Ferdows hot mineral spring in Iran: purification and biochemical characterization. *Int. J. Biol. Macromol.* 46, 289–297. doi: 10.1016/j.ijbiomac.2010.01.013
- Asodeh, A., Emtenani, S., Emtenani, S., Jalal, R., and Housaindokht, M. R. (2014). Molecular cloning and biochemical characterization of a thermoacidophilic, organic-solvent tolerant α -amylase from a Bacillus strain in Escherichia coli. *J. Molecular Cat. B: Enzymatic.* 99, 114–120. doi: 10.1016/j.molcatb.2013.10.025
- Auernik, K. S., Cooper, C. R., and Kelly, R. M. (2008). Life in hot acid: pathway analyses in extremely thermoacidophilic archaea. *Curr. Opin. Biotechnol.* 19, 445–453. doi: 10.1016/j.copbio.2008.08.001
- Babu, K. R., and Satyanarayana, T. (1993). Extracellular calcium-inhibited alpha-amylase of Bacillus coagulans B 49. *Enz. Microb. Technol.* 15, 1066–1069. doi: 10.1016/0141-0229(93)90056-8
- Babu, K. R., and Satyanarayana, T. (1995). α -Amylase production by thermophilic Bacillus coagulans in solid state fermentation. *Process Biochem.* 30, 305–309. doi: 10.1016/0032-9592(95)87038-5
- Bai, Y., Huang, H., Meng, K., Shi, P., Yang, P., Luo, H., et al. (2012). Identification of an acidic α -amylase from Alicyclobacillus sp. A4 and assessment of its application in the starch industry. *Food Chem.* 131, 1473–1478. doi: 10.1016/j.foodchem.2011.10.036
- Baker-Austin, C., and Dopson, M. (2007). Life in acid: pH homeostasis in acidophiles. *Trends Microbiol.* 15, 165–171. doi: 10.1016/j.tim.2007.02.005
- Bhattacharyya, S., Chakrabarty, B. K., Das, A., Jundu, P., and Banerjee, P. C. (1990). Acidiphilium symbioticum sp. nov., an acidophilic heterotrophic bacterium from Thiobacillus ferrooxidans cultures isolated from Indian mines. *Can. J. Microbiol.* 37, 78–85. doi: 10.1139/m91-012
- Boel, E., Brady, L., Brzozowski, A. M., Derewenda, Z., Dodson, G. G., Jensen, V. J., et al. (1990). Calcium binding in α -amylases: an x-ray diffraction study at 2.1 Å resolution of two enzymes from Aspergillus. *Biochemistry* 29, 6244–6249. doi: 10.1021/bi00478a019
- Brock, T. D., Brock, K. M., Belly, R. T., and Weiss, R. L. (1972). Sulfolobus: a new genus of sulphur-oxidizing bacteria living at low pH and high temperature. *Arch. Mikrobiol.* 84, 54–68. doi: 10.1007/BF00408082
- Brown, S. H., Costantino, H. R., and Kelly, R. M. (1990). Characterization of amyolytic enzyme activities associated with the hyperthermophilic archaeobacterium Pyrococcus furiosus. *Appl. Environ. Microbiol.* 56, 1985–1991.
- Chen, L., Brügger, K., Skovgaard, M., Redder, P., She, Q., Torarinsson, E., et al. (2005). The genome of Sulfolobus acidocaldarius, a model organism of the Crenarchaeota. *J. Bacteriol.* 187, 4992–4999. doi: 10.1128/JB.187.14.4992-4999.2005
- Clark, D. A., and Norris, P. R. (1996). Acidimicrobium ferrooxidans gen. nov., sp. nov.: mixed-culture ferrous iron oxidation with Sulfolobus species. *Microbiology* 142, 785–790. doi: 10.1099/00221287-142-4-785
- Crabb, W. D., and Mitchinson, C. (1997). Enzymes involved in the processing of starch to sugars. *Trends Biotechnol.* 15, 349–352. doi: 10.1016/S0167-7799(97)01082-2
- Cregg, J. M., Tolstorukov, I., Kusari, A., Sunga, J., Madden, K., and Chappell, T. (2009). Expression in the yeast Pichia pastoris. *Methods Enzymol.* 463, 169–189. doi: 10.1016/S0076-6879(09)63013-5
- Darland, G., Brock, T. D., Samsonoff, W., and Conti, S. F. (1970). A thermophilic, acidophilic mycoplasma isolated from a coal refuse pile. *Science* 170, 1416–1418. doi: 10.1126/science.170.3965.1416
- Demirjian, D. C., Moris-Varas, F., and Cassidy, C. S. (2001). Enzymes from extremophiles. *Curr. Opin. Chem. Biol.* 5, 144–151. doi: 10.1016/S1367-5931(00)00183-6
- Dong, G., Vieille, C., Savchenko, A., and Zeikus, J. G. (1997). Cloning, sequencing, and expression of the gene encoding extracellular alpha-amylase from Pyrococcus furiosus and biochemical characterization of the recombinant enzyme. *Appl. Environ. Microbiol.* 63, 3569–3576.
- Eichler, J. (2001). Biotechnological uses of archaeal extremozymes. *Biotechnol. Advances.* 19, 261–278. doi: 10.1016/S0734-9750(01)00061-1
- Elleuche, S., Schröder, C., Sahm, K., and Antranikian, G. (2014). Extremozymes-biocatalysts with unique properties from extremophilic microorganisms. *Curr. Opin. Biotechnol.* 29, 116–123. doi: 10.1016/j.copbio.2014.04.003
- Fan, Z., Wagschal, K., Lee, C. C., Kong, Q., Shen, K. A., Maiti, I. B., et al. (2009). The construction and characterization of two xylan-degrading chimeric enzymes. *Biotechnol. Bioeng.* 102, 684–692. doi: 10.1002/bit.22112
- Ferrer, M., Golyshina, O., Belouqui, A., and Golyshin, P. N. (2007). Mining enzymes from extreme environments. *Curr. Opin. Microbiol.* 10, 207–214. doi: 10.1016/j.mib.2007.05.004
- Fuchs, T., Huber, H., Teiner, K., Burggraf, S., and Stetter, K. O. (1996). Metallosphaera prunae, sp. nov., a novel metal-mobilizing, thermoacidophilic archaeum, isolated from a uranium mine in Germany. *Syst. Appl. Microbiol.* 18, 560–566. doi: 10.1016/S0723-2020(11)80416-9
- Gangadharan, D., Sivaramakrishnan, S., Nampoothiri, K. M., and Pandey, A. (2006). Solid culturing of Bacillus amyloliquefaciens for alpha amylase production. *Food Technol. Biotechnol.* 44, 269–274.
- Golovacheva, R. S., Golyshina, O. V., Karavaiko, G. I., Dorofeev, A. G., and Cheruykh, N. A. (1992). The new iron-oxidizing bacterium Leptospirillum thermoferrooxidans sp. nov. *Microbiology* 61, 744–750.

- Golovacheva, R. S., and Karavaiko, G. I. (1978). A new genus of thermophilic spore-forming bacteria, *Sulfobacillus*. *Microbiology* 47, 658–665.
- González-Toril, E., Llobet-Brossa, E., Casamayor, E. O., Amann, R., and Amils, R. (2003). Microbial ecology of an extreme acidic environment, the Tinto River. *App. Environ. Microbiol.* 69, 4853–4865. doi: 10.1128/AEM.69.8.4853-4865.2003
- Grogan, D., Palm, P., and Zillig, W. (1990). Isolate B12 which harbors a viruslike element represents a new species of the archaeobacterial genus *Sulfolobus*, *Sulfolobus shibatae* new species. *Arch. Microbiol.* 154, 594–599. doi: 10.1007/BF00248842
- Grzybowski, B., Szweida, P., and Synowiecki, J. (2004). Cloning of the thermostable α -amylase gene from *Pyrococcus woesei* in *Escherichia coli*. *Mol. Biotechnol.* 26, 101–109. doi: 10.1385/MB:26:2:101
- Harrison, A. P. (1981). *Acidiphilium cryptum* gen. nov., sp. nov., heterotrophic bacterium from acidic mineral environments. *Int. J. Syst. Bacteriol.* 31, 327–332. doi: 10.1099/00207713-31-3-327
- Hedrich, S., and Johnson, D. B. (2013). *Acidithiobacillus ferridurans* sp. nov., an acidophilic iron-, sulfur- and hydrogen-metabolizing chemolithotrophic gamma proteobacterium. *Int. J. Syst. Evol. Microbiol.* 63, 4018–4025. doi: 10.1099/ijs.0.049759-0
- Hough, D. W., and Danson, M. J. (1999). Extremozymes. *Curr. Opin Chem. Biol.* 3, 39–46. doi: 10.1016/S1367-5931(99)80008-8
- Huang, Y., Krauss, G., Cottaz, S., Driguez, H., and Lipps, G. (2005). A highly acid-stable and thermostable endo- β -glucanase from the thermoacidophilic archaeon *Sulfolobus solfataricus*. *Biochem. J.* 385, 581–588. doi: 10.1042/BJ20041388
- Huber, G., Spinnler, C., Gambacorta, A., and Stetter, K. O. (1989). *Metallosphaera sedula* gen. and sp. nov. represents a new genus of aerobic, metal-mobilizing, thermoacidophilic archaeobacteria. *Syst. Appl. Microbiol.* 12, 38–47. doi: 10.1016/S0723-2020(89)80038-4
- Huber, G., and Stetter, K. O. (1991). *Sulfolobus metallicus*, sp. nov., a novel strictly chemolithoautotrophic thermophilic archaeal species of metal mobilizers. *Syst. Appl. Microbiol.* 14, 372–378. doi: 10.1016/S0723-2020(11)80312-7
- Itoh, T., Yoshikawa, N., and Takashina, T. (2007). *Thermogymnomonas acidicola* gen. nov., sp. nov., a novel thermoacidophilic, cell wall-less archaeon in the order Thermoplasmatales, isolated from a solfataric soil in Hakone, Japan. *Int. J. Syst. Evol. Microbiol.* 57, 2557–2561. doi: 10.1099/ijs.0.65203-0
- Johnson, D. B. (1998). Biodiversity and ecology of acidophilic microorganisms. *FEMS Microbiol. Ecol.* 27, 307–317. doi: 10.1111/j.1574-6941.1998.tb00547.x
- Joshi, S., and Satyanarayana, T. (2015). *In vitro* engineering of microbial enzymes with multifarious applications: prospects and perspectives. *Biores. Technol.* 176, 273–283. doi: 10.1016/j.biortech.2014.10.151
- Joyet, P., Guerinéau, M., and Heslot, H. (1984). Cloning of a thermostable α -amylase gene from *Bacillus licheniformis* and its expression in *Escherichia coli* and *Bacillus subtilis*. *FEMS Microbiol. Lett.* 21, 353–358. doi: 10.1016/0378-1097(84)90097-1
- Kanno, M. (1986). α -Amylase production by *Bacillus acidocaldarius*, *B. stearothermophilus* and their D-cycloserine resistant mutants. *Agric. Biol. Chem.* 50, 2633–2635. doi: 10.1271/abb1961.50.2633
- Karavaiko, G. I., Golyshina, O. V., Troitskii, A. V., Valiehoroman, K. M., Golovacheva, R. S., and Pivovarova, T. A. (1994). *Sulfurococcus yellowstonii* sp. nov., a new species of iron oxidizing and sulphur oxidizing thermoacidophilic archaeobacteria. *Microbiology* 63, 379–387.
- Koch, R., Spreinat, A., Lemke, K., and Antranikian, G. (1991). Purification and properties of a hyperthermoactive α -amylase from the archaeobacterium *Pyrococcus woesei*. *Arch. Microbiol.* 155, 572–578. doi: 10.1007/BF00245352
- Kumar, P., and Satyanarayana, T. (2007). Optimization of culture variables for improving glucoamylase production by alginate-entrapped *Thermomucor indiciae*-seudaticae using statistical methods. *Biores. Technol.* 98, 1252–1259. doi: 10.1016/j.biortech.2006.05.019
- Kurosawa, N., Itoh, Y. H., Iwai, T., Sugai, A., Uda, I., Kimura, N., et al. (1998). *Sulfurisphaera ohwakuensis* gen. nov., sp. nov., a novel extremely thermophilic acidophile of the order Sulfolobales. *Int. J. Syst. Bacteriol.* 48, 451–456. doi: 10.1099/00207713-48-2-451
- Laderman, K. A., Davis, B. R., Krutzsch, H. C., Lewis, M. S., Griko, Y. V., Privalov, P. L., et al. (1993). The purification and characterization of an extremely thermostable alpha-amylase from the hyperthermophilic archaeobacterium *Pyrococcus furiosus*. *J. Biol. Chemistry* 268, 24394–24301.
- Linden, A., Mayans, O., Meyer-Klaucke, W., Antranikian, G., and Wilmanns, M. (2003). Differential regulation of a hyperthermophilic α -amylase with a novel (Ca, Zn) two-metal center by zinc. *J. Biol. Chem.* 278, 9875–9884. doi: 10.1074/jbc.M211339200
- Liu, Y. H., Hu, B., Xu, Y. J., Bo, J. X., Fan, S., Wang, J. L., et al. (2012). Improvement of the acid stability of *Bacillus licheniformis* α -amylase by error-prone PCR. *J. App. Microbiol.* 113, 541–549. doi: 10.1111/j.1365-2672.2012.05359.x
- Liu, Y. H., Lu, F. P., Li, Y., Wang, J. L., and Gao, C. (2008a). Acid stabilization of *Bacillus licheniformis* alpha-amylase through introduction of mutations. *App. Microbiol. Biotechnol.* 80, 795–803. doi: 10.1007/s00253-008-1580-5
- Liu, Y. H., Lu, F. P., Li, Y., Yin, X. B., Wang, Y., and Gao, C. (2008b). Characterization of mutagenised acid-resistant alpha-amylase expressed in *Bacillus subtilis* WB600. *Appl. Microbiol. Biotechnol.* 78, 85–94. doi: 10.1007/s00253-007-1287-z
- Lobos, J. H., Chisholm, T. E., Bopp, L. H., and Holmes, D. S. (1986). *Acidiphilium organovororum* sp. nov., an acidophilic heterotroph isolated from a *Thiobacillus ferrooxidans* culture. *Int. J. Syst. Bacteriol.* 36, 139–144. doi: 10.1099/00207713-36-2-139
- Matsubara, H., Goto, K., Matsumura, T., Mochida, K., Iwaki, M., Niwa, M., et al. (2002). *Alicyclobacillus acidiphilus* sp. nov., a novel thermo-acidophilic, omega-acyclic fatty acid-containing bacterium isolated from acidic beverages. *Int. J. Syst. Evol. Microbiol.* 52, 1681–1685. doi: 10.1099/00207713-52-5-1681
- Matzke, J., Schwermann, B., and Bakker, E. P. (1997). Acidostable and acidophilic proteins: the example of the α -amylase from *Alicyclobacillus acidocaldarius*. *Comparative Biochem. Physiol. Part A Physiol.* 118, 475–479.
- Mavromatis, K., Sikorski, J., Lapidus, A., Del Rio, T. G., Copeland, A., Tice, H., et al. (2010). Complete genome sequence of *Alicyclobacillus acidocaldarius* type strain (104-IA T). *Standards Genomic Sci.* 2, 9–21. doi: 10.4056/signs.591104
- Mehta, D., and Satyanarayana, T. (2013). Biochemical and molecular characterization of recombinant acidic and thermostable raw-starch hydrolysing α -amylase from an extreme thermophile *Geobacillus thermoleovorans*. *J. Mol. Catal. B Enz.* 85, 229–238. doi: 10.1016/j.molcatb.2012.08.017
- Mehta, D., and Satyanarayana, T. (2014). Domain C of thermostable α -amylase of *Geobacillus thermoleovorans* mediates raw starch adsorption. *App. Microbiol. Biotechnol.* 98, 4503–4519. doi: 10.1007/s00253-013-5459-8
- Mehta, D., and Satyanarayana, T. (2016). Bacterial and archaeal α -amylases: diversity and amelioration of the desirable characteristics for industrial applications. *Front. Microbiol.* 7:1129. doi: 10.3389/fmicb.2016.01129
- Murakami, S., Nishimoto, H., Toyama, Y., Shimamoto, E., Takenaka, S., Kaulpiboon, J., et al. (2007). Purification and characterization of two alkaline, thermotolerant α -amylases from *Bacillus halodurans* 38C-2-1 and expression of the cloned gene in *Escherichia coli*. *Biosci. Biotechnol. Biochem.* 71, 2393–2401. doi: 10.1271/bbb.60666
- Nielsen, J. E., Beier, L., Otzen, D., Borchert, T. V., Frantzen, H. B., Andersen, K. V., et al. (1999). Electrostatics in the active site of an α -amylase. *FEBS J.* 264, 816–824. doi: 10.1046/j.1432-1327.1999.00664.x
- Nielsen, J. E., and Borchert, T. V. (2000). Protein engineering of bacterial α -amylases. *Biochim. Biophys. Acta* 1543, 253–274. doi: 10.1016/S0167-4838(00)00240-5
- Nonaka, T., Fujihashi, M., Kita, A., Hagihara, H., Ozaki, K., Ito, S., et al. (2003). Crystal structure of calcium-free α -amylase from *Bacillus* sp. strain KSM-K38 (AmyK38) and its sodium ion binding sites. *J. Biol. Chem.* 278, 24818–24824. doi: 10.1074/jbc.M212763200
- Norris, P. R., Clark, D. A., Owen, J. P., and Waterhouse, S. (1996). Characteristics of *Sulfobacillus acidophilus* sp. nov. and other moderately thermophilic mineral-sulfide-oxidizing bacteria. *Microbiology* 142, 775–783. doi: 10.1099/00221287-142-4-775
- Orr, R. V., Shewfelt, R. L., Huang, C. J., Tefera, S., and Beuchat, L. R. (2000). Detection of guaiacol produced by *Alicyclobacillus acidoterrestris* in apple juice by sensory and chromatographic analyses, and comparison with spore and vegetative cell populations. *J. Food Protec.* 63, 1517–1522. doi: 10.4315/0362-028X-63.11.1517
- Parashar, D., and Satyanarayana, T. (2016a). Production of Ca²⁺-independent and acidstable Recombinant α -amylase of *Bacillus acidicola* extracellularly and its applicability in generating maltoooligosaccharides. *Mol. Biotechnol.* 58, 707–717. doi: 10.1007/s12033-016-9970-x

- Parashar, D., and Satyanarayana, T. (2016b). Enhancing the production of recombinant acidic α -amylase and phytase in *Pichia pastoris* under dual promoters [constitutive (GAP) and inducible (AOX)] in mixed fed batch high cell density cultivation. *Process Biochem.* 51, 1315–1322. doi: 10.1016/j.procbio.2016.07.027
- Parashar, D., and Satyanarayana, T. (2016c). A chimeric α -amylase engineered from *Bacillus acidicola* and *Geobacillus thermoleovorans* with improved thermostability and catalytic efficiency. *J. Ind. Microbiol. Biotechnol.* 43, 473–484. doi: 10.1007/s10295-015-1721-7
- Parashar, D., and Satyanarayana, T. (2017a). Production of chimeric acidic α -amylase by the recombinant *Pichia pastoris* and its applications. *Front. Microbiol.* 8:493. doi: 10.3389/fmicb.2017.00493
- Parashar, D., and Satyanarayana, T. (2017b). Engineering a chimeric acid-stable α -amylase-glucoamylase (Amy-Glu) for one step starch saccharification. *Int. J. Biol. Macromol.* 99, 274–281. doi: 10.1016/j.ijbiomac.2017.02.083
- Pedersen, H., and Nielsen, J. (2000). The influence of nitrogen sources on the α -amylase productivity of *Aspergillus oryzae* in continuous cultures. *App. Microbiol. Biotechnol.* 53, 278–281. doi: 10.1007/s002530050021
- Peng, S., Chu, Z., Lu, J., Li, D., Wang, Y., Yang, S., et al. (2016). Co-expression of chaperones from *P. furiosus* enhanced the soluble expression of the recombinant hyperthermophilic α -amylase in *E. coli*. *Cell Stress Chaperones* 21, 477–484. doi: 10.1007/s12192-016-0675-7
- Priyadarshini, R., Manoharan, S., Hemalatha, D., and Gunasekaran, P. (2010). Repeated random mutagenesis of α -amylase from *Bacillus licheniformis* for improved pH performance. *J. Microbiol. Biotechnol.* 20, 1696–1701. doi: 10.4014/jmb.1008.08013
- Prokofeva, M. I., Miroshnichenko, M. L., Kostrikin, N. A., Chernykh, N. A., Kuznetsov, B. B., Tourova, T. P., et al. (2000). *Acidilobus aceticus* gen. nov., sp. nov., a novel anaerobic thermoacidophilic archaeon from continental hot vents in Kamchatka. *Int. J. Syst. Evol. Microbiol.* 50, 2001–2008. doi: 10.1099/00207713-50-6-2001
- Raddadi, N., Cherif, A., Daffonchio, D., Neifar, M., and Fava, F. (2015). Biotechnological applications of extremophiles, extremozymes and extremolytes. *Appl. Microbiol. Biotechnol.* 99, 7907–7913. doi: 10.1007/s00253-015-6874-9
- Ranjan, B., and Satyanarayana, T. (2016). Recombinant HAP phytase of the thermophilic mold *Sporotrichum thermophile*: expression of the codon-optimized phytase gene in *Pichia pastoris* and applications. *Mol. Biotechnol.* 58, 137–147. doi: 10.1007/s12033-015-9909-7
- Reed, C. J., Lewis, H., Trejo, E., Winston, V., and Evilia, C. (2013). Protein adaptations in archaeal extremophiles. *Archaea* 13, 1–14. doi: 10.1155/2013/373275
- Ren-Long, J., Wu, J., Chaw, S. M., Tsai, C. W., and Tsen, S. D. (1999). A novel species of thermoacidophilic archaeon, *Sulfolobus yangmingensis* sp. nov. *Int. J. Syst. Bacteriol.* 49, 1809–1816. doi: 10.1099/00207713-49-4-1809
- Ribeiro, L. F., Furtado, G. P., Lourenzoni, M. R., Costa-Filho, A. J., Santos, C. R., Nogueira, S. C., et al. (2011). Engineering bifunctional laccase-xylanase chimeras for improved catalytic performance. *J. Biol. Chem.* 286, 43026–43038. doi: 10.1074/jbc.M111.253419
- Sajedi, R. H., Naderi-Manesh, H., Khajeh, K., Ahmadvand, R., Ranjbar, B., Asoodeh, A., et al. (2005). A Ca-independent α -amylase that is active and stable at low pH from the *Bacillus* sp. K1R-8104. *Enz. Microb. Technol.* 36, 666–671. doi: 10.1016/j.enzmictec.2004.11.003
- Savchenko, A., Vieille, C., Kang, S., and Zeikus, J. G. (2002). *Pyrococcus furiosus* α -amylase is stabilized by calcium and zinc. *Biochemistry* 41, 6193–6201. doi: 10.1021/bi012106s
- Schäfer, K., Magnusson, U., Scheffel, F., Schiefner, A., Sandgren, M. O., Diederichs, K., et al. (2004). X-ray structures of the maltose–maltodextrin-binding protein of the thermoacidophilic bacterium *Alicyclobacillus acidocaldarius* provide insight into acid stability of proteins. *J. Mol. Biol.* 335, 261–274. doi: 10.1016/j.jmb.2003.10.042
- Schleper, C., Puehler, G., Holz, L., Gambacorta, A., Janekovic, D., and Santarius, U. (1995). *Picrophilus* gen. nov., fam. Nov. a novel aerobic, heterotrophic, thermoacidophilic genus and family comprising archaea capable of growth around pH 0. *J. Bacteriol.* 177, 7050–7059.
- Schwermann, B., Pfau, K., Liliensiek, B., Schleyer, M., Fischer, T., and Bakker, E. P. (1994). Purification, properties and structural aspects of a thermoacidophilic α -amylase from *Alicyclobacillus acidocaldarius* ATCC 27009. *FEBS J.* 226, 981–991.
- Seeger, A., Langworthy, T. A., and Stetter, K. O. (1988). *Thermoplasma acidophilum* and *Thermoplasma volcanium* sp. nov. from solfatara fields. *Syst. Appl. Microbiol.* 10, 161–171.
- Seeger, A., Neuner, A., Kristjansson, J. K., and Stetter, K. O. (1986). *Acidianus infernus* gen. nov., sp. nov., and *Acidianus brierleyi* comb. nov.: facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeobacteria. *Int. J. Syst. Evol. Microbiol.* 36, 559–564. doi: 10.1099/00207713-36-4-559
- Sharma, A., Kawarabayasi, Y., and Satyanarayana, T. (2012). Acidophilic bacteria and archaea: acid-stable biocatalysts and their potential applications. *Extremophiles* 16, 1–19. doi: 10.1007/s00792-011-0402-3
- Sharma, A., Parashar, D., and Satyanarayana, T. (2016). “Acidophilic Microbes: Biology and Applications,” in *Biotechnology of Extremophiles*, ed P. H. Rampelotto (Cham: Springer International Publishing), 215–241.
- Sharma, A., and Satyanarayana, T. (2010). High maltose-forming, Ca²⁺-independent and acid-stable α -amylase from a novel acidophilic bacterium, *Bacillus acidicola*. *Biotechnol. Lett.* 32, 1503–1507. doi: 10.1007/s10529-010-0322-9
- Sharma, A., and Satyanarayana, T. (2011). Optimization of medium components and cultural variables for enhanced production of acidic high maltose-forming and Ca²⁺-independent α -amylase by *Bacillus acidicola*. *J. Biosci. Bioeng.* 111, 550–553. doi: 10.1016/j.jbiosc.2011.01.004
- Sharma, A., and Satyanarayana, T. (2012a). Cloning and expression of acid-stable, high maltose-forming, Ca²⁺-independent α -amylase from an acidophile *Bacillus acidicola* and its applicability in starch hydrolysis. *Extremophiles* 16, 515–522. doi: 10.1007/s00792-012-0451-2
- Sharma, A., and Satyanarayana, T. (2012b). Production of acid-stable and high maltose-forming α -amylase of *Bacillus acidicola* by solid-state fermentation and immobilized cells and its applicability in baking. *Appl. Biochem. Biotechnol.* 168, 1025–1034. doi: 10.1007/s12010-012-9838-x
- Sharma, A., and Satyanarayana, T. (2013a). Microbial acid-stable α -amylases: characteristics, genetic engineering and applications. *Process Biochem.* 48, 201–211. doi: 10.1016/j.procbio.2012.12.018
- Sharma, A., and Satyanarayana, T. (2013b). Structural and biochemical features of acidic α -amylase of *Bacillus acidicola*. *Int. J. Biol. Macromol.* 61, 416–423. doi: 10.1016/j.ijbiomac.2013.08.003
- Shaw, A., Bott, R., and Day, A. G. (1999). Protein engineering of α -amylase for low pH performance. *Curr. Opin. Biotechnol.* 10, 349–352. doi: 10.1016/S0958-1669(99)80063-9
- Shima, S., and Suzuki, K. I. (1993). *Hydrogenobacter acidophilus* sp. nov., a thermoacidophilic, aerobic, hydrogen-oxidizing bacterium requiring elemental sulfur for growth. *Int. J. Syst. Evol. Microbiol.* 43, 703–708. doi: 10.1099/00207713-43-4-703
- Shirokizawa, O., Akiba, T., and Horikoshi, K. (1990). Nucleotide sequence of the G6-amylase gene from alkalophilic *Bacillus* sp. H-167. *FEMS Microbiol. Lett.* 70, 131–135. doi: 10.1111/j.1574-6968.1990.tb13965.x
- Spohner, S. C., Müller, H., Quitmann, H., and Czermak, P. (2015). Expression of enzymes for the usage in food and feed industry with *Pichia pastoris*. *J. Biotechnol.* 202, 118–134. doi: 10.1016/j.jbiotec.2015.01.027
- Uma Maheswar Rao, J. L., and Satyanarayana, T. (2003). Statistical optimization of a high maltose-forming, hyperthermostable and Ca²⁺-independent alpha-amylase production by an extreme thermophile *Geobacillus thermoleovorans* using response surface methodology. *J. App. Microbiol.* 95, 712–718. doi: 10.1046/j.1365-2672.2003.02036.x
- Verma, D., and Satyanarayana, T. (2012). Molecular approaches for ameliorating microbial xylanases. *Biores. Technol.* 117, 360–367. doi: 10.1016/j.biortech.2012.04.034
- Wang, L., Zhou, Q., Chen, H., Chu, Z., Lu, J., Zhang, Y., et al. (2007). Efficient solubilization, purification of recombinant extracellular α -amylase from *Pyrococcus furiosus* expressed as inclusion bodies in *Escherichia coli*. *J. Ind. Microbiol. Biotechnol.* 34, 187–192. doi: 10.1007/s10295-006-0185-1
- Wang, P., Wang, P., Tian, J., Yu, X., Chang, M., Chu, X., et al. (2016). A new strategy to express the extracellular α -amylase from *Pyrococcus furiosus* in *Bacillus amyloliquefaciens*. *Sci. Rep.* 6, 222–229. doi: 10.1038/srep22229

- Widersten, M. (2014). Protein engineering for development of new hydrolytic biocatalysts. *Curr. Opin. Chem. Biol.* 21, 42–47. doi: 10.1016/j.cbpa.2014.03.015
- Wind, R. D., Uitdehaag, J. C., Buitelaar, R. M., Dijkstra, B. W., and Dijkhuizen, L. (1998). Engineering of cyclodextrin product specificity and pH optima of the thermostable cyclodextrin glycosyltransferase from *Thermoanaerobacterium thermosulfurigenes* EM1. *J. Biological. Chem.* 273, 5771–5779. doi: 10.1074/jbc.273.10.5771
- Xiang, X., Dong, X., and Huang, L. (2003). *Sulfolobus tengchongensis* sp. nov., a novel thermoacidophilic archaeon isolated from a hot spring in Tengchong, China. *Extremophiles* 7, 493–498. doi: 10.1007/s00792-003-0355-2
- Yang, H., Liu, L., Shin, H. D., Chen, R. R., Li, J., Du, G., et al. (2013). Structure-based engineering of histidine residues in the catalytic domain of α -amylase from *Bacillus subtilis* for improved protein stability and catalytic efficiency under acidic conditions. *J. Biotechnol.* 164, 59–66. doi: 10.1016/j.jbiotec.2012.12.007
- Zhang, R. Y., Xia, J. L., Peng, J. H., Zhang, Q., Zhang, C. G., Nie, Z. Y., et al. (2010). A new strain *Leptospirillum ferriphilum* YTW315 for bioleaching of metal sulfides ores. *Trans. Nonferrous Metals Soc. China* 20, 135–141. doi: 10.1016/S1003-6326(09)60110-2
- Zhu, H., Reynolds, L. B., and Menassa, R. (2017). A hyper-thermostable α -amylase from *Pyrococcus furiosus* accumulates in *Nicotiana tabacum* as functional aggregates. *BMC Biotechnol.* 17:53. doi: 10.1186/s12896-017-0372-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Parashar and Satyanarayana. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.