



CRISPR/Cas12a Coupled With Recombinase Polymerase Amplification for Sensitive and Specific Detection of *Aphelenchoides besseyi*

Anpeng Zhang^{1,2†}, Bin Sun^{1†}, Jianming Zhang¹, Can Cheng¹, Jihua Zhou¹, Fuan Niu¹, Zhongyong Luo³, Luzhen Yu⁴, Cui Yu⁴, Yuting Dai¹, Kaizhen Xie¹, Qiyang Hu¹, Yue Qiu¹, Liming Cao^{1,2*} and Huangwei Chu^{1,2*}

OPEN ACCESS

Edited by:

Xiaoming Zhou,
South China Normal University, China

Reviewed by:

Ulhas S. Kadam,
Gyeongsang National University,
South Korea
Hongxing Liu,
First Affiliated Hospital of Guangzhou
Medical University, China

*Correspondence:

Liming Cao
caoliming@saas.sh.cn
Huangwei Chu
chuhuangwei@saas.sh.cn

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Biosensors and Biomolecular
Electronics,
a section of the journal
Frontiers in Bioengineering and
Biotechnology

Received: 05 April 2022

Accepted: 17 May 2022

Published: 30 June 2022

Citation:

Zhang A, Sun B, Zhang J, Cheng C,
Zhou J, Niu F, Luo Z, Yu L, Yu C, Dai Y,
Xie K, Hu Q, Qiu Y, Cao L and Chu H
(2022) CRISPR/Cas12a Coupled With
Recombinase Polymerase
Amplification for Sensitive and Specific
Detection of *Aphelenchoides besseyi*.
Front. Bioeng. Biotechnol. 10:912959.
doi: 10.3389/fbioe.2022.912959

¹Institute of Crop Breeding and Cultivation, Shanghai Academy of Agricultural Sciences, Shanghai, China, ²Shanghai Key Laboratory of Agricultural Genetics and Breeding, Shanghai Academy of Agricultural Sciences, Shanghai, China, ³Shanghai Agricultural Science and Technology Seed Co., Ltd., Shanghai, China, ⁴Technical Center for Animal Plant and Food Inspection and Quarantine, Shanghai Customs, Shanghai, China

Aphelenchoides besseyi (*A. besseyi*), a seed-borne parasitic nematode, is the causal agent of rice white tip disease (RWTD), which may result in a drastic loss of rice yield. Seed treatments are currently considered to be the most effective means of preventing the spread of RWTD. Therefore, the rapid, highly specific, and accurate detection of *A. besseyi* from rice seeds is crucial for the surveillance, prevention, and control of RWTD. Here, we describe a novel detection assay that combines recombinase polymerase amplification (RPA) and CRISPR/Cas12a to detect *A. besseyi* (termed RPA-Cas12a-Ab), with a low limit of detection (LOD) of 1 copy/ μ l of plasmid or 1:10⁷ diluted DNA extracted from individual nematodes. To improve the user-friendliness, lateral flow strip assay (LFA) was adopted to visualize the detection result. The LOD of the RPA-Cas12a-Ab LFA assay was 1,000 copies/ μ l plasmid or 1:10 diluted DNA extracted from individual nematodes. The assay developed in this study was able to identify *A. besseyi* in 45 min with high accuracy and sensitivity without cross reaction with three closely related non-*A. besseyi* species. Thus, RPA-Cas12a-Ab is a rapid, sensitive, and specific detection system that requires no sophisticated equipment and shows promise for on-site surveillance of *A. besseyi*.

Keywords: CRISPR/Cas12a, RPA, *Aphelenchoides besseyi*, rice, molecular identification, detection

INTRODUCTION

Aphelenchoides besseyi (*A. besseyi*), also known as the rice white tip nematode, has a broad host range of more than 200 plants (Cheng et al., 2013), including rice, that is, the staple food of more than half of the world's population. *A. besseyi* is widely distributed in almost all rice-growing areas in the world, and severe infestations of *A. besseyi* may result in yield losses of up to 50% (Duncan and Moens, 2006; Cheng et al., 2013; Sánchez-Monge et al., 2015). *A. besseyi* is a seed-borne nematode, and infected seeds are considered the main source of initial inoculum for rice white tip disease (RWTD) (Devran and Gökür, 2020). It is difficult to implement a good means for thorough control

of RWTD once it outbreaks (Devran and Gökür, 2020). Therefore, there is an urgent need to develop reliable and rapid detection and identification tools for *A. besseyi* (Buonicontrò et al., 2018).

The conventional morphologic diagnostic method is difficult to be widely adopted due to the morphological comparability between the numerous nominal species in the *Aphelenchoides* genus (Bai et al., 2017). Therefore, molecular techniques have been the preferred methods for the identification of *A. besseyi* because of the higher sensitivity and specificity (Rybarczyk-Mydłowska et al., 2012). Currently, molecular methods for *A. besseyi* detection are mostly derived from polymerase chain reaction (PCR)-based technologies, such as loop-mediated isothermal amplification (LAMP) (Bai et al., 2017; Yang and Yu, 2019), PCR (Ibrahim et al., 1994; Devran et al., 2017; Sánchez-Monge et al., 2017), and real-time PCR (qPCR) (Rybarczyk-Mydłowska et al., 2012; Buonicontrò et al., 2018; Çelik and Devran, 2019). The sensitivity of the LAMP method is similar to that of the routine PCR method, which could specifically detect target DNA levels as low as 10^3 copies/ μ l (Yang and Yu, 2019), but significantly lower than that of qPCR (Buonicontrò et al., 2018; Çelik and Devran, 2019). Although qPCR is the most sensitive method for detecting low concentrations of target DNA, the utilization of the qPCR method in the identification of *A. besseyi* is hampered as it is time-consuming, costly, and dependent on expensive equipment. Therefore, the development of an easy, cost-effective, and sensitive molecular method, which does not require specialized technical knowledge and auxiliary equipment, is desired (Chang et al., 2020).

Cas12a, a class II type V-A endonuclease of the CRISPR/Cas effector family, is characterized as a dual nuclease catalyzing its own CRISPR RNA (crRNA) maturation, target-site recognition, and DNA cleavage (Zetsche et al., 2015). Cas12a can detect the presence of target DNA under the guidance of crRNA. Despite mediating target-site sequence-specific DNA cleavage, Cas12a has non-specific single-stranded DNase activity (*trans*-cleavage activity) upon the formation of the Cas12a/crRNA/target DNA ternary complex (Zetsche et al., 2015; Li et al., 2018a; Chen et al., 2018). Utilizing the *trans*-cleavage activity of Cas12a, a nucleic acid detection method, named one-Hour Low-cost Multipurpose Highly Efficient System (HOLMES), was first reported in 2018 (Li et al., 2018a; Li et al., 2018b). In parallel, another similar nucleic acid detection assay, named DNA endonuclease-targeted CRISPR *trans* reporter (DETECTR), was invented by another group (Chen et al., 2018). By coupling preamplification of the target nucleic acid with PCR or isothermal amplification, the HOLMES or DETECTR assays were able to detect target DNA at a concentration as low as aM (Li et al., 2018b; Chen et al., 2018; Gootenberg et al., 2018).

The CRISPR/Cas12a-based nucleic acid detection technology has been successfully applied to detect biallelic mutants (Xiao et al., 2020), genetically modified organisms (Zhang et al., 2020; Liu et al., 2021), and various pathogens, such as the high-risk HPV16 or HPV18 virus (ZIKV) (Tsou et al., 2019), the *pseudorabies* virus (Li L. et al., 2019), the *human papillomavirus* (Gootenberg et al., 2018), the cancer-related virus (Huang et al., 2018), the white spot syndrome virus (Chaijarasphong et al., 2019), the African swine fever (Bai et al., 2019), and *Magnaporthe oryzae* (Zhang et al., 2020). However, to the

best of our knowledge, no study has reported the use of a CRISPR/Cas12a-based assay for the molecular detection of *A. besseyi*.

In this study, we developed a novel detection assay that combines the CRISPR/Cas12a method with recombinase polymerase amplification (RPA) to detect *A. besseyi*. This assay has three steps, including 1) 20 min of preamplification of target DNA at 37°C, 2) 20 min of Cas12a-reaction at 37°C, and 3) readout detection results using a fluorescence analyzer or the lateral flow strip assay (LFA) within 5 min (Figure 1). Our new molecular assay (RPA-Cas12a-Ab) has the potential to detect *A. besseyi* with high accuracy, sensitivity, and specificity without the need for complicated equipment, providing a convenient and simple method that can be utilized on-site for rapid detection of *A. besseyi*.

MATERIALS AND METHODS

Materials

The nematodes used in this study are listed in Table 1. Two rice grain samples, AB1 and AB2, infected by *A. besseyi* were collected from paddy fields in Shanghai and Nanjing, respectively. The related *Aphelenchoides* spp., *A. fragariae* (AF), and *A. subtenuis* (AS), were intercepted in Shanghai ports and saved in Shanghai Entry-Exit Inspection and Quarantine Bureau. *M. incognita* (MI) were kindly provided by Prof. Xuan Wang (Nanjing Agricultural University, Nanjing, China) (Wang et al., 2018).


RPA primers, crRNA, and ssDNA reporters were synthesized by Sangon Biotech (Shanghai, China). The RPA assay kit (TwistAmp Basic kit, TABAS03KIT) was purchased from TwistDx Ltd. (TwistDx Ltd., United Kingdom). The LbCas12a (cpf1) nuclease (32108-03) was purchased from Tolo Biotech (Tolo Biotech, Anhui, China). The lateral flow strip (JY0301) was purchased from Warbio Biotech (Nanjing, China).

Recombinase Polymerase Amplification Primer, CRISPR RNA, and ssDNAs Reporter Design

The internal transcribed spacer region (ITS) sequence of the *A. besseyi* 18S ribosomal RNA (18S rRNA) gene was retrieved from GenBank under accession number JF826519. According to the design manual of the manufacturer, RPA primers should be 30–35 bases long, and the T_m values are not the critical factor for its performance. Based on these principles, a pair of specific RPA primers with a 237 bp fragment length was designed (Table 2, Supplementary Figure S1). Eight crRNA sequences targeting the ITS of *A. besseyi* were designed based on the PAM site of Cas12a (Table 2, Supplementary Figure S1). In addition, for target detection using Cas12a *trans*-cleavage activity, the HEX-BHQ1-labeled ssDNA reporter (HB-reporter) or FAM-Biotin labeled ssDNA reporter (FB-reporter) was designed for the fluorescence assay or LFA assay, respectively (Table 2).

Construction and Purification of Reference Plasmid pUC57-18S

In order to determine the sensitivity of the RPA-Cas12a-Ab assays, a reference plasmid was constructed by cloning the

**TABLE 1** | Plant-parasitic nematode species used in this study.

Sample No.	Species	Host	Origin
AB1	<i>A. besseyi</i>	<i>Oryza sativa</i>	Shanghai, China
AB2	<i>A. besseyi</i>	<i>Oryza sativa</i>	Jiangsu, China
AF	<i>A. fragariae</i>	<i>Paeonia albiflora</i>	Netherlands
AS	<i>A. Subtenuis</i>	<i>Allium giganteum</i>	Netherlands
MI	<i>M. incognita</i>	Tomato	Jiangsu, China

237 bp RPA fragment of 18S rRNA ITS into pUC57. The resultant plasmid, pUC57-18S, was verified by restriction enzyme digests and sequencing. The concentration of the purified pUC57-18S

was determined using a NanoDrop 2000 Spectrophotometer (Thermo Scientific, United States) and calculated in copy numbers per μl and then diluted in TE buffer to a concentration of 10^{10} copies/ μl .

Nematode Extraction and DNA Isolation

The nematodes were extracted using the modified Baermann funnel method (Yang and Yu, 2019). In brief, 200 husked rice grains or other 5-g chopped plant materials infected with nematodes were wrapped with gauze and placed into a funnel filled with sterile water. After soaking for 24 h, 1,000 μl of the nematode suspension was collected from the bottom of the funnel

TABLE 2 | RPA primer, crRNA, and ssDNA reporter sequences.

Name	Sequences (5'-3')
RPA-F	GTTAATAAGCACGAATTACAGATATTACGAG
RPA-R	AGTACTAACAACGTAATCACAGGCATACTC
CrRNA1	AAUUUCUACUGUUGUAGAUCGAUUGCAUUAUUGCGUCGU
CrRNA2	AAUUUCUACUGUUGUAGAUCGAUUGCAUUAUUGCGUCGUCGGGU
CrRNA3	AAUUUCUACUGUUGUAGAUCGAUGGAAAGGGAAGCCAAA
CrRNA4	AAUUUCUACUGUUGUAGAUCGAUGGAAAGGGAAGCCAAAUGCAU
CrRNA5	AAUUUCUACUGUUGUAGAUCGAUUCGAAUCGUCGACUC
CrRNA6	AAUUUCUACUGUUGUAGAUCGAUUCGAAUCGUCGACUCUCC
CrRNA7	AAUUUCUACUGUUGUAGAUGGGCGUAUUCUUCGGAGU
CrRNA8	AAUUUCUACUGUUGUAGAUGGGCGUAUUCUUCGGAGUAUGCC
HB-reporter	HEX-TTTTTTTTTT-BHQ1
FB-reporter	6FAM-TTTTTTTTTT-Biotin

into a 1.5 ml centrifuge tube and centrifuged for 3 min at a speed of 6,000 r/min. The supernatant was then discarded.

DNA was extracted from individual nematodes. A single nematode was transferred to a separate microtube containing 16 μ l of ddH₂O and 2 μ l of 10 \times PCR buffer, followed by three freeze-thaw cycles of 1 min freezing in liquid nitrogen and 2 min thawing at 95°C. 2 μ l of 10 mg/ml proteinase K was added to the mixture, incubated at 65°C for 90 min, and then incubated at 95°C for 10 min to deactivate proteinase K. The supernatant was used as the DNA template. Genomic DNA (gDNA) of rice leaves was extracted from fresh leaves using the CTAB method (Borges et al., 2009).

Recombinase Polymerase Amplification Reactions

RPA reactions were performed with a TwistAmp Basic kit (TABAS03KIT, TwistDx Ltd., United Kingdom), following the manufacturer's instructions. In brief, the RPA reaction mixture was prepared, containing 29.5 μ l of TwistAmp RPA buffer, 2.4 μ l each of 10 μ M RPA primer, 1 μ l of the DNA template, and 12.2 μ l of sterile ddH₂O. After gentle mixing, the reaction mixture was added to the TwistAmp Basic reaction tube containing lyophilized enzyme powder. Then, 2.5 μ l of 280 mM of magnesium acetate was added and mixed well, followed by incubation at 37°C for 20 min. In no template control (NTC) reaction, sterile ddH₂O was used in place of the DNA template.

Establishment of the RPA-Cas12a-Ab Fluorescence Assay

The Cas12a-mediated cleavage reaction was performed in a total volume of 20 μ l with 2 μ l 10 \times Cas12a reaction buffer, 250 nM crRNA, 50 nM LbCas12a, 10 U RNase inhibitor, 250 nM HB-reporter (labeled with HEX at the 5'-end and BHQ1 at the 3'-end), and 2 μ l of the RPA products. The mixture was placed in a PCR plate and then incubated at 37°C for up to 70 min in a Real-Time PCR system (Applied Biosystems, QuantStudio 6 Flex), with fluorescence signals measured every 30 s. Alternatively, the 96-well microplate was incubated at 37°C for 20 min in a fluorescence plate reader (Tecan, Infinite 200Pro) and with excitation set at 530 nm and fluorescence signals measured at 565 nm.

RPA-Cas12a-Ab Assay With the Lateral Flow Strip Assay

The Cas12a-mediated cleavage reaction mix for LFA detection was similar as above, except that FB-reporter (labeled with 6-FAM at the 5'-end and biotin at the 3'-end) was used instead of HB-reporter (Table 2). After 20 min incubation at 37°C, the reaction products were diluted to 100 μ l, after which lateral flow strips were dipped into the resultant reactions. The results were visualized after 5 min of incubation at room temperature.




FIGURE 2 | Evaluation of the performance of RPA primers. M, DL2000 DNA marker; 1–8, RPA products amplified at 25°C, 28°C, 32°C, 35°C, 37°C, 39°C, 42°C, and 45°C, respectively.

RESULTS


Establishing the RPA-Cas12a-Ab Fluorescence Detection Assay

In order to evaluate the performance of RPA primers, RPA experiments were carried out using purified reference plasmid DNA as a template at eight different temperatures including 25°C, 28°C, 32°C, 35°C, 37°C, 39°C, 42°C, and 45°C. All the temperatures were able to amplify a clear and intense 237bp DNA band visualized using agarose electrophoresis (Figure 2). For the convenience of experimental operation and the minimal requirement of equipment, 37°C was chosen for subsequent RPA reactions as it is also the optimal temperature for the activity of Cas12a.


crRNA is the main factor affecting the *trans*-cleavage activity of Cas12a (Creutzburg et al., 2020; Zhang et al., 2020; Liu et al., 2021; Ma et al., 2021). To obtain a better *trans*-cleavage activity of Cas12a, eight crRNAs targeting the 18S rRNA ITS of *A. besseyi* based on the PAM motif of Cas12a were designed (Table 2, Supplementary Figure S1), and the RPA-Cas12a fluorescence assay was performed to select the optimal crRNA for further experiments (Figure 1). The fluorescence signals were recorded at 30 s intervals with a QuantStudio 6 Flex Real-Time PCR System (Applied Biosystems). The fluorescence signals were detectable within 10 min for all crRNAs, among which crRNA5 generated the highest fluorescence intensity (Figures 3A–C), implying the high activity of crRNA5 for target DNA detection. The Cas12a reactions triggered by crRNA5 were completed within about 20 min, and the fluorescence kinetic curves plateaued thereafter (Figure 3A). Therefore, crRNA5 was chosen as the optimal crRNA for the RPA-Cas12a-Ab detection assay, which only takes 20 min to achieve ideal results from Cas12a reactions.

Evaluation of the Specificity of the RPA-Cas12a-Ab Fluorescence Assay

Five nematode samples, including AB1 and AB2, which were confirmed as *A. besseyi* isolates using the species-specific primers AbF5/AbR5 (Devran et al., 2017), and three non-*A. besseyi* (AF, AS, and MI) were used as templates to evaluate the specificity evaluation of the RPA-Cas12a-Ab system (Table 1,




Supplementary Figure S2). RPA results show that the 237bp DNA fragment could only be amplified from *A. besseyi* samples AB1 and AB2, and not from AF, AS, and MI samples, as well as the rice gDNA and negative control (**Figure 4A**). Furthermore,



using RPA-Cas12a-Ab fluorescence assays, AB1 and AB2 were found to have positive readings with significant fluorescence signals comparable with the plasmid positive control and did not cross-react with three non-*A. besseyi* (AF, AS, and MI) samples (**Figure 4B**). These results demonstrated the specificity of the RPA-Cas12a-Ab fluorescence assay.

Evaluation of the Sensitivity of the RPA-Cas12a-Ab Fluorescence Assay

The sensitivity of the RPA-Cas12a-Ab assay was evaluated and compared with that of PCR and real-time qPCR using a serial 10-fold dilution of plasmid pUC57-18S. The limit of detection (LOD) of PCR and real-time qPCR was 10^3 and 10^1 copies/ μ l, respectively (**Figures 5A,B**), while the LOD of the RPA-Cas12a-Ab assay reached 1 copy/ μ l target DNA (**Figure 5C**). The RPA-



Cas12a-Ab assay is thus more sensitive than PCR and qPCR in *A. besseyi* detection.

Furthermore, we used a serial 10-fold dilution of individual *A. besseyi* gDNA to assess the sensitivity of the RPA-Cas12a-Ab fluorescence assay. As shown in **Figure 5D**, $1:10^7$ dilution of individual *A. besseyi* gDNA was still able to generate a significant fluorescence signal as compared with the control using the RPA-Cas12a-Ab fluorescence assay (**Figure 5D**). The concentration of *A. besseyi* gDNA extracted from individual nematodes is about 9.6 ng/ μ l. Thus, the LOD of the RPA-Cas12a-Ab fluorescence assay for detection of *A. besseyi* gDNA is as low as 9.6 fg/ μ l.


Establishing the RPA-Cas12a-Ab Lateral Flow Strip Assay

For on-site detection, we combined the RPA-Cas12a with LFA technology to establish the RPA-Cas12a-Ab LFA assay (**Figure 1**). Five nematode samples (AB1, AB2, AF, AS, and MI, **Table 1**) were tested with the RPA-Cas12a-Ab LFA assay, of which *A. besseyi* samples (AB1 and AB2) showed a clear test band like that of the positive control, and no cross reactions were observed for non-*A. besseyi* samples (AF, AS, and MI). These results confirmed the specificity of the RPA-Cas12a-Ab LFA assay (**Figure 6A**).

A serial 10-fold dilution of reference plasmid pUC57-18S was used to evaluate the sensitivity of the RPA-Cas12a-Ab LFA assay. The LOD of the RPA-Cas12a-Ab LFA assay was 10^3 copies/ μ l (**Figure 6B**), which was similar to that of the PCR assay (**Figure 5A**). We then tested the RPA-Cas12a-Ab LFA system in terms of the detection of *A. besseyi* samples. The initial gDNA extracted individual *A. besseyi* was about 9.6 ng/ μ l, and then the DNA was serially diluted from $1/10^0$ to $1/10^6$. The results showed that $1/10^0$ and $1/10^1$ dilutions developed a clear positive test band (**Figure 6C**), implying that the detection limit of the RPA-Cas12a-Ab LFA assay was 0.96 ng/ μ l gDNA of *A. besseyi*.

DISCUSSION

RWTD is a typical seed-borne disease in rice (Devran and Göknur, 2020), which is caused by the plant parasitic nematode *A. besseyi* (Jones et al., 2013). *A. besseyi* is only 0.5–1 mm in length and composed of 1000–2000 cells. It has been reported that about 92% of the rice grains had nematodes in the panicles that were infected by *A. besseyi*, and the numbers of nematodes were up to 2014 nematodes/100 grains (Lin et al.,



2005). *A. besseyi* gathered inside of the glume axis of maturing grains and slowly dehydrated along with kernel drying. They become dormant and are able to survive for 8 months to 3 years during seed storage (Huang and Huang, 1974). After seed sowing, dormant *A. besseyi* rapidly reacts and embarks on a new life cycle. Therefore, accurate and rapid detection of *A. besseyi* is crucial to facilitating the quarantine of rice seeds and stopping the spread of RWTD. Due to the remarkably constrained morphology of nematodes from the same genus, molecular methods such as PCR or qPCR, have been proved helpful in detecting *A. besseyi*. However, most current PCR-based detection assays have a trade-off in performance metrics such as sensitivity and specificity and over-reliance on well-established laboratories with sophisticated facilities or well-trained operators (Li Y. et al., 2019).

In this study, we developed a new assay, the RPA-Cas12a-Ab assay, by combining RPA with CRISPR/Cas12a to detect *A. besseyi* (Figure 1). There are advantages for using this CRISPR/Cas12a-based new assay. First, unlike conventional assays, RPA-Cas12a-Ab does not rely on sophisticated equipment. Both RPA amplification and CRISPR/Cas12a reaction could be completed at body temperature, and only one fluorescence analyzer is needed to complete rapid detection. For on-site detection, we couple the RPA-Cas12a-Ab assay with LFA, allowing for readout of results without any equipment at room temperature. Second, in contrast to the time-consuming nature of conventional qPCR or PCR-based detection assays, which require 1–3 h, the newly developed RPA-Cas12a-Ab assay takes only 45 min, significantly reducing the amount of time it takes to obtain the detection results. Finally, the RPA-Cas12a-Ab fluorescence detection assay is shown to have higher detection sensitivity than conventional assays. This assay had an LOD of 1 copy/ μ l,

which is significantly more sensitive than that of PCR and qPCR assays (Figures 5A–C). Even though the adoption of the LFA technique to readout the RPA-Cas12a-Ab detection result will increase the LOD to 10^3 copies/ μ l (Figure 6B), this new assay is still robust enough to detect the presence of *A. besseyi* with high accuracy and specificity within 45 min (Figures 6A,C).

The crRNA, which plays a role in guiding Cas12a to recognize the target DNA, is critical for Cas12a-based genome editing or molecular detection. Strict selection of the target of crRNA by avoiding the homologous sites in the genome could minimize the “off-target” in genome editing (Kadam et al., 2018). Similarly, avoiding the conserved sequence in crRNA target selection is crucial for alleviating the false positive of Cas12a-based molecular detection. In this study, the ITS region sequence of the *A. besseyi* 18S rRNA gene was selected as the target sequence in RPA-Cas12a-Ab assays. The ITS region evolves much faster compared to the regions of the rRNA gene and has been widely applied as a diagnostic marker at the species level in nematodes (Powers, 2004). The specificity of the RPA-Cas12a-Ab assays was validated using three closely related non-*A. besseyi* species (Figures 4, 6A). The results showed that the assay accurately detected *A. besseyi* DNA without false-positive signals from other nematodes, confirming the specificity of RPA-Cas12a-Ab assays for *A. besseyi*.

In conclusion, we developed a new *A. besseyi* molecular detection assay that combines CRISPR/Cas12a with RPA preamplification, with readout using a fluorescence analyzer or LFA. This new assay could be completed within 45 min with no need for complicated instruments and skilled operators. The LOD of the new RPA-Cas12a-Ab assay could reach 1 copy/ μ l for the fluorescence detection assay and 10^3 copies/ μ l for the LFA

detection assay. Thus, the new RPA-Cas12a-Ab is a promising molecular method for *A. besseyi* detection with high accuracy, sensitivity, and specificity.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**; further inquiries can be directed to the corresponding authors.

AUTHOR CONTRIBUTIONS

AZ and BS conceptualized, designed the research, and performed the experiments. JZ, CC, and JZ, isolated nematode samples. FN and ZL performed data curation. LY and CY isolated *A. Subtenuis* and *A. fragariae* samples. YD, KX, QH, and YQ investigated and validated data. LC and HC analyzed data and wrote the manuscripts. All authors have read and approved the final manuscript.

REFERENCES

- Bai, J., Lin, H., Li, H., Zhou, Y., Liu, J., Zhong, G., et al. (2019). Cas12a-based On-Site and Rapid Nucleic Acid Detection of African Swine Fever. *Front. Microbiol.* 10, 2830. doi:10.3389/fmicb.2019.02830
- Bai, Z., Qin, M., Zhao, L., Han, Y., Wang, D., Xu, C., et al. (2017). Loop-mediated Isothermal Amplification Assay for Rapid Diagnosis of *Aphelenchoides Besseyi*. *Chin. J. Rice Sci.* 31, 432–440. doi:10.16819/j.1001-7216.2017.7001
- Borges, A., Rosa, M. S., Recchia, G. H., de Queiroz-Silva, J. R., Bressan, E. d. A., and Veasey, E. A. (2009). Ctab Methods for Dna Extraction of Sweetpotato for Microsatellite Analysis. *Sci. Agric. (Piracicaba, Braz.)* 66, 529–534. doi:10.1590/S0103-90162009000400015
- Buonincontro, D. S., Roberts, D. M., Oliveira, C. M. G., Blok, V. C., Neilson, R., and Oliveira, R. D. L. (2018). A Rapid Diagnostic for Detection of *Aphelenchoides Besseyi* and *a. Fujianensis* Based on Real-Time PCR. *Plant Dis.* 102, 519–526. doi:10.1094/PDIS-08-17-1160-RE
- Çelik, E. S., and Devran, Z. (2019). Identification and Quantification of *Aphelenchoides Besseyi* from Rice Using Qpcr. *Eur. J. Plant Pathol.* 154, 691–703. doi:10.1007/s10658-019-01692-4
- Chaijarasphong, T., Thammachai, T., Itsathitphaisarn, O., Sritunyalucksana, K., and Suebsing, R. (2019). Potential Application of Crispr-Cas12a Fluorescence Assay Coupled with Rapid Nucleic Acid Amplification for Detection of White Spot Syndrome Virus in Shrimp. *Aquaculture* 512, 734340. doi:10.1016/j.aquaculture.2019.734340
- Chang, Y., Deng, Y., Li, T., Wang, J., Wang, T., Tan, F., et al. (2020). Visual Detection of Porcine Reproductive and Respiratory Syndrome Virus Using CRISPR-Cas13a. *Transbound. Emerg. Dis.* 67, 564–571. doi:10.1111/tbed.13368
- Chen, J. S., Ma, E., Harrington, L. B., Da Costa, M., Tian, X., Palefsky, J. M., et al. (2018). Crispr-cas12a Target Binding Unleashes Indiscriminate Single-Stranded Dnase Activity. *Science* 360, 436–439. doi:10.1126/science.aar6245
- Cheng, X., Xiang, Y., Xie, H., Xu, C.-L., Xie, T.-F., Zhang, C., et al. (2013). Molecular Characterization and Functions of Fatty Acid and Retinoid Binding Protein Gene (Ab-far-1) in *Aphelenchoides Besseyi*. *Plos One* 8, e66011. doi:10.1371/journal.pone.0066011
- Creutzburg, S. C. A., Wu, W. Y., Mohanraju, P., Swartjes, T., Alkan, F., Gorodkin, J., et al. (2020). Good Guide, Bad Guide: Spacer Sequence-dependent Cleavage Efficiency of Cas12a. *Nucleic Acids Res.* 48, 3228–3243. doi:10.1093/nar/gkz1240

FUNDING

This work was supported by the Shanghai Sailing Program (grant number: 21YF1441000), the Agriculture Research System of Shanghai, China (Grant No. 2022003), and the SAAS Program for Excellent Research Team (grant number: 2022-A006).

ACKNOWLEDGMENTS

We would like to thank Dr. Jing-han Hong for valuable critiques and comments on the manuscript and Dr. Xuan Wang (Nanjing Agricultural University) for kindly providing *A. besseyi* and *M. incognita* samples.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fbioe.2022.912959/full#supplementary-material>

- Devran, Z., and Göknur, A. (2020). Development and Validation of a SNP-Based KASP Assay for Rapid Identification of *Aphelenchoides Besseyi* Christie, 1942. *Crop Prot.* 136, 105235. doi:10.1016/j.cropro.2020.105235
- Devran, Z., Tülek, A., Mıstanoğlu, İ., Çiftçigil, T. H., and Özalp, T. (2017). A Rapid Molecular Detection Method for *Aphelenchoides Besseyi* from Rice Tissues. *Australas. Plant Pathol.* 46, 43–48. doi:10.1007/s13313-016-0452-1
- Duncan, L. W., and Moens, M. (2006). “Migratory Endoparasitic Nematodes,” in *Plant Nematology: CAB International Publishing, Wallingford* Editor R. N. Perry and M. Moens, 123–152. doi:10.1079/9781845930561.0123
- Gootenberg, J. S., Abudayyeh, O. O., Kellner, M. J., Joung, J., Collins, J. J., and Zhang, F. (2018). Multiplexed and Portable Nucleic Acid Detection Platform with Cas13, Cas12a, and Csm6. *Science* 360, 439–444. doi:10.1126/science.aaq0179
- Huang, C.-H., Lee, K.-C., and Doudna, J. A. (2018). Applications of Crispr-Cas Enzymes in Cancer Therapeutics and Detection. *Trends Cancer* 4, 499–512. doi:10.1016/j.trecan.2018.05.006
- Huang, C. S., and Huang, S. P. (1974). Dehydration and the Survival of Rice White Tip Nematode, *Aphelenchoides Besseyi*1. *Nematol* 20, 9–18. doi:10.1163/187529274x00528
- Ibrahim, S. K., Perry, R. N., Burrows, P. R., and Hooper, D. J. (1994). Infectivity and Population Dynamics of the Rice Stem Nematode, *Ditylenchus Angustus*. *Nematol* 40, 412–422. doi:10.1163/003525994x00292
- Jones, J. T., Haegeman, A., Danchin, E. G. J., Gaur, H. S., Helder, J., Jones, M. G. K., et al. (2013). Top 10 Plant-Parasitic Nematodes in Molecular Plant Pathology. *Mol. Plant Pathol.* 14, 946–961. doi:10.1111/mpp.12057
- Kadam, U. S., Shelake, R. M., Chavhan, R. L., and Suprasanna, P. (2018). Concerns Regarding ‘off-Target’ Activity of Genome Editing Endonucleases. *Plant Physiology Biochem.* 131, 22–30. doi:10.1016/j.plaphy.2018.03.027
- Li, L., Li, S., Wu, N., Wu, J., Wang, G., Zhao, G., et al. (2019). Holmesv2: a Crispr-Cas12b-Assisted Platform for Nucleic Acid Detection and Dna Methylation Quantitation. *ACS Synth. Biol.* 8, 2228–2237. doi:10.1021/acssynbio.9b00209
- Li, S.-Y., Cheng, Q.-X., Liu, J.-K., Nie, X.-Q., Zhao, G.-P., and Wang, J. (2018a). Crispr-cas12a Has Both Cis- and Trans-cleavage Activities on Single-Stranded Dna. *Cell Res.* 28, 491–493. doi:10.1038/s41422-018-0022-x
- Li, S.-Y., Cheng, Q.-X., Wang, J.-M., Li, X.-Y., Zhang, Z.-L., Gao, S., et al. (2018b). Crispr-cas12a-assisted Nucleic Acid Detection. *Cell Discov.* 4, 20. doi:10.1038/s41421-018-0028-z
- Li, Y., Li, S., Wang, J., and Liu, G. (2019). Crispr/cas Systems towards Next-Generation Biosensing. *Trends Biotechnol.* 37, 730–743. doi:10.1016/j.tibtech.2018.12.005

- Lin, M., Ding, X., Wang, Z., Zhou, F., and Lin, N. (2005). Description of *Aphelenchoides besseyi* from Abnormal Rice with Small Grains and Erect Panicles in China. *Rice Sci.* 12, 289–294. doi:10.16819/j.1001-7216.2005.04.014
- Liu, H., Wang, J., Zeng, H., Liu, X., Jiang, W., Wang, Y., et al. (2021). Rpa-cas12a-fs: a Frontline Nucleic Acid Rapid Detection System for Food Safety Based on Crispr-Cas12a Combined with Recombinase Polymerase Amplification. *Food Chem.* 334, 127608. doi:10.1016/j.foodchem.2020.127608
- Ma, Q.-N., Wang, M., Zheng, L.-B., Lin, Z.-Q., Ehsan, M., Xiao, X.-X., et al. (2021). Raa-cas12a-tg: a Nucleic Acid Detection System for *Toxoplasma gondii* Based on Crispr-Cas12a Combined with Recombinase-Aided Amplification (Raa). *Microorganisms* 9, 1644. doi:10.3390/microorganisms9081644
- Powers, T. (2004). Nematode Molecular Diagnostics: from Bands to Barcodes. *Annu. Rev. Phytopathol.* 42, 367–383. doi:10.1146/annurev.phyto.42.040803.140348
- Rybarczyk-Mydlowska, K., Mooyman, P., van Megen, H., van den Elsen, S., Vervoort, M., Veenhuizen, P., et al. (2012). Small Subunit Ribosomal Dna-Based Phylogenetic Analysis of Foliar Nematodes (*Aphelenchoides* spp.) and Their Quantitative Detection in Complex Dna Backgrounds. *Phytopathology* 102, 1153–1160. doi:10.1094/PHYTO-05-12-0114-R
- Sánchez-monge, A., Flores, L., Salazar, L., Hockland, S., and Bert, W. (2015). An Updated List of the Plants Associated with Plant-Parasitic *Aphelenchoides* (Nematoda: Aphelenchoididae) and its Implications for Plant-Parasitism within This Genus. *Zootaxa* 4013, 207–224. doi:10.11646/zootaxa.4013.2.3
- Sánchez-Monge, A., Janssen, T., Fang, Y., Couvreur, M., Karssen, G., and Bert, W. (2017). Mtcoi Successfully Diagnoses the Four Main Plant-Parasitic *Aphelenchoides* Species (Nematoda: Aphelenchoididae) and Supports a Multiple Origin of Plant-Parasitism in This Paraphyletic Genus. *Eur. J. Plant Pathol.* 148, 853–866. doi:10.1007/s10658-016-1141-1
- Tsou, J.-H., Leng, Q., and Jiang, F. (2019). A Crispr Test for Detection of Circulating Nuclei Acids. *Transl. Oncol.* 12, 1566–1573. doi:10.1016/j.tranon.2019.08.011
- Wang, X., Xue, B., Dai, J., Qin, X., Liu, L., Chi, Y., et al. (2018). A Novel Meloidogyne Incognita Chorismate Mutase Effector Suppresses Plant Immunity by Manipulating the Salicylic Acid Pathway and Functions Mainly during the Early Stages of Nematode Parasitism. *Plant Pathol.* 67, 1436–1448. doi:10.1111/ppa.12841
- Xiao, G., Liu, S., Liu, H., He, X., Zhang, S., Liang, Z., et al. (2020). Crispr/cas12a-based Biosensing Platform for Precise and Efficient Screening of Crispr/cas9-Induced Biallelic Mutants. *Talanta* 210, 120613. doi:10.1016/j.talanta.2019.120613
- Yang, J.-L., and Yu, G.-Y. (2019). A Loop-Mediated Isothermal Amplification Assay for the Plant-Parasitic Nematode *Aphelenchoides besseyi* in Rice Seedlings. *J. Nematology* 51, 1–11. doi:10.21307/jofnem-2019-080
- Zetsche, B., Gootenberg, J. S., Abudayyeh, O. O., Slaymaker, I. M., Makarova, K. S., Essletzbichler, P., et al. (2015). Cpf1 Is a Single Rna-Guided Endonuclease of a Class 2 Crispr-Cas System. *Cell* 163, 759–771. doi:10.1016/j.cell.2015.09.038
- Zhang, Y.-m., Zhang, Y., and Xie, K. (2020). Evaluation of Crispr/cas12a-Based Dna Detection for Fast Pathogen Diagnosis and Gmo Test in Rice. *Mol. Breed.* 40, 11. doi:10.1007/s11032-019-1092-2

Conflict of Interest: Author ZL was employed by the company Shanghai Agricultural Science and Technology Seed Co., Ltd.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Zhang, Sun, Zhang, Cheng, Zhou, Niu, Luo, Yu, Yu, Dai, Xie, Hu, Qiu, Cao and Chu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.