



OPEN ACCESS

EDITED AND REVIEWED BY
Hongcheng Sun,
Hangzhou Normal University, China

*CORRESPONDENCE

Ping Ru,
✉ ruping_rp@163.com
Xinghui Liu,
✉ sylvixh@163.com
Tianzhao Xu,
✉ tianzhaoxufan@163.com

[†]These authors have contributed equally to this work

RECEIVED 07 April 2023

ACCEPTED 13 April 2023

PUBLISHED 20 April 2023

CITATION

Peng T, Liu Q, Song H, Zhang C, Wang X, Ru P, Xu T and Liu X (2023), Corrigendum: Biodegradable hollowed mesoporous SeO₂ nanoplatform loaded with indocyanine green for simultaneous NIR II fluorescence imaging and synergistic breast carcinoma therapy. *Front. Bioeng. Biotechnol.* 11:1201751. doi: 10.3389/fbioe.2023.1201751

COPYRIGHT

© 2023 Peng, Liu, Song, Zhang, Wang, Ru, Xu and Liu. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Biodegradable hollowed mesoporous SeO₂ nanoplatform loaded with indocyanine green for simultaneous NIR II fluorescence imaging and synergistic breast carcinoma therapy

Tingwei Peng^{1†}, Qing Liu^{2†}, Hui Song^{2†}, Conghui Zhang²,
Xue Wang², Ping Ru^{3*}, Tianzhao Xu^{4*} and Xinghui Liu^{2*}

¹Shanghai Gongli Hospital, Ningxia Medical University, Shanghai, China, ²Department of Clinical Laboratory, Shanghai Gongli Hospital, The Second Military Medical University, Shanghai, China, ³Department of Obstetrics, Shanghai East Hospital, School of Medicine, Tongji University, Shanghai, China, ⁴Hospital Department, Shanghai University of Medicine and Health Sciences Affiliated to Zhoupu Hospital, Shanghai, China

KEYWORDS

hollowed mesoporous SeO₂, ICG precise delivery, NIR II fluorescent imaging, photothermal therapy, ROS mediated oxidative therapy

A Corrigendum on

Biodegradable hollowed mesoporous SeO₂ nanoplatform loaded with indocyanine green for simultaneous NIR II fluorescence imaging and synergistic breast carcinoma therapy

by Peng T, Liu Q, Song H, Zhang C, Wang X, Ru P, Xu T and Liu X (2023), *Front. Bioeng. Biotechnol.* 11:1151148. doi: 10.3389/fbioe.2023.1151148

In the published article, there was an error in [Figure 6A](#) as published. The photograph of nude mouse is wrong in hmSeO₂@ICG-RGD group. The corrected [Figure 6](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 6

(A) Representative photographs of breast carcinoma-bearing BALB/c nude mice 0, 7 and 15 days after various treatments. (B) Representative resected tumors from mice in the various groups 15 days after treatment administration. (C) Tumor volumes in mice following various treatments. (D) Tumor weights after 15 days of treatment. ****p < 0.01.** (E) Body weights of mice in each treatment group. (F) H&E and TUNEL-staining photographs of tumor tissues resected from subcutaneous tumor-bearing mice after various treatments for 7 days.