



OPEN ACCESS

EDITED BY

Mattia Galli,
Agostino Gemelli University Polyclinic
(IRCCS), Italy

REVIEWED BY

Plinio Cirillo,
University of Naples Federico II, Italy
Salvatore Giordano,
Magna Graecia University, Italy

*CORRESPONDENCE

José Luis Ferreiro
jferreiro@bellvitgehospital.cat

SPECIALTY SECTION

This article was submitted to
Coronary Artery Disease,
a section of the journal
Frontiers in Cardiovascular Medicine

RECEIVED 29 September 2022

ACCEPTED 03 November 2022

PUBLISHED 22 November 2022


CITATION

Marcano AL, Gracida M, Roura G,
Gomez-Lara J, Romaguera R,
Teruel L, Fuentes L, Muntané-Carol G,
Meroño O, Sosa SG,
Gómez-Hospital JA, Comin-Colet J
and Ferreiro JL (2022) Antiplatelet
efficacy of ticagrelor versus
clopidogrel in Mediterranean patients
with diabetes mellitus and chronic
coronary syndromes: A crossover
pharmacodynamic investigation.
Front. Cardiovasc. Med. 9:1057331.
doi: 10.3389/fcvm.2022.1057331

COPYRIGHT

© 2022 Marcano, Gracida, Roura,
Gomez-Lara, Romaguera, Teruel,
Fuentes, Muntané-Carol, Meroño,
Sosa, Gómez-Hospital, Comin-Colet
and Ferreiro. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Antiplatelet efficacy of ticagrelor versus clopidogrel in Mediterranean patients with diabetes mellitus and chronic coronary syndromes: A crossover pharmacodynamic investigation

Ana Lucrecia Marcano^{1,2}, Montserrat Gracida^{1,2},
Gerard Roura^{1,2}, Josep Gomez-Lara^{1,2}, Rafael Romaguera^{1,2},
Luis Teruel^{1,2}, Lara Fuentes^{1,2}, Guillem Muntané-Carol^{1,2},
Oona Meroño^{1,2,3}, Silvia Gabriela Sosa^{1,2},
Joan Antoni Gómez-Hospital^{1,2}, Josep Comin-Colet^{2,4} and
José Luis Ferreiro  2,4*

¹Department of Cardiology, Bellvitge University Hospital, L'Hospitalet de Llobregat, Barcelona, Spain, ²Bio-Heart Cardiovascular Diseases Research Group, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain, ³Faculty of Medicine, University of Vic-Central University of Catalonia, Barcelona, Spain, ⁴Department of Cardiology, Bellvitge University Hospital, Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares, L'Hospitalet de Llobregat, Barcelona, Spain

Introduction: Patients with diabetes mellitus (DM) have augmented platelet reactivity and diminished responsiveness to clopidogrel. Ticagrelor, a more potent P2Y₁₂ inhibitor, is clinically superior to clopidogrel in acute coronary syndromes, although its role in chronic coronary syndromes (CCS) is still the subject of debate. The aim of this investigation was to compare the pharmacodynamic effectiveness of ticagrelor and clopidogrel in Mediterranean DM patients with CCS.

Materials and methods: In this prospective, randomized, crossover study, patients ($n = 20$) were randomized (1:1) to receive, on top of aspirin therapy, either ticagrelor 180 mg loading dose (LD)/90 mg maintenance dose (MD) b.i.d. or clopidogrel 600 mg LD/75 mg MD o.d. for 1 week in a crossover fashion with a 2–4 week washout period between regimens. Platelet function measurements were performed at 4 timepoints in each period (baseline, 2 h and 24 h after LD, and 1 week), including light transmission aggregometry (LTA, primary endpoint), VASP assay, Multiplate and VerifyNow P2Y₁₂.

Results: The ticagrelor LD achieved greater platelet inhibitory effect than clopidogrel LD, assessed with LTA (20 μ M ADP as agonist), at 2 h ($34.9 \pm 3.9\%$ vs. $63.6 \pm 3.9\%$; $p < 0.001$) and 24 h ($39.4 \pm 3.5\%$ vs. $52.3 \pm 3.8\%$; $p = 0.014$).

After 1 week of therapy, platelet reactivity was again significantly inferior with ticagrelor compared to clopidogrel ($30.7 \pm 3.0\%$ vs. $54.3 \pm 3.0\%$; $p < 0.001$). The results were consistent with the other platelet function assays employed.

Conclusion: In Mediterranean patients with DM and CCS, ticagrelor provides a more potent antiplatelet effect than clopidogrel after the LD and during the maintenance phase of therapy.

Clinical trial registration: [ClinicalTrials.gov], identifier [NCT02457130].

KEYWORDS

ticagrelor, chronic coronary syndrome, antiplatelet therapy, high platelet reactivity, diabetes mellitus

Introduction

Subjects with diabetes mellitus (DM) have a higher risk of developing cardiovascular disease and experiencing atherothrombotic events, which have poorer prognosis than those occurring in patients without DM (1). One of the factors involved in the augmented atherothrombotic risk of DM patients with coronary artery disease (CAD) is a hyper-reactive platelet phenotype, which contributes to an impaired responsiveness to antiplatelet drugs, mainly to clopidogrel (2, 3). Therefore, the augmented ischemic risk among DM patients with CAD clearly emphasizes the need to optimize platelet inhibition in this population with the goal of ameliorating clinical outcomes (4).

The use of more potent and less variable P2Y₁₂ receptor antagonists such as prasugrel or ticagrelor has demonstrated a reduction in adverse ischemic events when compared to clopidogrel in patients suffering an acute coronary syndrome (ACS) (5, 6). However, the observed clinical superiority of ticagrelor or prasugrel over clopidogrel in ACS patients has not been replicated in patients with stable CAD or undergoing elective percutaneous coronary intervention (PCI) (7, 8). In fact, clopidogrel is still widely used in real-life clinical practice as part of dual antiplatelet therapy (DAPT), e.g., in patients undergoing elective PCI or in those with stabilized symptoms after an ACS following a strategy of DAPT de-escalation. It is well established that clopidogrel has a large interindividual variability in response with genetic factors, such as polymorphisms of cytochrome P450 (CYP) isoforms (mainly CYP2C19), playing a key role in this phenomenon (9, 10). Evidently, the prevalence of genetic polymorphisms may vary greatly among races and, therefore, it is relevant that pharmacodynamic (PD) investigations take into consideration ethnicity when evaluating antiplatelet agents.

Since the evidence regarding the PD effectiveness of clopidogrel compared to ticagrelor in DM patients with a chronic coronary syndrome (CCS) is relatively scarce (11, 12), we designed the Comparison of Ticagrelor and clopidogrel



in patients with Coronary artery disease and type 2 Diabetes Mellitus (TICS-DM) study, with the aim of assessing the platelet inhibitory effects of these two P2Y₁₂ inhibitors in a Mediterranean population with a comprehensive panel of platelet function assays.

Materials and methods

Subject population and study design

This was a prospective, open-label, two-sequence, two-period, randomized, crossover study conducted in Mediterranean (Spanish nationality) type 2 DM patients with 18–75 years of age and known stable CAD (angiographically documented) on a background of aspirin therapy (NCT 02457130). The World Health Organization criteria were used to define DM status. Exclusion criteria included: known allergies to clopidogrel or ticagrelor, blood dyscrasia or bleeding diathesis, any recent acute coronary event (<1 year), hemodynamic instability, recent treatment with any other antiplatelet agent (<14 days) with the exception of aspirin, oral anticoagulation with a coumarin derivative, any active bleeding or malignancy, history of stroke (<6 months prior to inclusion) or any intracranial bleeding, platelet count $<100 \times 10^6/\mu\text{l}$, severe chronic kidney disease (creatinine clearance $<30 \text{ ml/min}$) and pregnant females.

Subjects were randomized in a 1:1 fashion to ticagrelor [180-mg loading dose (LD) followed by 90-mg maintenance dose] or clopidogrel (600-mg loading dose followed by 75-mg daily maintenance dose) for 1 week (Figure 1). All patients were on chronic aspirin therapy (100 mg o.d.), that was maintained at the same dose throughout the study. Patients crossed-over treatment regimen after a 2 to 4-week washout period. Blood sampling for platelet function measurements were performed at the two phases of the study at the following timepoints: (1) baseline, (2) 2 h after LD, (3) 24 h after LD, and (4) 7 days (in the morning, with last dose of study drug administered the



previous day). The washout periods were included in order to minimize carryover effects. A follow-up visit was performed at least 2 weeks after the last dose of the study drug to verify the absence of adverse events. Patient compliance was assessed by pill counting and interview.

The study was performed in compliance with the Declaration of Helsinki and was approved by the institutional Ethics Committee. All subjects included provided written informed consent.

Sample collection and platelet function assays

Blood samples for platelet function assessment were collected at the scheduled time points from an antecubital vein; the first 2–4 ml of blood were discarded in order to avoid spontaneous platelet activation. Samples were processed by trained laboratory personnel (blinded to allocated treatment). Platelet function tests (PFT) included light

transmission aggregometry (LTA), flow cytometric analysis of the phosphorylation status of the vasodilator-stimulated phosphoprotein (VASP), multiple electrode aggregometry (MEA) and VerifyNow P2Y₁₂ (VN-P2Y₁₂) assay.

Light transmission aggregometry

Light transmission aggregometry (a schematic example is shown in **Figure 2**) was performed according to standard protocols (13). Briefly, platelet aggregation was assessed using platelet-rich plasma (PRP) and platelet-poor plasma (PPP) by the turbidometric method in a two-channel aggregometer (Chrono-Log 490 Model, Chrono-Log Corp., Havertown, PA, USA). PRP was obtained as a supernatant after centrifugation of citrated blood at 100 g for 10 min and PPP was obtained by a second centrifugation of the blood fraction at 1500 g for 15 min. Light transmission was adjusted to 0% for PRP and to 100% for PPP for each measurement. Maximal platelet aggregation (MPA) was stimulated by 20 and 5 μmol/L adenosine diphosphate (ADP) as agonists. High on-treatment platelet reactivity (HPR) was defined as MPA > 64.5% and >42.9% with ADP 20 and 5 μmol/L, respectively (14).

Vasodilator-stimulated phosphoprotein assay

Vasodilator-stimulated phosphoprotein-phosphorylation (VASP-P) is a marker of the P2Y₁₂ receptor reactivity and, therefore, P2Y₁₂ inhibitors-induced inhibition. VASP was assessed according to standard protocols (15). Adding ADP to PGE₁-stimulated platelets diminishes PGE₁-induced VASP-P levels. If P2Y₁₂ receptors are successfully inhibited, the addition of ADP will not decrease the PGE₁-stimulated VASP-P levels. VASP-P levels were quantified with labeled monoclonal antibodies by flow cytometry with the Platelet VASP-FCM kit (Biocytex Inc., Marseille, France). The platelet reactivity index (PRI) was calculated once measured the VASP-P levels after stimulation with PGE₁ (MFI PGE₁) and also PGE₁ + ADP (MFI PGE₁ + ADP) with the following formula: $PRI = ([MFI\ PGE_1] - [MFI\ PGE_1 + ADP]) / [MFI\ PGE_1] \times 100\%$. A reduced PRI indicates a greater inhibition of the P2Y₁₂ signaling pathway, and a cut-off point of ≥50% PRI was utilized to define low responsiveness (16).

Multiple electrode aggregometry

Multiple electrode aggregometry (MEA) was assessed in whole blood with the Multiplate analyzer (Roche Diagnostics, Basel, Switzerland), which measures the change in impedance caused by platelets adhesion onto silver-covered electrodes working as sensor units (17). Curves were recorded for 6 min and platelet aggregation was determined as area under the curve of arbitrary aggregation units (AU*min) using 6.4 μmol/L ADP as agonist. The cut-off value used to define HPR was >468 AU*min (16).

VerifyNow P2Y₁₂ assay

The VerifyNow System is a turbidimetric based optical detection system which measures platelet induced aggregation as an increase in light transmittance (Accumetrics, San Diego, CA, USA) and was utilized according to manufacturer's instructions (18). The VerifyNow P2Y₁₂ Assay measures changes in platelet function specific to P2Y₁₂ inhibition by combining ADP + PGE₁ stimuli. The reagents are incorporated into the assay channel to induce platelet activation and light transmittance increases as activated platelets bind and aggregate fibrinogen-coated beads. The device then measures this change in optical signal and reports results in P2Y₁₂ Reaction Units (PRU). A cut-off point of >208 PRUs was used to define HPR (16).

Study endpoints and sample size calculation

The primary endpoint of the present study was the comparison of MPA measured with LTA (20 μM ADP as agonist) and achieved after 1 week of therapy with ticagrelor or clopidogrel using the treatment regimens described above. An initial sample size of 30 patients was planned, but a mid-course recalculation of the sample size due to an overestimation of the standard deviation was performed and specified in an amendment to the protocol. The revised calculation of the sample size was as follows: assuming a standard deviation of MPA of 13 (19, 20), a difference between treatment groups of 10 with 90% power and 2-sided alpha = 0.05 will be detected with 18 completed subjects per regimen group. Randomization of a total of 25 subjects was allowed, considering an approximate dropout of 25%, in order to ensure that complete data from 18 subjects would be available for analysis.

Other secondary end points included: (a) evaluation of platelet reactivity between clopidogrel and ticagrelor with all the PFT after 1 week of treatment; (b) comparison of the 2 treatment regimens at 2 and 24 h after LD with all the PFT; and (c) determination at the different time points assessed of the proportion of patients with HPR (measured with all tests).

Statistical analysis

Baseline continuous variables are expressed as mean ± SD, while categorical variables are reported as frequencies and percentages. Only those subjects who successfully completed the two treatment periods were considered for analysis. All statistical comparisons of platelet reactivity for the primary and secondary endpoints were performed using linear mixed-effects models with treatment, sequence, period, and treatment*period (treatment by period interaction to test for carryover effects) as fixed effects, subject as a random effect, and the baseline

value of each corresponding platelet function test (MPA, PRI, AU*min, or PRU) as a covariate. Results are reported as least-squares mean (LSM) ± standard error of the mean (SEM). Comparisons between HPR rates were conducted using the McNemar test or the binomial exact test. All the analyses performed were evaluated with a 2-tailed probability value <0.05 to indicate a statistically significant difference. Statistical analysis was performed using SPSS version 18.0 software (SPSS Inc., Chicago, IL, USA).

Results

Among 580 patients screened for eligibility, 95 met inclusion and exclusion criteria. Of these, 25 patients agreed to participate and were randomized to start with ticagrelor (*n* = 13) or clopidogrel (*n* = 12). Following randomization, four patients withdrew consent and one patient discontinued ticagrelor treatment due to side effects (dyspnea). Therefore, 20 patients successfully completed the two periods of the study and were included in the analysis. The flow chart of the study is illustrated in **Figure 1**, whereas baseline demographics and clinical variables are reported in **Table 1**. No significant dissimilarities were found between patients that initiated with either ticagrelor or clopidogrel. Among patients that completed the two phases of the study, 4 (20%) developed mild and transient dyspnea on ticagrelor therapy whereas no patient on clopidogrel therapy developed dyspnea. No patient experienced any ischemic or bleeding event during the study.

Pharmacodynamic effects of ticagrelor vs. clopidogrel

At baseline, there were no statistical differences between the two regimens studied. After 1 week of treatment, MPA (using 20 μM ADP as agonist, the primary endpoint of the present investigation) was significantly lower (**Figure 3**) with ticagrelor compared to clopidogrel (MPA: 30.7 ± 3.0% vs. 54.3 ± 3.0%; *p* < 0.001). When assessing the PD efficacy of the LD, ticagrelor also provided greater platelet inhibition than clopidogrel both at 2 h (MPA: 34.9 ± 3.9% vs. 63.6 ± 3.9; *p* < 0.001) and 24 h (MPA: 39.4 ± 3.5% vs. 52.3 ± 3.8%; *p* = 0.014), as shown in **Figure 3**. No statistically significant differences were found by sequence, period, or the treatment-by-period interaction, which suggest no carryover effect. Similar findings were observed with 5 μM ADP and the other platelet function tests employed, showing greater inhibition of platelet aggregation at 2 h, 24 h, and 1 week in the ticagrelor group compared with the clopidogrel group (**Figure 4**). Of note, no differences in clopidogrel- or ticagrelor- mediated platelet inhibition were found when comparing patients with or without insulin therapy (data not shown).

TABLE 1 Baseline characteristics.

	<i>n</i> = 20
Age, mean ± SD	65.45 ± 4.88
Male gender, n (%)	16 (80)
BMI, median [IQR]	29.7 [27.4–32.5]
Cardiovascular risk factors	
Active smoking, n (%)	1 (5)
Hypertension, n (%)	16 (80)
Dyslipidemia, n (%)	18 (90)
Peripheral artery disease, n (%)	3 (15)
Chronic kidney disease, n (%)	2 (10)
Prior stroke, n (%)	0
DM complications*, n (%)	8 (40)
Insulin treatment, n (%)	7 (35)
Oral antidiabetics, n (%)	20 (100)
Cardiovascular history	
Prior myocardial infarction, n (%)	14 (70)
Diseased vessels, mean ± SD	2.15 ± 0.75
Prior PCI, n (%)	17 (85)
Prior CABG, n (%)	4 (20)
LVEF, mean ± SD	58.5 ± 9.0
Laboratory measurements	
HbA1c, median [IQR]	6.8 [6.4–7.9]
Hb, mean ± SD	13.62 ± 1.66
Platelet count (×10 ³), mean ± SD	228 ± 51
MPV, mean ± SD	11.45 ± 1.10

*Complications of DM: Neuropathy, nephropathy, retinopathy, or vasculopathy. BMI, body mass index; CABG, coronary artery bypass grafting; DM, diabetes mellitus; LVEF, left ventricular ejection fraction; MPV, mean platelet volume; PCI, percutaneous coronary intervention.

High platelet reactivity rates according to treatment

Ticagrelor HPR rates ranged from 17.6 to 35.3% at 2 h, from 0 to 28.6% at 24 h, and from 0 to 12.5% at 1 week depending on the platelet function assay employed, whereas HPR rates with clopidogrel were higher, ranging from 29.4 to 93.8% at 2 h, from 23.1 to 81.8% at 24 h, and from 15.0 to 75.0% at 1 week, reaching statistical significance in most of the comparisons (**Figure 5**).

Discussion

The present study was specifically designed to compare the antiplatelet effect of ticagrelor and clopidogrel in Mediterranean patients with DM and CCS, consisting on stable patients with prior ACS or coronary revascularization. The main finding of this investigation is that in such patients the PD benefit of ticagrelor over clopidogrel is maintained. Indeed, a LD of ticagrelor 180 mg has a faster and greater effect on platelet inhibition compared to the LD of clopidogrel 600 mg, an effect




FIGURE 3

Platelet reactivity across study time points. Comparison of platelet reactivity over time measured with LTA and using 20 μmol ADP as agonists (primary endpoint). Values are expressed as least-squares means. Error bars indicate standard errors of the mean. * $p < 0.001$; § $p < 0.05$. ADP, adenosine diphosphate; LTA, Light transmission aggregometry; MPA, maximal platelet aggregation.





FIGURE 4

Platelet function measurements across study time points. (A) Light transmittance aggregometry using 5 μM adenosine diphosphate (ADP) as agonist. (B) Flow cytometric VASP analysis. (C) Multiple electrode aggregometry using ADP as agonist. (D) VerifyNow P2Y₁₂ assay. Values are expressed as least-squares means. Error bars indicate standard errors of the mean. * $p < 0.001$; § $p < 0.05$. LTA, light transmission aggregometry; MPA, maximal platelet aggregation; VASP, vasodilator-stimulated phosphoprotein.



that is seen as soon as 2 h after intake of the LD of the drug. These outcomes were also consistently observed during the maintenance phase of therapy. This PD effect translated into ticagrelor achieving significantly lower rates of HPR at any time point of the study and with all platelet function tests employed.

Compelling data from previous PD investigations have demonstrated a greater, and also faster, inhibition of platelet reactivity achieved with ticagrelor compared with clopidogrel (19, 21). However, very few studies have addressed this issue in DM patients, a subpopulation at high risk of recurrent ischemic events. Of note, the available studies addressing this issue are actually *post-hoc* analyses and, thus, are not

exclusively performed in DM patients (11, 22). In addition, it is quite relevant to consider ethnicity when evaluating responsiveness to antiplatelet agents (11, 23). In fact, the prevalence of loss-of-function alleles of the CYP2C19 isoform varies greatly among races (9), which has a huge impact on clopidogrel responsiveness. This investigation is, to the best of our knowledge, the first to specifically compare the antiplatelet efficacy of ticagrelor vs. clopidogrel in a Mediterranean Caucasian population with DM and provides a valid confirmation of the PD superiority of ticagrelor over clopidogrel irrespective of ethnicity.

DM patients have augmented platelet reactivity, leading to greater rates of HPR to clopidogrel than non-DM subjects, which is clearly associated with poorer clinical outcomes (2–4). This problem has incited the evaluation of more potent antiplatelet regimens in this high-risk population. The PD effectiveness in DM patients of other P2Y₁₂ inhibition strategies, more potent than clopidogrel, has been compared among them in a number of mechanistic studies. For instance, in the CLOTILDA study, ticagrelor displayed a greater platelet inhibitory effect than high-dose clopidogrel (150 mg daily) in stable patients at least 1 month after PCI (12). More importantly, a number of PD investigations have compared the platelet inhibitory efficacy of ticagrelor vs. prasugrel specifically in DM patients (24–26), although results were not completely consistent. Briefly, two studies have suggested separately a slightly greater antiplatelet efficacy of ticagrelor, although no differences in the rates of HPR to both agents were observed in any of these studies (24, 25). However, in the comprehensive OPTIMUS-4 investigation, the platelet inhibitory effectiveness of both agents were similar with most of the platelet function assays employed to evaluate the LD and MD regimens (26). In line with these findings, Galli et al. (27) observed in a recent investigation a similar PD efficacy of ticagrelor and prasugrel, after switching from clopidogrel, both in patients with and without DM; of note, despite an important increase in platelet inhibition after escalation of antiplatelet agents, platelet reactivity persisted higher among DM patients compared to those without DM.

Whether there is a clinical advantage of one of the two more potent P2Y₁₂ antagonists, prasugrel, or ticagrelor, in DM patients is yet to be determined. In fact, in a prespecified analysis of patients with DM of the ISAR-REACT 5 trial, conducted in ACS patients with planned invasive therapy, no differences in ischemic or hemorrhagic events were seen between prasugrel and ticagrelor (28). The latter is in contrast with the somewhat surprising findings of the main trial, in which prasugrel significantly reduced the rates of the primary efficacy outcome, a composite of death, myocardial infarction and stroke (29, 30).

The favorable PD profile of ticagrelor in CAD patients with DM may contribute to explain the consistent benefit in terms

of reduction of atherothrombotic outcomes observed in large-scale clinical trials that have evaluated different antiplatelet regimens with ticagrelor in several scenarios across the CAD spectrum. In the DM subgroup of the pivotal PLATO trial, dual antiplatelet therapy (DAPT) with ticagrelor diminished ischemic events compared to DAPT with clopidogrel in ACS patients at moderate to high ischemic risk, without differences in major bleedings (31). Nevertheless, the relative benefit achieved with ticagrelor in DM patients, although consistent with the global trial results, was somewhat attenuated (17 vs. 12% relative risk reduction of ischemic events in non-DM and DM patients, respectively), since a numerical (although not statistically significant) reduction of the occurrence of the primary efficacy endpoint was observed. In a different clinical setting, the addition of ticagrelor on top of aspirin as secondary prevention in patients with a prior myocardial infarction, which was evaluated in the PEGASUS-TIMI 54 trial, led to a significant reduction of recurring ischemic events with ticagrelor (pooled doses of 60 and 90 mg b.i.d.) compared to the control arm (aspirin monotherapy), including both cardiovascular and coronary heart disease mortality in the DM subgroup, although with the counterpart of a heightened risk of major bleeding (32, 33). Interestingly, a platelet function substudy of this trial showed a similar platelet inhibition of ticagrelor 60 mg and 90 mg b.i.d. doses regardless of diabetes status (34). More recently, the THEMIS trial, conducted in stable DM patients with CAD and without a history of myocardial infarction or stroke, showed that adding ticagrelor to aspirin resulted in a reduction of ischemic cardiovascular events albeit at the cost of a higher rate of major bleedings, when compared to aspirin monotherapy (35). Overall, these findings underline the need for carefully addressing the ischemic and bleeding risk of each and every patient in order to decide the most suitable antiplatelet strategy.

Clopidogrel is the preferred P2Y₁₂ antagonist in patients with stable CAD undergoing PCI but it is also commonly prescribed in ACS patients deemed not suitable for potent DAPT due to increased bleeding risk. Moreover, the results of recent trials have suggested that a de-escalation of dual antiplatelet therapy (DAPT) strategy by reducing the intensity of DAPT through switching from more potent P2Y₁₂ inhibitors (i.e., prasugrel or ticagrelor) to clopidogrel, could be useful to reduce hemorrhagic events in ACS patients at high risk of bleeding without losing efficacy in terms of preventing ischemic events (36–38). For these reasons among others, clopidogrel is still widely utilized in real-life clinical practice as part of DAPT (39, 40). However, the superior platelet inhibitory effect of prasugrel or ticagrelor compared to clopidogrel, as shown in the present study and other abovementioned investigations, suggest that high-risk subgroups such as DM patients may obtain a greater benefit from maintaining more potent antiplatelet regimens. Noteworthy, recent evidence points toward a potential benefit of personalized antiplatelet

therapy using platelet function of genetic assessment (e.g., guided escalation of P2Y₁₂ inhibitors) in the PCI setting, which may be of particular relevance in DM patients due to the heightened platelet reactivity and the high rates of clopidogrel suboptimal response that characterize this population (41, 42). Indeed, an individualized approach taking into consideration the balance between ischemic and bleeding risks is certainly recommendable before deciding the P2Y₁₂ inhibition strategy in CAD patients.

Limitations

We acknowledge several limitations of the present investigation, such as the open-label design and the relatively small sample size. Further, no pharmacokinetic or genetic (e.g., loss-of-function CYP2C19 alleles) assessments were done, which could have provided important insights on the mechanisms contributing to the differences observed in platelet reactivity between clopidogrel and ticagrelor. However, prior investigations in DM patients used a single platelet function assay to compare the PD effectiveness of ticagrelor vs. clopidogrel (10, 20), whereas four different assays were employed in the present study to evaluate the LD and MD effect, which yields a great consistency to the results obtained. Ultimately, the ticagrelor 90 mg b.i.d. regimen is not routinely employed in long-term secondary prevention and our results cannot be extrapolated to the 60 mg b.i.d. dose of ticagrelor, which is approved in this scenario due to the results obtained in the PEGASUS-TIMI 54 trial.

Conclusion

In Mediterranean DM patients with CCS, ticagrelor yields a more potent platelet inhibition than clopidogrel, which is detected promptly after the loading dose and is maintained after 1 week of treatment. This PD benefit results in significantly lower HPR rates with ticagrelor compared to clopidogrel both with the load and maintenance doses. Of note, ticagrelor HPR rates are almost negligible after 1 week of therapy. The present investigation is a valid confirmation of the consistent and favorable PD profile of ticagrelor among different high-risk subgroups, such as patients with DM.

Data availability statement

The data analyzed in this study are not publicly available due to internal policy. Any requests can be directed to the

corresponding author. Requests to access the datasets should be directed to JF, jlferreiro@bellvitgehospital.cat.

Ethics statement

The studies involving human participants were reviewed and approved by CEIm Hospital Universitario de Bellvitge. The patients/participants provided their written informed consent to participate in this study.

Author contributions

AM and JF contributed to the conception and design of the study, analysis and interpretation of data. AM acquired data for the work and drafted the manuscript, which was critically revised for important intellectual content by the other authors. All authors approved the final version submitted.

Funding

This study was supported by a grant from AstraZeneca (ISSBRIL0162) to the Sociedad Española de Cardiología. The funder had no role in the conduct of the study, data collection, statistical analysis, interpretation of data, drafting the manuscript and decision to submit it for publication.

References

1. Patti G, Cavallari I, Andreotti F, Calabrò P, Cirillo P, Denas G, et al. Prevention of atherothrombotic events in patients with diabetes mellitus: from antithrombotic therapies to new-generation glucose-lowering drugs. *Nat Rev Cardiol.* (2019) 16:113–30. doi: 10.1038/s41569-018-0080-2
2. Ferreiro JL, Angiolillo DJ. Diabetes and anti-platelet therapy in acute coronary syndrome. *Circulation.* (2011) 123:798–813. doi: 10.1161/CIRCULATIONAHA.109.913376
3. American Diabetes Association. 10. Cardiovascular disease and risk management: standards of medical care in diabetes-2020. *Diabetes Care.* (2020) 43(Suppl. 1):S111–34. doi: 10.2337/dc20-S010
4. Ferreiro JL, Angiolillo DJ. Challenges and perspectives of antiplatelet therapy in patients with diabetes mellitus and coronary artery disease. *Curr Pharm Des.* (2012) 18:5273–93. doi: 10.2174/138161212803251916
5. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* (2009) 361:1–13. doi: 10.1056/NEJMoa0904327
6. Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, et al. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* (2007) 357:2001–15. doi: 10.1056/NEJMoa0706482
7. Mehilli J, Baquet M, Hochholzer W, Mayer K, Tesche C, Aradi D, et al. Randomized comparison of intensified and standard P2Y12-receptor-inhibition before elective percutaneous coronary intervention: the SASSICAIA trial. *Circ Cardiovasc Interv.* (2020) 13:e008649. doi: 10.1161/CIRCINTERVENTIONS.119.008649

Acknowledgments

We thank CERCA Programme/Generalitat de Catalunya for institutional support.

Conflict of interest

JF reports: Speaker fees from Eli Lilly Co., Daiichi Sankyo, Inc., AstraZeneca, Roche Diagnostics, Pfizer, Abbott, Ferrer, Rovi, Boehringer Ingelheim, and Bristol-Myers Squibb; consulting fees from AstraZeneca, Eli Lilly Co., Ferrer, Boston Scientific, Pfizer, Boehringer Ingelheim, Daiichi Sankyo, and Bristol-Myers Squibb; and research grant from AstraZeneca.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

8. Silvain J, Lattuca B, Beygui F, Rangé G, Motovska Z, Dillinger JG, et al. Ticagrelor versus clopidogrel in elective percutaneous coronary intervention (ALPHEUS): a randomised, open-label, phase 3b trial. *Lancet.* (2020) 396:1737–44. doi: 10.1016/S0140-6736(20)32236-4
9. Nguyen AB, Cavallari LH, Rossi JS, Stouffer GA, Lee CR. Evaluation of race and ethnicity disparities in outcome studies of CYP2C19 genotype-guided antiplatelet therapy. *Front Cardiovasc Med.* (2022) 9:991646. doi: 10.3389/fcvm.2022.991646
10. Angiolillo DJ, Ferreiro JL. Platelet adenosine diphosphate P2Y12 receptor antagonism: benefits and limitations of current treatment strategies and future directions. *Rev Esp Cardiol.* (2010) 63:60–76. doi: 10.1016/s1885-5857(10)70010-4
11. Clavijo LC, Maya J, Carlson G, Angiolillo DJ, Teng R, Caplan R, et al. Platelet inhibition with ticagrelor versus clopidogrel in hispanic patients with stable coronary artery disease with or without diabetes mellitus. *Cardiovasc Revasc Med.* (2015) 16:450–4. doi: 10.1016/j.carrev.2015.08.007
12. Mangiacapra F, Panaioli E, Colaiori I, Ricottini E, Pantano AL, Pozzilli P, et al. Clopidogrel versus ticagrelor for antiplatelet maintenance in diabetic patients treated with percutaneous coronary intervention: results of the CLOTILDIA study (clopidogrel high dose versus ticagrelor for antiplatelet maintenance in diabetic patients). *Circulation.* (2016) 134:835–7. doi: 10.1161/CIRCULATIONAHA.116.023743
13. Ferreiro JL, Homs S, Berdejo J, Roura G, Gómez-Lara J, Romaguera R, et al. Clopidogrel pretreatment in primary percutaneous coronary intervention: prevalence of high on-treatment platelet reactivity and impact on preprocedural patency of the infarct-related artery. *Thromb Haemost.* (2013) 110:110–7. doi: 10.1160/TH13-01-0057

14. Breet NJ, van Werkum JW, Bouman HJ, Kelder JC, Ruven HJ, Bal ET, et al. Comparison of platelet function tests in predicting clinical outcome in patients undergoing coronary stent implantation. *JAMA*. (2010) 303:754–62. doi: 10.1001/jama.2010.181
15. Ferreiro JL, Ueno M, Tello-Montoliu A, Tomasello SD, Capodanno D, Capranzano P, et al. Effects of cangrelor in coronary artery disease patients with and without diabetes mellitus: an in vitro pharmacodynamic investigation. *J Thromb Thrombolysis*. (2013) 35:155–64. doi: 10.1007/s11239-012-0846-z
16. Sibbing D, Aradi D, Alexopoulos D, Ten Berg J, Bhatt DL, Bonello L, et al. Updated expert consensus statement on platelet function and genetic testing for guiding P2Y12 receptor inhibitor treatment in percutaneous coronary intervention. *JACC Cardiovasc Interv*. (2019) 12:1521–37. doi: 10.1016/j.jcin.2019.03.034
17. Malinin A, Pokov A, Spergling M, Defranco A, Schwartz K, Schwartz D, et al. Monitoring platelet inhibition after clopidogrel with the VerifyNow-P2Y12(R) rapid analyzer: the VERIFY thrombosis risk Assessment (VERITAS) study. *Thromb Res*. (2007) 119:277–84. doi: 10.1016/j.thromres.2006.01.019
18. Sibbing D, Schulz S, Braun S, Morath T, Stegherr J, Mehilli J, et al. Antiplatelet effects of clopidogrel and bleeding in patients undergoing coronary stent placement. *J Thromb Haemost*. (2010) 8:250–6. doi: 10.1111/j.1538-7836.2009.03709.x
19. Storey RF, Angiolillo DJ, Patil SB, Desai B, Ecol R, Husted S, et al. Inhibitory effects of ticagrelor compared with clopidogrel on platelet function in patients with acute coronary syndromes: the PLATO (PLATElet inhibition and patient Outcomes) PLATELET substudy. *J Am Coll Cardiol*. (2010) 56:1456–62. doi: 10.1016/j.jacc.2010.03.100
20. Cho JR, Rollini F, Franchi F, DeGroat C, Bhatti M, Dunn EC, et al. Pharmacodynamics, pharmacokinetics, and safety of the oral reversible P2Y12 antagonist AZD6140 with aspirin in patients with atherosclerosis: a double-blind investigation. *JACC Cardiovasc Interv*. (2015) 8:1075–83.
21. Husted S, Emanuelsson H, Heptinstall S, Sandset PM, Wickens M, Peters G. Pharmacodynamics, pharmacokinetics, and safety of the oral reversible P2Y12 antagonist AZD6140 with aspirin in patients with atherosclerosis: a double-blind comparison to clopidogrel with aspirin. *Eur Heart J*. (2006) 27:1038–47. doi: 10.1093/eurheartj/ehi754
22. Sweeney JM, Angiolillo DJ, Franchi F, Rollini F, Waksman R, Raveendran G, et al. Impact of diabetes mellitus on the pharmacodynamic effects of ticagrelor versus clopidogrel in troponin-negative acute coronary syndrome patients undergoing ad hoc percutaneous coronary intervention. *J Am Heart Assoc*. (2017) 6:e005650. doi: 10.1161/JAHA.117.005650
23. Waksman R, Maya J, Angiolillo DJ, Carlson GE, Teng R, Caplan RJ, et al. Ticagrelor versus clopidogrel in black patients with stable coronary artery disease: prospective, randomized, open-label, multiple-dose, crossover pilot study. *Circ Cardiovasc Interv*. (2015) 8:e002232. doi: 10.1161/CIRCINTERVENTIONS.114.002232
24. Laine M, Frère C, Toesca R, Berbis J, Barnay P, Pansieri M, et al. Ticagrelor versus prasugrel in diabetic patients with an acute coronary syndrome. A pharmacodynamic randomised study. *Thromb Haemost*. (2014) 111:273–8. doi: 10.1160/TH13-05-0384
25. Alexopoulos D, Xanthopoulou I, Mavronasiou E, Stavrou K, Siapika A, Tsoni E, et al. Randomized assessment of ticagrelor versus prasugrel antiplatelet effects in patients with diabetes. *Diabetes Care*. (2013) 36:2211–6. doi: 10.2337/dc12-2510
26. Franchi F, Rollini F, Aggarwal N, Hu J, Kureti M, Durairaj A, et al. Pharmacodynamic comparison of prasugrel versus ticagrelor in patients with type 2 diabetes mellitus and coronary artery disease: the OPTIMUS (optimizing antiplatelet therapy in diabetes mellitus)-4 study. *Circulation*. (2016) 134:780–92. doi: 10.1161/CIRCULATIONAHA.116.023402
27. Galli M, Rollini F, Been L, Zenni MM, Angiolillo DJ, Franchi F. Impact of diabetes mellitus on the pharmacodynamic effects of prasugrel and ticagrelor after switching from clopidogrel in patients with coronary artery disease. *J Thromb Thrombolysis*. (2022) 54:461–9. doi: 10.1007/s11239-022-02696-4
28. Ndrepepa G, Kastrati A, Menichelli M, Neumann FJ, Wöhrle J, Bernlochner I, et al. Ticagrelor or prasugrel in patients with acute coronary syndromes and diabetes mellitus. *JACC Cardiovasc Interv*. (2020) 13:2238–47. doi: 10.1016/j.jcin.2020.07.032
29. Schüpke S, Neumann FJ, Menichelli M, Mayer K, Bernlochner I, Wöhrle J, et al. Ticagrelor or prasugrel in patients with acute coronary syndromes. *N Engl J Med*. (2019) 381:1524–34. doi: 10.1056/NEJMoa1908973
30. Crea F, Thiele H, Sibbing D, Barthélémy O, Bauersachs J, Bhatt DL, et al. Debate: prasugrel rather than ticagrelor is the preferred treatment for NSTEMI-ACS patients who proceed to PCI and pretreatment should not be performed in patients planned for an early invasive strategy. *Eur Heart J*. (2021) 42:2973–85. doi: 10.1093/eurheartj/ehab277
31. James S, Angiolillo DJ, Cornel JH, Erlinge D, Husted S, Kontny F, et al. Ticagrelor vs. clopidogrel in patients with acute coronary syndromes and diabetes: a substudy from the PLATElet inhibition and patient outcomes (PLATO) trial. *Eur Heart J*. (2010) 31:3006–16. doi: 10.1093/eurheartj/ehq325
32. Bonaca MP, Bhatt DL, Cohen M, Steg PG, Storey RF, Jensen EC, et al. PEGASUS-TIMI 54 steering committee and investigators. Long-term use of ticagrelor in patients with prior myocardial infarction. *N Engl J Med*. (2015) 372:1791–800. doi: 10.1056/NEJMoa1500857
33. Bhatt DL, Bonaca MP, Bansilal S, Angiolillo DJ, Cohen M, Storey RF, et al. Reduction in ischemic events with ticagrelor in diabetic patients with prior myocardial infarction in PEGASUS-TIMI 54. *J Am Coll Cardiol*. (2016) 67:2732–40. doi: 10.1016/j.jacc.2016.03.529
34. Thomas MR, Angiolillo DJ, Bonaca MP, Ajjan RA, Judge HM, Rollini F, et al. Consistent platelet inhibition with ticagrelor 60 mg twice-daily following myocardial infarction regardless of diabetes status. *Thromb Haemost*. (2017) 117:940–7. doi: 10.1160/TH16-09-0703
35. Steg PG, Bhatt DL, Simon T, Fox K, Mehta SR, Harrington RA, et al. Ticagrelor in patients with stable coronary disease and diabetes. *N Engl J Med*. (2019) 381:1309–20. doi: 10.1056/NEJMoa1908077
36. Cuisset T, Deharo P, Quilici J, Johnson TW, Deffarges S, Bassez C, et al. Benefit of switching dual antiplatelet therapy after acute coronary syndrome: the TOPIC (timing of platelet inhibition after acute coronary syndrome) randomized study. *Eur Heart J*. (2017) 38:3070–8. doi: 10.1093/eurheartj/ehx175
37. Sibbing D, Aradi D, Jacobshagen C, Gross L, Trenk D, Geisler T, et al. Guided de-escalation of antiplatelet treatment in patients with acute coronary syndrome undergoing percutaneous coronary intervention (TROPICAL-ACS): a randomised, open-label, multicentre trial. *Lancet*. (2017) 390:1747–57. doi: 10.1016/S0140-6736(17)32155-4
38. Kim CJ, Park MW, Kim MC, Choo EH, Hwang BH, Lee KY, et al. Unguided de-escalation from ticagrelor to clopidogrel in stabilised patients with acute myocardial infarction undergoing percutaneous coronary intervention (TALOS-AMI): an investigator-initiated, open-label, multicentre, non-inferiority, randomised trial. *Lancet*. (2021) 398:1305–16. doi: 10.1016/S0140-6736(21)01445-8
39. Ferreiro JL, Vivas D, De La Hera JM, Marcano AL, Lugo LM, Gómez-Polo JC, et al. High and low on-treatment platelet reactivity to P2Y12 inhibitors in a contemporary cohort of acute coronary syndrome patients undergoing percutaneous coronary intervention. *Thromb Res*. (2019) 175:95–101. doi: 10.1016/j.thromres.2019.01.021
40. Marcucci R, Patti G, Calabrò P, Gori AM, Grossi G, Cirillo P, et al. Antiplatelet treatment in acute coronary syndrome patients: real-world data from the START-antiplatelet Italian registry. *PLoS One*. (2019) 14:e0219676. doi: 10.1371/journal.pone.0219676
41. Galli M, Benenati S, Capodanno D, Franchi F, Rollini F, D'Amario D, et al. Guided versus standard antiplatelet therapy in patients undergoing percutaneous coronary intervention: a systematic review and meta-analysis. *Lancet*. (2021) 397:1470–83. doi: 10.1016/S0140-6736(21)00533-X
42. Galli M, Ortega-Paz L, Franchi F, Rollini F, Angiolillo DJ. Precision medicine in interventional cardiology: implications for antiplatelet therapy in patients undergoing percutaneous coronary intervention. *Pharmacogenomics*. (2022) 23:723–37. doi: 10.2217/pgs-2022-0057