



## OPEN ACCESS

APPROVED BY  
Frontiers Editorial Office,  
Frontiers Media SA, Switzerland

\*CORRESPONDENCE  
Hui-Nam Pak  
✉ hnpak@yuhs.ac

RECEIVED 11 May 2023  
ACCEPTED 22 May 2023  
PUBLISHED 02 June 2023

## CITATION

Kim D, Yu HT, Kwon O-S, Kim T-H, Uhm J-S,  
Joung B, Lee M-H and Pak H-N (2023)  
Corrigendum: Effect of epicardial fat volume on  
outcomes after left atrial posterior wall isolation in  
patients with persistent atrial fibrillation.  
Front. Cardiovasc. Med. 10:1220717.  
doi: 10.3389/fcvm.2023.1220717

## COPYRIGHT

© 2023 Kim, Yu, Kwon, Kim, Uhm, Joung, Lee  
and Pak. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The use,  
distribution or reproduction in other forums is  
permitted, provided the original author(s) and  
the copyright owner(s) are credited and that the  
original publication in this journal is cited, in  
accordance with accepted academic practice.  
No use, distribution or reproduction is  
permitted which does not comply with these  
terms.

# Corrigendum: Effect of epicardial fat volume on outcomes after left atrial posterior wall isolation in addition to pulmonary vein isolation in patients with persistent atrial fibrillation

Daehoon Kim, Hee Tae Yu, Oh-Seok Kwon, Tae-Hoon Kim,  
Jae-Sun Uhm, Boyoung Joung, Moon-Hyoung Lee and  
Hui-Nam Pak\*

Division of Cardiology, Department of Internal Medicine, Yonsei University Health System, Seoul, Republic of Korea

## KEYWORDS

atrial fibrillation, epicardial adipose tissue, posterior wall isolation, catheter ablation-atrial fibrillation, rhythm outcome

## A Corrigendum on

**Effect of epicardial fat volume on outcomes after left atrial posterior wall isolation in addition to pulmonary vein isolation in patients with persistent atrial fibrillation**

by Kim D, Yu HT, Kwon OS, Kim TH, Uhm JS, Joung B, Lee MH, Pak HN. (2022) Front. Cardiovasc. Med. 9:1005760. doi: 10.3389/fcvm.2022.1005760

In the published article, there was an error. Incorrect numbers were used.

A correction has been made to **Results**, *Additional posterior wall box isolation in patients with a larger epicardial adipose tissue volume*, Paragraph 2. This sentence previously stated:

“After overlap weighting, the early recurrence was less frequent in the additional POBI group (38.6%) than in the CPVI alone group (52.4%) ( $P=0.001$ ) (Table 3).”

The corrected sentence appears below:

“After overlap weighting, the early recurrence was less frequent in the additional POBI group (39.4%) than in the CPVI alone group (54.1%) ( $P=0.001$ ) (Table 3).”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.