



OPEN ACCESS

EDITED AND REVIEWED BY
Matteo Cameli,
University of Siena, Italy

*CORRESPONDENCE

Jiří Plášek
✉ jiri.plasek@fno.cz

RECEIVED 27 February 2024

ACCEPTED 19 March 2024

PUBLISHED 03 April 2024

CITATION

Plášek J, Dodulík J, Lazárová M, Stejskal D, Švagera Z, Chobolová N, Šulc P, Evin L, Purová D and Václavík J (2024) Corrigendum: Secretoneurin levels are higher in dilated cardiomyopathy than in ischaemic cardiomyopathy: preliminary results. *Front. Cardiovasc. Med.* 11:1392484. doi: 10.3389/fcvm.2024.1392484

COPYRIGHT

© 2024 Plášek, Dodulík, Lazárová, Stejskal, Švagera, Chobolová, Šulc, Evin, Purová and Václavík. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Secretoneurin levels are higher in dilated cardiomyopathy than in ischaemic cardiomyopathy: preliminary results

Jiří Plášek^{1,2*}, Jozef Dodulík¹, Marie Lazárová¹, David Stejskal^{3,4}, Zdeněk Švagera^{3,4}, Nela Chobolová³, Patrik Šulc¹, Lukáš Evin^{1,2}, Dana Purová⁵ and Jan Václavík^{1,2}

¹Department of Internal Medicine and Cardiology, University Hospital Ostrava, Ostrava, Czechia, ²Research Center for Internal and Cardiovascular Diseases Faculty of Medicine, University of Ostrava, Ostrava, Czechia, ³Institute of Laboratory Medicine, University Hospital Ostrava, Ostrava, Czechia, ⁴Institute of Laboratory Medicine, University of Ostrava, Ostrava, Czechia, ⁵Social Health Institute, Palacky University Olomouc, Olomouc, Czechia

KEYWORDS

secretoneurin, heart failure, CaMKII, dilated cardiomyopathy, ischaemic cardiomyopathy

A Corrigendum on

Secretoneurin levels are higher in dilated cardiomyopathy than in ischaemic cardiomyopathy: preliminary results

By Plášek J, Dodulík J, Lazárová M, Stejskal D, Švagera Z, Chobolová N, Šulc P, Evin L, Purová D and Václavík J. (2023). *Front. Cardiovasc. Med.* doi: 10.3389/fcvm.2023.1297900

In the published article, there were multiple errors in Table 2 as published. The values in Table 2 as seen below have been corrected.

TABLE 2 Baseline characteristics of the healthy volunteers.

	Healthy volunteers	Males	Females	P value
	N = 34	N = 13	N = 21	
Age (years)	31 ± 7.1	30.8 ± 7.1	29.9 ± 6.9	0.596
Secretoneurin	50.7 ± 16.3	50.7 ± 15.3	50.7 ± 16.3	0.64
Body weight (kg)	69 ± 14	69 ± 14	67 ± 14	0.001
Body height (cm)	171.8 ± 8.9	171.8 ± 8.9	170.8 ± 8.9	<0.001
Body mass index (kg/m ²)	23.2 ± 3.6	23.1 ± 3.6	23 ± 3.6	0.477
NT-pro-BNP	55.5 ± 29.2	55.5 ± 29.2	57.5 ± 30	0.066
Creatinine	74.1 ± 12.45	74.1.1 ± 12.45	71.9 ± 10.6	<0.001
Natrium	138.2 ± 1.75	138.2 ± 1.75	138.2 ± 1.76	0.208
Potassium	4.1 ± 0.29	4.1 ± 0.29	4.1 ± 0.3	0.03
Ca ²⁺ (ionized)	1.17 ± 0.1	1.2 ± 0.1	1.17 ± 0.1	0.519
Total calcium	2.48 ± 0.13	2.48 ± 0.11	2.48 ± 0.13	0.079
Mg ²⁺	0.79 ± 0.1	0.79 ± 0.1	0.79 ± 0.05	0.341
hs-TnI	8.1 ± 29.4	8.1 ± 29.42	8.6 ± 31	0.195
Hs-CRP	1.8 ± 2.36	1.8 ± 2.4	2 ± 2.4	0.026
IL-6	3.1 ± 2.6	3.1 ± 2.6	3.2 ± 2.8	0.440
Leukocytes	7.52 ± 1.7	7.5 ± 1.7	7.5 ± 1.6	0.201
Thrombocytes	280 ± 59.1	280 ± 59.1	281.8 ± 59	0.013

Indices are shown as mean ± standard deviation and compared for males and females. Ca²⁺, ionized calcium; Hs-TnI, high-sensitivity Troponine I; IL-6, interleukin 6; Mg²⁺, ionized magnesium; NT-pro-BNP, N-terminal-pro-brain natriuretic peptide.

The corrected Table 2 and its caption appear below.

TABLE 2 Baseline characteristics of the healthy volunteers.

	Healthy volunteers	Males	Females	P value
	N = 34	N = 13	N = 21	
Age (years)	31 ± 7.1	30.8 ± 7.1	29.9 ± 6.9	0.596
Secretoneurin	50.3 ± 15	50.2 ± 15.1	50.7 ± 15.5	0.64
Body weight (kg)	69.1 ± 13.5	70 ± 14	66.6 ± 13	0.47
Body height (cm)	171.1 ± 8.9	172 ± 8.7	170.6 ± 8.8	0.52
Body mass index (kg/m ²)	23.2 ± 3.6	23.1 ± 3.6	23 ± 3.6	0.477
NT-pro-BNP	55.4 ± 28.7	58.1 ± 31.4	53.6 ± 28.9	0.066
Creatinine	73 ± 10.9	75.6 ± 10.1	72.3 ± 10.3	0.36
Natrium	138.2 ± 1.73	138 ± 1.8	138.1 ± 1.7	0.208
Potassium	4.1 ± 0.3	4.1 ± 0.29	4.1 ± 0.3	0.03
Ca ²⁺ (ionized)	1.17 ± 0.1	1.2 ± 0.1	1.17 ± 0.1	0.519
Total calcium	2.48 ± 0.13	2.48 ± 0.11	2.48 ± 0.13	0.079
Mg ²⁺	0.79 ± 0.1	0.79 ± 0.1	0.79 ± 0.05	0.341
hs-TnI	8.6 ± 30	10.2 ± 34.2	8.8 ± 30.1	0.9
Hs-CRP	1.9 ± 2.4	1.5 ± 1.7	2 ± 2.4	0.45
IL-6	3.1 ± 2.6	3.1 ± 2.6	3.2 ± 2.8	0.440
Leukocytes	7.52 ± 1.7	7.5 ± 1.7	7.5 ± 1.6	0.201
Thrombocytes	277.5 ± 59.8	271.3 ± 64	280.4 ± 59.8	0.69

Indices are shown as mean ± standard deviation and compared for males and females. Ca²⁺, ionized calcium; Hs-TnI, high-sensitivity Troponine I; IL-6, interleukin 6; Mg²⁺, ionized magnesium; NT-pro-BNP, N-terminal-pro-brain natriuretic peptide.

The authors apologize for this errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

In the published article, there was an error in assumptions about differences in males and females in the healthy volunteers group.

A correction has been made to **Results**. This sentence previously stated:

“Males and females in the healthy individuals group differed in anthropometric parameters, plasma creatinine level, hs-CRP, and thrombocyte count (Table 2)”.

The corrected sentence appears below:

“Males and females in the healthy individuals’ group did not differ in any parameters (Table 2)”.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.